

V. 15



LIBRARY OF
THE NEW YORK BOTANICAL GARDEN

E. D. Merrill Library
1954





QK3

M42

V.15

AN ENUMERATION OF PHILIPPINE FLOWERING PLANTS

BY

ELMER D. MERRILL

DIRECTOR AND BOTANIST, BUREAU OF SCIENCE

VOLUME I



MANILA
BUREAU OF PRINTING
1925

GOVERNMENT OF THE PHILIPPINE ISLANDS
DEPARTMENT OF AGRICULTURE AND NATURAL RESOURCES
BUREAU OF SCIENCE
MANILA

PUBLICATION No. 18, VOLUME 1

DATES OF ISSUE

- Fascicle 1, pages 1 to 128, January 3, 1923.
2, pages 129 to 240, January 3, 1923.
3, pages 241 to 368, December 29, 1924.
4, pages 369 to 463, April 7, 1925.

PREFACE

The object of the present work is to enumerate the known species of flowering plants occurring in the Philippines; to adjust the synonymy; to account for all binomials accredited to the Philippines in botanical literature; to give the most important literature references, especially to the special Philippine publications and to the important extra-Philippine ones; to record the numerous local names for the various species; to give the habitat, occurrence, altitudinal range, Philippine distributions, and approximate extra-Philippine distribution of each species admitted; to cite illustrative specimens when necessary or expedient, especially in critical groups; in brief, to summarize in convenient form for consultation our present knowledge of the Philippine flora.

When publication of this work was commenced in 1921, it was planned to issue it in three volumes, of four or five fascicles each. As publication progressed, it was found necessary to abandon the plan of issuing the work in three volumes and the inclusion of the extensive introductory data in the prefatory part of the present volume and to add a fourth volume to contain the general introduction, bibliography, and indices to local and scientific names. In the individual volumes of the actual enumeration merely brief indices have been included covering the family and generic names that appear in each volume. Complete indices to the entire work form a part of volume 4.

In the arrangement of families and of genera the sequence in general follows the Engler and Prantl system, but for convenience in indexing and in consulting the species have been arranged alphabetically under their respective genera. While an attempt has been made to compare the very numerous species within each genus with each other and also with extra-Philippine representatives of the same genera, so far as material has been available, much critical work remains to be done; and future studies must of necessity modify the enumeration, not only as to the validity of numerous admitted species but also as to questions of synonymy and geographic distribution. It should be remembered that most of the actual work has been accomplished on the basis of the material and the literature available in Manila,

and that the local botanical library and the herbarium date only from 1902. While the library is a very excellent working unit, especially for the Indo-Malaysian region, it is far from complete, although practically all the publications appertaining especially to the Philippine flora are represented. The herbarium, containing upward of 250,000 mounted specimens, is especially rich in Philippine material, and contains few local collections made previous to 1902, except a series of approximately 1,200 numbers from Cuming's collections, made between 1836 and 1840, and a fairly large set of duplicates representing Warburg's collections. However, nearly all species described from the Philippines of which the types are extant are represented in the herbarium by types, cotypes, fragments of authentic specimens, photographs or sketches of types, or by critically compared specimens. Types of none of the species described by Blanco in his *Flora de Filipinas*, few of those described by Llanos, or the few described by F.-Villar and Naves are extant. The Blanco and Llanos species have been critically considered in a special publication.*

The extra-Philippine collections are relatively extensive, the herbarium being rich in material from surrounding regions, received in exchange and by collection, from Formosa, China, Indo-China, India, Ceylon, Burma, Siam, the Malay Peninsula, Sumatra, Java, Borneo, Celebes, the Moluccas, New Guinea, New Caledonia, Australia, the Marianne and the Caroline Islands, and Polynesia.

Some idea of the extent and the value of the extra-Philippine botanical material available in the herbarium of the Bureau of Science may be had from the following data: From India, Burma, Siam, Indo-China, and the Malay Peninsula, more or less extensive collections are available from material secured by the following collectors and botanists: Burkill, Buchanan-Hamilton, Burn-Murdoch, Balansa, Bon, Cave, Clarke, Craib, Curtis, Debeaux, Foxworthy, Eberhardt, Harmand, Gamble, Godefroy, Hole, Hooker f. and Thomson, Jacquemont, Kurz, Kunstler, King, King's collectors, Kerr, Kloss, Meebold, Petelot, Pierre, Prazer, Rock, Robinson, Ridley, Shaik Mokim, Talmy, Thorel, Thwaites, Walker, Wallich, Wight, and Wray. The flora of China, Japan, and Formosa is well represented by collections made by the following individuals: Bodinier, Callery, Cavalerie, Chun, Chung,

* Merrill, E. D., *Species Blancoanae: A critical revision of the Philippine species of plants described by Blanco and by Llanos*, Bur. Sci. Publ. 12 (1918) 1-423, map.

Clemens, Dunn, Ducloux, Delavay, Farges, Faurie, Ford, Groff, Henry, Hu, Ito, Kanehira, Katsumada, Kawakami, Levine, Nakahira, Merrill, McClure, Maire, Steward, Sasaki, Taquet, Ts'oong, Tutcher, Wilson, and Wichura. The flora of some parts of the Malay Archipelago is particularly well represented by botanical material from the following collectors and botanists: Achmad, Agama, Amdjah, Backer, Backhuizen, Blume, Beguin, Beccari, Bartlett, Bünnenmeyer, Clemens, Elmer, Forbes, Foxworthy, Gjellerup, Hallier, Haniff, Hose, Kloss, Koorders, Korthals, C. King, La Rue, Lörzing, Mousset, Moulton, Nieuwenhuis, Ramos, Robinson, Raciborski, Schlechter, Teysmann, Topping, Versteeg, Villamil, Warburg, Winkler, Wood, and Yates. The flora of some parts of Polynesia is fairly well represented by collections made by the following individuals: Heller, Hombron, Le Batard, Le Guillou, McGregor, Nelson, Setchell, Thompson, Vaupel, Vesco, and Volkens. The flora of New Caledonia is particularly well represented by an extensive series of collections made by the following individuals: Balansa, Cribbs, Deplanche, Franc, Germain, Lécard, Le Rat, Pancher, and Vieillard.

The duplicates of botanical material secured by the older collectors have in large part been obtained in exchange from European botanical institutions, notably the British Museum (Natural History), the Edinburgh Botanic Garden, the Paris Museum of Natural History, the Berlin Botanic Garden, and the Rijks Herbarium, Leyden; thus, from the Berlin institution more than 8,000 specimens have been secured; from the Paris Museum, more than 12,000 specimens; from the British Museum, about 6,000 specimens; and from the Edinburgh Botanic Garden, 3,500 specimens. An extensive series of exchanges has been carried on with botanic institutions in different parts of India, Malaysia, and Australia; more than 12,000 specimens from the Malay Archipelago have been received from the Buitenzorg Botanic Garden in Java, about 5,000 specimens from the Malay Peninsula have been received from the Botanic Garden at Singapore, and the Australian flora is represented by somewhat over 10,000 specimens secured in exchange with various botanic institutions in Australia. The large exchanges received from the Paris Museum have been particularly rich in material from Indo-China, China, and New Caledonia; the material from the British Museum has been largely from British India, with miscellaneous collections from China and New Guinea; the material received from the Edinburgh Botanic Garden has been largely from India and China. The large amount of extra-Philippine material from

regions surrounding the Philippines has been invaluable for purposes of direct comparison in the prosecution of our work on the Philippine flora.

Botanical exploration under the auspices of the Bureau of Science has been confined largely to the Philippines, but members of our staff have prosecuted field work in China (Merrill), Indo-China (Robinson), Borneo (Foxworthy, Ramos), Amboina (Robinson), and Guam (McGregor). Large collections have been submitted for identification from various parts of China, Formosa, Hainan, Indo-China, Burma, Siam, the Malay Peninsula, Sumatra, Java, Borneo, New Guinea, Guam, Samoa, and Tahiti.

No attempt has been made to cite all the collections and numbers, as this procedure would have added enormously to the extent of the present publication and would have added very little to its value. In general, well-known and widely distributed species are merely enumerated with their synonyms, occurrence, distribution, etc., without the citation of specimens. In critical groups selected representative specimens only are cited. Generally speaking, when only one or two collections are cited under a species these represent the only collections extant. It should be noted that in approximately 1,800 cases Philippine records for individual species are based solely on single collections, while in about 900 additional cases individual species records are based on two collections only. This will give some idea of the great amount of work still remaining to be done before we can hope to have a reasonably complete knowledge of the very rich Philippine flora. Collections cited in parentheses—for example, (*Warburg 14231*), (*Loher 4250*)—indicate specimens not examined by me during the actual preparation of this work, although most of these were seen by me in various European herbaria in 1907 and 1908.

While most of the identifications on which this enumeration is based have been made by me, in certain groups much of this work in earlier years was done by former botanists connected with the Bureau of Science; notably, the late Dr. C. B. Robinson (Myrtaceae, Urticaceae, and certain groups of Euphorbiaceae, especially) and Dr. F. W. Foxworthy (Dipterocarpaceae and Taxaceae). Mr. Oakes Ames has made practically all of the identifications of the Orchidaceae since 1903. Many of the Pandanaceae have been determined by Dr. U. Martelli, while most of the Palmae were identified by the late Dr. O. Beccari. Many identifications in the Gramineae were made by Dr. E. Hackel, especially previous to 1907; while in the Cyperaceae the Bureau of Science has had the benefit of the extensive knowledge

of the late Mr. C. B. Clarke and of the Rev. G. Kükenthal. The Araceae have been largely identified by Dr. A. Engler, and the Dioscoreaceae by Sir David Prain and Mr. I. H. Burkill. The Piperaceae have largely been identified by the late Dr. C. de Candolle. Dr. L. Diels has named many of the Menispermaceae. Dr. J. Perkins in addition to her general work on earlier collections in various families has identified many of the Monimiaceae. The late Mr. O. Focke has named or examined many of the *Rubus* specimens. Practically all of the Sapindaceae have been named or examined by Dr. L. Radlkofer. The *Impatiens* material was named by the late Sir J. D. Hooker. Many of the Symlocaceae were determined by Dr. A. Brand. Dr. R. Schlechter has identified or examined most of the species of Asclepiadaceae. Dr. F. Kränzlin accomplished considerable work on the Gesneriaceae. The late Mr. C. B. Clarke identified numerous representatives of the Acanthaceae. Numerous individual identifications and comparisons with type material have been made by other botanists in various European countries, the United States, Japan, Java, Singapore, India, and Australia.

AN ENUMERATION OF PHILIPPINE
FLOWERING PLANTS

By ELMER D. MERRILL

Director and Botanist, Bureau of Science

GYMNOSPERMAE

CYCADACEAE

1. CYCAS Linnaeus

CYCAS CAIRNSIANA F.-Muell. Fragm. 10 (1876-77) 63?*Cycas* sp. Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 151.

Culion, *Merrill 657*. In open grasslands at low altitudes. A curious species, of which only the leaves are known, perhaps representing the above Australian species.

CYCAS REVOLUTA Thunb. Fl. Jap. (1784) 229; A. DC. Prodr. 16² (1868) 526; F.-Vill. Novis. App. (1880) 212; Vidal Cat. Pl. Prov. Manila (1880) 46; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 151; Merr. Fl. Manila (1912) 65.

A native of China and Japan, introduced into the Philippines for ornamental purposes and now occasionally cultivated in Manila and in other large towns in the Archipelago. It never produces flowers or fruits in the Philippines so far as our records show.

Local name: Olíva (Sp.).

CYCAS RUMPHII Miq. in Bull. Soc. Phys. Nat. Néerl. (1839) 45; F.-Vill. Novis. App. (1880) 212; Merr. Interpret. Herb. Amb. (1917) 74, Sp. Blancoanae (1918) 52.

Cycas circinalis Blanco Fl. Filip. (1837) 745, ed. 2 (1845) 513, ed. 3, 3 (1879) 146; F.-Vill. Novis. App. (1880) 212; Vidal Sinopsis Atlas (1883) 43, t. 99; Warb. Monsunia (1900) 178; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 24, 3 (1908) Bot. 394, Fl. Manila (1912) 65; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 151 (non Linn.?).

Cycas riuminiana Porte ex Regel in Gartenflora 12 (1863) 16, fig., Ill. Hort. 28 (1881) 32; A. DC. Prodr. 16² (1868) 528; F.-Vill. Novis. App. (1880) 212.

Batan Islands and northern Luzon to Palawan and Mindanao. Chiefly along or near the seashore, sometimes in forests; of rather local occurrence. Malaya to Polynesia.

The common coastal form may not be specifically distinct from *C. circinalis* Linn. A form from the forests of Mt. Mariveles (*Merrill 3257*, *Whitford 269*, *1325*, etc.), with small ♂ cones and small fruits may be specifically distinct. A second form from Mindoro and Batangas with small fruits may also be distinct (*B. S. 22391 Ramos*).

Local names: Báit (Sul.); báyit (Yak.); bitógo (Tag.); oliva (Sp.); patúbo (Tag.); patúgo (Tag.); pitógo (Tag.); pitúgo (P. Bis.); saúang (Ilk.); uliba (Tag.).

CYCAS sp. Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 152, t. 27.

Palawan, *F. B. 3842 Curran*. Leaves like those of *C. rumphii* Miq., but the ♂ cones are entirely different. Probably an undescribed species.

TAXACEAE

1. PODOCARPUS L'Héritier

PODOCARPUS AMARUS Blume Enum. Pl. Jav. (1827) 88, Rumphia 3 (1847) 213, t. 170; Pilger in Engl. Pflanzenreich 18 (1903) 68, f. B, A-D, Warb. Monsunia (1900) 192; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 159.

Podocarpus euryntcha Miq. Fl. Ind. Bat. 2 (1859) 1074; F.-Vill. Novis. App. (1880) 211.

Luzon (Benguet, Lepanto), Mindoro, Mindanao (Davao), *Elmer 11682, Merrill 5703, F. B. 10895, 10951 Curran, 18356 Alvarez, 18029 Merritt*. In the mossy forest, altitude 1,600 to 2,400 m. Java, Sumatra, Lombok; also reported from tropical Australia.

Local names: Pasuig (Ig.); tumpís (Bag.).

PODOCARPUS BLUMEI Endl. Syn. (1847) 208; Parl. in DC. Prodr. 16² (1868) 508; Pilger in Engl. Pflanzenreich 18 (1903) 60; Merr. in Govt. Lab. (Philip.) Publ. 17 (1904) 5, Philip. Journ. Sci. 1 (1906) Suppl. 24, 2 (1907) Bot. 258; Foxw. op. cit. 6 (1911) Bot. 158.

Podocarpus latifolius Blume Enum. Pl. Jav. (1827) 89; C. B. Rob. in Bull. Torr. Bot. Club. 35 (1908) 63, non R. Br. (1825).

Luzon (Cagayan, Apayao, Bataan), Mindoro, Sibuyan, Panay, Samar. *Whitford 1353, Williams 329, 753, 1035, Merrill 5728, Elmer 12360, B. S. 28348 Fénix, F. B. 23937, 23918, 23940 Cortes & Knapp, 25949 Knapp, 16738, 17200, 17616 Curran, 20888 Hinoban, 194, 147 Barnes*. In primary forests, altitude 300 to 1,800 m. Java, Moluccas, Celebes, New Guinea.

Local name: Makopulá (P. Bis.).

PODOCARPUS BREVIFOLIUS (Stapf) Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 160, t. 29, f. 2.

Podocarpus nerifolius Don var. *brevifolia* Stapf in Trans. Linn. Soc. Bot. 4 (1894) 249; Pilger in Engl. Pflanzenreich 18 (1903) 93.

Luzon (Zambales), Mindanao (Bukidnon), *B. S. 5002 Ramos, 38500 Ramos & Edaña, F. B. 9511 Curran & Merritt*. In elfinwood on exposed ridges, altitude 2,000 m. Borneo.

PODOCARPUS COSTALIS Presl. Epim. (1849) 236, p.p. quoad spec. Haenke; Walp. Ann. 3 (1852-53) 448; F.-Vill. Novis. App. (1880) 211; Miq. Fl. Ind. Bat. 2 (1859) 1074; Ceron Cat. Pl. Herb. (1892) 187; Warb. Monsunia (1900) 193; Pilger in Engl. Pflanzenreich 18 (1903) 161; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 161.

Podocarpus sp. Vidal Sinopsis Atlas (1883) 43, t. 97, f. D.

Luzon (Laguna, Tayabas), *Haenke, Merrill 7506, Loher 2140, 7128, B. S. 19581 Ramos, 2393, 2423 Foxworthy, F. B. 7913 Curran & Merritt*. In the mossy forest on Mt. Banahao, altitude 1,700 to 2,200 m. Endemic.

PODOCARPUS GLAUCUS Foxw. in Philip. Journ. Sci. 2 (1907) Bot. 258, 6 (1911) Bot. 159, t. 29, f. 1.

Mindoro, Merrill 5672. Border of an open heath on Mt. Halcon, altitude about 2,400 m. Endemic.

PODOCARPUS JAVANICUS (Burm. f.) Merr. in Philip. Journ. Sci. 19 (1921) 338. = *P. imbricatus* Blume!

(No *Thuja* ? *javanica* Burm. f. Fl. Ind. (1768) 202, t. 64, f. 3.)

Podocarpus imbricatus Blume Enum. Pl. Jav. (1827) 89; Pilger in Engl. Pflanzenreich 18 (1903) 56; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 157; W. H. Br. op. cit. 12 (1917) Bot. 317, t. 17.

Podocarpus cupressina R. Br. ex Mirb. in Mém. Mus. Paris 13 (1825) 75, *nomen nudum*; Benn. Pl. Jav. Rar. (1838) 35, t. 10; Miq. Fl. Ind. Bat. 2 (1859) 1074; F.-Vill. Novis. App. (1880) 211; Vidal Sinopsis Atlas (1883) 43, t. 97, f. B, Phan. Cuming. Philip. (1885) 160, Rev. Pl. Vasc. Filip. (1886) 295; Warb. Monsunia (1900) 191.

Podocarpus cumingii Parl. in DC. Prodr. 16² (1868) 521.

Nageia cumingii O. Kuntze Rev. Gen. Pl. 2 (1891) 800.

Podocarpus imbricatus Blume var. *cumingii* Pilger in Engl. Pflanzenreich 18 (1903) 56, Perk. Frag. Fl. Philip. (1904) 44; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 324.

Luzon (Abra, Lepanto, Bontoc, Benguet, Zambales, Tayabas), Mindoro, Negros, Panay, Mindanao. Common on the higher mountains in the mossy forest, altitude 1,300 to 2,500 m. Burma to Indo-China, Sumatra, Java, Borneo, Celebes, Timor, and New Guinea.

Local names: Apilum (Buk.); baginsóloi (Ting.); banyás (Mang.); dionái (Sub.); haluko (Ig.); iguem (Ig.); igum (Ig.); rorogóngon (Lan.); sarambrum (Bag.).

PODOCARPUS NERIIFOLIUS Don in Lamb. Pin. (1824) 21; Parl. in DC. Prodr. 16² (1868) 514; Pilger in Engl. Pflanzenreich 18 (1903) 80, Perk. Frag. Fl. Philip. (1904) 44; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 24, 2 (1907) Bot. 258; Foxw. op. cit. 6 (1911) Bot. 162 (incl. var. *brevipes* Blume).

Luzon (Abra, Benguet, Tayabas), Polillo, Mindoro, Sibuyan, Mindanao, Jolo, Merrill 1992, 5768, 5615, F. B. 25285 Somonte, 26834 Hirro, 10894, 10826 Curran, 14422 Darling, B. S. 25522 Yates, 10779 McGregor. In primary forests at low and medium altitudes. India to southern China through Malaya to New Guinea.

Local names: Dilang-butikí (Tag.); pasuik (Ig.).

PODOCARPUS PHILIPPINENSIS Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 163, t. 30.

Podocarpus rumphii Foxw. op. cit. 6 (1911) Bot. 164, non Blume.

Luzon (Ilocos Sur, Bataan, Pampanga), Mindoro, B. S. 1660, 5174, 10844 Foxworthy, F. B. 27786, 13253 Paraiso, 17523, 17594, 6326, 6325, 6537, 7512, 8987, 17664, 17723 Curran, 2743 Borden, Merrill 5553, 3917. In primary forests, altitude 400 to 800 m. Endemic.

Local name: Kasírai (Ilk.).

PODOCARPUS PILGERI Foxw. in Philip. Journ. Sci. 2 (1907) Bot. 259, 6 (1911) Bot. 160.

Podocarpus celebicus Warb. Monsunia (1900) 192; Pilger in Engl. Pflanzenreich 18 (1903) 78, non Hemsl. (1896).

Mindoro, Negros, Mindanao (Agusan, Misamis), *Merrill 5754, Phil. Pl. 241, Elmer 14086, F. B. 4673 Mearns & Hutchinson*. In the mossy forest on the higher mountains, altitude 1,400 to 2,500 m. Celebes.

Local names: Lúbang-lúbang (Mbo.); tambiayang (Bag.).

PODOCARPUS POLYSTACHYUS R. Br. ex Mirb. in *Mém. Mus. Paris* 13 (1825) 75; Endl. *Syn.* (1847) 215; Parl. in *DC. Prodr.* 16² (1868) 515; Pilger in *Engl. Pflanzenreich* 18 (1903) 79; Merr. in *Philip. Journ. Sci.* 3 (1908) Bot. 394; Foxw. *op. cit.* 6 (1911) Bot. 161.

Batan Islands, Luzon (Ilocos Norte), Palawan, Bucas, *B. S. 27146 Ramos, 26902 Edaño, 3586 Félix, 20771 Escritor, 13202 Foxworthy & Ramos, 904 Foxworthy, F. B. 24264 Bernardo, 3854 Curran, Merrill 5268*. In forests at low and medium altitudes, sometimes along the seashore. Malay Peninsula, Sumatra, Java, Borneo.

Local name: Bantígi (Neg.).

2. DACRYDIUM Solander

DACRYDIUM ELATUM (Roxb.) Wall. in *Hook. Journ. Bot.* 2 (1843) 144, *t. 2*; Blume *Rumphia* 3 (1847) 221, *t. 172 B, f. 1, t. 172 C, f. 2*; de Boer *Conif. Archip. Ind.* (1868) 29; Parl. in *DC. Prodr.* 16² (1868) 494; Rendle in *Journ. Bot.* 34 (1896) 355; Merr. in *Philip. Journ. Sci.* 2 (1907) Bot. 257; Pilger in *Engl. Pflanzenreich* 18 (1903) 51; Foxw. in *Philip. Journ. Sci.* 6 (1911) Bot. 154.

Juniperus elata Roxb. *Fl. Ind. ed. 2, 3* (1832) 838.

Luzon (Nueva Ecija), Mindoro, Negros, Panay, Mindanao, *B. S. 26510, 34497 Ramos & Pascasio, F. B. 4547, 4548, 4731 Mearns & Hutchinson, 4227 Everett, 4419, 8527 Merritt, 13612, 13621 Curran & Foxworthy, Merrill 5789*. Chiefly in the mossy forest, altitude 1,200 to 2,400 m. on the east coast of Mindanao occurring as low as 200 m. Indo-China, Malay Peninsula, Sumatra, Borneo, Celebes.

Local names: Hamo-hamo (Neg.); hinubáyan (Buk.); lokínai (Sub.); maglíging (Mbo.).

DACRYDIUM FALCIFORME (Parl.) Pilger in *Engl. Pflanzenreich* 18 (1903) 45; Merr. in *Philip. Journ. Sci.* 2 (1907) Bot. 257; Foxw. *op. cit.* 6 (1911) Bot. 153, *t. 28, f. 1*.

Podocarpus falciformis Parl. in *DC. Prodr.* 16² (1868) 685; Rendle in *Journ. Bot.* 34 (1896) 355; Warb. *Monsunia* (1900) 193.

Nageia falciformis O. Kuntze *Rev. Gen. Pl.* 2 (1891) 800.

Luzon (Nueva Ecija), Mindoro, Panay, Mindanao (Davao, Surigao), *F. B. 27746 Cruz, 4425 Merritt, 26478 Mallonga, 26394 Ramos & Edaño, Merrill 5744*. Chiefly in the mossy forest, altitude 1,000 to 2,000 m. Borneo, Malay Peninsula, Lingga.

Local name: Bináton (Mbo.); magíbus (C. Bis.); maradayónit (Neg.).

3. PHYLLOCLADUS L. C. and A. Richard

PHYLLOCLADUS HYPOPHYLLUS Hook. *f. Ic.* 9 (1852) *t. 889*; Carr. *Conif.* (1867) 706; Parl. in *DC. Prodr.* 16² (1868) 499; Pilger in *Engl. Pflanzenreich* 18 (1903) 99.

Phyllocladus protractus Pilger in *Engl. Pflanzenreich* 18 (1903) 99, *Perk. Frag. Fl. Philip.* (1904) 44; Foxw. ex Merr. in *Philip. Journ. Sci.* 2 (1907) Bot. 259, 6 (1911) Bot. 165, *t. 31*.

Phyllocladus hypophyllus Hook. f. var. *protracta* Warb. Monsunia (1911) 194.

Luzon (Isabela, Abra, Ifugao, Lepanto, Bontoc, Benguet), Mindoro, Mindanao (Bukidnon, Lanao, Misamis, Davao), *Merrill 4753, 5788, Lohr 5203, Elmer 11463, F. B. 27235 Niebert, 18364, 18567 Alvarez, 10957 Curran, 14587 Darling, 4679 Mearns & Hutchinson, B. S. 37757, 38738 Ramos & Edaño*. In the mossy forest on the higher mountains, altitude 1,200 to 2,400 m. Borneo. *Batjan. New Guinea*

The supposed differences between *P. protractus* Pilger and *P. hypophyllus* Hook. f. break down entirely on comparison of a large series of Kinabalu and Philippine specimens.

Local names: Aransisĩngit (Ig.); dálung (Ting.); galingkinga (Sub.); salumayag (Bag.).

4. TAXUS Linnaeus

TAXUS WALLICHIANA Zucc. in Abh. Bayr. Akad. Wiss. 3 (1843) 803, t. 5; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 166.

Taxus baccata Linn. subsp. *wallichiana* (Zucc.) Pilger in Engl. Pflanzenreich 18 (1903) 112, Perk. Frag. Fl. Philip. (1904) 44.

Taxus baccata Merr. in Philip. Journ. Sci. 5 (1910) Bot. 324, non Linn.

Cephalotaxus mannii Rendle in Journ. Bot. 34 (1896) 355, non Hook. f.

Luzon (Benguet, Lepanto, Laguna, Tayabas). In the mossy forest, altitude 2,000 to 2,600 m; south of the Mountain Province known in the Philippines only from Mt. Banahao; represented by numerous collections. India to Burma, Sumatra and Celebes.

Local name: Amugáuen (Ig.).

PINACEAE

1. AGATHIS Salisbury

AGATHIS ALBA (Lam.) Foxw. in Philip. Journ. Sci. 5 (1910) Gen. Sci. 173, 6 (1911) Bot. 167; Merr. Interpret. Herb. Amb. (1917) 76, Sp. Blancoanae (1918) 52.

Dammara alba Lam. Encycl. 2 (1786) 259; Parl. in DC. Prodr. 16² (1868) 374.

Agathis loranthifolia Salisb. in Trans. Linn. Soc. Bot. 8 (1807) 311; Blanco Fl. Filip. ed. 2 (1845) 528, ed. 3, 3 (1879) 170; F.-Vill. Novis. App. (1880) 211; Vidal Sinopsis Atlas (1883) 43, t. 98, f. A, Phan. Cuming. Philip. (1885) 160, Rev. Pl. Vasc. Filip. (1886) 295.

Dammara rumphii Presl Epim. (1851) 236.

Podocarpus philippeanus Benth. ex Parl. in DC. Prodr. 16² (1868) 375, in syn.

Agathis philippinensis Warb. Monsunia (1900) 185, Perk. Frag. Fl. Filip. (1904) 36; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 24, 2 (1907) Bot. 257.

Agathis borneensis Warb., *beccarii* Warb., *celebica* Warb. et *macrostachys* Warb. Monsunia (1900) 184, 185.

Babuyan Islands and northern Luzon to Palawan and Mindanao, in most islands and provinces. Always in the primary forest, chiefly at medium and higher altitudes; represented by over 60 individual collections. Indo-China through the Malay Peninsula and Archipelago to the Moluccas.

Local names: Adiāngau (Bik.); alinságo (Ig.); almáciga (Sp.); alin-tágau (Ig.); anínga (Ig.); anano (S. L. Bis.); anting (Neg.); arínga (Klg.); bagtík (Kuy.); bálau (C. Bis.); baltík (Tagb.); bidiāngau (P. Bis.); badiāngau (P. Bis.); biáyo (Bis.); bunsóg (Ig.); buntóg (Ig.); dadiāngau (C. Bis., Tag.); dadúñgoi (Bik.); dinar (Bag.); gala-gála (Tag., Tagb.); ladiāngau (Bik., Tag.); makáu (C. Bis.); olinságo (Ig.); sálang (Neg.); sáleng (Neg.); sálong (Tag., Bik.); titau (Ting.); uli (Sbl.); uniñgát (Ilk.).

2. PINUS Linnaeus

PINUS INSULARIS Endl. Syn. Conif. (1847) 157; Presl Epim. (1851) 37; Parl. in DC. Prodr. 16² (1868) 390; F.-Vill. Novis. App. (1880) 212; Vidal Sinopsis Atlas (1883) 43, t. 98, f. C, Phan. Cumming. Philip. (1885) 160, Rev. Pl. Vasc. Filip. (1886) 296; Merr. in Forest. Bur. (Philip.) Bull. 1 (1903) 15, Govt. Lab. Publ. (Philip.) 6 (1904) 6, Philip. Journ. Sci. 5 (1910) Bot. 325, Sp. Blancoanae (1918) 53; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 170; Shaw Gen. Pinus (1914) 60, t. 23, f. 208-210.

Pinus taeda Blanco Fl. Filip. (1837) 767, ed. 2 (1845) 528, ed. 3, 3 (1879) 169, t. 453, non Linn.

Pinus insularis × *merkusii* Perk. Frag. Fl. Philip. (1904) 35.

Pinus kasya Royle ex Parl. in DC. Prodr. 16² (1868) 390; F.-Vill. Novis. App. (1880) 212; Merr. in Forest. Bur. (Philip.) Bull. 1 (1903); 15 (*khasya*).

Luzon (Ifugao, Bontoc, Lepanto, Benguet, Nueva Vizcaya, Nueva Ecija). The dominant species in the Mountain Province, its altitudinal range being from about 1,000 to 2,700 m. Burma and Indo-China.

Local names: Alal (Ig.); balibo (Ig.); bariat (Bon.); bata (Bon.); batáng (Bon.); bebe (Bon.); bolbol (Ig.); bubu (Ig.); bulhúl (Ig.); lolol (Ig.); parúa (Ilk.); sáhing (Tag.); sáleng (Bon., Ilk., Ig., Ting.); sálit (Sbl.); talang (Gad.); tapúlau (Sbl.).

PINUS MERKUSII Jungh. & DeVr. in Pl. Nov. Ind. Bat. Orient. ⁵ (1845) t. 2, Bot. Zeit. 4 (1846) 13; Miq. Fl. Ind. Bat. 2 (1859) 1069; F.-Vill. Novis. App. (1880) 211; Vidal Sinopsis Atlas (1883) 43, t. 98, f. B, Rev. Pl. Vasc. Filip. (1886) 296; Perk. Frag. Fl. Philip. (1904) 35; Merr. in Forest. Bur. (Philip.) Bull. 1 (1903) 15, Govt. Lab. Publ. (Philip.) 6 (1904) 6; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 169; Shaw Gen. Pinus (1914) 58, t. 23, f. 198-200.

Luzon (Zambales), Mindoro. Very similar to *P. insularis* Endl. but with only two leaves in a fascicle instead of three; in places it occurs as low as 100 m altitude. Burma, Indo-China, Sumatra, ? Borneo.

Local names: Agú (Mang.); agúu (Mang.); sálit (Sbl.); sálong (Sbl.); tapúlau (Sbl.).

The following species of Pinaceae have been credited to the Philippines, but are all introduced and but rarely cultivated and have no claims to be considered as representatives of the Philippine flora:

ARAUCARIA EXCELSA R. Br.; F.-Vill. Novis. App. (1880) 211.

This does not thrive in the Philippines and never reaches maturity here.

CUPRESSUS TORULOSA Don; F.-Vill. l. c.

This is rarely cultivated and is commonly known by its Spanish name *cipres*.

JUNIPERUS OCCIDENTALIS Hook.; Usteri Beitr. Ken. Philip. Veg. (1905) 134.
Cultivated according to Usteri; I have seen no specimens.

GNETACEAE

1. GNETUM Linnaeus

GNETUM ARBOREUM Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 174, t. 32.

Luzon (Rizal, Nueva Ecija, Tayabas), *B. S. 9439 Robinson, 29635, 26388 Ramos & Edaño*. In the mossy forest, altitude about 900 m. Endemic.

Additional collections show this to be a vine, not a tree, as Robinson and Foxworthy supposed.

Local name: Lapiñgat (Neg.).

GNETUM GNEMON Linn. Mant. 1 (1867) 125; Blume Rumphia 4 (1848) 3, t. 176, 178 B, f. 5; Parl. in DC. Prodr. 16² (1868) 349; Blanco Fl. Filip. (1837) 747, ed. 2 (1845) 514, ed. 3, 3 (1879) 147; F.-Vill. Novis. App. (1880) 210; Vidal Phan. Cuming. Philip. (1885) 160, Rev. Pl. Vasc. Filip. (1886) 295; Karsten in Ann. Jard. Bot. Buitenz. 11 (1893) 203; Merr. Forestry Bur. (Philip.) Bull. 1 (1903) 16, Philip. Journ. Sci. 1 (1906) Suppl. 24, Sp. Blancoanae (1918) 53; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 173.

Gnetum vinosum Elm. Leaf. Philip. Bot. 7 (1915) 2673.

Luzon (Ilocos Norte, Bataan, Batangas, Laguna, Tayabas, Camarines), Mindoro, Palawan, Sibuyan, Panay, Samar, Leyte, Siargao, Bucas Grande, Mindanao. In forests at low and medium altitudes. Tropical Africa and Asia through Malaya to New Guinea.

Local names: Babayong (Sub.); bágo (Buk., Bik., Bag., C. Bis., Mbo., Tag.); bágo-sili (C. Bis.); banágo (Bis.); kugitis (Mbo.); kúman (Mand.); lamparan (Tag.); magatúnḡal (Mag.); nábo (Bik.).

GNETUM GNEMONOIDES Brongn. in Duperry Voy. Bot. (1829) 12; Merr. Interpret. Herb. Amb. (1917) 78.

Gnetum rumphianum Becc. Malesia 1 (1877) 182.

Gnetum verrucosum Karst. in Ann. Jard. Bot. Buitenz. 11 (1893) 216.

Funis gnemoniformis Rumph. Herb. Amb. 5 (1747) 12, t. 8.

Luzon (Tayabas, east coast), Basilan, *B. S. 15496 Reillo, 29096 Ramos & Edaño*. In forests at low or medium altitudes. Moluccas, New Guinea.

GNETUM INDICUM (Lour.) Merr. Interpret. Herb. Amb. (1917) 77, Sp. Blancoanae (1918) 53.

Abutua indica Lour. Fl. Cochinch. (1790) 630.

Gnetum funiculare Brongn. in Duperry Voy. Bot. (1829) 12; Miq. Fl.

Ind. Bat. 2 (1859) 1068; F.-Vill. Novis. App. (1880) 211.

Gnetum latifolium Blume Fam. Nov. (1833) 30, Rumphia 4 (1848) 5, t. 174; Presl Epim. (1851) 236; Miq. Fl. Ind. Bat. 2 (1859) 1067; Parl. in DC. Prodr. 16² (1868) 350; F.-Vill. Novis. App. (1880) 210; Vidal Phan. Cuming. Philip. (1885) 160, Rev. Pl. Vasc. Filip. (1886) 295, Sinopsis Atlas (1883) 42, t. 97, f. A; Warb. Monsunia (1900) 195; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 25; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 174.

Thoa pendula Blanco Fl. Filip. (1837) 746.

Thoa edulis Blanco Fl. Filip. ed. 2 (1845) 514, ed. 3, 3 (1879) 146, non Willd.

Gnetum philippinense Warb. *Monsunia* (1900) 196.

Gnetum scandens F.-Vill. *Novis. App.* (1880) 211, non Roxb.

Gnetum laxifrutescens Elm. *Leaf. Philip. Bot.* 4 (1912) 1478.

Luzon (Cagayan to Sorsogon), Mindoro, Palawan, Sibuyan, Panay, Leyte, Samar, Siargao, Mindanao. In forests at low and medium altitudes; distinctly variable in vegetative characters. Represented by very numerous collections. India to southern China and Malaya.

Local names: Biás (Tag.); bias-biás (Tag.); bulso (Buk.); dadaotum (Mbo.); kadiát (Itñ.); kalát (Ig., Ilk.); kaliát (Ilk.); kandiát (Ig.); konjât (Ig.); koliát (Tag.); kuliád (Ibn.); kuliát (Bik., Tag., Pamp., Mang.); lamparahan (Tag.); malaígot (S. L. Bis.); nonok (P. Bis.); tobal (Tag.).

GNETUM MINUS Foxw. in *Philip. Journ. Sci.* 6 (1911) *Bot.* 176, t. 33.

Luzon (Benguet), *B. S.* 2513 *Mearns*. In forests or thickets, altitude probably about 1,500 m. Endemic.

EXCLUDED SPECIES

GNETUM NEGLECTUM Blume; F.-Vill. *Novis. App.* (1880) 211.

ANGIOSPERMAE

MONOCOTYLEDONAE

TYPHACEAE

1. TYPHA Linnaeus

TYPHA CAPENSIS Rohrb. in *Verh. Bot. Ver. Brandenb.* 11 (1869) 96;

Hallier f. in *Lorenz Nova Guinea* 8 (1913) 911.

Typha angustifolia Blanco *Fl. Filip.* (1837) 687, ed. 2 (1845) 477, ed. 3, 3 (1879) 91; *Naves Novis. App.* (1882) 297, non Linn.

Typha angustifolia Linn. subsp. *javanica* Graebn. in *Perk. Frag. Fl. Philip.* (1904) 42, *Merr. Fl. Manila* (1912) 66, *Sp. Blancoanae* (1918) 53, non Schnizl.

Luzon (Laguna), Mindoro, Negros, *Merrill* 895, *Sp. Blancoanae* 33, *Piper* 74, *B. S.* 9881 *Robinson*, *F. B.* 9775 *Merritt*. In fresh water swamps at low altitudes. Africa and Madagascar, Philippines, New Guinea.

Local names: Balangót (Tag., S. L. Bis.); homai-homai (Bis.); kaidkéd (Pang.); lampakánai (C. Bis.); tubal-tubal (C. Bis.).

TYPHA DOMINGENSIS Pers. *Syn.* 2 (1807) 532; Graebn. in *Engl. Pflanzenreich* 2¹ (1900) 14; Hallier f. in *Lorenz Nova Guinea* 8 (1913) 912, in nota.

Mindoro, Panay, Negros, Cebu, *F. B.* 13371 *Curran*, 25893 *Tamesis & Azurin*, *B. S.* 11125 *Ramos*. In swamps at low altitudes. Tropical America; Java, Timor.

Local names: Balangót (P. Bis.); buhai-búhai (P. Bis.); lampakánai (C. Bis.).

TYPHA SHUTTLEWORTHII Koch & Sonder in *Koch Syn. ed.* 2 (1844) 786; Hallier f. in *Lorenz Nova Guinea* 8 (1913) 913, in nota.

Typha angustifolia Vidal *Rev. Pl. Vasc. Filip.* (1886) 280, non Linn.

Typha orientalis Presl *Epim.* (1851) 239; C. B. Rob. in *Bull. Torr.*

Bot. Club 35 (1908) 63; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 394.

Typha shuttleworthii Koch & Sond. subsp. *orientalis* Graebn. in Engl. Pflanzenreich 2¹ (1900) 10, quoad syn. Presl.

Luzon (Benguet, Bontoc, Sorsogon), Cebu, Mindanao (Bukidnon, Lanao), *Cuming 1767, Elmer 6382, 16244, Clemens 856, B. S. 24984 Fénix*. In swamps at low and medium altitudes, ascending to 1,400 m. Europe to China.

Local names: Anibung (Bon.); badok-badok (Ilk.); dosi-dosi (Ig.); palabog (Buk.).

I have arbitrarily followed Hallier f. in this classification of the Philippine forms but am by no means satisfied that the identifications are correct.

PANDANACEAE

1. FREYCINETIA Gaudichaud

FREYCINETIA ACUTIFOLIA Merr. in Philip. Journ. Sci. 13 (1918) Bot. 267.

Luzon (Camarines), Catanduanes, *B. S. 30319 Ramos, 33561 Ramos & Edaño*. In damp primary forests at low altitudes. Endemic.

FREYCINETIA ANGULATA C. B. Rob. ex Martelli in Webbia 3 (1910) 16, 309.

Luzon (Albay), Samar, Panay, *B. S. 6221 Robinson, 24813 Edaño, 32393 McGregor*. In forests at low altitudes. Endemic.

FREYCINETIA APAYAOENSIS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 269.

Luzon (Apayao, Ilocos Norte), *B. S. 28034 Fénix, 33111 Ramos*. In forests along streams at low altitudes. Endemic.

FREYCINETIA APOENSIS Martelli in Leaf. Philip. Bot. 3 (1911) 1111.

Mindanao (Davao), *Elmer 10781*. In forests along streams, altitude about 1,200 m. Endemic.

Local name: Rasrás (Bag.).

FREYCINETIA ATOCENSIS Martelli in Webbia 3 (1910) 28, 309.

Luzon (Benguet), (*Loher 1570*). Probably in forests, altitude at or above 1,000 m. Endemic.

FREYCINETIA AURICULATA Merr. in Philip. Journ. Sci. 3 (1908) Bot. 312; Martelli in Webbia 3 (1910) 15, 309.

Palawan, *B. S. 876 Foxworthy*. In forests at low or medium altitudes. Endemic.

FREYCINETIA BANAHAENSIS Elm. Leaf. Philip. Bot. 1 (1907) 215; Martelli in Webbia 3 (1910) 29, 309.

Luzon (Laguna, Tayabas), *Elmer 7902, 18424*. In primary forests, altitude 800 to 2,000 m. Endemic.

FREYCINETIA BATANENSIS Martelli in Webbia 3 (1910) 29, 309..

Batan Islands, Luzon (Ilocos Norte), *B. S. 3806 Fénix, 33301 Ramos*. In forests, altitude about 1,000 m. Endemic.

Local name: Uyód (Iv.).

FREYCINETIA BOTULIFORMIS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 268.

Luzon (Sorsogon), *B. S. 23363 Ramos*. In primary forests at medium altitudes. Endemic.

FREYCINETIA BULUSANENSIS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 268.

Luzon (Sorsogon), Panay, *B. S. 23686 Ramos, 32571 McGregor, Elmer 14978, 16294* as *F. vulcanica* Elm. Apparently in the mossy forest, altitude at or above 900 m. Endemic.

FREYCINETIA CUERNOSENSIS Martelli in Webbia 3 (1910) 26, 310.

Luzon (Sorsogon), Negros, *Elmer 9677, 16314, 14914*. In forests at medium altitudes. Endemic.

FREYCINETIA CUMINGIANA Gaudich. Bot. Voy. Bonite (1839-46), t. 37, f. 12-14, t. 60; C. B. Rob. in Bull. Torr. Bot. Club 35 (1908) 64; Martelli in Webbia 3 (1910) 19, 310.

Freycinetia luzonensis Presl Epim. (1851) 238 (incl. var. *heterophylla* Presl); Walp. Ann. 3 (1852-53) 494; Miq. Fl. Ind. Bat. 3 (1856) 172; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 280; Naves Novis. App. (1882) 286; Warb. in Engl. Pflanzenreich 3 (1900) 35, f. 10, D-E; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 311; C. B. Rob. op. cit. 6 (1911) Bot. 193.

Luzon (Cagayan, Camarines, Sorsogon), *Cuming 1455, Weber 1585, Elmer 15394* as *F. membranifolia* Elm., *15930, 16016* as *F. leuteocarpa* Elm., *B. S. 13874 Ramos, F. B. 11361 Curran*. In forests at low and medium altitudes. Endemic.

Local name: Alapúnti (C. Bis.).

FREYCINETIA CURRANII Merr. in Philip. Journ. Sci. 3 (1908) Bot. 312; Martelli in Webbia 3 (1910) 14, 310.

Luzon (Camarines, Sorsogon), Catanduanes, Panay, Camiguin de Misamis, *F. B. 11359 Curran, 23789 Alvarez, B. S. 35735 Martelino & Edaña, 14642, 30267, 30317 Ramos, Elmer 16590*. In primary forests at low and medium altitudes. Endemic.

FREYCINETIA DISCOIDEA Martelli in Leaf. Philip. Bot. 3 (1911) 1113.

Sibuyan, *Elmer 12197*. In forests, altitude about 300 m. Endemic.

FREYCINETIA ENSIFOLIA Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 5, Philip. Journ. Sci. 1 (1906) Suppl. 25, 3 (1908) Bot. 315; Martelli in Webbia 3 (1910) 32, 310; Elm. Leaf. Philip. Bot. 3 (1911) 1118.

Luzon (Bontoc, Benguet, Bataan, Pampanga), Mindoro, Negros, *Merrill 3242, Elmer 6840, Whitford 329, B. S. 1944 Foxworthy, F. B. 6285 Curran, 1347 Borden, 2624 Meyer, B. S. 39472 Ramos*. In the mossy forest, altitude 1,200 to 1,600 m. Endemic.

Local names: Kanġingai (Ig.); kakaag (Bon.).

FREYCINETIA FEROX Warb. in Engl. Pflanzenreich 3 (1900) 33; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 310; Martelli in Webbia 3 (1910) 13, 310.

Freycinetia lucbanensis Elm. Leaf. Philip. Bot. 1 (1907) 212; Martelli in Webbia 3 (1910) 14, 312.

Luzon (Tayabas), *Warburg* in herb. Berol., *Elmer* 8230. In primary forests, altitude about 500 m. Endemic.

FREYCINETIA GITINGIANA Martelli in Leaf. Philip. Bot. 3 (1911) Bot. 1112.

Sibuyan, *Elmer* 12358. In forests, altitude about 500 m. Endemic.

FREYCINETIA INSIPIDA Martelli in Leaf. Philip. Bot. 3 (1911) 1114.

Sibuyan, *Elmer* 12426. In forests along streams, altitude 500 to 600 m. Endemic.

Local name: Bagánot (P. Bis.).

FREYCINETIA JAGORII Warb. in Engl. Pflanzenreich 3 (1900) 39; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 313; Martelli in Webbia 3 (1910) 25, 311.

Luzon (Sorsogon), Biliran, Samar, Leyte, Mindanao (Agusan, Lanao, Zamboanga), *Elmer* 15378, 16210, 15057, 13717, *Wenzel* 306, *Merrill* 8300, *Clemens*, B. S. 18852 *McGregor*, 17395 *Ramos*, 37032 *Ramos & Edaño*. In forests at low and medium altitudes. Endemic.

Local name: Magalop (Sub.).

FREYCINETIA LEPTOPHYLLA Martelli in Webbia 3 (1910) 20, 312.

Luzon (Rizal, Laguna), (*Loher* 5450), B. S. 15006 *Ramos*. In forests at medium altitudes. Endemic.

FREYCINETIA LOHERI Martelli in Webbia 3 (1910) 15, 312.

Luzon (Lepanto, Rizal), (*Loher* 1577, 1578, 5469). Probably from forests at or above 1,200 m altitude. Endemic.

FREYCINETIA MAXIMA Merr. in Philip. Journ. Sci. 3 (1908) Bot. 310; Martelli in Webbia 3 (1910) 13, 312, Leaf. Philip. Bot. 3 (1911) 1111.

Luzon (Laguna, Tayabas, Camarines, Albay, Sorsogon), Samar, *F. B.* 10754, 1238 *Curran*, 25374 *Mabesa*, B. S. 23343, 17444 *Ramos*, 6345 *Robinson*, *Elmer* 17530, 14697, 18152. In primary forests at low and medium altitudes, ascending to 500 m. Endemic.

FREYCINETIA MEGACARPA Merr. in Philip. Journ. Sci. 3 (1908) Bot. 314; Martelli in Webbia 3 (1910) 31, 312, Leaf. Philip. Bot. 3 (1911) 1117.

Mindanao (Agusan, Lanao), *Clemens*, *Elmer* 14242. In forests, altitude about 650 m. Endemic.

FREYCINETIA MERRILLII Elm. Leaf. Philip. Bot. 1 (1907) 216; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 314; Martelli in Webbia 3 (1910) 31, 313.

Luzon (Cagayan, Tayabas), *Elmer* 9010, B. S. 10727 *McGregor*. In forests along streams, altitude about 600 m. Endemic.

FREYCINETIA MONOCEPHALA Elm. Leaf. Philip. Bot. 1 (1906) 78 bis, (1907) 218; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 314; Martelli in Webbia 3 (1910) 31, 313; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 193.

Luzon (Rizal, Laguna, Tayabas), *Elmer* 7380, 9012, *Whitford* 971, *Merrill Phil. Pl.* 423, *F. B.* 13094 *Curran*, B. S. 29372 *Ramos & Edaño*. In forests, altitude about 700 m. Endemic.

FREYCINETIA MONTALBANICA Martelli in Webbia 3 (1910) 18, 313; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 193.

Luzon (Rizal), (*Loher s. n.*, 5451). Probably in forests at medium altitudes. Of this I have seen only a photograph of *Loher* 5451. Endemic.

FREYCINETIA MULTIFLORA Merr. in Philip. Journ. Sci. 2 (1907) Bot. 259, 3 (1908) Bot. 312; Elm. Leaf. Philip. Bot. 1 (1907) 212; Martelli in Webbia 3 (1910) 17, 313.

Freycinetia luzonensis Naves in Blanco Fl. Filip. ed. 3 (1877-83) t. 437, non Presl.

Luzon (Laguna, Bataan), Mindoro, Mindanao, *Merrill* 5647, *Elmer* 8039, 9009, 17974, *Clemens* 1028, *B. S.* 16857 *Serviñas*, 15117, 2092 *Ramos*, 11285 *McGregor*. In primary forests up to 900 m altitude. Endemic.

FREYCINETIA NEGROSENSIS Merr. in Philip. Journ. Sci. 3 (1908) Bot. 313; Martelli in Webbia 3 (1910) 28, 313.

Luzon (Laguna, Sorsogon), Leyte, Negros, Mindanao (Agusan), *Whitford* 1541, *Elmer* 16161, 13766, as *F. urdanetensis* Elm., *F. B.* 26216 *Franco*, *B. S.* 16637 *Ramos*. In forests at medium altitudes, ascending to 1,200 m. Endemic.

Local name: Pandán-magamoi (C. Bis.).

FREYCINETIA OBLONGIFOLIA Merr. in Philip. Journ. Sci. 3 (1908) Bot. 310; Martelli in Webbia 3 (1910) 16, 313.

Panay, Mindanao (Surigao), *Bolster* 249, 342, *B. S.* 32533 *McGregor*, 31060 *Ramos* & *Edaño*. In primary forests at low altitudes. Endemic.

FREYCINETIA PALAWANENSIS Merr. ex Elm. Leaf. Philip. Bot. 1 (1907) 216, Philip. Journ. Sci. 3 (1908) Bot. 313; Martelli in Webbia 3 (1910) 23, 313.

Freycinetia elmeri Martelli in Webbia 3 (1910) 22, 310, Leaf. Philip. Bot. 3 (1911) 1117.

Luzon (Tayabas, Sorsogon), Palawan, Sibuyan, Panay, Mindanao (Agusan, Lanao), *B. S.* 706 *Foxworthy*, 23707 *Ramos*, 32527, 32231 *McGregor*, 35753 *Martelino* & *Edaño*, *Elmer* 12072, 7810, 9386, 14030, *Clemens* 796. In primary forests, altitude 600 to 1,000 m. Endemic.

FREYCINETIA PERIPIEZOCARPA Martelli in Webbia 3 (1910) 314.

Luzon (Ifugao, Bontoc, Benguet), Negros, *F. B.* 18020 *Merritt*, 13638 *Curran* & *Foxworthy*, 10858 *Curran*, *Merrill Phil. Pl.* 727. On ridges in thickets and forests, altitude 1,200 to 1,600 m. Endemic.

FREYCINETIA PHILIPPINENSIS Hemsl. in Kew Bull. (1896) 165; Warb. in Engl. Pflanzenreich 3 (1900) 40; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 313; Martelli in Webbia 3 (1910) 25, 314.

Luzon (Cagayan, Tayabas, Camarines), Samar, Biliran, Mindanao (Bukidnon), *Gregory* 117, *Cuming* 1898, *B. S.* 18787 *McGregor*, 26028 *Fénix*, 22088, 22167 *Ramos*. In primary forests at low and medium altitudes. Endemic.

FREYCINETIA PLATYPHYLLA Merr. in Philip. Journ. Sci. 13 (1918) Bot. 267.

Samar, *B. S.* 24349 *Ramos*. In forests along streams at low altitudes. Endemic.

FREYCINETIA POLYSTACHYA Martelli in Webbia 3 (1910) 14, 314.

Luzon (Ilocos Norte, Rizal, Tayabas), (*Loher 5454*), *B. S. 26642 Ramos & Edaño, 33299 Ramos, 25540 Yates*. In forests at medium altitudes. Endemic.

FREYCINETIA RIGIDA Elm. Leaf. Philip. Bot. 1 (1908) 362; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 313; Martelli in Webbia 3 (1910) 26, 314.

Freycinetia hemsleyi Elm. Leaf. Philip. Bot. 1 (1907) 214, non Warb.

Luzon (Tayabas), *Elmer 7847*. In primary forests, altitude about 800 m. Endemic.

FREYCINETIA ROBINSONII Merr. in Philip. Journ. Sci. 3 (1908) Bot. 311; Martelli in Webbia 3 (1910) 17, 314.

Luzon (Ilocos Norte, Bataan, Pampanga), Leyte, *F. B. 752, 2466, 3037 Borden, 2194 Meyer, B. S. 2006 Foxworthy, 27256 Ramos, Whitford 1311, Merrill 3791*. In forests at low and medium altitudes. Endemic.

Var. **LATIFOLIA** Martelli in Webbia 3 (1910) 18, 314.

Luzon (Benguet, Laguna, Rizal), Negros, *Elmer 6196*. In forests at low and medium altitudes. Endemic.

Local names: Malabaling-uái (Tag.); malapandán (Tag.); maloran (Pamp.).

FREYCINETIA ROBUSTA Elm. Leaf. Philip. Bot. 8 (1919) 3097.

Luzon (Laguna), *Elmer 18026*. In primary forests at medium altitudes. Endemic.

FREYCINETIA ROSTRATA Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 177, 3 (1908) Bot. 315; Martelli in Webbia 3 (1910) 34, 314, Leaf. Philip. Bot. 3 (1911) 1118.

Samar, Negros, Siargao, Mindanao (Lanao), *Merrill 5235, Clemens s. n., B. S. 24787 Edaño, F. B. 19097 Curran*. In forests at low and medium altitudes. Endemic.

Local names: Rasrás (Bag.); sagatap (P. Bis.).

FREYCINETIA SCABRIPES Warb. in Engl. Pflanzenreich 3 (1900) 41; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 313, 394; Martelli in Webbia 3 (1910) 27, 314.

Freycinetia dilatata Merr. ex Elm. Leaf. Philip. Bot. 1 (1907) 214, Philip. Journ. Sci. 3 (1908) Bot. 313; Martelli in Webbia 3 (1910) 310.

Luzon (Cagayan, Bataan, Rizal, Tayabas, Sorsogon), *F. B. 4529 Maule, B. S. 99 Foxworthy, Elmer 9008, Merrill 2301*. In forests at low and medium altitudes. Endemic.

Local name: Palindán (Tag.).

FREYCINETIA SPHAEROCEPHALA Gaudich. Bot. Voy. Bonite (1843) t. 52; Solms-Laub. in Linnaea 42 (1878) 96; Warb. in Engl. Pflanzenreich 3 (1900) 35; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 314; Martelli in Webbia 3 (1910) 33, 315.

Freycinetia strobilacea Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 280; F.-Vill. Novis. App. (1882) 285, non Blume.

Freycinetia globosa Merr. in Philip. Journ. Sci. 2 (1907) Bot. 260; Elm. Leaf. Philip. Bot. 1 (1907) 217.

Luzon (Ilocos Norte, Apayao, Laguna, Camarines, Albay, Sorsogon), Mindoro, *Cuming* 839, *Merrill* 5791, *Elmer* 14920, *Wenzel* 511, *B. S.* 16593, 22016, 20510 *Ramos*. In forests at low and medium altitudes, ascending to 1,000 m. Endemic.

Local name: Looi-lóoi (Bik.).

The figure of *Freycinetia strobilacea* Blume given by Vidal, Sinopsis Atlas (1883) 42, t. 95, f. B, was copied from Blume's original in Rumphia fide Vidal l. c. Blume's species does not occur in the Philippines.

FREYCINETIA SUBFLAGELLATA Elm. Leaf. Philip. Bot. 8 (1919) 3098.

Luzon (Laguna), *Elmer* 17967. In primary forests at medium altitudes. Endemic.

FREYCINETIA SUPERBA Martelli in Leaf. Philip. Bot. 3 (1911) 1115.

Sibuyan, *Elmer* 12257. In forests, altitude 300 to 600 m. Endemic.

FREYCINETIA VIDALII Hemsl. in Kew Bull. (1896) 166; Warb. in Engl.

Pflanzenreich 3 (1900) 36; Merr. in Philip. Journ. Sci. 3 (1908) Bot.

311; Martelli in Elm. Leaf. Philip. Bot. 3 (1910) 21, 315.

Freycinetia confusa Elm. Leaf. Philip. Bot. 1 (1907) 213.

Luzon (Cagayan, Apayao, Nueva Ecija, Rizal, Tayabas), *Vidal* 3964, *Elmer* 9007, *B. S.* 16600, 13861 *Ramos*, 28434 *Fénix*, *F. B.* 19573 *Curran*, 22192 *Alvarez*. In primary forests at low and medium altitudes. Endemic.

Local name: Takiakad (Ig.).

FREYCINETIA WARBURGII Elm. Leaf. Philip. Bot. 1 (1907) 218; Merr.

in Philip. Journ. Sci. 3 (1908) Bot. 315; Martelli in Webbia 3 (1910)

34, 315; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 193.

Luzon (Laguna, Tayabas, Sorsogon), *Elmer* 8229, 15833, as *F. sorsogonensis* Elm., *B. S.* 14988, 16604, 20608, 10925 *Ramos*. In forests at medium altitudes. Endemic.

FREYCINETIA WILLIAMSII Merr. in Philip. Journ. Sci. 3 (1908) Bot.

315, 394; Martelli in Webbia 3 (1910) 32, 315.

Batan Islands, Luzon (Ifugao, Bontoc, Benguet, Rizal, Pangasinan, Laguna, Tayabas), Panay, *Williams* 1013, *Elmer* 5857, 18006, *Merrill* 8611, *B. S.* 3786, 29862 *Fénix*, 29737 *Ramos & Edaño*, 22233 *Catalan*, 19755 *McGregor*, 9380, 17120 *Robinson*. In forests, extending into the mossy forest, altitude 500 to 1,600 m. Endemic.

Local names: Pandán (Ig.); vayasúbas (Iv.).

DOUBTFUL AND EXCLUDED SPECIES

FREYCINETIA GRAMINEA Blume; Naves Novis. App. (1882) 286; Usteri Beitr. Ken. Philip. Veg. (1905) 130.

This Javan species is not known from the Philippines, both of the above records being undoubtedly based on erroneously identified specimens.

FREYCINETIA INSIGNIS Blume; Vidal Cat. Pl. Prov. Manila (1880) 46; Naves Novis App. (1882) 286.

This Javan species certainly does not occur in the Philippines; from the local name cited by Vidal his specimens probably were *Pandanus sabotan* Blanco.

TILLANDSIA PSEUDO-ANANAS Blanco Fl. Filip. (1837) 853, ed. 2 (1845) 162, ed. 3, 1 (1877) 292.

This was reduced by Naves to *Freycinetia insignis* Blume, a species that does not extend to the Philippines. I now consider that Blanco's description applies to *Freycinetia*, but further identification of the species is impossible from the very imperfect description given by Blanco. It is possibly the same as *Pandanus copelandii* Merr.

2. PANDANUS Linnaeus

PANDANUS ACLADUS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 265.

Catanduanes, Mindanao (Bukidnon), *B. S. 30462 Ramos, 39172 Ramos & Edaño*. Along streams in forests, altitude about 150 m. Endemic. Local name: Ulañgón (Buk.).

PANDANUS APICULATUS Merr. in Philip. Journ. Sci. 17 (1920) 240.

Mindanao (Surigao), *B. S. 34572 Ramos & Pascasio*. On ridges, altitude about 650 m. Endemic.

PANDANUS APOENSIS Martelli in Leaf. Philip. Bot. 3 (1911) 1129, Webbia 4¹ (1913) 6, t. 13, f. 12-14.

Mindanao (Davao), *Elmer 11313*. In primary forests, altitude about 1,200 m. Endemic.

Local names: Baráia (Bag.); maráia (Bag.).

PANDANUS BARAI Martelli in Leaf. Philip. Bot. 3 (1911) 1126, Webbia 4¹ (1913) 7, t. 38, f. 10-13.

Mindanao (Davao), *Elmer 10924*. In damp forests, altitude about 1,050 m. Endemic.

Local name: Barai (Bag.).

PANDANUS BILIRANENSIS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 266.

Biliran, *B. S. 18895 McGregor*. In forests, altitude about 300 m. Endemic.

PANDANUS BOTRYOIDES Martelli in Philip. Journ. Sci. 3 (1908) Bot. 66, Webbia 4¹ (1913) 8, t. 4, f. 14-16; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 193.

Polillo, Mindanao (Surigao, Agusan, Lanao), *Merrill 5442, Weber 1212, Clemens 984, B. S. 34570 Ramos & Pascasio, 10473 Robinson*. In forests along tidal streams, and at Lake Lanao along the lake shore, altitude about 650 m. Endemic.

Specifically distinct from *P. radicans* Blanco?

PANDANUS BREVISPATHUS Martelli in Philip. Journ. Sci. 3 (1908) Bot. 69, Webbia 4¹ (1913) 8, t. 26, f. 3-5.

Pandanus polycephalus Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 8, non Lour.

Mindanao (Davao), *Copeland 442*. In thickets near the seashore. Endemic.

PANDANUS BREVISTIPES Martelli in Webbia 4¹ (1913) 8, 4² (1914) 421, t. 38, f. 4-9.

Luzon (Batangas), *F. B. 7719 Curran & Merritt*. On ridges in forests, altitude about 800 m. Endemic.

Local name: Pandán (Tag.).

PANDANUS CALCEIFORMIS Martelli in *Leaf. Philip. Bot.* 3 (1911) 1127, *Webbia* 4¹ (1913) 9, t. 35, f. 3-6.

Mindanao (Bukidnon, Davao), *Elmer 11654, B. S. 38580, 39170 Ramos & Edaño*. On forested ridges, altitude about 1,350 m. Endemic.

Local name: Baráui (Bag.).

PANDANUS CAMARINENSIS Merr. in *Philip. Journ. Sci.* 14 (1919) 367.

Luzon (Camarines), *B. S. 33635 Ramos & Edaño*. In forests at low altitudes. Endemic.

PANDANUS CAUDATUS Merr. in *Govt. Lab. Publ. (Philip.)* 29 (1905) 6; Martelli in *Philip. Journ. Sci.* 3 (1908) Bot. 64, *Webbia* 4¹ (1913) 9, t. 4, f. 1-3.

Luzon (Benguet), *Elmer 6143*. In thickets or forests, altitude about 1,000 m. Endemic.

Local name: Pañgíhang (Ig.).

PANDANUS CLEMENTIS Merr. in *Philip. Journ. Sci.* 1 (1906) Suppl. 178, t. 1, 3 (1908) Bot. 68; Martelli in *Webbia* 4¹ (1913) 10, t. 27, f. 6-10.

Panay, Mindanao (Lanao), *Clemens 436, Merrill 6708*. In forests, altitude 100 to 700 m. Endemic.

PANDANUS COPELANDII Merr. in *Govt. Lab. Publ. (Philip.)* 17 (1904) 7, *Philip. Journ. Sci.* 3 (1908) Bot. 71; Martelli op. cit. 71, *Leaf. Philip. Bot.* 3 (1911) 1132, *Webbia* 4¹ (1913) 11, t. 31, f. 5-7.

Pandanus muricatus Elm. *Leaf. Philip. Bot.* 1 (1906) 76.

Luzon (Cagayan, Apayao, Benguet, Pangasinan, Nueva Vizcaya, Nueva Ecija, Zambales, Tayabas, Camarines, Sorsogon), Polillo, Catanduanes, Samar, Bucas Grande, Panay, Leyte, Mindanao (Bukidnon, Zamboanga, Surigao). A very characteristic species, represented by more than 30 collections. In forests at low and medium altitudes, ascending to 1,000 m. Endemic.

Tillandsia pseudo-ananas Blanco is possibly a synonym of this rather than a *Freycinetia*; see page 15.

Local names: Baléo (Bis.); baléu (P. Bis.); balíku (C. Bis.); balói (Mbo.); bariéu (Bag.); baríu (Bik.); barói (Mbo.); lagutlut (Tag.); pandán (Tag., Ibn.); pangdán (Ilk., Pang.); patága (Ibn.); sirí (Ibn.).

PANDANUS CUMINGIANUS Martelli in *Philip. Journ. Sci.* 3 (1908) Bot. 70, *Webbia* 4¹ (1913) 11, t. 25, f. 13-15.

Negros, (*Cuming s. n.* in herb. Mus. Brit.). I have seen only a photograph of the syncarps. The leaves are unknown. Endemic.

PANDANUS DECIPIENS Martelli in *Webbia* 4¹ (1913) 12, 4² (1914) 402, t. 37, f. 7-9.

Palawan, *Elmer 12593, Merrill 9246*. In thickets and forests at low altitudes, often back of the mangrove swamps. Endemic.

Local name: Olañgó (Tagb.).

PANDANUS DINAGATENSIS Merr. in Philip. Journ. Sci. 17 (1920) 240.

Dinagat, B. S. 35183 Ramos & Pascasio. In forests along streams at low altitudes. Endemic.

PANDANUS DUBIUS Spreng. Syst. 3 (1826) 897; Miq. Fl. Ind. Bat. 3 (1855) 159; Naves Novis. App. (1882) 285; Warb. in Engl. Pflanzenreich 3 (1900) 50; Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 8; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 67, Leaf. Philip. Bot. 3 (1911) 1122, Webbia 4¹ (1913) 12.

? *Pandanus latifolius* Perr. in Mém. Soc. Linn. Paris 3 (1824) 134; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 72; C. B. Rob. op. cit. 306; Warb. in Engl. Pflanzenreich 3 (1900) 88.

Mindanao (Davao), *Copeland* 613. In thickets or forests along the seashore; Java, Borneo, Celebes, Moluccas, New Guinea to the New Hebrides, Caroline, and Marianne Islands.

Local names: Bákong (P. Bis.); báuang (P. Bis.); taboán (Bik.).

PANDANUS ESCULENTUS Martelli in Webbia 4¹ (1913) 13, 4² (1914) 403, t. 25, f. 1-3.

Luzon (Cagayan, Apayao, Ilocos Norte, Bulacan), *F. B.* 17813, 17817 *Curran*, 14009 *Merritt & Darling*, B. S. 28352 *Fénix*, 34089 *Ramos & Edaño*, 22208 *Ramos*. In forests at low and medium altitudes, ascending to 1,000 m. Endemic.

Local name: Lígít (Ibn.).

PANDANUS EXALTATUS Blanco Fl. Filip. (1837) 778, ed. 2 (1845) 536, ed. 3, 3 (1879) 183; Miq. Fl. Ind. Bat. 3 (1856) 163; Warb. in Engl. Pflanzenreich 3 (1900) 84; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 64, Webbia 4¹ (1913) 13, t. 10, f. 1-3, 9-11; Merr. Sp. Blancoanae (1918) 54.

Pandanus arayatensis Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 7, t. 3, Philip. Journ. Sci. 1 (1906) Suppl. 25; Martelli in Webbia 4¹ (1913) 6.

Pandanus banahaensis Elm. Leaf. Philip. Bot. 1 (1906) 79; Martelli in Webbia 4¹ (1913) 7 (*banahaoensis*).

Pandanus vidalii Martelli in Philip. Journ. Sci. 3 (1908) Bot. 65, Webbia 4¹ (1913) 36, t. 8, f. 11-13.

Pandanus fascicularis Naves Novis. App. (1882) 284, non Lam.

Luzon (Apayao, Benguet, Nueva Vizcaya, Pampanga, Rizal, Bataan, Tayabas), Catanduanes, *Whitford* 1325, *Elmer* 7574, 9074, *Merrill* 3832, *Sp. Blancoanae* 421, B. S. 30498 *Ramos*, 28256, 13546 *Fénix*, 11226 *McGregor*, 2701 *Ahern's collector*. In primary forests at medium altitudes, ascending to 800 m. Endemic.

Local names: Alasás (Tag.); bambán (Ibn.); pandán (Tag., Pamp., Ibn.); pangdán (Ilk.).

Forma **AHERNIANUS** Martelli in Philip. Journ. Sci. 3 (1908) Bot. 64, Webbia 4¹ (1913) 13, t. 10, f. 4-8, 12-15.

Luzon (Cagayan, Rizal, Pampanga), *Merrill Dec. Philip. Forest Fl.* 296, *F. B.* 17264, 17722 *Curran*. Habitat of the species from which it is scarcely distinguishable. Endemic.

PANDANUS GLAUCIPHYLLUS C. B. Rob. in Bull. Torr. Bot. Club 35 (1908) 64, 74; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 68, Leaf. Philip. Bot. 3 (1911) 1124, Webbia 4¹ (1913) 15, t. 27, f. 14-16.

Negros, Romblon, Siargao, Mindanao (Zamboanga), *Williams 2423*, *Copeland, F. B. 17372 Curran, B. S. 34855 Ramos & Pascasio*. In forests at low altitudes. Endemic.

Local name: Pandán (P. Bis.).

PANDANUS GRACILIS Blanco Fl. Filip. (1837) 778, ed. 2 (1845) 536, ed. 3, 3 (1879) 182, t. 446; Warb. in Engl. Pflanzenreich 3 (1900) 84; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 89, Philip. Journ. Sci. 2 (1907) Bot. 436, Sp. Blancoanae (1918) 56; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 68, Webbia 4¹ (1913) 16, t. 25, f. 16-24, 25-26.

Pandanus whitfordii Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 8, Philip. Journ. Sci. 1 (1906) Suppl. 25.

Luzon (Nueva Vizcaya, Zambales, Bataan, Rizal, Laguna, Tayabas, Camarines), Mindoro, Mindanao (Bukidnon, Davao), *Whitford 351*, *Merrill 7612*, *Sp. Blancoanae 890, 896*, *B. S. 23860, 22469, 39493 Ramos, 15783 Fénix, 34844 Ramos & Pascasio, F. B. 20188 Aguilar, 2944 Borden, 6003 Curran*. In primary forests at medium altitudes, ascending to 1,000 m. Endemic.

Local names: Ayasás (Tag.); lagúloi (Buk.); pandán (Tag.).

PANDANUS JULIFERUS Martelli in Webbia 4¹ (1913) 18, 4² (1914) 428, t. 43, f. 10-12.

Camiguin de Misamis, *B. S. 14670, 14671 Ramos*. Probably in forests, altitude about 1,000 m. Endemic.

This seems to be very near *Pandanus polyglossus* Martelli.

PANDANUS LATERALIS Martelli in Philip. Journ. Sci. 3 (1908) Bot. 68, Leaf. Philip. Bot. 3 (1911) 1125, Webbia 4¹ (1913) 20, t. 27, f. 31-33.

Mindanao (Davao), *Copeland 336, Elmer 11029*. In forests at low or medium altitudes. Endemic.

Local names: Bakúling (Bag.); pandán (C. Bis., Sul.).

PANDANUS LINNAEI Gaudich. forma **PHILIPPINENSIS** Martelli in Webbia 4¹ (1913) 21, t. 6, f. 1-2, Philip. Journ. Sci. 3 (1908) Bot. 60 (as *P. linnaei* Gaudich.).

Pandanus exaltatus Merr. in Philip. Journ. Sci. 2 (1907) Bot. 436, non Blanco.

Semirara, Busuanga, *Merrill 4140, B. S. 41264 Ramos*. In thickets at low altitudes. The species in Celebes, Key, and Aru Islands.

PANDANUS LOHERIANUS Martelli in Webbia 4¹ (1913) 22, 4² (1914) 423, t. 27, f. 17-21.

Luzon (Rizal, Laguna), *B. S. 20549 Ramos, 13192 Curran*. In forests along streams at low or medium altitudes. Endemic.

PANDANUS LUZONENSIS Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 6, t. 1, 2, Philip. Journ. Sci. 1 (1906) Suppl. 25; Martelli op. cit. 3 (1908) Bot. 67, Webbia 4¹ (1913) 22, t. 15, f. 5-9.

Pandanus calicarpus Martelli in Webbia 1 (1905) 365, 4¹ (1913) 9.

Luzon (Pangasinan, Zambales, Bulacan, Rizal, Bataan, Laguna). Chiefly in secondary forests at low altitudes; represented by numerous collections, *Merrill Phil. Pl.* 1931 and 759 are typical. Endemic.

Local names: Alasás (Tag.); dasá (Tag.); pandán (Tag., Ilk., Ibn.).

PANDANUS MAPOLA Martelli in Leaf. Philip. Bot. 3 (1911) 1124, *Webbia* 4¹ (1913) 22, t. 41, f. 14-17.

Sibuyan, *Elmer* 12139. In thickets or forests at low altitudes; the local name cited, *mapulá*, merely means red in reference to the color of the fruits. Endemic.

PANDANUS MARTELLII Elm. Leaf. Philip. Bot. 1 (1908) 272; Martelli in *Webbia* 4¹ (1913) 23, t. 11, f. 4-7.

Luzon (Tayabas), *Elmer* 9082, 9083, 17646, *B. S.* 23025 *McGregor*, 23829 *Ramos*. In forests, altitude about 650 m. Endemic.

PANDANUS MERRILLII Warb. in Perk. Frag. Fl. Philip. (1904) 50; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 71, *Webbia* 4¹ (1913) 23, t. 32, f. 35-37.

Palawan, *Merrill* 840, 7250, 9274, 9375, *B. S.* 232 *Bermejos*. In primary forests, sea level to 500 m altitude. Endemic.

Local name: Pandán (Tagb.).

PANDANUS MINDANAENSIS Martelli in Elm. Leaf. Philip. Bot. 3 (1911) 1121, *Webbia* 4¹ (1913) 24, t. 36, f. 4-7 (*mindanaoensis*).

Mindanao (Davao), *Elmer* 10823. In primary forests, altitude about 1,200 m. Endemic.

PANDANUS MULTIBRACTEATUS Merr. in Philip. Journ. Sci. 17 (1920) 241.

Mindanao (Surigao), *B. S.* 34571, 34819 *Ramos & Pascasio*. In forests along streams, altitude about 650 m. Endemic.

PANDANUS OCCULTUS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 265.

Palawan, *Merrill* 9361. In primary swamp forests at low altitudes. Endemic.

PANDANUS PALLIDUS Merr. in Govt. Lab. Publ. (Philip.) 29 (1905) 5; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 68, *Webbia* 4¹ (1913) 26, t. 27, f. 28-30.

Luzon (Benguet, Nueva Ecija), *Elmer* 5840, *Williams* 917, *B. S.* 26270 *Ramos & Edaño*, *F. B.* 5106, 5069 *Curran*. In forested stream depressions up to 1,500 m altitude. Endemic.

Local names: Pangchán (Ig.); tachiáng (Ig.).

PANDANUS PALOENSIS Elm. Leaf. Philip. Bot. 1 (1906) 75; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 68, Leaf. Philip. Bot. 3 (1911) 1123, *Webbia* 4¹ (1913) 26, t. 27, f. 22-27.

Luzon (Cagayan, Apayao, Camarines, Sorsogon), Leyte, Mindanao (Davao), *Elmer* 7343, 14858, 11224, *Wenzel* 1304, *Weber* 1568, *B. S.* 33556 *Ramos & Edaño*, 28072 *Fénix*, 13921 *Ramos*, *F. B.* 19569 *Curran*. In forests at low altitudes. Endemic.

Local name: Baróai-latong (Bag.).

PANDANUS PANAYENSIS Merr. in Philip. Journ. Sci. 14 (1919) 366.

Panay, *B. S. 30796 Ramos & Edaño, 35740 Martelino & Edaño*. In primary forests at low or medium altitudes. Endemic.

PANDANUS PHILIPPINENSIS Merr. in Philip. Journ. Sci. 13 (1918) Bot. 264.

Luzon (Kalinga, Nueva Vizcaya), *B. S. 11289 McGregor, 37495 Ramos & Edaño*. On forested slopes at medium altitudes. Endemic.

PANDANUS POLYGLOSSUS Martelli in Leaf. Philip. Bot. 3 (1911) 1130, Webbia 4¹ (1913) 28, t. 39, f. 16-18.

Mindanao (Davao), *Elmer 11335*. In primary forests, altitude about 1,350 m. Endemic.

PANDANUS RADICANS Blanco Fl. Filip. (1837) 780, ed. 2 (1845) 537, ed. 3, 3 (1879) 184; Miq. Fl. Ind. Bat. 3 (1856) 166; Warb. in Engl. Pflanzenreich 3 (1900) 85; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 89, Philip. Journ. Sci. 2 (1907) Bot. 436, Sp. Blancoanae (1918) 54; Elm. Leaf. Philip. Bot. 1 (1906) 74; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 66, Webbia 4¹ (1913) 29, t. 6, f. 8-10, *Pandanus olango* Blanco ex Espejo & Garcia Suppl. Cat. Seem. Hort. Bot. Manila (1876) 14, *nomen nudum*.

Luzon (Sorsogon), Leyte, *Elmer 7200, 14892*. In forests at low altitudes. Endemic.

Local names: Olañgó (S. L. Bis.); ouañgó (P. Bis.); oyañgó (Bik.); uañgó (P. Bis.); uyañgó (Mbo.).

PANDANUS RAMOSII Merr. in Philip. Journ. Sci. 17 (1920) 242.

Bucas Grande, Dinagat, *B. S. 35136, 35198 Ramos & Pascasio*. In dry forests at low altitudes. Endemic.

PANDANUS RECLINATUS Martelli in Webbia 4¹ (1913) 29, 4² (1914) 432, t. 27, f. 3, 4.

Palawan, *Elmer 12731*. In forests at low altitudes. Endemic.

PANDANUS SABOTAN Blanco Fl. Filip. (1837) 779, ed. 2 (1845) 537, ed. 3, 3 (1878) 184; Miq. Fl. Ind. Bat. 3 (1856) 166; Warb. in Engl. Pflanzenreich 3 (1900) 84; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 89, Sp. Blancoanae (1918) 56; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 72, Webbia 4¹ (1913) 30.

Luzon (Laguna, Rizal), *Merrill Sp. Blancoanae 185, F. B. 10099 Curran, B. S. 6532, 11934 Robinson, 11239 McGregor*. A planted pandan of which the fruits and flowers are unknown. The Filipinos state that it never produces fruit. It is perhaps but a form of *P. tectorius* Solander, and Martelli has so identified some of the specimens cited above. Endemic?

Local names: Parauan (Gad.); sabotán (Tag.); sibután (Tag.).

PANDANUS SIBUYANENSIS Martelli in Leaf. Philip. Bot. 3 (1911) 1120, Webbia 4¹ (1913) 31, t. 34, f. 4-6 (*sibujanensis*).

Sibuyan, *Elmer 12146*. In forests, altitude about 350 m. Endemic.

Local name: Pandan-bugás (P. Bis.).

PANDANUS SIMPLEX Merr. in Govt. Lab. Publ. (Philip.) 29 (1905) 6; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 67, Webbia 4¹ (1913) 31, t. 14, f. 11-14.

Pandanus utilisimus Elm. Leaf. Philip. Bot. 1 (1906) 80; Martelli

in Philip. Journ. Sci. 3 (1908) Bot. 67, *Webbia* 4¹ (1913) 36, t. 14, f. 3-5; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 193.

Luzon (Ifugao, Laguna, Tayabas, Sorsogon), Catanduanes, *Elmer* 7379, 16366, *Whitford* 782, *F. B.* 20166 *Aguilar*, 8507 *Curran*, *B. S.* 20033 *McGregor*, 6542 *Robinson*. In forests at low and medium altitudes; also planted. A species remarkable for its enormous syncarps. Endemic.

Local names: Bankoang (Tag.); ilohan (Tag.); kalagímai (Tag.); karagómoi (Bik.); karagúmoi (Bik., Bis., Tag.); pandán (Tag.); pandan-totóo (Tag.).

PANDANUS SUBACaulis Merr. in Philip. Journ. Sci. 13 (1918) Bot. 263.

Catanduanes, *B. S.* 30489 *Ramos*. In forests at low or medium altitudes. Endemic.

PANDANUS TECTORIUS Solander in Parkins. Journ. Voy. H. M. S. Endeavour (1773) 46; Warb. in Engl. Pflanzenreich 3 (1900) 46; Perk. Frag. Fl. Philip. (1904) 50; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 25, 3 (1908) Bot. 394, Fl. Manila (1912) 66, Sp. Blancoanae (1918) 55; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 61, Leaflet. Philip. Bot. 3 (1911) 118, *Webbia* 4¹ (1913) 33.

Pandanus odoratissimus Linn. f. Suppl. (1781) 424; Vidal Sinopsis Atlas (1883) 42, t. 95, f. A; F.-Vill. Novis. App. (1882) 284.

Pandanus malatensis Blanco Fl. Filip. (1837) 777, ed. 2 (1845) 535, ed. 3, 3 (1879) 181; Warb. in Engl. Pflanzenreich 3 (1900) 84; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 89; ? Martelli in Philip. Journ. Sci. 3 (1908) Bot. 70, *Webbia* 4¹ (1913) 22, t. 26, f. 22.

Pandanus spiralis Blanco Fl. Filip. (1837) 777, ed. 2 (1845) 535, ed. 3, 3 (1879) 181; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 89, non R. Br.

Pandanus blancoi Kunth Enum. 3 (1841) 583.

Pandanus tectorius Soland. var. *surigaensis* Martelli in Philip. Journ. Sci. 3 (1908) Bot. 64, *Webbia* 4¹ (1913) 34, t. 8, f. 3-7 (*surigaensis*).

Pandanus tectorius Soland. var. *sinensis* Warb. in Engl. Pflanzenreich 3 (1900) 48; Martelli in Philip. Journ. Sci. 3 (1908) Bot. 62, *Webbia* 4¹ (1913) 34, t. 9, f. 2-10.

Pandanus tectorius Soland. var. *spiralis* Martelli in Philip. Journ. Sci. 3 (1908) Bot. 62, *Webbia* 4¹ (1913) 34, t. 9, f. 1-7, t. 12, f. 8.

Pandanus tectorius Soland. var. *douglasii* Martelli forma *philippinensis* Martelli in Philip. Journ. Sci. 3 (1908) Bot. 63, *Webbia* 4¹ (1913) 33, t. 8, f. 8-10.

Pandanus vidalii Martelli in Philip. Journ. Sci. 3 (1908) Bot. 65, *Webbia* 4¹ (1913) 36, p. p., quoad syn. Vidal.

Pandanus coronatus Martelli in Philip. Journ. Sci. 3 (1908) Bot. 65 (incl. f. *minor*), *Webbia* 4¹ (1913) 11, t. 4, f. 8-13.

Along the seashore throughout the Philippines, abundant and forming characteristic thickets back of the beach. Indo-Malayan and Polynesian regions generally, extending to southern China and tropical Australia.

Local names: Bároi (Bik.); laha (Sul.); ohañgó (Iv.); padán (Iv.); pandán (Tag., P. Bis., C. Bis., Sub.); pandan-dágat (Tag.); pangdán (Ilk., Pang.); panglán (Ilk., Sbl.); panhakad (Bis.); uhañgó (Iv.).

PANDANUS TENUIPEDUNCULATUS Merr. in Philip. Journ. Sci. 17 (1920) 243.

Siargao, Mindanao (Surigao), *B. S. 34739, 34842 Ramos & Pascasio*. In open forests at low altitudes. Endemic.

PANDANUS ZAMBOANGENSIS Martelli in Webbia 4¹ (1913) 37, 4² (1914) 426, t. 42, f. 7-10.

Mindanao (Zamboanga, Bukidnon, Lanao), *Merrill 8051, 8065, F. B. 9494 Whitford & Hutchinson, B. S. 15696 Fénix, 37428, 37429 Ramos & Edaño*. In thickets and secondary forests at low altitudes. Endemic.

Local names: Balíu (Sub.); bariú (Sul.); pandán (Sul.).

DOUBTFUL AND EXCLUDED SPECIES

PANDANUS INERMIS Blanco Fl. Filip. ed. 2 (1845) 537, ed. 3, 3 (1879) 184, non Roxb.

This is unquestionably a *Pleomele* (*Dracaena*) and is referable to *Pleomele angustifolia* (Roxb.) N. E. Br.

PANDANUS KAFU Martelli in Webbia 4¹ (1913) 19, t. 19, f. 1-3.

"Is. Filippine" is an error, the type being from Guam, Marianne Islands.

PANDANUS PORTEANUS Lescuy. ex Hérincq Hort. Franc. 1 (1866) 16; Warb. in Engl. Pflanzenreich 3 (1900) 90; Martelli in Webbia 4¹ (1913) 28.

A *nomen nudum* based on a cultivated plant of Philippine origin.

Pandanus bagea Miq., *P. caricosus* Rumph., *P. ceramicus* Rumph., *P. drupaceus* Thouars., *P. elegans* Thouars., *P. furcatus* Roxb., *P. humilis* Rumph., *P. latifolius* Rumph., *P. littoralis* Jungh., *P. montanus* Miq., *P. moschatus* Rumph., *P. repens* Rumph., *P. spurius* Rumph., and *P. sylvestris* Rumph.

The above fourteen species, several of them of entirely doubtful status, were credited to the Philippines by Naves, Novis. App. (1882) 284, 285. With the exception of *Pandanus caricosus* Rumph. = *Scirpiodendron ghaeri* (Gaertn.) Merr. none of them are known to occur in the Philippines.

3. SARARANGA Hemsley

SARARANGA PHILIPPINENSIS Merr. in Govt. Lab. Publ. (Philip.) 29 (1905) 5; Elm. Leaf. Philip. Bot. 3 (1911) 1110.

Luzon (Tayabas, Camarines, Albay, Sorsogon), Sibuyan, Samar, Dinagat, Mindanao (Surigao, Bukidnon), *Merrill 5230, Whitford 749, Elmer 17206, B. S. 6401 Robinson, 35181, 34680 Ramos & Pascasio, 33710 Ramos & Edaño, F. B. 26530 Rola*. Chiefly in secondary forests, sea level to an altitude of 750 m. Endemic.

A genus represented by one other known species in the Solomon Islands.

Local names: Bagáas (P. Bis.) balúi (Buk.); balur (Buk.); nagsalalusáu (Bik.).

POTAMOGETONACEAE

1. POTAMOGETON Linnaeus

POTAMOGETON ANGUSTIFOLIUS Berchtold & Presl Rostlinar 2 (1821) 19; A. Benn. in Philip. Journ. Sci. 9 (1914) Bot. 340.

Luzon (Benguet), *Loher 1597*. In shallow water, altitude about 1,400 m. Europe, Asia, and North America.

POTAMOGETON MAACKIANUS A. Benn. in Journ. Bot. 42 (1904) 74, Philip. Journ. Sci. 9 (1914) Bot. 342.

Mindanao (Lanao), *Clemens 216*. In shallow water of Lake Lanao, altitude about 650 m. Northern Asia, Korea, China, Japan, and Formosa.

POTAMOGETON MALAINUS Miq. Ill. Fl. Archip. Ind. (1871) 46 (*malaina*); Aschers. & Graebn. in Engl. Pflanzenreich 31 (1907) 83; A. Benn. in Philip. Journ. Sci. 9 (1914) Bot. 341; Merr. Fl. Manila (1912) 67.

Potamogeton mucronatus Presl Epim. (1851) 245; Walp. Ann. 3 (1852-53) 504, non Schrad.

Potamogeton lucens Naves Novis. App. (1882) 297; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 282, non Linn.

Luzon (Rizal, Laguna, Sorsogon), Mindanao (Lanao), *Merrill 5107*, *Williams 2041*, *Elmer 18113*, *B. S. 22797* *McGregor*, *11892* *Robinson & Ramos*. In shallow water of lakes at low and medium altitudes. India to Korea and Japan southward to Borneo, Java, and Celebes. The Philippine form has been somewhat confused with *Potamogeton gaudichaudii* C. & S. which is a form of *P. lucens* Linn.

Local name: Damúng-kulút (Tag.).

POTAMOGETON PERVERSUS A. Benn. in Philip. Journ. Sci. 9 (1914) Bot. 343.

Luzon (Bontoc), *Vanoverbergh 249, 2684*. In shallow slow streams and in rice paddies, altitudes about 1,300 m. India to China, Manchuria, Korea, and Formosa.

Local name: íbas (Bon.).

POTAMOGETON PHILIPPINENSIS A. Benn. in Philip. Journ. Sci. 9 (1914) Bot. 342, hybr. *P. malainus* × *P. maackianus*.

Potamogeton nipponicus Ostenfeld in Philip. Journ. Sci. 9 (1914) Bot. 260, non Makino.

Mindanao (Lanao), *Clemens*. In shallow water of Lake Lanao, altitude about 650 m. Endemic.

POTAMOGETON PUSILLUS Linn. Sp. Pl. (1753) 127; A. Benn. in Philip. Journ. Sci. 9 (1914) Bot. 342.

Luzon (Benguet), *Elmer 5951*. In shallow water, altitude about 1,300 m. Widely distributed in Europe, Asia, Africa, and North America.

POTAMOGETON TEPPERI A. Benn. in Journ. Bot. 25 (1887) 178; Philip. Journ. Sci. 9 (1914) Bot. 340.

Luzon (Benguet), *Merrill Phil. Pl. 1722*, *Elmer 5952*, *Topping 45*, *B. S. 14110* *Robinson*. In shallow water, altitude about 1,300 m. Yunnan and Australia.

EXCLUDED SPECIES

POTAMOGETON CRISPUS Linn.; Naves Novis. App. (1882) 297.

POTAMOGETON NATANS Linn.; Naves l. c.

POTAMOGETON JAVANICUS Linn.; Naves l. c.

2. *RUPPIA* Linnaeus

RUPPIA MARITIMA Linn. Sp. Pl. (1753) 127; Naves Novis. App. (1882) 297; Aschers. & Graebn. in Engl. Pflanzenreich 31 (1907) 142; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 228, Fl. Manila (1912) 68.

Luzon (Manila Bay), Panay, *Merrill Phil. Pl. 799, Loher 1594, B. S. 20660 Serviñas*. In shallow brackish water. In salt or brackish water in most warm and tropical regions; the Philippine form is referable to the var. *rostrata* Agardh. (*R. maritima* Linn. subsp. *rostellata* Koch).

Local names: Buñgüt-buñgüt (P. Bis.); damong-palai (Tag.).

3. *CYMODOCEA* Koenig

CYMODOCEA ROTUNDATA (Ehrb. & Hempr.) Aschers. & Schweinf. in Sitzber. Ges. Naturf. Freunde Berlin (1870) 84; Graebner in Engl. Pflanzenreich 31 (1907) 147; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 2.

Phucagrostis rotundata Ehrb. & Hempr. ex Aschers. in Boiss. Fl. Orient. 5 (1884) 21.

Palawan, Cavilli, Mindanao, *Merrill 7180a, Phil. Pl. 1264, B. S. 15559 Fénix*. On sand about coral reefs in shallow water. In shallow seas from the Red Sea to Australia.

CYMODOCEA SERRULATA (R. Br.) Aschers. & Magnus in Sitzber. Ges. Naturf. Freunde Berlin (1870) 84; Aschers. & Graebn. in Engl. Pflanzenreich 31 (1907) 147.

Caulinia serrulata R. Br. Prodr. (1810) 339.

Mindanao, fide *Ascherson & Graebner*, l. c. In shallow seas from the Red Sea and western Africa to Australia.

Cymodocea aequorea Koenig = *C. nodosa* (Ucria) Aschers.! was erroneously credited to the Philippines by Naves Novis. App. (1882) 297.

4. *DIPLANTHERA* Thouars

DIPLANTHERA UNINERVIS (Forst.) Aschers. in Engl. & Prantl Nat. Pflanzenfam. 2¹ (1897) 37; Graebn. in Engl. Pflanzenreich 31 (1907) 152; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 3.

Zostera uninervis Forsk. Fl. Aeg.-Arab. (1775) 159.

Luzon (Manila Bay), Palawan, Cavilli, *Merrill 7179, 9223, Phil. Pl. 1263, Shaw 1148*. In sand about reefs. In shallow seas from the Red Sea and western Africa to Australia and Polynesia.

The family Aponogetonaceae has no known representatives in the Philippines; *Aponogeton monostachyus* Linn. f. and *Ouvirandra fenestralis* Poir., credited to the Archipelago by Naves, Novis. App. (1882) 298, 299, must be excluded.

NAJADACEAE

1. *NAJAS* Linnaeus

NAJAS GRAMINEA Delile Fl. Aegypt. (1812) 282, t. 50, f. 3; Naves Novis. App. (1882) 298; Rendle in Trans. Linn. Soc. Bot. 5 (1899) 424, 443, t. 32, f. 192-201, Engl. Pflanzenreich 7 (1901) 18, f. 5, Q V; Merr. Fl. Manila (1912) 68.

Northern Luzon to Mindanao. In shallow stagnant fresh water, at low and medium altitudes; often common. Northern Africa to Japan southward to New Caledonia and Australia.

Local names: Aragán (Ilk.); bangbángi (Bon.); lábui (Ig.).

NAJAS PALUSTRIS Blanco Fl. Filip. (1837) 660; Rendle in Trans. Linn. Soc. Bot. 5 (1899) 427, Engl. Pflanzenreich 7 (1901) 18; Merr. Sp. Blancoanae (1918) 57.

Najas lobata Blanco Fl. Filip. ed. 2 (1845) 459, ed. 3, 3 (1879) 65.

Najas foveolata A. Br. ex Magnus Beitr. (1870) 7; Rendle in Trans. Linn. Soc. Bot. 5 (1899) 416, t. 31, f. 139-144, Engl. Pflanzenreich 7 (1901) 15.

Najas tenuifolia Naves Novis. App. (1880) 297, non R. Br.

Luzon (Union, Rizal, Laguna, Tayabas), *Merrill 5109, Sp. Blancoanae 519, Elmer 14420, Whitford 837, B. S. 24079 Ramos*. In shallow stagnant fresh water at low altitudes. India to Japan southward to Timor.

Local names: Aragán-tamnái (Ilk.); rigmáu (Tag.).

DOUBTFUL AND EXCLUDED SPECIES

NAJAS FALCICULATA A. Br. in Journ. Bot. 2 (1864) 278, f. 4; Rendle in Trans. Linn. Soc. Bot. 5 (1899) 417, t. 32, f. 145-151, Engl. Pflanzenreich 7 (1901) 15.

Manila, *Mertens* fide A. Braun l. c. The type is an Indian specimen and I strongly suspect that the Philippine specimen referred here by A. Braun is *Najas palustris* Blanco.

NAJAS INDICA Cham.; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 425.

The record was based on a specimen from Laguna de Bay, collected by Meyen. The species is known only from India.

ALISMACEAE

1. SAGITTARIA Linnaeus

SAGITTARIA SAGITTAEFOLIA Linn. Sp. Pl. (1753) 993; Naves Novis. App. (1882) 298; Micheli in DC. Monog. Phan. 3 (1881) 66; Buchenau in Engl. Pflanzenreich 16 (1903) 46; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 421, Sp. Blancoanae (1918) 58.

Alisma sagittifolium Llanos Frag. Pl. Filip. (1851) 69; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 51, non Willd.

Luzon (Pangasinan, Nueva Ecija, Tayabas, Camarines, Albay, Sorsogon), Catanduanes, Samar, Leyte, Mindanao (Lanao), *Merrill Phil. Pl. 982, Elmer 14339, Wenzel 1228, 1521, Clemens 888, B. S. 30161 Ramos, 10703 McGregor*. In fresh-water swamps, rice paddies, etc., at low and medium altitudes. Europe through Asia to Japan, southward to Java.

Local names: Gauai-gáuai (S. L. Bis.); tikóg (Bik.).

EXCLUDED GENERA

LIMNOPHYTON OBSTUSIFOLIUM Miq.; Naves Novis. App. (1882) 298.

LOPHIOCARPUS GUYANENSIS Micheli; Naves Novis. App. (1882) 298.

HYDROCHARITACEAE

1. HALOPHILA Thouars

HALOPHILA BECCARII Aschers. in Nuov. Giorn. Bot. Ital. 3 (1871) 302; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 228, Fl. Manila (1912) 70.

Luzon (Rizal), *Merrill 8045*. In brackish fish ponds. Borneo.

HALOPHILA OVALIS (R. Br.) Hook. f. Fl. Tasm. 2 (1860) 45; Naves Novis. App. (1882) 298; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 2.

Caulinia ovalis R. Br. Prodr. (1810) 339.

Palawan, Bancoran, Biliran, Merrill 7155, Phil. Pl. 1223, B. S. 18755 McGregor. In sand on coral reefs. On sandy bottoms of shallow seas. India to Australia and Polynesia.

HALOPHILA OVATA Gaudich. Bot. Freycinet's Voy. (1826) 430, t. 40, f. 1; Ostenfeld in Philip. Journ. Sci. 4 (1909) Bot. 68; Merr. Fl. Manila (1912) 70.

Luzon (Manila Bay), Merrill 4112, Phil. Pl. 1098, B. S. 9899 Robinson. In shallow water on rather muddy bottom. Marianne Islands.

HALOPHILA SPINULOSA (R. Br.) Aschers. in Neumayer Anleit. Wissenschaft. Beob. (1875) 368, ed. 2 (1888) 396; Benth. Fl. Austral. 7 (1878) 183; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 2.

Caulinia spinulosa R. Br. Prodr. (1810) 339.

Luzon (Camarines), Mindanao, F. B. 12275 Curran. In shallow seas on reefs. In shallow seas, eastern and northern coasts of Australia; Java, and probably in other parts of Malaya.

2. HYDRILLA L. C. Richard

HYDRILLA VERTICILLATA (Roxb.) Royle Ill. Bot. Himal. (1839) t. 376; Presl Bemerk. (1842) 112; Casp. in Pringsh. Bot. Jahrb. 1 (1858) 494; Naves Novis. App. (1880) 214; Merr. & Rolfe in Philip. Journ. Sci. 3 (1908) Bot. 96; Merr. Fl. Manila (1912) 69, Sp. Blancoanae (1918) 58.

Serpicula verticillata Linn. f. Suppl. (1781) 416; Roxb. Pl. Coromandel 2 (1798) t. 164.

Udora verticillata Spreng. Syst. 1 (1825) 170; Llanos Frag. Pl. Filip. (1851) 101; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 78.

Northern Luzon to Mindanao, in most or all islands and provinces. In shallow fresh water at low and medium altitudes; often abundant. Central Europe to Asia southward to Australia.

Local names: Gínga (Ibn.); ináta (Tag.); lomotlomótan (Tag.); lúsai (Mag.).

3. VALLISNERIA Linnaeus

VALLISNERIA GIGANTEA Graebn. in Engl. Bot. Jahrb. 49 (1912) 68; Merr. Fl. Manila (1912) 69; Hallier f. Nov. Guinea 8 (1913) 915.

Vallisneria spiralis Blanco Fl. Filip. (1837) 781, ed. 2 (1845) 538, ed. 3, 3 (1879) 187; Naves Novis. App. (1880) 214; Vidal Phan. Cuming. Philip. (1885) 148, non Linn.

Luzon (Pangasinan, Bulacan, Nueva Ecija, Laguna, Rizal, Camarines). In lakes and slow streams at low altitudes. Enormously variable in size, apparently depending largely on the depth of the water in which it grows. New Guinea, New South Wales.

Local names: Baláiba (Ilk.); lanténg (Ibn.); mariu-báriu (Bik.); palá-ipa (Ilk.) sabotán-buáia (Tag.); sintas-sintásan (Tag., from Sp. cinta = ribbon).

4. BLYXA Thouars

BLYXA OCTANDRA (Roxb.) Planch. ex Thwaites Enum. Pl. Ceyl. (1859-64) 332; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 73.

Vallisneria octandra Roxb. Pl. Coromand. 2 (1798) 34, t. 165.

Blyxa roxburghii Rich. in Mém. Inst. France (1811) 77, t. 5; Naves Novis. App. (1880) 214.

Luzon (Bontoc), Palawan, *Merrill Phil. Pl. 1244, Vanoverbergh 664*. In shallow lakes, rice paddies, etc., from sea level to 1,250 m altitude; apparently very local. Tropical Asia to Malaya, Australia, and the Caroline Islands.

Local name: Pangpangdán (Bon.).

5. ENHALUS L. C. Richard

ENHALUS ACOROIDES (Linn. f.) L. C. Rich. ex Steud. Nomencl. ed. 2, 1 (1840) 554; Chatin Anat. Pl. Aquat. (1862) 15, t. 6; Merr. Interpret. Herb. Amb. (1917) 84, Sp. Blancoanae (1918) 59.

Stratiotes acoroides Linn. f. Suppl. (1781) 268.

Enhalus koenigii L. C. Rich. in Mém. Inst. France 2 (1811) 78; Naves Novis. App. (1880) 214.

Vallisneria sphaerocarpa Blanco Fl. Filip. (1837) 780, ed. 2 (1845) 538, ed. 3, 3 (1879) 186.

Northern Luzon to Palawan and Mindanao, *Merrill 8070, Sp. Blancoanae 383, B. S. 18723 McGregor, 19001, 15558 Fénix*. In shallow water of sheltered bays along the seashore; of general occurrence where favorable habitats are found. Ceylon through Malaya to tropical Australia.

Local names: Iaaí (Tag.); lamón (Tag., Bik.); mariu-báriu (Bik.); paláipat-baibái (Ilk.); bariau-bariau (Kuy.).

6. THALASSIA Banks

THALASSIA HEMPRICHII (Ehrenb.) Aschers. in Engl. & Prantl Nat. Pflanzenfam. 2¹ (1889) 254; Merr. Fl. Manila (1912) 71, Philip. Journ. Sci. 10 (1915) Bot. 2.

Schizotheca hemprichii Ehrenb. in App. Akad. Berl. 1 (1832) 429.

Luzon (Manila Bay), Alabat, Palawan, Cavilli. In tide pools, about coral reefs, etc., probably of general Philippine distribution. In shallow seas, sandy bottom, India to Polynesia.

7. BOOTTIA Wallich

BOOTTIA RENIFOLIA Merr. in Philip. Journ. Sci. 4 (1909) Bot. 247.

Luzon (Camarines), Mindanao (Cotabato), *F. B. 12274 Curran, B. S. 11593 Robinson*. In swamps and slow streams at low altitudes. Endemic.

Local names: Butid-na-ubal (Mag.); gabi-gábi (Bik.).

Boottia cordata Wall., reported from the Philippines by Naves, Novis. App. (1880) 213, was unquestionably admitted on the basis of an erroneous identification.

8. OTTELIA Persoon

OTTELIA ALISMOIDES (Linn.) Pers. Syn. 1 (1805) 400; Blanco Fl. Filip. (1837) 461, ed. 2 (1845) 321, ed. 3, 2 (1878) 230; Naves Novis. App. (1880) 213; Vidal Phan. Cuming. Philip. (1885) 148; Merr. Fl. Manila (1912) 70, Sp. Blancoanae (1918) 59.

Stratiotes alismoides Linn. Sp. Pl. (1753) 535.

Damasonium lancifolium Presl Rel. Haenk. 1 (1827) 89.

Ottelia ensiformis Blanco Fl. Filip. (1837) 460, ed. 2 (1845) 320, ed. 3, 2 (1878) 229; Naves Novis. App. (1880) 214.

Ottelia philippinensis Ostenfeld in Philip. Journ. Sci. 9 (1914) Bot. 259.

Northern Luzon to Mindanao, in most islands and provinces. In shallow fresh water of lakes, pools, and slow streams. India through Malaya to tropical Australia.

A species presenting great variation in its vegetative characters, which largely depends on the depth of the water in which it grows, whether stagnant or not, and whether the plants are crowded or scattered. Two additional synonyms are *Ottelia javanica* Miq. and *O. ovalifolia* L. C. Rich., credited to the Philippines by Naves, Novis. App. (1880) 214, as distinct species.

Local names: Damong-ilálím (Tag.); espáda (Sp.); kalabúa (Tag.); lantíng (Bik.); lantén-sápa (Tag.); maraduyúk (Ib.); maradúyug (It.); tarabáng (Ilk.).

EXCLUDED GENERA

HYDROCHARIS MORSUS-RANAE Linn.; Naves Novis. App. (1880) 213.

NECHAMANDRA ROXBURHII Planch.; Naves Novis. App. (1880) 215 = *Lagarosiphon roxburghii* Planch.

TRIURIDACEAE

1. SCIAPHILA Blume

SCIAPHILA CLEMENSAE Hemsl. in Hook. Ic. IV 9 (1907) t. 2850, f. 7-14; Fedde Repert. 5 (1908) 265.

Mindanao (Lanao), *Clemens* 775. In damp primary forests at low altitudes. Borneo.

This species seems to be closely allied to *S. tenella* Blume. Dr. J. J. Smith states that *B. S. 6347 Robinson*, *F. B. 9350, 9449 Whitford & Hutchinson*, and *Clemens 775 p. p.* are very similar to Blume's species.

SCIAPHILA MACULATA Miers in Trans. Linn. Soc. Bot. 21 (1852) 48, t. 6, f. 19-36; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 282.

Luzon (Bataan), Mindanao (Butuan, Zamboanga), *F. B. 5537 Curran, 9126 Whitford & Hutchinson, Weber 1462*. In damp primary forests at low and medium altitudes. Endemic.

In addition to these two species what may be *Sciaphila tenella* Blume occurs in our collections, as noted above, as well as fragmentary material representing two other distinct species.

GRAMINEAE

1. ZEA Linnaeus

ZEA MAYS Linn. Sp. Pl. (1753) 971; Blanco Fl. Filip. (1837) 686, ed. 2 (1845) 476, ed. 3, 3 (1879) 90, t. 279; F.-Vill. Novis. App. (1882) 314; Usteri Beitr. Ken. Philip. Veg. (1905) 133; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 320, Fl. Manila (1912) 74, Sp. Blancoanae (1918) 59.

Extensively cultivated in most parts of the Philippines, in some islands and provinces a staple article of food. Introduced from America.

Local names: Gahilang (If.); igi (Bon.); maís (Sp.); maít (It.); mañgi (Ibn.); tigi (Bon.).

Teosinte, *Euchlaena luxurians* Schrad., a native of Mexico, was introduced into the Philippines in the late years of the Spanish occupation and again in the early years of the American occupation and was cultivated experimentally for forage purposes. It has not persisted in cultivation.

2. POLYTOCA R. Brown

POLYTOCA HETEROCLITA (Roxb.) Koord. in Eksksfl. Java 1 (1911) 99;

Merr. in Philip. Journ. Sci. 10 (1915) Bot. 288.

Coix heteroclita Roxb. Fl. Ind. ed. 2, 3 (1832) 572.

Polytoxa bracteata R. Br. in Benn. Pl. Jav. Rar. (1838) 20, t. 5.

Busuanga, Mindanao (Bukidnon), *Baker 3616*, *B. S. 39226*, *38484*, *41267* *Ramos & Edaño*. In open grasslands at medium altitudes. India to southern China and Java.

3. CHIONACHNE R. Brown

CHIONACHNE BIAURITA Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 263; Merr. op. cit. 320.

Luzon (Bontoc, Benguet), *Merrill 4232*, *Vanoverbergh 934*. On fresh talus slopes, untilled soil, etc., altitude about 1,200 m. Endemic.

EXCLUDED SPECIES

CHIONACHNE BARBATA R. Br.; F.-Vill. Novis. App. (1882) 314.

4. COIX Linnaeus

COIX LACHRYMA-JOBI Linn. Sp. Pl. (1753) 972; Pilg. ex Perk. Frag. Fl. Philip. (1904) 137; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 320, 3 (1908) Bot. 394, 5 (1910) Bot. 325, Fl. Manila (1912) 75, Sp. Blancoanae (1918) 59; Usteri Beitr. Ken. Veg. Philip. (1905) 132 (*lachrymae-jovis*).

Coix lachryma Linn. Syst. ed. 10 (1759) 1261; Blanco Fl. Filip. (1837) 688, ed. 2 (1845) 478, ed. 3, 3 (1879) 92, t. 188; Miq. Fl. Ind. Bat. 3 (1857) 476; F.-Vill. Novis. App. (1882) 314; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288.

Coix agrestis Lour. Fl. Cochinch. (1790) 551; F.-Vill. Novis. App. (1882) 314.

Coix exaltata Jacq. ex Spreng. Syst. 1 (1825) 228; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Throughout the Philippines in the settled areas at low and medium altitudes, in most or all islands and provinces; common. A native of the Old World, now pantropic.

Local names: Abúkai (Iv.); adlái (Bis.); agágai (Iv.); agdá (Ig.); agláí (C. Bis.); alimúdias (P. Bis.); apági (Ig.); atákai (Ilk., Bon.); balantákan (Pamp.); barubaióko (Bik.); bintíkai (Bik.); bitógan (Bag.); dalái (Sub.); damáu (C. Bis.); glias (Sub.); kalabúgau (Buk.); kambot (Ting.); katáyan (Ig.); katigbí (C. Bis.); kibaoang (If.); koldásan (Bik.); kudlásan (Tag.); lamúdias (P. Bis.); liás (Sub.); paiás (P. Bis., Bag.); paliás (P. Bis.); pintáka (C. Bis.); tidbí (S. L. Bis.); tigbí (Bik., Tag.); tighíkai (Bik.); tiguas (Sul.); tikaian (Bon.).

Var. **MA-YUEN** (Roman.) Stapf in Hook. f. Fl. Brit. Ind. 7 (1897) 100; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 321.

Coix ma-yuen Roman. in Bull. Soc. Acclim. Paris III 8 (1881) 442.

Luzon (Rizal), Mindanao (Zamboanga), Jolo, *Merrill 5326*, *F. B. 3394 Ahern's collector*. Occasionally cultivated for its edible grain or for use in making fermented drinks. Tropical Asia.

Local name: Ilás (Tag.).

5. DIMERIA R. Brown

DIMERIA CILIATA Merr. in Philip. Journ. Sci. 9 (1914) Bot. 262.

Palawan, *Merrill 9320*. In low open grasslands just above sea level. Endemic.

DIMERIA ORNITHOPODA Trin. Fund. Agrost. (1820) 167, *t. 14*, var.

TENERA (Trin.) Hack. in DC. Monog. Phan. 6 (1889) 81; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 321, Fl. Manila (1912) 75.

Dimeria tenera Trin. in Mém. Acad. Sci. St. Pétersb. VI 2 (1833) 335; F.-Vill. Novis. App. (1882) 315.

Luzon (Bontoc, Benguet, Pangasinan, Zambales, Bulacan, Rizal, Camarines), Mindoro, Mindanao (Davao). In open grasslands, on rice-paddy banks, etc., sea level to an altitude of 1,500 m, locally abundant. India to Japan and Malaya.

Haplachne pilosissima Presl and *Andropogon chloridiformis* Gaudich. are included in the Novissima Appendix by F.-Villar as distinct species. Both are synonyms of *Dimeria chloridiformis* K. Schum. & Lauterb. (*D. pilosissima* Triñ.) a very characteristic species known only from the Marianne Islands.

6. IMPERATA Cyrillo

IMPERATA CYLINDRICA (Linn.) Beauv. Agrost. (1912) expl. *pl. 5*, *t. 5*, *f. 1*, var. **KOENIGII** (Retz.) Benth. ex Pilger in Perk. Frag. Fl. Philip. (1904) 137; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 322, 3 (1908) Bot. 395, 5 (1910) Bot. 325, Fl. Manila (1912) 76, Sp. Blancoanae (1918) 60.

Saccharum koenigii Retz. Obs. 5 (1789) 16.

Imperata koenigii Beauv. Agrost. (1812) 165; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 100.

Imperata arundinacea Cyr. Pl. Rar. Ic. 2 (1792) 26, *t. 11*; Miq. Fl. Ind. Bat. 3 (1857) 514; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 289; F.-Vill. Novis. App. (1882) 316.

Imperata arundinacea Cyr. var. *koenigii* Benth.; Hack. in DC. Monog. Phan. 6 (1889) 94; Usteri Beitr. Ken. Phil. Veg. (1905) 132.

Saccharum spicatum Presl Rel. Haenk. 1 (1828) 346; Kunth Enum. 1 (1833) 476; Miq. Fl. Ind. Bat. 3 (1857) 513; F.-Vill. Novis. App. (1882) 317; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 52, *t. 1*.

Throughout the Philippines, on open slopes, often gregariously occupying enormous areas, forming characteristic *kogonales*, the name of these areas being taken from its almost universal local name *kogon*. It ascends to 2,300 m altitude. Tropical Asia and Africa to Australia and Polynesia.

Local names: Buchid (Iv.); bulum (If.); gáon (Ig.); gógon (Bik.); góon (Bon.); ilib (Pamp.); kógon (Ilk., Tag., Bik.); panáu (Ilk.); parrang (Sul.).

IMPERATA EXALTATA Brongn. Voy. Coq. Bot. (1829) 101, excl. syn. Roxb.; Hack. in DC. Monog. Phan. 6 (1889) 98; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 322; Pilg. in Perk. Frag. Fl. Philip. (1904) 137.

Saccharum confertum Presl Rel. Haenk. 1 (1828) 346; Kunth Enum. 1 (1833) 476; F.-Vill. Novis. App. (1882) 317.

Imperata ramosa Anders. in Oefvers. Vet. Akad. Forhandl. Stockh. (1855) 158, Rolfe in Journ. Bot. 23 (1885) 216; Vid. Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 289.

Saccharum alopecurus Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 100; Walp. Ann. 3 (1853) 792; Miq. Fl. Ind. Bat. 3 (1857) 513; F.-Vill. Novis. App. (1882) 317.

Saccharum negrosense Steud. Syn. Pl. Glum. 1 (1854) 407.

Central Luzon to Palawan and Mindanao. In habitats similar to the preceding species, but much less common. Java to the Moluccas; a variety in tropical America as Hackel interprets it.

Subsp. **MERRILLII** Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 264; Merr. op. cit. 322.

Luzon (Ifugao, Benguet), Negros, *Merrill 4813*, *Phil. Pl. 582*, *B. S. 2768* *Mearns*, *B. S. 19627* *McGregor*. On exposed ridges, altitude 1,500 to 2,250 m. Endemic.

Local names: Gógon (Bik.); kógon (Tag., Bis., Sul.); padang-lubuk (Sul.); paraugutau (Sub.).

7. **MISCANTHUS** Andersson

MISCANTHUS DEPAUPERATUS Merr. in Philip. Journ. Sci. 5 (1910) Bot. 170.

Negros, *Merrill Phil. Pl. 583*. In open places on the crater of Canlaon Volcano, altitude 1,800 to 2,000 m. Endemic.

MISCANTHUS JAPONICUS (Thunb.) Anders. in Oefv. Vet. Akad. Forhandl. Stockh. (1855) 166; Hack. in DC. Monog. Phan. 6 (1889) 107; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 322, 3 (1908) Bot. 395.

Saccharum japonicum Thunb. in Trans. Linn. Soc. 2 (1794) 328.

Saccharum praegrande Steud. Syn. Pl. Glum. 1 (1854) 408.

Miscanthus luzoniensis Anders. in Oefv. Vet. Akad. Forhandl. Stockh. (1855) 166; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 289.

Saccharum densum Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 100; Walp. Ann. 3 (1853) 792.

Babuyan Islands to Mindanao, in most islands and provinces. In thickets, on slopes, and along streams at low and medium altitudes, ascending to 1,200 m. Japan to the Moluccas and Polynesia.

Local names: Búyau (If.); Ioni (Bag.); tanlák (Tag.); tubás (Tag.); rúnò (Pang., Ig., Ilk.); víau (Iv.).

MISCANTHUS SINENSIS Anders. in Oefv. Vet. Akad. Forhandl. Stockh. (1855) 166; Hack. in DC. Monog. Phan. 6 (1889) 105; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 323, 2 (1907) Bot. 260, 3 (1908) Bot. 395, 5 (1910) Bot. 325.

Eulalia japonica Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 333; Miq. Fl. Ind. Bat. 3 (1857) 518; F.-Vill. Novis. App. (1882) 316, non *Miscanthus japonicus* Anders.

Miscanthus japonicus Pilg. in Perk. Frag. Fl. Philip. (1904) 137, non Anders.

Batan Islands, Luzon (Bontoc, Benguet, Lepanto, Pampanga, Tayabas, Bataan, Zambales), Mindoro, Negros. In open places, on slopes, and in ravines, altitude 800 to 2,700 m, often abundant. Japan to southern China and Celebes.

Local names: Bigáho (Bik.); bígau (Bik.); bídu (Ig.); bílau (Ig., If.); gáho (Bik.); gísa (Bik.); lólo (Bon.); padéd (Ig.); pao (Bon.); rúno (Ilk.); víau (Iv.).

8. SACCHARUM Linnaeus

SACCHARUM ARUNDINACEUM Retz. Obs. 6 (1791) 14; Hack. in DC.

Monog. Phan. 6 (1889) 117; Hook. f. Fl. Brit. Ind. 7 (1897) 119.

Erianthus sp. Pilger in Perk. Frag. Fl. Philip. (1904) 137.

Luzon (Bontoc, Nueva Vizcaya, Tayabas), Mindoro, Samar, Mindanao, Merrill 1794, 5448, Clemens 809, Williams 2625, B. S. 13279 Ramos, 14207 McGregor, 20906 Escritor. Along river banks at low and medium altitudes. India to southern China.

Local names: Bagi'-unás (Ilk.); bahí-onás (Isn.); gathó (Tag.); lakbó (Tag.); tanig (Bon.); zaigbau (Sub.).

SACCHARUM OFFICINARUM Linn. Sp. Pl. (1753) 54; Blanco Fl. Filip.

(1837) 42; ed. 2 (1845) 29, ed. 3, 1 (1877) 55, t. 18; F.-Vill. Novis.

App. (1882) 317; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 323,

Fl. Manila (1912) 78, Sp. Blancoanae (1918) 60.

Saccharum violaceum F.-Vill. Novis. App. (1882) 317, non Tussac.

Cultivated throughout the Philippines, in some islands and provinces very extensively, being one of the major crops of the Philippines. Not a native of the Archipelago. Sugar cane.

Local names: Agbó (Ibn.); caña dulce (Sp.); tubó (Tag., Bik.); tubu (Sul.); uná (Ibn.); unás (Ilk.); unát (It.).

SACCHARUM SPONTANEUM Linn. Mant. 2 (1771) 183 subsp. **INDICUM**

Hack. in DC. Monog. Phan. 6 (1889) 117; Pilger in Perk. Frag. Fl.

Philip. (1904) 137; Merr. in Philip. Journ. Sci. 1 (1906) Suppl.

323, 5 (1910) Bot. 325, Fl. Manila (1912) 77, Sp. Blancoanae (1918) 60.

Saccharum spontaneum Linn.; Miq. Fl. Ind. Bat. 3 (1857) 512; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 289.

Saccharum spontaneum Linn. subsp. *luzonicum* Hack. in DC. Monog. Phan. 6 (1889) 117.

Anthistiria gigantea Blanco Fl. Filip. (1837) 49, ed. 2 (1845) 33, ed. 3, 1 (1877) 62, non Cav.

Saccharum insulare Brongn. var. *amplum* et var. *depauperatum* Anders. in Oefv. Vet. Akad. Forhandl. Stockh. 12 (1855) 157.

Throughout the Philippines, in open areas at low and medium altitudes, ascending to 1,500 m, often gregarious and almost exclusively occupying large areas. India to southern China through Malaya to Polynesia.

Local names: Bogáng (C. Bis.); bugáng (C. Bis.); liddá (Ilk.); sálin (Bon.); síkal (Ibn.); taláhib (Tag.); tibáyo (Bag.); tigbáu (S. L. Bis., P. Bis.).

9. POLLINIA Trinius

POLLINIA ARGENTEA (Brongn.) Trin. in Mém. Acad. St. Pétersb. VI 4 (1836) 90; Hack. in DC. Monog. Phan. 6 (1889) 162; Pilger in Perk. Frag. Fl. Philip. (1904) 138; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 265; Merr. op. cit. 325, Fl. Manila (1912) 77.
Eulalia argentea Brongn. Voy. Coqu. Bot. (1829) 92.

Luzon (Cagayan, Isabela, Bontoc, Nueva Vizcaya, Pangasinan, Rizal), Semirara, Merrill 4152, 108, Elmer 6474, B. S. 17350 Otanes, 14216 McGregor, 7815, 8092, 7831 Ramos. In open grasslands at low and medium altitudes. India to Malaya and Australia.

Var. **LAGOPUS** Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 79; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 325.

Pollinia speciosa Pilger in Perk. Frag. Fl. Philip. (1904) 138, non Hack.

Luzon (Ilocos Norte, Lepanto, Bontoc, Benguet, Pampanga), Mindoro, Palawan, Merrill 4222, 3902, 4365, 4535, 472, B. S. 32753, 32779, 5849 Ramos. In open grasslands at low and medium altitudes, ascending to 1,600 m. Endemic.

POLLINIA CONTORTA (Brongn.) comb. nov.

Pogonatherum contortum Brongh. Bot. Voy. Coqu. Bot. (1829) 90, t. 17.

Andropogon koretrostachys Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 273; Steud. Syn. Pl. Glum. 1 (1854) 380; Miq. Fl. Ind. Bat. 3 (1857) 484; F.-Vill. Novis. App. (1882) 316.

Pollinia articulata Trin. in Mém. Acad. St. Pétersb. VI 4 (1836) 90; F.-Vill. Novis. App. (1882) 315; Hack. in DC. Monog. Phan. 6 (1889) 153 (subsp. *fragilis* Hack. var. *setifolia* Hack.); Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 324; Pilger in Perk. Frag. Fl. Philip. (1904) 138.

Pollinia setifolia Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 88; Miq. Fl. Ind. Bat. 3 (1857) 521; F.-Vill. Novis. App. (1882) 315; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 290.

Andropogon asthenostachys Steud. Syn. Pl. Glum. 1 (1854) 381; Miq. Fl. Ind. Bat. 3 (1857) 485.

Eulalia contorta O. Kuntze Rev. Gen. Pl. (1891) 775.

Luzon (Cagayan, Isabela, Abra, Lepanto, Nueva Ecija, Nueva Vizcaya), Mindanao (Bukidnon), Merrill 317, B. S. 20141 McGregor, 7035, 8121, 7809, 4737 Ramos, F. B. 5875 Curran, 22128 Alvarez, 16460 Bacani. In open grasslands chiefly at medium altitudes. Burma, southern China, Moluccas, Australia.

Local name: Sañgumati (Buk.).

POLLINIA FULVA (R. Br.) Benth. Fl. Austral. 7 (1878) 526.

Saccharum fulvum R. Br. Prodr. (1810) 203.

Pollinia cumingii Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850)

98; Steud. Syn. Pl. Glum. 1 (1854) 373; Walp. Ann. 3 (1853) 803; Miq. Fl. Ind. Bat. 3 (1857) 522; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 290; Hack. in DC. Monog. Phan. 6 (1889) 167; Pilger in Perk. Frag. Fl. Philip. (1904) 138; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 325.

Andropogon aureo-fulvus Steud. Syn. Pl. Glum. 1 (1854) 373.

Pollinia aurea F.-Vill. Novis. App. (1882) 315, non Benth.

Eulalia fulva O. Kuntze Rev. Gen. Pl. (1891) 775.

Luzon (Ilocos Norte, Amburayan, Lepanto, Benguet, Pangasinan, Nueva Vizcaya), Mindoro, Cuming 1538, Merrill 260, 4293, B. S. 33254, 4827 Ramos. In open grasslands, sandy river bars, etc., at low and medium altitudes, ascending to 1,200 m. Bengal, southern China, Timor, Java, Australia.

POLLINIA GLABRATA (Brongn.) Trin. in Mém. Acad. St. Pétersb. VI 4 (1836) 90; Hack. in DC. Monog. Phan. 6 (1889) 179.

Eulalia glabrata Brongn. Voy. Coqu. Bot. (1829) 93, t. 19.

Andropogon glabratus Steud. Nomencl. ed. 2, 1 (1840) 91.

Nemastachys taitensis Steud. Syn. Pl. Glum. 1 (1854) 357.

Pollinia monantha Nees, var. *leptathera* Hack. in Philip. Journ. Sci. 3 (1908) Bot. 167.

Luzon (Pangasinan, Rizal, Bataán, Laguna, Tayabas), Mindoro, Samar, Panay, Mindanao, Balut, Merrill 5212, 8272, 7323, 5627, 5426, Phil. Pl. 151, B. S. 31411, 29010 Ramos & Edaño, 21993, 13784 Ramos. In open ravines, thickets, and old clearings at low and medium altitudes. New Caledonia to the Marianne Islands and Polynesia.

Local names: Gulos (Sub.); kapong (Bon.); lusimat (Sub.).

Var. **ELMERI** (Hack.) comb. nov.

Pollinia monantha Nees var. *elmeri* Hack. in Philip. Journ. Sci. 3 (1908) Bot. 167.

Luzon (Bontoc, Benguet), Elmer 6524, Merrill 7784, Vanoverbergh 3665. In thickets in damp ravines, altitude 1,200 to 1,500 m. Endemic.

POLLINIA IRRITANS (R. Br.) Hack. in DC. Monog. Phan. 6 (1889) 155, Philip. Journ. Sci. 1 (1906) Suppl. 265; Merr. op. cit. 325.

Saccharum irritans R. Br. Prodr. (1810) 203.

Erianthus irritans Kunth Rev. Gram. 1 (1829) 160.

Eulalia irritans O. Kuntze Rev. Gen. Pl. (1891) 775.

Luzon (Bontoc, Benguet), Merrill 4388, Vanoverbergh 3838. In open grasslands, altitude about 1,000 m. Tropical Australia.

POLLINIA MARITIMA Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 326.

Lumbucan, Merrill 5277, Phil. Pl. 112. On sandy seashores. Closely allied to *P. fulva* Benth. Endemic.

POLLINIA NUDA Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 307; Hack. in DC. Monog. Phan. 6 (1889) 178; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 265.

Eulalia nuda O. Kuntze Rev. Gen. Pl. (1891) 775.

Luzon (Benguet), Merrill 4727, 4842. In openings in the mossy forest, altitude 2,000 to 2,400 m. India to China and Japan.

POLLINIA QUADRINERVIS Hack. in DC. Monog. Phan. 6 (1889) 158, Govt. Lab. Publ. (Philip.) 35 (1906) 79; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 325, 5 (1910) Bot. 325.

Eulalia quadrinervis O. Kuntze Rev. Gen. Pl. (1891) 775.

Luzon (Bontoc, Benguet), Merrill 4354, 4560, 6519, Elmer 5783, B. S. 31980 Santos, F. B. 16183, 16208 Curran, Merritt, & Zschokke. On open slopes in the pine region, altitude 1,400 to 2,200 m. India to southern China.

Local names: Galakgákan (Ig.); taaó (Ig.).

POLLINIA TENUIS Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 307; F.-Vill. Novis App. (1882) 315; Miq. Fl. Ind. Bat. 3 (1857) 521; Hack. in DC. Monog. Phan. 6 (1889) 181; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 326.

Luzon (Ilocos Sur, Bontoc, Benguet, Bataan, Rizal), Merrill 4310, 4382, Kneucker Gram. Exsic. 721, B. S. 5310 Ramos. On dry open slopes, on and about cliffs, etc., from sea level to 1,200 m altitude. Caroline Islands.

POLLINIA VIMINEA (Trin.) comb. nov.

Andropogon vimineus Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 268.

Pollinia imberbis Nees in Steud. Syn. Pl. Glum. 1 (1854) 410, var. *willdenowiana* Hack. in DC. Monog. Phan. 6 (1889) 178, forma *monostachya* Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 265; Merr. op. cit. 326.

Luzon (Bontoc, Benguet, Rizal), Merrill 4707, Williams 1960, Vanoverbergh 309, 991, 3810. On open wet slopes and in the mossy forest, altitude 1,200 to 2,200 m. India to southern China and Japan.

10. POLYTRIAS Hackel

POLYTRIAS AMAUREA (Büse) O. Kuntze Rev. Gen. Pl. (1891) 788; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 327.

Andropogon amaurus Büse in Miq. Pl. Jungh. (February, 1854) 320.

Andropogon diversiflorus Steud. in Zoll. Syst. Verzeich. (1854) 58, *nomen nudum*, Syn. Pl. Glum. 1 (July or August, 1854) 370.

Pollinia praemorsa Nees in Steud. Syn. Pl. Glum. 1 (1854) 409.

Polytrias praemorsa Hack. in DC. Monog. Phan. 6 (1889) 189, t. 1, f. 13; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 7; Pilger in Perk. Frag. Fl. Philip. (1904) 138.

Polytrias diversiflora Nash in Torreya 5 (1905) 110; Merr. in Philip. Journ. Sci. 11 (1916) Bot. 253.

Luzon (Rizal, Sorsogon), Jolo, Merrill 386, Phil. Pl. 106, Kneucker Gram. Exsic. 602. In open grasslands in and about towns at low altitudes, certainly introduced. Malay Peninsula, Borneo, Java, introduced in the West Indies.

11. POGONATHERUM Beauvois

POGONATHERUM PANICEUM (Lam.) Hack. in Allg. Bot. Zeitschr. 12 (1906) 178; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 395, Fl. Manila (1912) 76, Sp. Blancoanae (1918) 60.

Saccharum paniceum Lam. Encycl. 1 (1785) 595, t. 40, f. 1.

Pogonatherum saccharoideum Beauv. Agrost. (1812) 9, t. 11, f. 7;

- F.-Vill. Novis. App. (1882) 316; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 290; Hack. in DC. Monog. Phan. 6 (1889) 192 (var. *monandrum* Hack.); Pilger in Perk. Frag. Fl. Philip. (1904) 138; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 327.
- Pogonatherum polystachyum* R. & S. Syst. 2 (1825) 497; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.
- Pogonopsis tenera* Presl Rel. Haenk. 1 (1830) 333, t. 46; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 52, t. 2.
- Pogonatherum crinitum* Kunth Enum. 1 (1833) 478; Miq. Fl. Ind. Bat. 3 (1857) 516; Usteri Beitr. Ken. Veg. Philip. (1905) 133.
- Cinna filiformis* Llanos Frag. Pl. Filip. (1851) 9; F.-Vill. & Naves in Blanco Fl. Filip. 3, 4¹ (1880) 3.
- Deyeuxia quadriseta* F.-Vill. Novis. App. (1882) 319, non Benth.
- Ischaemum crinitum* Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 298; Presl Repert. 1 (1833) 296.

Throughout the Philippines, common on banks of ravines, along streams on cliffs and steep banks, etc., at low and medium altitudes, ascending to 2,000 m. India to China through Malaya to the New Hebrides.

Local names: Dumenéi (Bon.); kisáng (Buk.); lagitlit (C. Bis.); mala-kauáyan (S. L. Bis.); obán-obán (Mbo.); panaguáyan (C. Bis.); silsilig (Bon.).

12. APLUDA Linnaeus

- APLUDA MUTICA** Linn. Sp. Pl. (1753) 82; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 192, Hook. Journ. Bot. Kew Miscel. 2 (1850) 99; Miq. Fl. Ind. Bat. 3 (1857) 501; F.-Vill. Novis. App. (1882) 318; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 332, Fl. Manila (1912) 79.
- Calamina humilis* Presl Rel. Haenk. 1 (1828) 344.
- Apluda humilis* Kunth Enum. 1 (1833) 517; F.-Vill. Novis. App. (1882) 318.
- Apluda cumingii* Büse in De Vriese Pl. Ind. Bat. Or. (1856) 105; Miq. Fl. Ind. Bat. 3 (1857) 501; F.-Vill. Novis. App. (1882) 318.
- Apluda varia* Hack. var. *humilis* Hack. in DC. Monog. Phan. 6 (1889) 196.
- Apluda varia* Hack. var. *mutica* Hack. f. *intermedia* Hack. in DC. Monog. Phan. 6 (1889) 197.
- Apluda mutica* Linn. forma *intermedia* Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 332.
- Apluda varia* Hack. var. *major* Hack. in DC. Monog. Phan. 6 (1889) 198.
- Apluda mutica* Linn. var. *major* Hack. in Allgem. Bot. Zeitschr. 12 (1906) 179.

In thickets and open grasslands throughout the Philippines at low and medium altitudes, ascending to 1,800 m. India to southern China, through Malaya to Australia, New Caledonia, and Polynesia.

Local names: Kauakauáyan (Tag.); kolokauáyan (Tag.); kurukauáyan (Bik); magkauáyan (C. Bis.).

- Var ARISTATA** (Linn.) Rendle ex Forbes & Hemsl. in Journ. Linn. Soc. Bot. 36 (1904) 279; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 332, Fl. Manila (1912) 80.

Apluda aristata Linn. Amoen. Acad. 4 (1759) 303; F.-Vill. Novis. App. (1882) 318; Usteri Beitr. Ken. Philip. Veg. (1905) 132.

Luzon (Rizal), *B. S.* 2041 *Ramos*, Merrill 629, *Silva* 391. Habitat and range of the species.

13. ISCHAEMUM Linnaeus

ISCHAEMUM ANGUSTIFOLIUM (Trin.) Hack. ex Oliv. in Hook. Ic. III 8 (1888) t. 1773, DC. Monog. Phan. 6 (1889) 241; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 331.

Spodiopogon angustifolius Trin. in Mém. Acad. St. Petersb. VI 2 (1833) 300; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 290.

Andropogon obvallatus Steud. Syn. Pl. Glum. 1 (1854) 373.

Ischaemum notopogon Nees in Miq. Fl. Ind. Bat. 3 (1857) 500; F.-Vill. Novis. App. (1882) 315.

Luzon (Cagayan, Ilocos Norte, Bontoc, Nueva Vizcaya, Pangasinan), Cuming 1002, Merrill 172, *Phil. Pl.* 171, *B. S.* 7748, 7919 *Ramos*, 14214 *McGregor*, *F. B.* 16527 *Curran & Merritt*. In open grasslands at low and medium altitudes, ascending to 1,400 m. India to southern China and Formosa.

Local names: Kabóot (Ilk., Ibn.); poen (Bon.); pueng (Bon.); puenig (Ig.); sangumáti (Buk.).

ISCHAEMUM ARISTATUM Linn. Sp. Pl. (1753) 1049 var. **GIBBUM** (Trin.) Hack. in DC. Monog. Phan. 6 (1889) 204; Pilg. in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 330, Fl. Manila (1912) 80.

Ischaemum gibbum Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 295; Miq. Fl. Ind. Bat. 3 (1857) 498; F.-Vill. Novis. App. (1882) 315.

Andropogon gibbus Steud. Syn. Pl. Glum. 1 (1854) 376.

Luzon (Isabela, Ilocos Norte, Nueva Ecija, Nueva Vizcaya, Zambales, Bulacan, Rizal, Camarines), Culion. In open grasslands at low and medium altitudes; often common. Well represented by *Merrill Phil. Pl.* 179, 1977, and *Kneucker Gram. Exsic.* 605. The variety endemic, the species India to China.

ISCHAEMUM ARUNDINACEUM F.-Muell. ex Benth. Fl. Austral. 7 (1878) 519 var. **RADICANS** Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 266; Merr. op. cit. 331.

Luzon (Benguet, Bontoc), *Merrill* 4264, *Phil. Pl.* 1000, *Williams* 1197, *Vanoverbergh* 861. On open grassy slopes in the pine regions, altitude 1,200 to 1,600 m. The variety endemic, the species in tropical Australia.

ISCHAEMUM CILIARE Retz. Obs. 6 (1791) 36; F.-Vill. Novis. App. (1882) 315; Hack. in DC. Monog. Phan. 6 (1889) 255 (var. *malachophyllum* Hack.); Merr. in Philip. Journ. Sci. 3 (1908) Bot. 395.

Batan Islands (Batan), *B. S.* 3169 *Mearns*. In open grasslands. India to China and Formosa.

ISCHAEMUM GLAUDESCENS Merr. in Philip. Journ. Sci. 9 (1914) Bot. 263.

Palawan, *Merrill* 9453. In shallow water, borders of Lake Manguao, at low altitudes. Endemic.

ISCHAEMUM INTERMEDIUM Brongn. in Duperr. Voy. Coqu. (1829) Bot. 73; Pilg. in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 331, Fl. Manila (1912) 80.

Luzon (Rizal, Laguna, Sorsogon), Polillo, Leyte, Mindanao (Surigao), Merrill 68, 5222, 5213, Phil. Pl. 102, 1453, Elmer 14326, 18318, 17408, 16936, B. S. 34749 Ramos & Pascasio, 12103, 12022 Ramos. In open wet grassy places at low altitudes. Caroline Islands.

ISCHAEMUM MERRILLII Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 266; Merr. op. cit. 331.

Luzon (Lepanto, Benguet), Merrill 4622, B. S. 12917 *Fénix*. In open rather wet places, altitude about 1,500 m. Endemic.

Local names: Kugañgán (Ig.); sinamon (Ig.).

ISCHAEMUM MUTICUM Linn. Sp. Pl. (1753) 1049; Miq. Fl. Ind. Bat. 3 (1857) 496; F.-Vill. Novis. App. (1882) 315; Hack. in DC. Monog. Phan. 6 (1889) 212; Pilg. in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 330, 3 (1908) Bot. 395. *Ischaemum glabratum* Presl Rel. Haenk. 1 (1828) 328; Kunth Enum. 1 (1833) 513; Miq. Fl. Ind. Bat. 3 (1857) 498; F.-Vill. Novis. App. (1882) 315.

Andropogon relictus Steud. Syn. Pl. Glum. 1 (1854) 377.

Ischaemum repens Roxb. Fl. Ind. 1 (1820) 325; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 100.

Along sandy seashores from northern Luzon to Palawan and Mindanao, in most or all islands. India through Malaya to tropical Australia, New Caledonia, and western Polynesia.

Local name: Karukauáyan (Bik.).

ISCHAEMUM PUBESCENS Merr. in Philip. Journ. Sci. 9 (1914) Bot. 264.

Dumaran, B. S. 21639 *Escritor*. In open grasslands at low altitudes. Endemic.

ISCHAEMUM RUGOSUM Salisb. Ic. (1791) 1, t. 1 var. **DISTACHYUM** (Cav.) Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 330, 2 (1907) Bot. 436, Fl. Manila (1912) 80, Sp. Blancoanae (1918) 61.

Colladoa distachya Cav. Ic. 5 (1799) 37, t. 460; Willd. Sp. Pl. 4² (1805) 496.

Andropogon ramosus Blanco Fl. Filip. (1837) 37, ed. 2 (1845) 25, ed. 3, 1 (1877) 48; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 91, non Forsk.

Ischaemum colladoa Spreng. Syst. 1 (1825) 298; Kunth Enum. 1 (1833) 513; Miq. Fl. Ind. Bat. 3 (1857) 498; F.-Vill. Novis. App. (1882) 315.

Ischaemum rugosum Salisb. var. *segetum* Hack. in DC. Monog. Phan. 6 (1889) 206; Pilg. in Perk. Frag. Fl. Philip. (1904) 139.

Meoschium elegans Nees & Arn. in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 199; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 100.

Ischaemum rugosum F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 105; Miq. Fl. Ind. Bat. 3 (1857) 496; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 290; Rolfe in Journ. Bot. (1885) 216, vix Salisb.

Luzon (Ilocos Norte, Lepanto, Bontoc, Pangasinan, Rizal, Bataan), Polillo, *Cuming 565*, *Merrill 1550*, *Phil. Pl. 592*, *Sp. Blancoanae 165*. In open usually rather wet grasslands. India to China and Malaya.

Local names: Dava (Sub.); gulonglapas (Pang.); tinitrigo (Tag.).

ISCHAEMUM TODAYENSE Elm. Leaf. Philip. Bot. 7 (1915) 2678.

Mindanao (Davao), *Elmer 10521*. In ravines in secondary forests, altitude about 900 m. Endemic.

Local name: Halopos (Bag.).

EXCLUDED SPECIES

ISCHAEMUM POLYSTACHYUM Presl; F.-Vill. Novis. App. (1882) 315.

The type of this was from the Marianne Islands and F.-Villar does not really enter the species as Philippine. It is a synonym of *Ischaemum digitatum* Brongn.

ISCHAEMUM TIMORENSE Kunth; F.-Vill. Novis. App. (1882) 315.

ISCHAEMUM MINUS Presl Rel. Haenk. 1 (1828) 329; F.-Vill. Novis. App. (1882) 315; Kunth Enum. 1 (1833) 514; Miq. Fl. Ind. Bat. 3 (1857) 498.

"Hab. in insulis Philippinis." This is an American species erroneously localized by Presl. It is *Ischaemum urvilleanum* Kunth; see Hack. in DC. Monog. Phan. 6 (1889) 218. *Andropogon minor* Steud. Syn. Pl. Glum. 1 (1854) 377 is a synonym.

13a. EREMOCHLOA Büse

EREMOCHLOA CILIARIS (Linn.) Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 331.

Nardus ciliaris Linn. Sp. Pl. (1753) 53.

Ischaemum leersiioides Munro in Proc. Amer. Acad. 4 (1864-65) 363.

Eremochloa leersiioides Hack. in DC. Monog. Phan. 6 (1889) 264;

Pilger in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 6.

Luzon (Cagayan, Isabela), *Merrill 138*, *B. S. 7813*, *7832*, *8109 Ramos*. In open grasslands at low altitudes. Southern China to Tonkin and Pegu.

13b. ROTTBOELLIA Linnaeus f.

ROTTBOELLIA EXALTATA Linn. f. Suppl. (1781) 114; Nees in Nov.

Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 173, Hook. Journ. Bot.

Kew Miscel. 2 (1850) 100; Miq. Fl. Ind. Bat. 3 (1857) 407; F.-Vill.

Novis. App. (1882) 314; Vidal Phan. Cuming. Philip. (1885) 158,

Rev. Pl. Vasc. Filip. (1886) 290; Hack. in DC. Monog. Phan. 6

(1889) 293; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 328, Fl.

Manila (1912) 78.

Rottboellia setosa Presl Rel. Haenk. 1 (1828) 329; Kunth Enum. 1 (1833) 466.

Rottboellia denudata Steud. Syn. Pl. Glum. 1 (1854) 362; Miq. Fl.

Ind. Bat. 3 (1857) 408; F.-Vill. Novis. App. (1882) 314.

Rottboellia coelorrhachis F.-Vill. Novis. App. (1882) 314, non Forst.

Manisuris exaltata O. Kuntze Rev. Gen. Pl. (1891) 779.

Throughout the Philippines in open places, borders of thickets, etc.,

at low and medium altitudes; common. India to China, tropical Africa, through Malaysia to tropical Australia.

Local names: Agíngai (Tag.); annárai (Iv.); bódo (Bon.); bukal (C. Bis.); gaho (Bik.); girun (Bag.); nagéi (Bon.); sagisi (Ilk.).

ROTTBOELLIA GLANDULOSA Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 250; Hack. in DC. Monog. Phan. 6 (1889) 302; Pilger in Perk. Frag. Fl. Philip. (1904) 138; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 328.

Coelorhachis muricata Brongn. Bot. Voy. Coqu. 1 (1829) 65, t. 14.

Rottboellia muricata Miq. Fl. Ind. Bat. 3 (1857) 407; F.-Vill. Novis.

App. (1882) 314; Vidal Phan. Cuming. Philip. (1885) 158, Rev.

Pl. Vasc. Filip. (1886) 290, non Retz.

Manisuris glandulosa O. Kuntze Rev. Gen. Pl. (1891) 780.

Culion, Palawan, Balabac, Siargao, Mindanao, *Merrill 505*; *Phil. Pl. 1375*, *B. S. 34988 Ramos & Pascasio, 491 Mangubat*. In thickets, old clearings, etc., at low altitudes. Burma to Java.

Local name: Murabon (Sul.).

ROTTBOELLIA OPHIUROIDES Benth. Fl. Austral. 7 (1878) 514; Hack. in DC. Monog. Phan. 6 (1889) 303; Pilger in Perk. Frag. Fl. Philip. (1904) 138; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 328, 5 (1910) Bot. 326.

Ischaemum rottboellioides R. Br. Prodr. (1910) 205.

Manisuris rottboellioides O. Kuntze Rev. Gen. Pl. (1891) 779.

Andropogon rottboellioides Steud. Syn. Pl. Glum. 1 (1854) 382.

Luzon (Cagayan, Bontoc, Benguet, Nueva Vizcaya, Tarlac, Pampanga, Pangasinan, Zambales, Rizal), Masbate, Panay, Bohol. In open grasslands at low and medium altitudes, ascending to 2,000 m, often common. New Guinea, tropical Australia.

Local names: Gilon (C. Bis.); katalon (Ig.); lagopak (S. L. Bis.); talaño (Tag., Sbl.); tanlo (Ig.).

Var. **INTERMEDIA** Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 265, Merr. op. cit. 329.

Luzon (Benguet), *Merrill 4667*, *Elmer 6393*. On dry slopes in the pine region, altitude 1,000 to 1,600 m. Endemic.

ROTTBOELLIA TRIFLORA Hubbard in Philip. Journ. Sci. 9 (1914) Bot. 257.

Palawan, *Ledesma*. Borders of thickets at low altitudes. Endemic.

EXCLUDED SPECIES

ROTTBOELLIA MYUROS Linn.; Ceron Cat. Pl. Herb. Manila (1892) 181.

This record was probably based on an erroneously identified specimen of *Ophiurus monostachyus* Presl.

ROTTBOELLIA MUTICA Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 99.

This is an evident misprint for *Rottboellia muricata* Retz., a species that does not occur in the Philippines.

14. MANISURIS Swartz

MANISURIS GRANULARIS Linn. f. Nov. Gram. Gen. (1779) 37, f. 4-7;
Miq. Fl. Ind. Bat. 3 (1857) 409; F.-Vill. Novis. App. (1882) 314;
Hack. in DC. Monog. Phan. 6 (1889) 314; Merr. in Philip. Journ.
Sci. 1 (1906) Suppl. 329, Fl. Manila (1912) 79.

Cenchrus granularis Linn. Mant. 2 (1771) 575.

Hackelochloa granularis O. Kuntze Rev. Gen. Pl. (1891) 776.

Rytidix granularis Skeels in U. S. Dept. Agric. Bur. Pl. Ind. Bull.
282 (1913) 20.

Batan Islands to Mindanao, in most islands and provinces. In disturbed soil, open waste places, old clearings, etc., at low and medium altitudes. Pantropic.

Local name: Gin̄getkaran (Bag.).

15. OPHIURUS Gaertner

OPHIURUS EXALTATUS (Linn.) O. Kuntze Rev. Gen. Pl. (1891) 780;
Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 329.

Aegilops exaltata Linn. Mant. 2 (1771) 575.

Ophiurus corymbosus Gaertn. Fruct. 3 (1805) 4, t. 181, f. A; F.-Vill.
Novis. App. (1882) 314; Pilger in Perk. Frag. Fl. Philip. (1904)
138.

Rottboellia tongcalingii Elm. Leaf. Philip. Bot. 7 (1915) 2680.

Luzon (Isabela, Nueva Vizcaya, Negros, Mindanao (Bukidnon, Davao),
Merrill 127 p. p., Williams 2617, Elmer 10317, 10521, 16486, B. S. 8076
Ramos. In open rather damp places at low and medium altitudes. India
to tropical Australia.

Local names: Girun (Bag.); talañgiu (Bon.)

OPHIURUS MONOSTACHYUS Presl Rel. Haenk. 1 (1830) 330; Kunth
Enum. 1 (1833) 464; Miq. Fl. Ind. Bat. 3 (1857) 405; F.-Vill.
Novis. App. (1882) 314; Vidal Phan. Cuming. Philip. (1885)
158, Rev. Pl. Vasc. Filip. (1886) 290; Hack. in DC. Monog. Phan.
6 (1889) 318; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 329,
Fl. Manila (1912) 79.

Ophiurus undatus Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850)
100; Walp. Ann. 3 (1853) 791.

Ophiurus undulatus Miq. Fl. Ind. Bat. 3 (1857) 405.

Luzon (Cagayan, Ilocos Norte, Lepanto, Ifugao, Rizal), Mindanao (Bukidnon, Zamboanga, Lanao), Cuming 1339, Merrill Phil. Pl. 124, B. S. 32781
Ramos, 19978 McGregor, 37450, 39203 Ramos & Edaño. In open grasslands at low and medium altitudes. Southern China, Indo-China.

16. ARTHRAXON Beauvois

ARTHRAOXON HISPIDUS (Thunb.) Merr. in Philip. Journ. Sci. 7 (1912)
Bot. 229.

Phalaris hispida Thunb. Fl. Jap. (1784) 44.

Arthraxon ciliaris Beauv. Agrost. (1812) 111, t. 11, f. 6; F.-Vill.
Novis. App. (1882) 315.

Pleuroplitis langsdorffii Trin. Fund. Agrost. (1820) 174, f. 16.

Arthraxon ciliaris Beauv. subsp. *langsdorffii* Hack. in DC. Monog.
Phan. 6 (1889) 354, Philip. Journ. Sci. 1 (1906) Suppl. 265;
Merr. op. cit. 333; Perk. Frag. Fl. Philip. (1904) 139.

Arthraxon ciliaris Beauv. subsp. *nudus* Hack. in DC. Monog. Phan. 6 (1889) 356, Philip. Journ. Sci. 1 (1906) Suppl. 266; Merr. op. cit. 333.

Luzon (Cagayan, Ifugao, Bontoc, Lepanto, Benguet, Nueva Vizcaya), Mindanao (Lanao), Merrill 102, 4427, 4628, 7676, Phil. Pl. 175, Vanoverbergh 3811, 1760, 204, Clemens 100, B. S. 19628 McGregor, 5534, 5926, 5970, 7021, 7912 Ramos. In damp places about streams, in old rice paddies, etc., at medium altitudes, ascending to 1,600 m. Japan, China, Formosa. Local name: Timi-timí (Bon.).

ARTHRAOXON MICROPHYLLUS (Trin.) Hochst. in Flora 39 (1856) 188; Hack. in DC. Monog. Phan. 6 (1889) 351, Philip. Journ. Sci. 1 (1906) Suppl. 265; Merr. op. cit. 333, 5 (1910) Bot. 326.

Andropogon microphyllus Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 275.

Pleuroplitis microphylla Regel in Bull. Acad. St. Pétersb. 5 (1866) 749.

Luzon (Ilocos Norte, Bontoc, Benguet), Merrill 4363, 4719, Phil. Pl. 193, Clemens 9233, Vanoverbergh 1732, F. B. 12487 Merritt & Darling. On open slopes in the pine region, altitude 1,200 to 2,000 m. Abyssinia, India, Ceylon.

ARTHRAOXON QUARTINIANUS (A. Rich.) Nash in N. Am. Flora 17 (1912, Sept. 18) 99; Merr. in Philip. Journ. Sci. 7 (1912, Sept. 30) Bot. 229, Fl. Manila (1912) 77.

Alectoridia quartiniana A. Rich. Tent. Fl. Abyss. 2 (1851) 448, t. 99.

Arthraxon ciliaris Beauv. subsp. *quartinianus* Hack. in DC. Monog. Phan. 6 (1889) 356, Philip. Journ. Sci. 1 (1906) Suppl. 266; Merr. op. cit. 333, 5 (1910) Bot. 326.

Luzon (Ifugao, Bontoc, Benguet, Rizal), Merrill 4704, 4677, 7370, 9656, 4272, 6514, Phil. Pl. 999, Williams 1189, B. S. 19709 McGregor. On open slopes, in old rice paddies, etc., chiefly at altitudes from 1,100 to 1,600 m, rarely from about sea level. Abyssinia, India, introduced in the West Indies.

17. ANDROPOGON Linnaeus

ANDROPOGON ACICULATUS Retz. Obs. 5 (1789) 22; Hack. in DC. Monog. Phan. 6 (1889) 562; Pilger in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 338, Fl. Manila (1912) 83, Sp. Blancoanae (1918) 162.

Chrysopogon aciculatus Trin. Fund. Agrost. (1820) 188; Miq. Fl. Ind. Bat. 3 (1857) 490; F.-Vill. Novis. App. (1882) 316; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291.

Rhaphis trivialis Lour. Fl. Cochinch. (1790) 553; Blanco Fl. Filip. (1837) 45.

Andropogon acicularis Willd. Sp. Pl. 4² (1805) 906; Blanco Fl. Filip. ed. 2 (1845) 26, ed. 3, 1 (1877) 49.

Andropogon subulatus Presl Rel. Haenk. 1 (1828) 341; Kunth Enum. 1 (1833) 505.

Chrysopogon subulatus Miq. Fl. Ind. Bat. 3 (1857) 491; F.-Vill. Novis. App. (1882) 317.

Rhaphis javanica Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.

Throughout the Philippines in open places at low and medium altitudes; a veritable pest on account of its spikelets adhering to fur, clothing, etc. India to China southward through Malaya to tropical Australia and Polynesia.

Local names: Amor-seco (Sp.); dalekédek (Bon.); dalukdúk (Bon.); lakut-lapas (Sul.); marisékos (Tag.); marískos (Tag.); pagippi (Ibn.); tinlói (Tag.).

ANDROPOGON BAILEYI F.-Muell. in Vict. Nat. 7 (1891) 16; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 297; Merr. op. cit. 337.

Sorghum laxiflorum Bailey Syn. Queensl. Fl. Suppl. 3 (1890) 84, Queensl. Fl. 6 (1902) 1869; Rept. Exped. Bellend. Ker Range 25, non *Andropogon laxiflorus* Steud.

Luzon (Bontoc, Lepanto, Benguet, Rizal), Merrill 4399, Vanoverbergh 2131, 3339, B. S. 24082 Ramos. On open grassy slopes at medium altitudes, ascending to 1,200 m. Tropical Australia.

ANDROPOGON BREVIFOLIUS Sw. Prodr. (1788) 26; Hack. in DC. Monog. Phan. 6 (1889) 363; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291; Pilger in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 335.

Luzon (Cagayan, Ifugao, Bontoc, Benguet, Nueva Vizcaya, Bulacan, Zambales, Bataan, Laguna), Mindanao (Lanao, Bukidnon, Davao). In open grasslands at low and medium altitudes. Pantropic.

Local name: Bung-bung (Bag.).

ANDROPOGON CINCTUS Steud. Syn. Pl. Glum. 1 (1854) 398.

Chrysopogon micranthus Benth. var. *spicigerus* Benth. Fl. Austral. 7 (1878) 538.

Andropogon micranthus Kunth var. *spicigerus* Hack. in DC. Monog. Phan. 6 (1889) 489; Merr. in Philip. Journ. Sci. 1 (1906) 336, 3 (1908) Bot. 395.

Batan and Babuyan Islands, Luzon (Bontoc, Benguet), Merrill 4336, Elmer 5918, F. B. 15612, 4854 Curran, B. S. 5367 Ramos, 3726, 3918, 4040 Fénix. In open grasslands up to 1,800 m altitude. China, Australia, New Caledonia.

I believe this to be specifically distinct from *Andropogon micranthus* Kunth.

ANDROPOGON CITRATUS DC. Cat. Hort. Monsp. (1813) 78; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 167, Fl. Manila (1912) 82, Sp. Blancoanae (1918) 62.

Andropogon schoenanthus Blanco Fl. Filip. (1837) 39, ed. 2, (1845) 27, ed. 3, 1 (1877) 50; F.-Vill. Novis. App. (1882) 316; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 339, 2 (1907) Bot. 436, non Linn.

Cymbopogon citratus Stapf in Kew Bull. (1906) 257.

Planted here and there about houses in most or all parts of the Philippines for its fragrant leaves, but nowhere spontaneous; I have seen only one flowering specimen, Jolo, Link 215. A native of tropical Asia, now widely distributed in cultivation.

Local names: Balióko (Bis.); baráni (Ilk.); paja de méca (Sp.); tang-lád (Tag., Bik., Bis.); sálai (Tag.); sái (Mbo., Mand., Sul.); sálaid (Tag.).

ANDROPOGON CONTORTUS Linn. Sp. Pl. (1753) 1045; Blanco Fl. Filip. ed. 1 (1837) 38, ed. 2 (1845) 26, ed. 3, 1 (1877) 49; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 338, Fl. Manila (1912) 83, Sp. Blancoanae (1918) 61.

Heteropogon contortus Beauv. ex R. & S. Syst. 2 (1817) 836; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99; F.-Vill. Novis. App. (1882) 315; Miq. Fl. Ind. Bat. 3 (1857) 493; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291.

Heteropogon polystachyus F.-Vill. Novis. App. (1882) 315, non Schultes.

Heteropogon roxburghii Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99, haud Arn.

Luzon (Ilocos Norte, Union, Lepanto, Nueva Ecija, Nueva Vizcaya, Pangasinan, Pampanga, Rizal, Zambales), Mindoro, Mindanao (Bukidnon, Misamis). In open grasslands at low and medium altitudes, often very abundant. Certainly an introduced plant in the Philippines. Pantropic.

Local name: Sibat-sibátan (Tag.).

The Philippine form is mostly referable to the var. *hispidissimus* Hack. in DC. Monog. Phan. 6 (1889) 587.

ANDROPOGON FASTIGIATUS Sw. Prodr. (1788) 26; Hack. in DC. Monog. Phan. 6 (1889) 393, Philip. Journ. Sci. 1 (1906) Suppl. 267; Merr. op. cit. 335.

Luzon (Lepanto, Bontoc, Pangasinan), Merrill 4461, B. S. 30055 Ramos. On dry open grassy slopes at low and medium altitudes. Pantropic.

ANDROPOGON FESTUCOIDES Presl Rel. Haenk. 1 (1828) 340; Kunth Enum. 1 (1833) 500; Miq. Fl. Ind. Bat. 3 (1857) 489; F.-Vill. Novis. App. (1882) 316; Merr. Sp. Blancoanae (1918) 61.

Andropogon anis Llanos Frag. Pl. Filip. (1851) 29; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 20; F.-Vill. Novis. App. (1882) 316.

Andropogon muricatus Retz. var. *aristatus* Büse in De Vriese Pl. Ind. Bat. Orient. (1856) 104.

Andropogon nigritanus Benth. in Hook. Niger Fl. (1849) 573.

Andropogon squarrosus Linn. f. var. *nigritanus* Hack. in DC. Monog. Phan. 6 (1889) 544.

Vetiveria zizanioides Stapf var. *nigrimana* Camus in Bull. Mus. Hist. Nat. Paris 25 (1919) 674.

Luzon (Pampanga, Nueva Ecija), Merrill 4231, 4240, Phil. Pl. 389, B. S. 22238 Santos. In open level lands, rice-paddy banks, etc., at low altitudes. Tropical Africa, Java.

Local name: Aniás (Pamp.).

The roots are entirely odorless in striking contrast to those of *Andropogon zizanioides* Urban to which it has been erroneously reduced.

ANDROPOGON FILIPENDULUS Hochst. in Flora 29 (1846) 115, var. **LACHNATHERUS** (Benth.) Hack. in DC. Monog. Phan. 6 (1889) 635, Philip. Journ. Sci. 1 (1906) Suppl. 267; Merr. op. cit. 339. *Andropogon lachnatherus* Benth. Fl. Austral. 7 (1878) 534.

Luzon (Bontoc, Lepanto, Benguet), *Merrill 7799, 4398, Elmer 6392*. On open grassy slopes, altitude 1,100 to 1,600 m.

Forma **BISPICULATA** Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 269; Merr. op. cit. 339.

Luzon (Benguet), *Merrill 4298*. On open slopes, altitude about 1,350 m. The var. *lachnatherus* Hack. in tropical Australia, other varieties in tropical Africa and Ceylon.

Local names: Páuai (Ig.); taáu (Bon.).

ANDROPOGON FRAGILIS R. Br. Prodr. (1810) 202, var. **LUZONIENSIS** Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 267; Merr. op. cit. 335.

Luzon (Bontoc, Benguet), *Merrill 4468, 4386*. In dry open grasslands at medium altitudes.

Var. **MALAYANUS** Merr. var. nov. A typo differt spiculis pedicellisqu omnino glabris.

Andropogon fragilis Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 335, Fl. Manila (1912) 82, haud R. Br.

Luzon (Bontoc, Ifugao, Pangasinan, Rizal), Mindanao (Lanao), *Merrill 8012, Phil. Pl. 156, Loher 1864, B. S. 19977 McGregor*. In dry open grasslands at low and medium altitudes. New Guinea, *King 1002*. The species in southern China, New Guinea, and in tropical Australia, a variety also in China.

ANDROPOGON GRYLLUS Linn. Amoen. Acad. 4 (1759) 332; Hack. in DC. Monog. Phan. 6 (1889) 550, var. **PHILIPPINENSIS** Merr. in Philip. Journ. Sci. 14 (1919) 368.

Chrysopogon gryllus Trin.; F.-Vill. Novis. App. (1882) 316 (prob.).

Panay, *B. S. 30964 Ramos & Edaña*. On bowlders and ledges along streams at low altitudes. The variety endemic, the species, in various forms, southern Europe to Australia.

ANDROPOGON HALEPENSIS (Linn.) Brot. Fl. Lusit. 1 (1804) 89, var. **PROPINQUUS** (Kunth) Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 336, Fl. Manila (1912) 84.

Andropogon propinquus Kunth Enum. 1 (1833) 502.

Andropogon affinis Presl Rel. Haenk 1 (1828) 343, non R. Br.

Sorghum halepense Pers. var. *muticum* Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.

Sorghum halepense Miq. Fl. Ind. Bat. 3 (1857) 503; F.-Vill. Novis. App. (1882) 317; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 184, haud Pers.

Andropogon sorghum Pers. subsp. *halepensis* Hack. var. *propinquus* Hack. in DC. Monog. Phan. 6 (1889) 503.

Batan Islands and northern Luzon to Mindanao, in most or all islands and provinces, often common. The species pantropic, the variety Ceylon to New Guinea.

Local names: Aroró (Bik.); baku (If.) batad-batádan (Tag.); dala-pak (Sub.); gingai (Sul.); giñgia (Bag.); riaría (Sul.); sikal (Ibn.); uginaí (Buk.).

- ANDROPOGON INTERMEDIUS** R. Br. Prodr. (1810) 202; Hack. in DC. Monog. Phan. 6 (1889) 485; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 335 var. **HAENKEI** (Presl) Hack. in DC. Monog. Phan. 6 (1889) 486; Pilger in Perk. Frag. Fl. Philip. (1904) 139; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 335, Fl. Manila (1912) 83.
Andropogon haenkei Presl Rel. Haenk. 1 (1828) 340; Kunth Enum. 1 (1833) 501; Miq. Fl. Ind. Bat. 3 (1857) 489; F.-Vill. Novis. App. (1882) 316.
Rhaphis stricta Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99; Walp. Ann. 3 (1853) 803.
Andropogon leptanthus Steud. Syn. Pl. Glum. 1 (1854) 391; Miq. Fl. Ind. Bat. 3 (1857) 389; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291.

Luzon (Cagayan, Ilocos Norte, Union, Nueva Vizcaya, Nueva Ecija, Pangasinan, Rizal, Batangas), Negros, Mindanao, *Cuming 1400, Merrill 6734, 313, 5456, Phil. Pl. 134, 1842, B. S. 30072 Fénix, 8246, 7830, 7658, 7918, 7916, 1426 Ramos*. In open grasslands at low and medium altitudes. The species in China and Australia, the variety in China and Ceylon, other varieties in Asia and Africa.

Local names: Sálai (Pang.); sálai-párang (Tag.).

- ANDROPOGON MICRANTHUS** Kunth Rev. Gram 1 (1829) 165; Enum. 1 (1833) 504; Hack. in DC. Monog. Phan. 6 (1889) 488 (var. *genuinus*); Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 336.
Andropogon alternans Presl Rel. Haenk. 1 (1828) 342.
Chrysopogon violascens Trin. in Mém. Acad. St. Pétersb. VI 2 (1833) 319.
Rhaphis villosulus Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102.
Rhaphis microstachys Nees l. c., var.
Chrysopogon villosulus Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291.
Andropogon parvispica Steud. Syn. Pl. Glum. 1 (1854) 397.
Chrysopogon parviflorus Benth. Fl. Austral. 7 (1878) 537; F.-Vill. Novis. App. (1882) 316.

Luzon (Ilocos Norte, Bontoc, Benguet, Nueva Vizcaya, Nueva Ecija, Zambales), *Cuming 980, 1397, Merrill 118, 4434, 4703, Phil. Pl. 176, Elmer 6593, Williams 1959, B. S. 32948 Ramos, 32010 Santos, 26038 Fénix, 14198 McGregor, 8241 Ramos*. In open grasslands chiefly at medium altitudes, ascending to 2,300 m. Abyssinia to China through Malaya to tropical Australia.

Local names: Sañgongŕoan (Bon.); sañguan (Bon.); sisamon (Ig.).

- ANDROPOGON NARDUS** Linn. Sp. Pl. (1753) 1046 var. **TORTILIS** (Presl) comb. nov.
Anthistiria tortilis Presl Rel. Haenk. 1 (1828) 347; Kunth Enum. 1 (1833) 483; F.-Vill. Novis. App. (1882) 317.
Andropogon hamatulus Nees in Hook. & Arn. Bot. Beechy's Voy. (1841) 244.
Andropogon martyni Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.

Andropogon nardus Linn. var. *hamatulus* Hack. in DC. Monog. Phan. 6 (1889) 606; Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 339, 3 (1908) Bot. 396.
Andropogon nardus Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291, non Linn.

Batan Islands, Luzon (Cagayan, Isabela, Ilocos Norte, Abra, Bontoc, Pangasinan, Zambales, Nueva Ecija), Cagayan Sulu, *Merrill 5307*, *Cuming 1000*, *B. S. 7893, 7793, 8047, 32888 Ramos, 10169 McGregor, F. B. 5858, 17037 Curran, 16461 Bacani*. In open grasslands at low and medium altitudes. The variety in southern China, the species in India, Africa, and Malaya.

Local name: Balaniu (Bon.).

ANDROPOGON NITIDUS (Vahl) Kunth Rev. Gram. 1 (1829) 166; Merr. Fl. Manila (1912) 84.

Holcus nitidus Vahl Symb. 2 (1791) 102.

Sorghum nitidum Pers. Syn. 1 (1805) 101.

Andropogon fuscus Presl Rel. Haenk. 1 (1828) 342; Kunth Enum. 1 (1833) 503.

Sorghum fuscum Miq. Fl. Ind. Bat. 3 (1857) 503; F.-Vill. Novis. App. (1882) 317; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291.

Sorghum tropicum Nees var. *muticum* Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.

Andropogon pedicellatus Steud. Syn. Pl. Glum. 1 (1854) 394; Miq. Fl. Ind. Bat. 3 (1857) 488; F.-Vill. Novis. App. (1882) 316.

Andropogon serratus Thunb. var. *nitidus* Hack. in DC. Monog. Phan. 6 (1889) 521; Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 337, 3 (1908) Bot. 396.

Batan Islands, northern Luzon (Cagayan) to Mindanao, in most islands and provinces. In open grasslands at low and medium altitudes.

Forma **HIRSUTA** Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 337.

Luzon (Isabela), *Merrill 137*. In open grasslands at low altitudes. The form endemic, the species India to southern China and Formosa southward to Java.

ANDROPOGON PHILIPPINENSIS Merr. in Philip. Journ. Sci. 14 (1919) 367.

Luzon (Ilocos Norte), *B. S. 32946, 32755 Ramos*. On dry open rocky slopes at low altitudes. Endemic.

ANDROPOGON SANGUINEUS (Retz.) Merr. in Philip. Journ. Sci. 12 (1917) Bot. 101; op. cit. 14 (1919) 369.

Rottboellia sanguinea Retz. Obs. 3 (1783) 25.

Thelepogon sanguineus Spreng. Syst. 1 (1825) 299.

Andropogon pseudograya Steud. Syn. Pl. Glum. 1 (1854) 465.

Luzon (Ilocos Norte), *Culion, B. S. 32777, 41310, 41329 Ramos*. On open slopes at low altitudes. Southern China, India, Ceylon, and Madagascar.

ANDROPOGON SERICEUS R. Br. Prodr. (1810) 201; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99; Hack. in DC. Monog. Phan. 6 (1889) 575; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 291; Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 338, Fl. Manila (1912) 82.

Andropogon tenuiculus Steud. Syn. Pl. Glum. 1 (1854) 371.

Heteropogon tenuiculus Miq. Fl. Ind. Bat. 3 (1857) 494; F.-Vill. Novis. App. (1882) 315.

Luzon (Benguet, Nueva Ecija, Pangasinan, Rizal), *Cuming 1398, Merrill 266, Phil. Pl. 163, B. S. 4824 Ramos*. In open grasslands at low altitudes. New Guinea, New Caledonia, Australia.

ANDROPOGON SERRATUS Thunb. Fl. Jap. (1784) 41; Hack. in DC. Monog. Phan. 6 (1889) 520; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 337, Fl. Manila (1912) 84.

Sorghum fulvum Beauv. Agrost. (1812) 164; F.-Vill. Novis. App. (1882) 317.

Luzon (Bontoc, Benguet, Rizal), Mindanao (Lanao, Davao, Bukidnon), *Merrill 4297, Phil. Pl. 593, Elmer 5958, Clemens 597, Piper 471, B. S. 5764 Ramos*. In open grasslands at low and medium altitudes. Japan and China through Malaya to Australia.

Local names: Girun (Bag.); taáu (Bon.); tígau (Mag.).

ANDROPOGON SORGHUM (Linn.) Brot. Fl. Lusit. 1 (1804) 88; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 336, Fl. Manila (1912) 84, Sp. Blancoanae (1918) 63 (var. *vulgaris* Hack.); Pilger in Perk. Frag. Fl. Philip. (1904) 140 (var. *sativus* Hack.); Usteri Beitr. Ken. Philip. Veg. (1905) 132 (var. *niger* Kunth).

Holcus sorghum Linn. Sp. Pl. (1753) 1047.

Holcus saccharatus Linn. Sp. Pl. (1753) 1047; Blanco Fl. Filip. (1837) 47, ed. 2 (1845) 32, ed. 3, 1 (1877) 58, t. 436.

Sorghum saccharatum Pers. Syn. 1 (1804) 101; F.-Vill. Novis. App. (1882) 317; Vidal Rev. Pl. Vasc. Filip. (1886) 291.

Sorghum vulgare Pers. Syn. 1 (1804) 101; Miq. Fl. Ind. Bat. 3 (1857) 502.

Andropogon sorghum Brot. subsp. *sativus* Hack. var. *saccharatus* Hack. in DC. Monog. Phan. 6 (1889) 505.

Several forms or varieties of sorghum are cultivated in the Philippines, but none are established. A native of Asia or Africa, now cultivated in all warm countries. Sorghum.

Local names: Bakau (Ibn.); bátad (Tag., Bik., Bis.); bátag (Tag.); bukákau (Ilk.); gau (If.); layagan (Sul.).

ANDROPOGON TRITICEUS R. Br. Prodr. (1810) 201; Hack. in DC. Monog. Phan. 6 (1889) 588; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 338.

Andropogon ischyranthus Steud. Syn. Pl. Glum. 1 (1854) 367.

Heteropogon ischyranthus Miq. Fl. Ind. Bat. 3 (1857) 493; F.-Vill. Novis. App. (1882) 315.

Heteropogon insignis Thwaites Enum. Pl. Zeyl. (1859-64) 437; F.-Vill. Novis. App. (1882) 315.

Luzon (Abra, Ilocos Norte, Pangasinan), Busuanga, *Cuming 1003, B. S. 7216, 4900, 41269 Ramos, F. B. 12477 Merritt & Darling*. In open grasslands at low or medium altitudes. Ceylon and Bruma to tropical Australia.

ANDROPOGON ZIZANIOIDES (Linn.) Urban Symb. Antil. 4 (1903) 79; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 228, Fl. Manila (1912) 83, Sp. Blancoanae (1918) 62.

Phalaris zizanioides Linn. Mant. 2 (1771) 183.

Andropogon squarrosus Hack. in DC. Monog. Phan. 6 (1889) 542; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 337, non Linn. f. (cf. Stapf in Kew Bull. (1906) 348.

Andropogon muricatus Retz. Obs. 3 (1783) 43, 5 (1789) 20; F.-Vill. Novis. App. (1882) 316.

Andropogon nardus Blanco Fl. Filip. (1837) 39, ed. 2 (1845) 27, ed. 3, 1 (1877) 51, non Linn.

Vetiveria zizanioides Stapf in Kew Bull. (1906) 346; Camus in Bull. Mus. Hist. Nat. Paris 25 (1919) 673.

Batan Islands, Luzon (Amburayan, Rizal, Cavite), *Merrill Sp. Blancoanae 355, Phil. Pl. 118, Loher 7213*. Planted about rice-paddy banks, often abundant; certainly an introduced plant in the Philippines. A native of tropical Asia, now pantropic. Vetiver.

Local names: Amóora (C. Bis.); amóras (Ilk.); anías de móras (Pamp.); giron (P. Bis.); filib (Pamp.); móra (Bik., Bis.); móras (Tag., Bis., Bik.); móro (Tag.); narawastu (Sul.); raíz de móras (Sp.); rimódas (P. Bis.); rimóra (Sbl.); rimóras (Bik.); tres-móras (Bis.).

EXCLUDED SPECIES

ANDROPOGON ERIOSTACHYS Presl Rel. Haenk. 1 (1830) 339; Kunth Enum. 1 (1833) 496; Miq. Fl. Ind. Bat. 3 (1857) 487; F.-Vill. Novis. App. (1882) 316.

"Hab. in insulis Philippinis" Presl. This is the tropical American *Andropogon ternatus* Nees var. *eristachys* (Presl) Hack. in DC. Monog. Phan. 6 (1889) 425. The type was one of the numerous Malaspina Expedition plants collected by Presl but erroneously localized as Philippine.

HETEROPOGON TENELLUS Schultes; F.-Vill. Novis. App. (1882) 315.

18. ASTHENOCHLOA Büse

ASTHENOCHLOA TENERA Büse in Miq. Pl. Jungh. (February, 1854) 367.

Andropogon leptos Steud. Syn. Pl. Glum. 1 (1854) 397; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 268, 337, 4 (1909) Bot. 250.

Garnotiella philippinensis Stapf in Hook. Ic. IV 5 (1896) t. 2494; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 374.

Garnotiella leptos Stapf in Kew Bull. (1910) 302.

Luzon (Benguet), Panay, *Merrill 4322, Phil. Pl. 154, B. S. 5318 Ramos*. On steep open slopes up to an altitude of 1,400 m. Originally described from "Andor," a locality I have not been able to locate; perhaps "Ind. or." intended. Java. A monotypic genus.

The alliance of the genus is with *Andropogon*, not with *Garnotia*.

19. THEMEDA Forskål

THEMEDA GIGANTEA (Cav.) Hack. in DC. Monog. Phan. 6 (1889) 670 (var. *genuina* Hack.); Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 340, 3 (1908) Bot. 396, 5 (1910) Bot. 326.

Anthistiria gigantea Cav. Ic. 5 (1799) 36, t. 458; Willd. Sp. Pl. 4² (1805) 902; F.-Vill. Novis. App. (1882) 317; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292.

Androscepiia gigantea Brongn. Bot. in Duperr. Voy. (1829) 78; Miq. Fl. Ind. Bat. 3 (1857) 506; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.

Perobachne secunda Presl Rel. Haenk. 1 (1828) 348, t. 48; Kunth Enum. 1 (1833) 485; Miq. Fl. Ind. Bat. 3 (1857) 507; F.-Vill. Novis. App. (1882) 317.

Batan Islands to Mindanao, in most or all islands and provinces. In open grasslands on dry steep slopes at low and medium altitudes. India to China, Malaya, New Caledonia, and Australia.

Local names: 'Girun (Bag.); maṅgambit (Sub.); sai-gaja (Sul.); salai-sálai (Tag.); tanháð (Iv.); tanik (Bon.); taurar (Ilk.); teldak (Pang.); tulaṅgo (Tag.); talauyo (Tag.).

Var. **VULPINA** (Anders.) Hack. in DC. Monog. Phan. 6 (1889) 670; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 340.

Anthistiria vulpina Anders. in Nov. Act. Upsal. III 2 (1856) 423.

Anthistiria arundinacea Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292, haud Roxb.

Luzon (Cagayan), *Cuming 1272* and numerous other specimens. India. Scarcely distinguishable from the species.

Subsp. **INTERMEDIA** Hack. var. **DUBIA** Hack. in DC. Monog. Phan. 6 (1889) 675; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 341.

Mindanao (Misamis), *Cuming 1609*.

The species is variable in its spikelet characters, a number of forms being represented in the Philippines.

THEMEDA TRIANDRA Forsk. Fl. Aegypt.-Arab. (1775) 178; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 340, Fl. Manila (1912) 81. *Themeda forskalii* Hack. in DC. Monog. Phan. 6 (1889) 659; Pilger in Perk. Frag. Fl. Philip. (1904) 140.

Anthistiria ciliata Cav. Ic. 5 (1799) 36, t. 459; F.-Vill. Novis. App. (1882) 317; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99 (var. *subglabra* Nees), non Linn. f.

Anthistiria depauperata Anders. in Nov. Act. Upsala III 2 (1856) 343; F.-Vill. Novis. App. (1882) 317.

Anthistiria barbata Presl Repert. 1 (1833) 305, non Desf.

Apluda barbata Llanos in Mém. Acad. Cienc. Madr. 4 (1858) 497; Fl. Filip. ed. 3, 4¹ (1880) 99 (misprint for *Anthistiria barbata*).

Batan Islands, Luzon (Cagayan, Isabela, Nueva Vizcaya, Ilocos Norte, Abra, Bontoc, Benguet, Pangasinan, Nueva Ecija, Rizal, Albay), Mindoro, Busuanga, Semirara, Mindanao. In open grasslands at low and medium

altitudes, ascending to 2,300 m. Frequently abundant and gregarious over large areas. Tropical Asia and Africa through Malaya to Australia.

Local names: Punáu (Bon.); samsamóng (Ilk.); táau (Ig.); usímau (Ilk.).

EXCLUDED SPECIES

ANTHISTIRIA JUNGHUHNIANA Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 99.

This is credited to the Philippines by Nees, on the basis of *Cuming* 2431, which was from Sumatra, not from the Philippines. It is a synonym of *Themeda frondosa* (R. Br.) Merr. (*T. arguens* Hack., not *Stipa arguens* Linn.), which does not occur in the Philippines.

20. PEROTIS Aiton

PEROTIS INDICA (Linn.) O. Kuntze Rev. Gen. Pl. (1891) 787; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 341.

Anthoxanthum indicum Linn. Sp. Pl. (1753) 28.

Perotis latifolia Ait. Hort. Kew. 1 (1789) 85; F.-Vill. Novis. App. (1882) 313.

Perotis glabrata Steud. Syn. Pl. Glum. 1 (1854) 186; Miq. Fl. Ind. Bat. 3 (1857) 479; F.-Vill. Novis. App. (1882) 313; Vid. Phan. Cuming. Philip. (1885) 158, Rev. Pl. Vasc. Filip. (1886) 289.

Perotis hordeiformis Nees in Steud. Nomencl. ed. 2, 2 (1841) 306, Hook. Journ. Bot. Kew Miscel. 2 (1850) 102.

Luzon (Cagayan, Nueva Ecija), Mindanao (Agusan), *Cuming* 1899, *Elmer* 13873, *B. S.* 7911 *Ramos*. In open grasslands at low and medium altitudes. Tropical Asia and Africa.

Local name: Bañgat (Mbo.).

PEROTIS RARA R. Br. Prodr. (1810) 172; Miq. Fl. Ind. Bat. 3 (1857) 479; F.-Vill. Novis. App. (1882) 314; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 341.

Xystidium barbatum Presl Rel. Haenk. 1 (1828) 228; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 55.

Xystidium maritimum Trin. Fund. Agrost. (1820) 102, t. 2.

Luzon (Ilocos Norte, Lepanto, Amburayan, Pangasinan, Zambales), Mindoro, *Merrill* 4447, *Phil. Pl.* 152, *F. B.* 9979 *Merritt*, 5859 *Curran*. On dry open slopes at low and medium altitudes. Australia.

21. ZOISIA Willdenow

(*Zoysia* auct.)

ZOISIA MATRELLA (Linn.) Merr. in Philip. Journ. Sci. 7 (1912) Bot. 230, Fl. Manila (1912) 85.

Agrostis matrella Linn. Mant. 2 (1771) 185.

Zoysia pungens Willd. in Ges. Naturf. Fr. Neue Schrift 3 (1801) 441; F.-Vill. Novis. App. (1882) 313; Pilger in Perk. Frag. Fl. Philip. (1904) 140; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 342.

Osterdamia matrella O. Kuntze Rev. Gen. Pl. (1891) 781.

Northern Luzon to Palawan, in most islands, occurring in open places chiefly along or near tidal streams. Tropical Asia and Africa through Malaya to Australia and western Polynesia.

Local name: Barit-baritan (Tag.).

22. ARUNDINELLA Raddi

ARUNDINELLA AGROSTOIDES Trin. Sp. Gram. Ic. (1836) 23, t. 265;
F.-Vill. Novis. App. (1882) 318; Merr. Fl. Manila (1912) 85.

Luzon (Ilocos Norte, Pangasinan, Rizal), *Merrill 7399, Phil. Pl. 594, Loher 7217, F. B. 12482 Merritt & Darling*. In open grasslands and on ledges along small streams at low altitudes. India to Burma.

Var. **CILIATA** (Roxb.) Hook. f. Fl. Brit. Ind. 7 (1897) 71; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 268; Merr. op. cit. 243.

Holcus ciliatus Roxb. Fl. Ind. 1 (1820) 321.

Arundinella ciliata Nees in Wall. Cat. (1829) No. 1668.

Luzon (Ilocos Sur, Bontoc, Benguet), *Merrill 4328, Williams 1961, F. B. 15940 Bacani, 15671 Merritt & Darling*. On open slopes, about cliffs, etc., altitude 800 to 1,600 m. Range of the species.

ARUNDINELLA MILIACEA Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; Vidal Phan. Cuming. Philip. (1885) 158.

Arundinella nepalensis F.-Vill. Novis. App. (1882) 318; Vidal Rev. Pl. Vasc. Filip. (1886) 289; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 343, non Trin.

Arundinella hispida Hack. subsp. *humilior* Hack. in Bull. Herb. Boiss. II 4 (1904) 527, Allgem. Bot. Zeitschr. 12 (1906) 179.

Arundinella filiformis Jan. ex Mez in Fedde Repert. 17 (1921) 85.

Arundinella caespitosa Jan. l. c.

Luzon (Apayao, Bulacan, Rizal, Laguna), Panay, *Merrill 2262, Phil. Pl. 161, Cuming 667, B. S. 30966 Ramos & Edaño, 6700 Robinson, 35453 Martelino & Edaño, 19286 Reillo, 28251 Fénix, 2101 Ramos, 30 Foxworthy, F. B. 7173 Curran, Kneucker Gram. Exsic. 606*. On ledges and banks along swift-running streams at low altitudes. Formosa.

The more-recent collections have been distributed as *A. hispida* (HBK.) O. Kuntze, following Hackel's identification.

ARUNDINELLA PUBESCENS Merr. & Hack. in Philip. Journ. Sci. 2 (1907) Bot. 419.

Palawan, *B. S. 856 Foxworthy*. On river banks at low altitudes. Endemic.

ARUNDINELLA SETOSA Trin. Diss. 2 (1824) 63, Gram. Pan. (1826) 245; F.-Vill. Novis. App. (1882) 318; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 342, 5 (1910) Bot. 326.

Arundinella stricta Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; Walp. Ann. 3 (1853) 804; Miq. Fl. Ind. Bat. 3 (1857) 520.

Danthonia luzonensis Steud. Syn. Pl. Glum. 1 (1854) 245; Miq. Fl. Ind. Bat. 3 (1857) 427; F.-Vill. Novis. App. (1882) 319; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292.

Arundinella setosa Trin. var. *stricta* Pilger in Perk. Frag. Fl. Philip. (1904) 140.

Luzon (Cagayan, Isabela, Bontoc, Benguet, Nueva Vizcaya, Zambales, Rizal), Mindoro, Mindanao (Bukidnon). In open grasslands, on cliffs, etc., at low and medium altitudes, ascending to 2,000 m. India to China through Malaya to New Guinea.

Local name: Salai-salai (Sbl.).

23. THYSANOLAENA Nees

THYSANOLAENA MAXIMA (Roxb.) O. Kuntze Rev. Gen. Pl. (1891) 794; Pilger in Perk. Frag. Fl. Philip. (1904) 141; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 343.

Agrostis maxima Roxb. Fl. Ind. 1 (1820) 319.

Thysanolaena agrostis Nees in Edinb. New Philos. Journ. 18 (1835) 180.

Thysanolaena acarifera Arn. & Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 181; Vidal Rev. Pl. Vasc. Filip. (1886) 289.

Luzon (Ilocos Norte, Apayao, Bontoc, Benguet, Nueva Vizcaya, Nueva Ecija, Bulacan, Zambales, Bataan, Laguna, Tayabas, Sorsogon), Mindoro, Palawan, Mindanao. On banks of ravines, slopes, about cliffs, etc., at low and medium altitudes, ascending to 1,600 m. India through Malaya to New Guinea.

Local names: Bugúbui (Pamp.); buybúi (Ilk.); eagadu (Bon.); gatbó (Bik.); lása (Tag.); tagádeu (Bon.); tagisa (Sub.); tambú (Tag.); tálankaran (Ig.).

24. DIGITARIA Scopoli

DIGITARIA CHINENSIS Nees in Hook. & Arn. Bot. Beechey Voy. (1841) 231.

Paspalum filiculme Nees ex Thwaites Enum. Pl. Zeyl. (1859-64) 358.

Digitaria violascens Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 347, 3 (1908) Bot. 396, Fl. Manila (1912) 88, non Link.

Batan Islands and northern Luzon to Palawan and Mindanao, in most islands and provinces, Merrill 4148, 9388, 7107, 7677, 4788, Phil. Pl. 174, B. S. 21715, 14524, 13939, 8250 Ramos. In open grasslands at low and medium altitudes. India to China and Malaya.

Local names: Timi (Bon.); tumi (Bon.); ualisíbis (Iv.).

DIGITARIA CORYMBOSA (Roxb.) comb. nov.

Panicum corymbosum Roxb. Hort. Beng. (1814) 7, *nomen nudum*, Fl. Ind. 1 (1820) 292.

Panicum microbachne Presl Rel. Haenk. 1 (1828) 298; Kunth Enum. 1 (1833) 81.

Digitaria consanguinea Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 347, 3 (1908) Bot. 396, Fl. Manila (1912) 88, Sp. Blancoanae (1918) 64, non Gaudich.

Panicum sanguinale Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 157, non Linn.

Digitaria lanosa Llanos Fragm. Pl. Filip. (1851) 28; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 19.

Panicum gaudichaudii F.-Vill. Novis. App. (1882) 311, non Kunth.

Panicum pruriens Trin. var. *glabrum* Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Throughout the Philippines at low and medium altitudes in open waste places, old clearings, etc., usually common, Merrill 4230, 5347, 7363, 9276, Sp. Blancoanae 787, Phil. Pl. 126, Elmer 18216, 16818, 16417, 9570, B. S. 12036, 13334, 32736, 14746 Ramos. India to southern China and Formosa through Malaya to Samoa.

Local names: Babanai (Sub.); balíli (C. Bis.); balisíbis (Iv.); balud-

gañgan (Tag.); barírid (Bag.); gasa (Kuy.); halos (Tag.); maya-máya (S. L. Bis.); maraséda (Ilk.); sagísi (Ilk.).

This species has been confused with *Digitaria sanguinalis* Scop. and by me with *D. consanguinea* Gaudich. It is distinguished by its almost obsolete first glume and short second glume, the latter rarely more than one-fourth as long as the spikelet.

DIGITARIA HETERANTHA (Nees & Mey.) var. **PACHYRHACHIS** (Hack.) comb. nov.

Panicum heteranthum Nees & Mey. var. *pachyrhachis* Hack. in Philip. Journ. Sci. 3 (1908) Bot. 167.

Palmas, Lumbucan, Merrill 5355, 5276. Along sandy seashores. The variety endemic, the species in the Malay Peninsula, Java, China, and Formosa.

DIGITARIA LONGIFLORA (Gmel.) Pers. Syn. 1 (1805) 85; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 347, 5 (1910) Bot. 326, Fl. Manila (1912) 88, Sp. Blancoanae (1918) 64.

Panicum longiflorum Gmel. Syst. (1788) 158.

Panicum parvulum Trin. in Mém. Acad. St. Pétersb. VI 3 (1835) 205; Mez in Perk. Frag. Fl. Philip. (1904) 142.

Paspalum brevifolium Flüggé Monogr. (1810) 150; F.-Vill. Novis. App. (1882) 310.

Paspalum fuscescens Presl Rel. Haenk. 1 (1828) 213, ex descr.

Syntherisma fuscescens Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 49, t. 10.

Paspalum fasciculatum Llanos Frag. Pl. Filip. (1851) 23; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 16, non Willd.

Northern Luzon to Mindanao, in most islands and provinces, *Elmer* 16733, *Merrill* 249, 331, 1129, 6518, *Phil. Pl.* 167, *Sp. Blancoanae* 1022, *B. S.* 32751, 33448 *Ramos*, 32009 *Fénix*, 10911 *McGregor*. In open grasslands, sea level to an altitude of 1,600 m. Warmer parts of the Old World.

Local name: Sasámon (Ig.).

DIGITARIA PEDICELLARIS (Trin.) Prain Bengal Pl. (1903) 1182; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 348, Fl. Manila (1912) 98.

Paspalum pedicellare Trin. ex Steud. Nomencl. ed. 2, 2 (1840) 272.

Panicum pedicellare Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 268.

Panicum puberulum Mez in Perk. Frag. Fl. Philip. (1904) 143, non (?) Kunth.

Luzon (Nueva Vizcaya, Nueva Ecija, Pampanga, Rizal, Bataan), Culion, *Merrill* 4225, 3164, 105, *Phil. Pl.* 577, *B. S.* 26273 *Ramos* & *Edaño*, *Williams* 93. In open grasslands at low and medium altitudes, ascending to 850 m. India to Burma.

DIGITARIA SANGUINALIS (Linn.) Scop. Fl. Carn. ed. 2, 1 (1772) 52; Vidal Rev. Pl. Vasc. Filip. (1886) 286; Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 346, Fl. Manila (1912) 88 (var. *australis* Thw.).

Panicum sanguinale Linn. Sp. Pl. (1753) 57.

Digitaria inaequalis Spreng. Syst. 1 (1825) 271.

Digitaria biformis Willd. Enum. (1809) 92.

Digitaria radicata Miq. Fl. Ind. Bat. 3 (1857) 437.

Panicum radicosum Presl Rel. Haenk. 1 (1828) 297; F.-Vill. Novis. App. (1882) 311.

Paspalum inaequale Link Enum. Hort. Berol. 1 (1821) 103; Kunth Enum. 1 (1833) 48; F.-Vill. Novis. App. (1882) 310.

Syntherisma sanguinalis Dulac Fl. Hautes Pyr. (1867) 77.

Widely scattered in the Philippines in open waste places, *B. S.* 27396, 7258, 7092 *Ramos*, 3681 *Fénix*, 18737 *Otanes*, *Merrill* 7258, *Phil. Pl.* 114, *Williams* 1172. All warm countries.

Local names: Dibutbut (Iv.); pagpagai (Bon.); saka-sáka (Ilk.).

Var. **FIMBRIATA** (Link.) Stapf in Prain Fl. Trop. Afr. 9 (1919) 440.

Digitaria fimbriata Link. Hort. Berol. 1 (1821) 226.

Digitaria ciliaris Merr. in Philip. Journ. Sci. 5 (1910) Bot. 168, Fl. Manila (1912) 87, non ? Pers.

Panicum ciliare Retz. Obs. 4 (1786) 16?

Paspalum sanguinale Lam. var. *ciliare* Hook. f. Fl. Brit. Ind. 7 (1897) 15.

Northern Luzon to Mindanao, *Merrill* 40, 330, *Phil. Pl.* 147, *B. S.* 7661 *Ramos*, 2249 *Mearns*. In open waste places. Pantropic.

Var. **AUSTRALIS** (Spreng.).

? *Panicum australe* Spreng. Syst. 1 (1825) 309.

Panicum sanguinale Linn. var. *australe* Thwaites Enum. Pl. Zeyl. (1864) 358.

? *Panicum striatum* R. Br. Prodr. (1810) 192, non Lam.

Throughout the Philippines at low and medium altitudes, in thickets, waste places, etc., *Merrill* 529, 89, 5359, 7259, 9787, *Phil. Pl.* 578, *Elmer* 5855, 15147. India through Malaya to Australia.

This identification follows Hackel, but I am by no means certain that our form is the same as the one originally described by R. Brown.

DOUBTFUL AND EXCLUDED SPECIES

PANICUM ELYTHROBLEPHARUM Steud.; F.-Vill. Novis. App. (1882) 311.

PANICUM STIPATUM Presl Rel. Haenk. 1 (1828) 297; F.-Vill. Novis. App. (1882) 311.

"Hab. in Mexico, Luzon?" Presl. This is the American *Digitaria setosa* Desv. fide Scribner in Rept. Mo. Bot. Gard. 10 (1899) 49.

PASPALUM FILIFORME Sw.; Presl Rel. Haenk. 1 (1828) 297; F.-Vill. Novis. App. (1882) 310.

"Hab. in Luzonia ? Mexico" Presl. This is the American *Digitaria filiforme* Delile fide Scribner in Rept. Mo. Bot. Gard. 10 (1899) 49.

PASPALUM MOLLE Presl Rel. Haenk. 1 (1828) 213 = *Paspalum mollicomum* Kunth Enum. 1 (1833) 310; Miq. Fl. Ind. Bat. 3 (1857) 433; F.-Vill. Novis. App. (1882) 310 = *Syntherisma molle* Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 50.

"Hab. in Luzonia" Presl. This was considered by Scribner to be a valid species, but Presl's description does not apply to any Philippine form known to me and I suspect that the type was an American, not a Philippine plant.

PASPALUM FUSCUM Presl Rel. Haenk. 1 (1828) 214; Kunth Enum. 1 (1833) 46; F.-Vill. Novis. App. (1882) 310 = *Syntherisma fusca* Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 49, t. 11.

"Hab. in Luzonia ? in Peruviae montanis huanoccensibus ? Mexico?" Presl. From the figure given by Scribner, drawn from a cotype of Presl's species, this does not appear to represent any known Philippine species. The type was probably from America.

25. PASPALUM Linnaeus

PASPALUM CONJUGATUM Berg. in Act. Helvet. 7 (1772) 129, t. 8; Mez in Perk. Frag. Fl. Philip. (1904) 141; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 28, Philip. Journ. Sci. 1 (1906) Suppl. 345, Fl. Manila (1912) 87.

Throughout the Philippines in the settled areas, about towns, along trails, streams, etc., often very abundant. A native of tropical America, now pantropic.

Local names: Bantótan (Mbo.); kauatkáuat (C. Bis.); laau-láau (Tag.).

PASPALUM LONGIFOLIUM Roxb. Hort. Beng. (1814) 7, *nomen nudum*, Fl. Ind. 1 (1820) 283; Mez in Perk. Frag. Fl. Philip. (1904) 141; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 345, Fl. Manila (1912) 86.

Paspalum flexuosum Klein in Presl Rel. Haenk. 1 (1828) 216; Kunth Enum. 1 (1833) 54; Miq. Fl. Ind. Bat. 3 (1857) 433; F.-Vill. Novis. App. (1882) 310.

Luzon (Cagayan, Benguet, Bontoc, Nueva Vizcaya, Bulacan, Rizal, Laguna, Albay), Polillo, Palawan, Bohol, Mindanao, *Weber 1063, Merrill 9303, Phil. Pl. 1956, B. S. 26002, 26675 Fénix, 21684, 14512, 29504, 10979 Ramos*. In open, usually rather wet lands at low and medium altitudes. India to Malaya.

Local name: Taltal-likód (Ilk.).

Var. TRICHOCOLEUM Hack. in Philip. Journ. Sci. 3 (1908) Bot. 167.

Mindoro, Mindanao (Surigao), *B. S. 941 Mangubat, Merrill 5455*. In open level grasslands at low altitudes. Endemic.

PASPALUM SCROBICULATUM Linn. Mant. 1 (1767) 29; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; F.-Vill. Novis. App. (1882) 310; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 286; Mez in Perk. Frag. Fl. Philip. (1904) 141; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 344, Fl. Manila (1912) 86, Sp. Blancoanae (1918) 63.

Paspalum cartilagineum Presl Rel. Haenk. 1 (1828) 216; Miq. Fl. Ind. Bat. 3 (1857) 432; F.-Vill. Novis. App. (1882) 310; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 49.

Paspalum kora Willd. Sp. Pl. 1 (1797) 332; Presl Rel. Haenk. 1 (1828) 216.

Paspalum villosum Blanco Fl. Filip. (1837) 40, ed. 2 (1845) 28, ed. 3, 1 (1877) 53, non Thunb.

Paspalum sumatrense Roth Nov. Pl. Sp. (1821) 35; Llanos Frag. Pl. Filip. (1851) 22; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 15; F.-Vill. Novis. App. (1882) 310.

Paspalum thunbergii Kunth ex Steud. Syn. Pl. Glum. 1 (1854) 28;
Mez in Perk. Frag. Fl. Philip. (1904) 141.

Throughout the Philippines, in open grasslands at low and medium altitudes, ascending to 1,500 m; common and somewhat variable. Pantropic.

Local names: Angangsúg (Ig.); bias-biásan (S. L. Bis.); bubulis' (Sub.); paragis (Tag.); sabuñg-sabuñgan (Tag.).

Var. PHILIPPINENSE Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 345.

Luzon (Nueva Vizcaya, Rizal), *Merrill 140a*, *B. S. 1449 Ramos*. In open, rather wet grasslands at medium altitudes. Endemic.

Var. BISPICATUM Hack. ex Merr. Fl. Manila (1912) 86.

Luzon (Ilocos Norte, Ilocos Sur, Benguet, Rizal, Bataan), Panay, *Merrill Phil. Pl. 576*, *B. S. 10083*, *32275 McGregor*, *27378 Ramos*. In open grasslands at low and medium altitudes. Endemic.

Var. AURICULATUM (Presl) Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 345, Fl. Manila (1912) 86.

Paspalum auriculatum Presl Rel. Haenk. 1 (1828) 217; Kunth Enum. 1 (1833) 54; Miq. Fl. Ind. Bat. 3 (1857) 432; F.-Vill. Novis. App. (1882) 310.

Luzon (Rizal), Palawan, Samar, Leyte, Mindanao, *Merrill 5361*, *7292*, *820*, *Phil. Pl. 153*, *B. S. 843 Foxworthy*, *24435 Ramos*. In open rather wet grasslands at low altitudes. Tropical Asia and Africa.

PASPALUM VAGINATUM Sw. Prodr. Veg. Ind. Occ. (1788) 21; Stapf in Prain Fl. Trop. Afr. 9 (1919) 570.

Paspalum distichum F.-Vill. Novis. App. (1882) 310; Usteri Beitr. Ken. Veg. Philip. (1905) 133; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 346, Fl. Manila (1912) 87, non Linn.

Luzon (Amburayan, Union, Pampanga, Rizal, Laguna), Panay, Samar, *Elmer 18140*, *Merrill 4253*, *5229*, *7116*. In open wet lands, usually in places subject to the influence of salt or brackish water. Pantropic.

Local names: Pagetpet (Ilk.); panluilui (Pamp.).

EXCLUDED SPECIES

PASPALUM ELEGANS Flügge; F.-Vill. Novis. App. (1882) 310.

PASPALUM CHRYSOTRICHUM Presl Rel. Haenk. 1 (1828) 211; Kunth Enum. 1 (1833) 64; Miq. Fl. Ind. Bat. 3 (1857) 431; F.-Vill. Novis. App. (1882) 310.

"Hab. in Luzonia." The localization is unquestionably erroneous. Presl's species is probably the same as *Paspalum aureum* HBK. of tropical America.

PASPALUM BORYANUM Presl Rel. Haenk. 1 (1828) 209; Kunth Enum. 1 (1833) 52; Miq. Fl. Ind. Bat. 3 (1857) 432; F.-Vill. Novis. App. (1882) 310.

"Hab. ad Sorzogon" Presl. This may prove to be a *Digitaria*, or again it may be an American species of *Paspalum*.

PASPALUM FILIFORME Sw.; F.-Vill. Novis. App. (1882) 310.

26. **ERIOCHLOA** Kunth

ERIOCHLOA RAMOSA (Retz.) O. Kuntze Rev. Gen. Pl. (1891) 775; Hack. in Bull. Acad. Int. Bot. 16 (1906) 19; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 348, Fl. Manila (1912) 85, Sp. Blancoanae (1918) 64.

Milium ramosum Retz. Obs. 6 (1791) 22.

Paspalus annulatus Flügge Monogr. (1810), 133; Trin. Gram. Pan. (1826) 117, Sp. Gram. 2 (1829) t. 133.

Eriochloa annulata Kunth Enum. 1 (1833) 73; F.-Vill. Novis. App. (1882) 311; Perk. Frag. Fl. Philip. (1904) 141.

Piptantherum annulatum Presl Rel. Haenk. 1 (1828) 221.

Helopus laevis Trin. in Spreng. Neue Entdeck. 2 (1821) 49; Miq. Fl. Ind. Bat. 3 (1857) 434.

Eriochloa punctata F.-Vill. Novis. App. (1882) 312, non Ham.

Milium zonatum Llanos Frag. Pl. Filip. (1851) 24; F.-Vill. and Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 16.

Throughout the Philippines at low and medium altitudes, in open usually damp grasslands at low and medium altitudes. Pantropic.

Var. **INVOLUCRATA** Hack. ex Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 349.

Mindoro, Bohol, *Merrill 3333*, *B. S. 1247* *McGregor*. Habitat of the species and scarcely distinguishable from it. Endemic.

27. **ISACHNE** R. Brown

ISACHNE BENECKEI Hack. in Oesterr. Bot. Zeitschr. 51 (1901) 459, Govt. Lab. Publ. (Philip.) 35 (1906) 79; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 350, 5 (1910) Bot. 327.

Luzon (Ifugao, Bontoc, Lepanto, Benguet, Bataan, Pampanga), *Merrill 4680, 5019, 3201, 4544, 4619, 4523, 4522*, *Phil. Pl. 1467*, *B. S. 19690, 1979*, *McGregor, 37736* *Ramos & Edaña, Williams 1193*. In open wet places, borders of cold pools and sphagnum swamps, and on narrow forested ridges, altitude 800 to 2,400 m. Java.

ISACHNE CONFERTA Merr. in Philip. Journ. Sci. 9 (1914) Bot. 261.

Luzon (Laguna), *B. S. 14914* *Ramos*. On boulders and ledges along streams at low or medium altitudes. Endemic.

ISACHNE DEBILIS Rendle ex Forbes & Hemsl. in Journ. Linn. Soc. Bot. 36 (1904) 322; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 268; Merr. l. c. 350.

Isachne monticola Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 79, Philip. Journ. Sci. 1 (1906) Suppl. 268, non Büse.

Luzon (Abra, Ifugao, Bontoc, Benguet, Nueva Vizcaya, Nueva Ecija, Bataan, Laguna), Mindanao (Davao), *Merrill 5315, 9680, 7675*, *Phil. Pl. 599*, *Elmer 5821, 18367*. Along streams in damp ravines and on damp forested ridges, altitude 1,000 to 2,000 m. Formosa.

This seems to be close to *I. myosotis* Nees.

ISACHNE DEPAUPERATA (Hack.) comb. nov.

Isachne beneckeii Hack. forma *depauperata* Hack. ex Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 350.

Luzon (Lepanto, Benguet), Negros, *Merrill 4545, 4489, 6978, 6504, B. S. 4249 Mearns*. In the mossy forest and in damp thickets, altitude 1,700 to 2,200 m. Endemic.

ISACHNE GLOBOSA (Thunb.) O. Kuntze Rev. Gen. Pl. (1891) 778.

Milium globosum Thunb. Fl. Jap. (1784) 49.

Isachne australis R. Br. Prodr. (1812) 196; F.-Vill. Novis. App. (1882) 321.

Mindanao (Lanao), *Clemens 870, 1114*. In swamps about Lake Lanao, altitude about 650 m. India to Japan south to Australia.

ISACHNE INCRASSATA (Hack.) Merr. in Philip. Journ. Sci. 5 (1910) Bot. 168.

Isachne debilis Rendle var. *incrassata* Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 268; Merr. op. cit. 350.

Negros, Mindanao (Davao), *Merrill 6977, De Vore & Hoover 358*. In crevices of ledges along mountain streams, altitude 1,200 to 1,500 m. Endemic.

ISACHNE MAGNA (Merr.) Merr. in Philip. Journ. Sci. 5 (1910) Bot. 327.

Isachne beneckeii Hack. var. *magna* Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 350.

Isachne apoensis Elm. Leaf. Philip. Bot. 7 (1915) 2676.

Luzon (Ifugao, Bontoc, Benguet), Mindanao (Davao), *Merrill 4541, 7672, 7678, 6569, Elmer 11578, B. S. 31686 Santos, 8870, 19731 McGregor, 4483, 4248 Mearns*. In the mossy forest, altitude 1,800 to 2,400 m. Endemic.

Local names: Buyo (Bag.); popoktót (Bon.).

ISACHNE MICRANTHA Merr. in Philip. Journ. Sci. 5 (1910) Bot. 168.

Luzon (Cagayan, Tayabas), *F. B. 16837, 16841 Curran, B. S. 28708 Ramos & Edaña*. On siopes, altitude 300 m and above. Endemic.

Closely allied to *I. myosotis* Nees.

ISACHNE MILIACEA Roth Nov. Pl. Sp. (1821) 58; Merr. Fl. Manila (1912) 89.

Panicum minutulum Gaudich. in Freyc. Voy. (1826) Bot. 410.

Isachne minutula Kunth Rev. Gram. 2 (1829) t. 117, Enum. 1 (1833) 137; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 172; F.-Vill. Novis. App. (1882) 321; Mez in Perk. Frag. Fl. Philip. (1904) 141; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 349.

Isachne pulchella Mez in Perk. Frag. Fl. Philip. (1904) 141 p. p. non Roth.

Panicum macilentum Presl Rel. Haenk. 1 (1828) 310; Kunth Enum. 1 (1833) 116; F.-Vill. Novis. App. (1882) 312.

Throughout the Philippines, in low open wet places, old rice paddies, etc., often abundant; *Kneucker Gram. Exsic. 609* from Manila well represents the species. India to Malaya and Polynesia.

Local names: Lagitlít (C. Bis.); matáng-lamók (Pang.); tagik-tagik (C. Bis.).

ISACHNE MYOSOTIS Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 98; Walp. Ann. 3 (1853) 720; Miq. Fl. Ind. Bat. 3 (1857) 462; F.-Vill. Novis. App. (1882) 321; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 286; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 349, 2 (1907) Bot. 261, 5 (1910) Bot. 327.

Panicum myosotis Steud. Syn. Pl. Glum. 1 (1854) 96.

Isachne pauciflora Hack. var. *hirsuta* Hack. in Philip. Journ. Sci. 3 (1908) Bot. 167.

Luzon (Abra, Benguet, Rizal, Pampanga), Negros, Mindoro, Mindanao (Zamboanga), *B. S.* 1765, 13768, 27029 *Ramos*, 29677 *Ramos & Edaño*, 1921 *Foxworthy*, *F. B.* 4405 *Merritt*, *Merrill* 6167, *Copeland* 162. In open places at medium and higher altitudes, ascending to 2,400 m. Reported also from Australia.

ISACHNE PANGERANGENSIS Z. & M. Syst. Verz. Zoll. Pflanz. (1845) 102, var. **HALCONENSIS** Hack. in Philip. Journ. Sci. 3 (1908) Bot. 167, Merr. op. cit. 5 (1910) Bot. 327.

Luzon (Benguet), Mindoro, *Merrill* 6221, 6203, *F. B.* 16182 *Curran*, *Merritt*, & *Zschokke*. In open heaths, altitude 2,400 to 2,600 m. The variety endemic, the species in Java.

ISACHNE PAUCIFLORA Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 80; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 350, 5 (1910) Bot. 327.

Luzon (Benguet, Nueva Vizcaya), *Elmer* 6486, *Merrill* 4292, 9770, 7673, 7674, *Phil. Pl.* 195, *B. S.* 31753 *Santos*, 14211 *McGregor*. In wet places, along streams, in rice paddies, etc., altitude 1,400 to 2,300 m. Endemic.

ISACHNE STRICTA Elm. Leaf. Philip. Bot. 2 (1908) 463.

Leyte, Negros, Mindanao (Lanao), *Elmer* 10425, *Wenzel* 781, *Clemens* 1135a. In crevices of ledges, on seepage slopes, etc., altitude 600 to 1,200 m. Endemic.

ISACHNE VULCANICA Merr. in Philip. Journ. Sci. 5 (1910) Bot. 169.

Negros, *Merrill* 6975, *Phil. Pl.* 585. On bare slopes of the old crater of Canlaon Volcano, altitude 1,800 to 2,100 m. Endemic.

EXCLUDED SPECIES

ISACHNE JAVANA Nees.

This species was credited to the Philippines by Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97, on the basis of *Cuming* 2288; this specimen was from Malacca, not from the Philippines. The species has not been found in the Archipelago.

ISACHNE PULCHELLA Roth.

This species has been credited to the Philippines by F.-Villar, Novis. App. (1882) 321, and by Mez ex Perk. Frag. Fl. Philip. (1904) 141, but one of the specimens cited by the latter is *Isachne miliacea* Roth, and another is *I. debilis* Rendle. Roth's species is *Sphaerocaryum pulchellum* (Roth) Merr. (*S. elegans* Nees) which has not been found in the Philippines.

28. **PANICUM** Linnaeus

PANICUM AMBIGUUM Trin. in Mém. Acad. St. Pétersb. VI 3² (1835) 243; Steud. Syn. Pl. Glum. 1 (1854) 61; Miq. Fl. Ind. Bat. 3 (1857) 447; F.-Vill. Novis. App. (1882) 311; Mez in Perk. Frag. Fl. Philip. (1904) 141; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 355.

Urochloa paspaloides Presl Rel. Haenk. 1 (1830) 318; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 54, t. 14, non Beauv.

Northern Luzon (Cagayan) to Mindanao and Palawan, *Merrill 9590*, *Phil. Pl. 580, 3332, 5369, B. S. 14511, 27128*. *Ramos, 31536 Ramos & Edaño*. Widely scattered in open more or less waste places, old clearings, etc., at low and medium altitudes. Burma to the Mascarene Islands, Malaya and Polynesia.

PANICUM AMPLEXICAULE Rudge Pl. Guian. (1805) 21, t. 27; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 356, Fl. Manila (1912) 92, Sp. Blancoanae (1918) 66.

Panicum myuros Kunth Enum. 1 (1833) 86; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27; Usteri Beitr. Ken. Philip. Veg. (1905) 133, non Lam.

Panicum polygonatum Llanos Frag. Pl. Filip. (1851) 41; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 30; non Kunth.

Panicum acutiglumum Steud. Syn. Pl. Glum. 1 (1854) 66.

Hymenachne serrulata Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 98, *nomen nudum*.

Central Luzon to Palawan and Mindanao, *Merrill 9468, Phil. Pl. 157, Sp. Blancoanae 967, Elmer 14561, F. B. 8881 Curran*. In open swamps and along swampy margins of lakes and streams at low altitudes. Pantropic. Local name: Lagtóm (Bik.).

PANICUM AURITUM Presl Rel. Haenk. 1 (1830) 305 (incl. var. *procerus* Presl); Kunth Enum. 1 (1833) 113; Steud. Syn. Pl. Glum. 1 (1854) 70; Miq. Fl. Ind. Bat. 3 (1857) 456; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 172, Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; F.-Vill. Novis. App. (1882) 312; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 46, t. 15; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 356, Fl. Manila (1912) 92.

Northern Luzon (Cagayan) to Palawan and Mindanao, in most or all islands and provinces. Common in open, rather wet places at low and medium altitudes. India to southern China and Malaya.

Local name: Mongoto (Sub.).

PANICUM BREVIFOLIUM Linn. Sp. Pl. (1753) 59.

Panicum ovalifolium Poir. in Lam. Encycl. Suppl. 4 (1816) 279; Hook. f. Fl. Brit. Ind. 7 (1897) 44.

Balabac, *Merrill 5386, Phil. Pl. 110, B. S. 442 Mangubat*. In thickets, along trails in shaded places, etc., at low altitudes. Tropical Africa and Asia to China and Malaya.

PANICUM CARINATUM Presl Rel. Haenk. 1 (1830) 309; Kunth Enum. 1 (1833) 112; Miq. Fl. Ind. Bat. 3 (1857) 452; F.-Vill. Novis. App. (1882) 312; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 46, t. 17; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 362.

Luzon (Ilocos Norte, Pangasinan, Rizal, Laguna, Batangas, Bataan, Camarines), Mindoro, Palawan, Culion, Balabac, Panay, Mindanao, Basilan, *Presl!* in Herb. Bernhardi, B. S. 30023 *Fénix*, 29483 *Ramos & Edaño*, 16887 *Serviñas*, Merrill 477, 10596, 9241, *Phil. Pl.* 178. In old clearings, thickets, etc., at low and medium altitudes. Borneo and probably other parts of Malaya.

Perhaps better considered as a variety of *P. patens* Linn.

Local names: Baile (Sul.); gamog-gamog (Sub.); hirabon (Sul.); ka-uang-káuang (Sul.); malakauáyan (Tag., P. Bis.); tuad-tuáran (Tag.).

PANICUM CAUDIGLUME Hack. in Oesterr. Bot. Zeitschr. 51 (1901) 428, Govt. Lab. Publ. (Philip.) 35 (1906) 80; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27, 360, Fl. Manila (1912) 92.

Luzon (Ilocos Sur, Bataan), Culion, Palawan, Merrill 832, 3307, 7436, 11607, *Kneucker Gram. Exsic.* 731, B. S. 29240, 24083, 41285 *Ramos*. In open waste places at low altitudes, apparently introduced. Java.

Local name: Kumut-palakâ (Tag.).

PANICUM COCCOSPERMUM Steud. Syn. Pl. Glum. 1 (1854) 62.

Panicum villosum Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 269; Merr. op. cit. 355, 5 (1910) Bot. 327, non Lam.

Luzon (Bontoc, Benguet, Nueva Vizcaya, Rizal), Culion, Mindanao (Lanao), Merrill 7692, 4281, 4396, 4459, 4360, 11606, *Phil. Pl.* 591, B. S. 5918, 8214, 41339 *Ramos*, 4492 *Mearns*. On open grassy slopes and banks chiefly at medium altitudes, ascending to 1,500 m. The identification of our material with Steudel's species was made by Mr. C. A. Backer of Buitenzorg, Java. India to Java.

Local name: Tumi (Bon.).

PANICUM COLONUM Linn. Syst. ed. 10 (1759) 870; Steud. Syn. Pl. Glum. 1 (1854) 46; F.-Vill. Novis. App. (1882) 311; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 286; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 354; Fl. Manila (1912) 91, Sp. Blancoanae (1918) 65.

Panicum colonum Linn. var. *pseudocolonum* Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 172, Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Panicum cumingianum Steud. Syn. Pl. Glum. 1 (1854) 58; Miq. Fl. Ind. Bat. 3 (1857) 447; F.-Vill. Novis. App. (1882) 311.

Echinochloa colona Link Hort. Berol. 2 (1833) 209.

Oplismenus colonus HBK. Nov. Gen. Sp. Pl. 1 (1815) 108; Presl Rel. Haenk. 1 (1830) 321.

Orthopogon dichotomus Llanos Frag. Pl. Filip. (1851) 38; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 28.

Throughout the Philippines at low and medium altitudes. In open, rather wet places. Pantropic.

Local names: Búlang (Tag.); dakayáng (Ilk.); dukayáng (Ilk.); tiri-búhan (Tag.); tumi (Bon.).

PANICUM CORDATUM Büse in Miq. Pl. Jungh. (1854) 376; Merr. in Philip. Journ. Sci. 4 (1909) Bot. 249.

Panicum luxurians Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 359, non Willd.

Panicum montanum Naves Novis. App. (1882) 312; Ceron Cat. Pl. Herb. Manila (1892) 180; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27, non Roxb.

Luzon (Bontoc, Union, Bataan, Bulacan), Culion, Palawan, Merrill 9348, 9318, 554, Phil. Pl. 146, Elmer 5595, 6735, Williams 134. In thickets, old clearings, etc., at low and medium altitudes. Java, Borneo.

PANICUM CRASSIAPICULATUM Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 356.

Panicum latifolium Hook. f. Fl. Brit. Ind. 7 (1897) 39, excl. syn. et var. *majus* Hook. f., non Linn.

Palawan, Balabac, Mindanao (Agusan), Merrill 5387, 7290, 9240, 9286, Phil. Pl. 111. In thickets and along trails in forests at low altitudes. Malay Peninsula, Java.

Closely allied to the American *P. oryzoides* Sw.

PANICUM CRUS-GALLI Linn. Sp. Pl. (1753) 56; F.-Vill. Novis. App. (1882) 311; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 354, 5 (1910) Bot. 327, Fl. Manila (1912) 91, Sp. Blancoanae (1918) 65.

Oplismenus limosus Presl Rel. Haenk. 1 (1830) 321; Kunth Enum. 1 (1833) 144.

Panicum limosum F.-Vill. Novis. App. (1882) 311.

Panicum hispidulum Lam. Ill. 1 (1791) 171; F.-Vill. Novis. App. (1882) 311.

Echinochloa crus-galli Beauv. Agrost. (1812) 53.

Orthopogon hispidus Spreng. Syst. 1 (1825) 307; Llanos Frag. Pl. Filip. (1851) 37; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 27.

Orthopogon subverticillatus Llanos op. cit. 38; F.-Vill. & Naves op. cit. 28.

Throughout the Philippines at low and medium altitudes, in open wet places, rice paddies, etc. All warm countries.

Local names: Baobao (If.); daua-dáua (Tag.); daua-dauáhan (Tag.); dauána (Mbo.); lagtom (Bik.); sabsablúg (Bon.); sabsábug (Bon.).

Var. **MUTICUM** Doell.; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 268; Merr. op. cit. 354.

Luzon (Bontoc, Benguet, Rizal), Balabac, Palmas, Merrill 5363, 4307, B. S. 460 Mangubat, 5975 Ramos. Habitat of the species. Pantropic.

PANICUM DISTACHYUM Linn. Mant. 1 (1767) 138; F.-Vill. Novis App. (1882) 311; Merr. Fl. Manila (1912) 91, Sp. Blancoanae (1918) 66.

Panicum remotum Retz. Obs. 4 (1786) 71; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 354.

Panicum petiveri Trin. Ic. Gram. t. 176, f. C.

Panicum ramosum Mez in Perk. Frag. Fl. Philip. (1904) 143, non (?) Linn.

? *Panicum umbrosum* F.-Vill. Novis. App. (1882) 311, non Retz.

Panicum miliiforme Presl Rel. Haenk. 1 (1830) 300; Kunth Enum. 1 (1833) 96; Miq. Fl. Ind. Bat. 3 (1857) 488; F.-Vill. Novis. App. (1882) 312; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 47, t. 20.

Northern Luzon to Palawan and Mindanao, in most or all islands and provinces. In open waste places in and about towns at low and medium altitudes; common. India to China through Malaya to tropical Australia.

PANICUM FLAVIDUM Retz. Obs. 4 (1786) 15; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; F.-Vill. Novis. App. (1882) 311; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 286; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27, 353, Fl. Manila (1912) 90.

Panicum floridum Usteri Beitr. Ken. Philip. Veg. (1905) 133, sphalm.

Northern Luzon to Palawan and Mindanao. A common weed in open waste places in and about towns at low altitudes. Tropical Asia, Africa, and Malaya.

Local names: Gíling (C. Bis.); sabuñg-sabuñgan (Tag.); bali itau (Sul.).

PANICUM HUMILE Nees ex Steud. Syn. Pl. Glum. 1 (1854) 84; Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 80; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 358.

Luzon (Bontoc, Lepanto, Tarlac, Pangasinan, Zambales, Rizal), Culion, Merrill 3623, 4466, 11608, B. S. 21998 Ramos, F. B. 5872 Curran. In open waste places, grasslands, etc., at low and medium altitudes. India to southern China and Malaya.

PANICUM INDICUM Linn. Mant. 1 (1767) 184; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; F.-Vill. Novis. App. (1882) 311; Vid. Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27, 357, Fl. Manila (1912) 92.

Hymenachne indica Büse ex Miq. Fl. Ind. Bat. 3 (1857) 458.

Sacciolepis indica Chase in Proc. Biol. Soc. Wash. 21 (1908) 8.

Northern Luzon to Palawan and Mindanao. Common in open wet grasslands at low and medium altitudes. India through Malaya to tropical Australia.

Local names: Buntót-pusá (Tag.); sabsablúg (Bon.); salai-máya (Tag.); sangumai (Tag.); sikuán (Ig.).

Var. **ANGUSTUM** (Trin.) Hook. f. Fl. Brit. Ind. 7 (1897) 42; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 357.

Panicum angustum Trin. Ic. Gram. 3 (1836) t. 334; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287.

Occasional, in the same habitat as the species. India.

PANICUM LUZONENSE Presl Rel. Haenk. 1 (1828) 308; Kunth Enum. 1 (1833) 121; Miq. Fl. Ind. Bat. 3 (1857) 457; F.-Vill. Novis. App. (1882) 312; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 358.

Luzon (Amburayan, Nueva Ecija, Bataan, Rizal, Cavite, Laguna), Palawan, Merrill 316, 3107, 4183, Phil. Pl. 1238, B. S. 4600, 21997 Ramos,

23060 *McGregor*. In open waste places, old clearings, etc., at low altitudes. Malay Peninsula.

Local name: Niknikan (Tag.).

PANICUM MALABARICUM (Linn.) Merr. in Philip. Journ. Sci. 4 (1909) Bot. 248.

Poa malabarica Linn. Sp. Pl. (1753) 69.

Panicum arnottianum Nees in Steud. Syn. Pl. Glum. 1 (1854) 59.

Mindoro, Palawan, Mindanao, Palmas, *F. B. 5349 Merritt, Merrill 5367, 9378, 9581, B. S. 36687 Ramos & Edaño*. In old clearings at low altitudes. India to Malaya.

Local name: Salid (Sub.).

PANICUM MAXIMUM Jacq. Ic. Pl. Rar. 1 (1781-86) t. 13; Merr. Fl. Manila (1912) 93.

Cultivated for forage purposes, occasional as an escape or persisting from former cultivation. Cultivated in all tropical countries. Guinea grass.

PANICUM MILIACEUM Linn. Sp. Pl. (1753) 58; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 358.

Very rarely cultivated in the Philippines, and nowhere spontaneous. Cultivated in most warm countries. Millet.

Local name: Kabug (C. Bis.).

PANICUM MINDANAENSE Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 360.

Luzon (Ilocos Sur, Rizal, Batangas, Zambales), Mindanao (Lanao), *Clemens 99, B. S. 27379, 5103, 29242 Ramos, Merrill Phil. Pl. 168, as P. caudiglume Hack*. In open places at low and medium altitudes. Endemic.

PANICUM MYOSUROIDES R. Br. Prodr. (1810) 189; Kunth Enum. 1 (1833) 77; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 357.

Panicum angustissimum Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 286, non ? Hochst.

Panicum contractum Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97, non ? Trin.

Luzon (Abra, Nueva Vizcaya), Panay, Mindanao (Misamis), *Cuming 1668, B. S. 31283 Ramos & Edaño, 14221 McGregor, F. B. 16459 Bacani*. In open wet lands at low and medium altitudes. Tropical Africa and Asia through Malaya to Australia.

PANICUM NODOSUM Kunth Enum. 1 (1833) 97; Miq. Fl. Ind. Bat. 3 (1857) 448; F.-Vill. Novis. App. (1882) 312; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 357, Fl. Manila (1912) 92, Sp. Blancoanae (1918) 66.

Panicum multinode Presl Rel. Haenk. 1 (1828) 303, non Lam.

Panicum violaceum Llanos Frag. Pl. Filip. (1851) 42; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 31.

Panicum ouonbiense Mez in Perk. Frag. Fl. Philip. (1904) 142, non Balansa.

Northern Luzon to Palawan and Mindanao, in most islands and provinces, Merrill 4228, 527, 7056, 127, 4182, 255, 5385, 5247, Phil. Pl. 113, Sp. Blancoanae 944. In thickets, old clearings, etc., at low and medium altitudes; common. India to China and Malaya.

Local names: Baníg-usá (Tag.); kauakauayánan (Tag.); kúmut-usá (Tag.).

PANICUM PALUDOSUM Roxb. Fl. Ind. 1 (1820) 310; Usteri Beitr. Ken. Philip. Veg. (1905) 133; Merr. in Philip. Journ. Sci. 4 (1909) Bot. 249, Fl. Manila (1912) 93.

Panicum proliferum Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 359, non Lam.

Luzon (Cagayan, Rizal, Laguna), Mindanao (Lanao), Merrill 97, Phil. Pl. 159, B. S. 23144 McGregor, 7429 Ramos. In shallow water along margins of streams and lakes at low and medium altitudes. India to southern China and Malaya.

PANICUM PATENS Linn. Sp. Pl. (1753) 86; Mez in Perk. Frag. Fl. Philip. (1904) 142; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 361, 3 (1908) Bot. 396.

Panicum radicans Retz. Obs. 4 (1786) 18; Steud. Syn. Pl. Glum. 1 (1854) 87; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Miq. Fl. Ind. Bat. 3 (1857) 453; F.-Vill. Novis. App. (1882) 312; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287; Mez in Perk. Frag. Fl. Philip. (1904) 143.

Northern Luzon to Palawan and Mindanao, in most or all islands and provinces. Common in thickets, some secondary forests, old clearings, etc., at low and medium altitudes. India to Malaya.

Local names: Ililit (Ig.); malaniknik (Tag.); púgad-lagúyo (Tag.).

PANICUM PILIPES Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Büse in Miq. Pl. Jungh. (1854) 376; Miq. Fl. Ind. Bat. 3 (1857) 453; Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27, 361, 3 (1908) Bot. 396, Fl. Manila (1912) 94.

Panicum hermaphroditum Steud. Syn. Pl. Glum. 1 (1854) 67; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287.

Panicum trigonum Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 172; F.-Vill. Novis. App. (1882) 312, non Retz.

Northern Luzon to Palawan and Mindanao, in most or all islands and provinces. In thickets, along streams, in old clearings, etc., at low and medium altitudes. India to the Mascarene Islands, Malaya, tropical Australia, and Polynesia.

Local names: Baitbút (Tagb.); baníg-usá (Tag.); baririt-pualas (Bag.); marikauáyan (Ig.).

PANICUM PSILOPODIUM Trin. Gram. Pan. (1826) 217; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 168, Fl. Manila (1912) 93, Sp. Blancoanae (1918) 66.

Panicum crispum Llanos Frag. Pl. Filip. (1851) 42; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 30.

Luzon (Benguet, Rizal), *Merrill 7079, Phil. Pl. 597, 1764, Elmer 5972, 6589*. In open wet places at low and medium altitudes, ascending to 1,400 m; local. India to Burma.

PANICUM PUNCTATUM Burm. f. Fl. Ind. (1768) 26; Hook. f. Fl. Brit. Ind. 7 (1897) 29; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 353, Fl. Manila (1912) 90.

Panicum fluitans Retz. Obs. 3 (1783) 8; Miq. Fl. Ind. Bat. 3 (1857) 455; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287.

Paspalum pluriracemosum Steud. Syn. Pl. Glum. 1 (1854) 27; Miq. Fl. Ind. Bat. 3 (1857) 431; F.-Vill. Novis. App. (1882) 310.

Panicum brizoides Nees in Hook. Journ. Bot. 2 (1850) 97, Miq. Fl. Ind. Bat. 3 (1857) 445, non Linn.

Luzon (Rizal, Laguna), Leyte, Mindanao (Misamis, Davao, Zamboanga, Agusan), *Merrill 7406, 8111, Phil. Pl. 131, Cuming 532, Wenzel 1780, Elmer 12019, B. S. 37364 Ramos & Edaño*. In shallow water and in open muddy places at low altitudes. Northern Africa to Mauritius, India, and Malaya.

Local names: Baririan (Bag.); lalabok (Sub.).

PANICUM REPENS Linn. Sp. Pl. ed. 2 (1762) 87; Hook. f. Fl. Brit. Ind. 7 (1897) 49; Ceron Cat. Pl. Herb. Manila (1892) 180; Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 359, 3 (1908) Bot. 396, Fl. Manila (1912) 93, Sp. Blancoanae (1918) 67.

Panicum ischaemoides Retz. Obs. 4 (1786) 17; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 175; Miq. Fl. Ind. Bat. 3 (1857) 450; F.-Vill. Novis. App. (1882) 312.

Panicum convolutum Beauv. ex Spreng. Syst. 1 (1825) 319; Presl Rel. Haenk. 1 (1828) 304; F.-Vill. Novis. App. (1882) 312; Steud. Syn. 1 (1854) 73.

Panicum tuberosum Llanos Frag. Pl. Filip. (1851) 40; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 29.

Panicum miliare Mez in Perk. Frag. Fl. Philip. (1904) 142, non Lam.

Throughout the Philippines in open low lands, especially near the sea, but also inland locally and ascending to 1,500 m; common. Pantropic. Local names: Kayana (Mbo.); luya-luyahan (Tag.).

PANICUM REPTANS Linn. Syst. ed. 10 (1759) 871; Hitchc. in Contr. U. S. Nat. Herb. 12 (1908) 119; Merr. in Philip. Journ. Sci. 4 (1909) Bot. 249.

Panicum prostratum Lam. Ill. 1 (1791) 171; Miq. Fl. Ind. Bat. 3 (1857) 446; F.-Vill. Novis. App. (1882) 311; Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 355.

Panicum calaccanzense Steud. Syn. Pl. Glum. 1 (1854) 65; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287.

Panicum barbatum Lam. Tabl. Encycl. 1 (1791) 171; Presl Rel. Haenk. 1 (1830) 299.

Panicum procumbens Nees Agrost. Bras. (1823) 109, Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Panicum luxurians Willd. ex Nees Agrost. Bras. (1829) 233, non *P. luxurians* Willd. ex Spreng. Syst. 1 (1825) 305.

Throughout the Philippines in open waste places, old clearings, thickets, etc., at low and medium altitudes. Pantropic.

Local names: Babáka-nalabága (Ilk.); marakauáyan (Ilk.).

PANICUM SARMENTOSUM Roxb. Fl. Ind. 1 (1820) 311; F.-Vill. Novis. App. (1882) 312; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287; Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 27, 360, 2 (1907) Bot. 261.

Panicum incomptum Trin. Gram. Pan. (1826) 200, Ic. Gram. (1828-36) t. 223; Kunth Enum. 2 (1833) 112; Miq. Fl. Ind. Bat. 3 (1857) 451; Nees in Mart. Fl. Bras. 2: 207.

Panicum extensum Steud. Syn. Pl. Glum. 1 (1854) 72.

Panicum vacillans Steud. op. cit. 75.

Panicum concinnum Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Walp. Ann. 3 (1853) 719.

Luzon (Bataan, Cavite, Zambales, Tayabas), Mindoro, Culion, Palawan, Balabac, Panay, Negros. In thickets and secondary forests at low and medium altitudes, often common. India to southern China and Malaya.

Local names: Kanubsúban (Tag.); kauakauáyan (Tag.); kauáyan-sauák (P. Bis.); kaykauáyan (P. Bis.); kuayan-kuáyan (Sul.).

PANICUM STAGNINUM Retz. Obs. 4 (1786) 17; F.-Vill. Novis. App. (1882) 311; Usteri Beitr. Ken. Philip. Veg. (1905) 133; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 354, 3 (1908) Bot. 396, Fl. Manila (1912) 91, Sp. Blancoanae (1918) 64.

Echinochloa stagnina Beauv. Agrost. (1812) 57; Miq. Fl. Ind. Bat. 3 (1857) 464.

Aegilops fluviatilis Blanco Fl. Filip. (1837) 47, ed. 2 (1845) 32, ed. 3, 1 (1879) 59.

Orthopogon loliaceus Llanos Frag. Pl. Filip. (1851) 36; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 26, non Spreng.

Panicum crus-galli Linn. var. *stagninum* O. Kuntze Rev. Gen. Pl. (1891) 783.

Babuyan Islands and northern Luzon to Mindanao, in most islands and provinces. In open wet places, swamps, borders of lakes and streams, at low and medium altitudes. India to Africa and Malaya.

Local names: Balili (Tag.); banágo (Sub.); lagtóm-na-pulá (Bik.); timsím (Tag.); urarói (Bik.).

PANICUM TRICHODES Sw. Prodr. Veg. Ind. Occ. (1788) 24; Hitchc. & Chase in Contr. U. S. Nat. Herb. 15 (1910) 129, f. 125.

Panicum capillaceum Lam. Tabl. Encycl. 1 (1791) 173.

Luzon (Ilocos Norte, Pangasinan), Culion, Merrill 458, B. S. 7618 Ramos, F. B. 15530 Merritt & Darling, Gates 5675. In thickets at low and medium altitudes. Similar in appearance to *Panicum brevifolium* Linn., but distinguished by its short first glume. I am indebted to Prof. A. S. Hitchcock for indicating to me the error in the original determinations. Widely distributed in tropical America, probably an introduced species in the Philippines.

PANICUM TRYPPERON Schultes Mant. 2 (1882) 244; Hook. f. Fl. Brit. Ind. 7 (1897) 47; Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 358.

Luzon (Cagayan, Lepanto), Semirara, Mindoro, Palawan, Mindanao (Davao), Merrill 678, 4136, 4478, Phil. Pl. 1281, Copeland 396, F. B. 16585 Curran. In open grasslands at low and medium altitudes. Tropical Africa through India to China and Malaya.

PANICUM TUBERCULATUM Presl Rel. Haenk. 1 (1830) 307; Kunth Enum. 1 (1833) 120; Miq. Fl. Ind. Bat. 3 (1857) 454; F.-Vill. Novis. App. (1882) 312.

Panicum caesium Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 358.

Panicum acroanthum Mez in Perk. Frag. Fl. Philip. (1904) 141, non Steud.

Luzon (Isabela, Nueva Ecija, Pampanga, Rizal, Laguna), Panay, Mindanao (Misamis), Cuming 1667, Merrill 123, 6707, 4229, 1469, Phil. Pl. 2057, B. S. 23081 McGregor, 8122 Ramos. In open waste places, old clearings, etc., at low and medium altitudes. India to Borneo and New Guinea.

PANICUM UNCINATUM Raddi Agrost. Bras. (1823) 41; Trin. Gram. Pan. (1826) 174, Ic. t. 216; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 362.

Mindanao (Davao), Clemens 266. In thickets or open lands, altitude about 650 m. Pantropic.

PANICUM WARBURGII Mez in Perk. Frag. Fl. Philip. (1904) 143; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 362, Fl. Manila (1912) 93.

Panicum patens Linn. var. *parvulum* Warb. l. c. in syn.

Luzon (Ilocos Norte, Ifugao, Lepanto, Nueva Vizcaya, Rizal, Zambales, Bataan, Laguna, Tayabas), Mindanao (Lanao), Merrill 303, Kneucker Gram. Exsic. 734, B. S. 28834 Ramos & Edaño, 27018, 27394 Ramos, 20007 McGregor. In thickets and forests, sea level to 1,200 m altitude; probably but a greatly reduced form of *P. carinatum* Presl, the latter being intermediate between the present species and *P. patens* Linn. Endemic.

EXCLUDED SPECIES

PANICUM BRIZAEFORME Presl Rel. Haenk. 1 (1830) 302; Kunth Enum. 1 (1833) 78; Miq. Fl. Ind. Bat. 3 (1857) 445; F.-Vill. Novis. App. (1882) 311; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 46, t. 16.

Panicum paspaloides Pers. Syn. 1 (1805) 81; Kunth Enum. 1 (1833) 77; F.-Vill. Novis. App. (1882) 311.

Hitchcock & Chase, in Contr. U. S. Nat. Herb. 15 (1910) 30, state that the Haenke specimen of *P. brizaeforme* in the Prague herbarium is labeled "Acapulco, Mexico," and is *Panicum geminatum* Forsk.

PANICUM ELATIUS Kunth; F.-Vill. Novis. App. (1882) 312.

PANICUM PALUDICOLA Nees Agrost. Bras. (1823) 179; Steud. Syn. Pl. Glum. 1 (1854) 65; Miq. Fl. Ind. Bat. 3 (1857) 455; F.-Vill. Novis. App. (1882) 312.

"Brazil . . . ins. Manil, Java" Steudel. This is a Brazilian species, *Panicum palustre* Trin.; see Chase in Proc. Biol. Soc. Wash. 21 (1908) 5, sub *Hymenachne palustris* (Trin.) Chase.

PANICUM MERTENSII Roth; F.-Vill. Novis. App. (1882) 312.

PANICUM LEUCOPHAEUM HBK.; Presl Rel. Haenk. 1 (1830) 299; F.-Vill. Novis. App. (1882) 311.

"Hab. in Mexico, in Luzonia" Presl. An American species to which some Australian forms have been referred. The Luzon record was based on an erroneously localized Malaspina Expedition collection.

PANICUM HELOPUS Trin.; F.-Vill. Novis. App. (1882) 311.

29. ICHNANTHUS Beauvois

ICHNANTHUS VICINUS (F. M. Bail.) comb. nov.

Panicum vicinum F. M. Bail. Syn. Queensl. Fl. Suppl. 3 (1890) 82.

Ichnanthus pallens Munro in Benth. Fl. Hongk. (1861) 414; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 261, 363; Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 80, non *Panicum pallens* Sw.

Panicum paludicolum Miq. Fl. Ind. Bat. 3 (1857) 454; F.-Vill. Novis. App. (1882) 312, non Nees.

Panicum nitens Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 8.

Luzon (Bontoc, Kalinga, Bataan, Laguna, Sorsogon), Catanduanes, Mindoro, Biliran, Mindanao, Merrill 3756, 3221, 5498, 5538, Phil. Pl. 109, Kneucker Gram. Exsic. 740, B. S. 30226, 23608 Ramos, 18502 McGregor, 37475, 38608 Ramos & Edaño. On forested slopes and ridges up to 1,400 m altitude. India to southern China and Formosa through Malaya to tropical Australia.

Local name: Dóot (Ilk.).

30. ALLOTEROPSIS Presl

ALLOTEROPSIS SEMIALATA (R. Br.) Hitchc. in Contr. U. S. Nat. Herb. 12 (1909) 210; Chase in Proc. Biol. Soc. Wash. 24 (1911) 159.

Panicum semialatum R. Br. Prodr. (1810) 192; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287.

Alloteropsis distachya Presl Rel. Haenk. 1 (1830) 344, t. 47; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 37, t. 23.

Holoseetum philippicum Steud. in Flora 33 (1850) 228, nomen, Syn. Pl. Glum. 1 (1854) 118; Miq. Fl. Ind. Bat. 3 (1857) 444.

Arundinella nervosa Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; Steud. Syn. Pl. Glum. 1 (1854) 115; Miq. Fl. Ind. Bat. 3 (1857) 519; F.-Vill. Novis. App. (1882) 318.

Coridochloa semialata Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Panicum philippicum F.-Vill. Novis. App. (1882) 312.

Urochloa semialata Kunth Rev. Gram. 1 (1829) 31; Pax in Perk. Frag. Fl. Philip. (1904) 144.

Axonopus semialatus Hook. f. Fl. Brit. Ind. 7 (1897) 64; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 367.

Luzon (Cagayan, Ilocos Norte, Bontoc, Lepanto, Benguet, Pangasinan, Nueva Vizcaya, Nueva Ecija, Zambales, Laguna). On open grassy slopes

at low and medium altitudes, ascending to 1,800 m. India to South Africa, China, and tropical Australia.

Local name: Timi (Bon.).

31. OPLISMENUS Beauvois

OPLISMENUS BURMANNII (Retz.) Beauv. Agrost. (1812) 54; F.-Vill. Novis. App. (1882) 312; Pilger in Perk. Frag. Fl. Philip. (1904) 144; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 364, Fl. Manila (1912) 94.

Panicum burmannii Retz. Obs. 3 (1783) 10.

Panicum hirtellum Burm. f. Fl. Ind. (1768) 24, t. 12, f. 1, non Linn.

Orthopogon burmannii R. Br. Prodr. (1810) 194; Miq. Fl. Ind. Bat. 3 (1857) 442.

Luzon (Pangasinan, Rizal, Bataan), Mindanao (Zamboanga), Merrill 3290, Phil. Pl. 196, Elmer 6645, Williams 135, B. S. 24084 Ramos. In thickets at low altitudes. Pantropic.

OPLISMENUS COMPOSITUS (Linn.) Beauv. Agrost. (1812) 54; F.-Vill. Novis. App. (1882) 312; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 364, 3 (1908) Bot. 397, Fl. Manila (1912) 94, Sp. Blancoanae (1918) 67.

Panicum compositum Linn. Sp. Pl. (1753) 57.

Panicum loliaceum Lam. Tabl. Encycl. 1 (1790) 170, Encycl. 4 (1798) 743.

Oplismenus loliaceus Beauv. Agrost. (1812) 168; Kunth Enum. 1 (1833) 140; F.-Vill. Novis. App. (1882) 312.

Orthopogon sylvaticus Miq. Fl. Ind. Bat. 3 (1857) 443.

Oplismenus sylvaticus R. & S. Syst. 2 (1817) 481; F.-Vill. Novis. App. (1882) 312.

Orthopogon compositus R. Br. Prodr. (1810) 194; Miq. Fl. Ind. Bat. 3 (1857) 443.

Oplismenus indicus R. & S. Syst. 2 (1817) 484; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288.

Orthopogon setarius Llanos Frag. Pl. Filip. (1851) 35, non Spreng.

Orthopogon hirtellus R. Br.; Llanos op. cit. 37.

Panicum lanceolatum Retz.; F.-Vill. Novis. App. (1882) 311.

Panicum sylvaticum Lam. Encycl. 4 (1798) 733; Steud. Syn. Pl. Glum. 1 (1854) 45.

Throughout the Philippines in all or most islands and provinces, in thickets, old clearings, etc., at low and medium altitudes; often common. Pantropic.

Local names: Balibátong (Sub.); balisibis (Iv.); baníg-usá (Tag.); kaukauáyan (Tag.); litlitun (Bon.); malakauáyan (Pang.); marikauáyan (Ig.); yamog-yamóg (P. Bis.); baillituganalu (Sul.).

Vor. **LASIORHACHIS** Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 81; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 364.

Oplismenus burmannii Mez in Perk. Frag. Fl. Philip. (1904) 144, non Beauv.

Palawan, Merrill 826. In open dry places at low altitudes. Endemic.

OPLISMENUS UNDULATIFOLIUS (Ard.) Beauv. Agrost. (1812) 54; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 364.

Panicum undulatifolium Ard. Anim. Bot. Sp. Alt. (1764) 14, t. 4.

Luzon (Ilocos Norte, Benguet, Bontoc, Ifugao, Union, Bataan, Laguna), Mindoro, Mindanao, B. S. 19708 McGregor, 27397 Ramos, 36693, 37488, 37108 Ramos & Edaña, F. B. 2547 Borden, Williams 1179, 1962, 144, Elmer 6987. In thickets and forests at low and medium altitudes. Europe to tropical Asia and Malaya.

Local names: Lulúpong (Sub.); malanikník (Tag.).

Var. **IMBECILLIS** (R. Br.) Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 81; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 364.

Panicum imbecille Trin. Ic. Gram. (1828-36) 16, t. 191.

Orthopogon imbecillis R. Br. Prodr. (1810) 194.

Oplismenus minus Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 9.

Luzon (Cagayan, Benguet, Bontoc, Rizal, Laguna, Tayabas, Bulacan, Pangasinan, Bataan, Sorsogon), Catanduanes, Leyte, Negros, Mindanao, Merrill 3203, 4511, 5485, 8276, 6976, Kneucker Gram. Exsic. 739, Elmer 17105, B. S. 29923 Fénix, 12281 Foxworthy, 30286, 19381 Ramos, 31692 Santos. Along streams in shaded places and on ridges in the mossy forest, at low and medium altitudes, ascending to 2,400 m. Australia.

32. SETARIA Beauvois

SETARIA GENICULATA (Lam.) Beauv. Agrost. (1812) 51, 178.

Panicum geniculatum Lam. Encycl. 4 (1798) 727 (err. typ. 737).

Panicum flavum Nees in Mart. Fl. Bras. 2 (1829) 180.

Setaria flava Kunth Rev. Gram. 1 (1829) 46; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 365, 5 (1910) Bot. 327, Fl. Manila (1912) 95, Sp. Blancoanae (1918) 68.

Panicum penicillatum Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 173, Hook. Journ. Bot. Kew Miscel. 2 (1850) 97, non Willd.

Panicum chrysanthum Steud. Nomen. ed. 2, 2 (1841) 254, Syn. Pl. Glum. 1 (1854) 50.

Panicum rubiginosum Steud. Syn. Pl. Glum. 1 (1854) 50.

Setaria aurea Hochst. ex A. Br. in Flora 24 (1841) 276.

Setaria pilifera Llanos Frag. Pl. Filip. (1851) 34; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 25, non Desv.

Setaria glauca F.-Vill. Novis. App. (1882) 312; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288, non Beauv.

Chaetochloa glauca Scribn. var. *aurea* W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 223.

Setaria glauca Beauv. var. *aurea* K. Schum. Fl. Deutsch. Schutzgeb. Südsee (1901) 180.

Setaria rubiginosa Miq. Fl. Ind. Bat. 3 (1857) 467; F.-Vill. Novis. App. (1882) 313.

Chaetochloa geniculata Millsp. & Chase in Field Mus. Bot. 3 (1903) 37; Hitchc. in Contr. U. S. Nat. Herb. 22 (1920) 168, f. 41.

Throughout the Philippines in open grasslands at low and medium altitudes, ascending to 1,800 m. Pantropic.

Local names: Buntót-púsa (Tag.); lagilikáran (Bag.); putiñgán (Ig.); sabsablúg (Bon.); samsamáng (Bon.); tirtirígo (Ilk.); daua-ambau (Sul.).

SETARIA ITALICA (Linn.) Beauv. Agrost. (1812) 51; Miq. Fl. Ind. Bat. 3 (1857) 467; F.-Vill. Novis. App. (1882) 312; Vidal. Phan. Cuming. Philip. (1885) 157; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 365, Sp. Blancoanae (1918) 68; Hubbard in Am. Journ. Bot. 2 (1915) 192 (subsp. *stramineo-fructa* Hubb. var. *brunneo-seta* Hubb.).

Panicum italicum Linn. Sp. Pl. (1753) 56; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Panicum miliaceum Blanco Fl. Filip. (1837) 39; ed. 2 (1845) 28; ed. 3, 1 (1877) 52, non Linn.

Panicum comosum Steud. Syn. Pl. Glum. 1 (1854) 53, ex descr.

Setaria comosa Miq. Fl. Ind. Bat. 3 (1857) 468; F.-Vill. Novis. App. (1882) 313.

Chaetochloa italica Scribn. in U. S. Dept. Agr. Div. Agrost. Bull. 4 (1897) 39.

Cultivated here and there throughout the Philippines, occasional as an escape in old clearings. Cultivated in all warm countries. Italian millet.

Local names: Bikákau (Ilk.); boróna (Pamp.); bukákau (Ilk.); dáua (C. Bis., P. Bis., Tag.); rautnokara (Iv.); sábug (Ig.) sammang (Bon.); turai (Sul.).

SETARIA LAXA Merr. in Philip. Journ. Sci. 1 Suppl. (1906) 366, Fl. Manila (1912) 96.

Chamaeraphis gracilis Hack. in Engl. Bot. Jahrb. 6 (1885) 236, non *Setaria gracilis* HBK.

Panicum chamaeraphoides Hack. ex Koord. Exkursionsfl. Java 1 (1911) 135.

Luzon (Rizal, Laguna), Merrill 7440, Phil. Pl. 595, Hallier. In thickets and fallow lands at low altitudes. Java.

SETARIA PALMIFOLIA (Koenig) Stapf in Journ. Linn. Soc. Bot. 42 (1914) 186.

Panicum palmaefolium Koenig in Naturf. 23 (1788) 208; Miq. Fl. Ind. Bat. 3 (1857) 449; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 361, 2 (1907) Bot. 261, 5 (1910) Bot. 327.

Panicum plicatum F.-Vill. Novis. App. (1882) 311; Mez in Perk. Frag. Fl. Philip. (1904) 143, non ? Lam.

Panicum amplissimum Steud. Syn. Pl. Glum. 1 (1854) 54; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 287.

Panicum nepalense Spreng. Syst. 1 (1825) 321; Rolfe in Journ. Bot. 23 (1885) 216.

Panicum neurodes Schultes Syst. Mant. 2 (1824) 321; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 287; Usteri Beitr. Ken. Philip. Veg. (1905) 133.

Panicum lene Steud. Syn. Pl. Glum. 1 (1854) 54.

Setaria lenis Miq. Fl. Ind. Bat. 3 (1857) 468; F.-Vill. Novis. App. (1882) 313.

Chamaeraphis setosa O. Kuntze var. *lenis* O. Kuntze Rev. Gen. Pl. (1891) 769.

Panicum kleinianum Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97.

Chaetochloa palmifolia Hitchc. & Chase in Contr. U. S. Nat. Herb. 22 (1920) 161, f. 38.

Northern Luzon to Palawan and Mindanao, in most or all islands and provinces. Common in thickets, old clearings, abacá plantations, etc., at low and medium altitudes, ascending to 200 m. India to China and Malaya, introduced in tropical America.

Local names: Agusáis (Bik.); asáhas (C. Bis.); baṅgas (Ig.); dumbug (Sul.); hagusáhis (Bik.); lalasa (Ig.); liahon (If.); lias (Bon.); yas (Ig.).

SETARIA VERTICILLATA (Linn.) Beauv. Agrost. (1812) 51; F.-Vill. Novis. App. (1882) 312; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 397.

Panicum verticillatum Linn. Sp. Pl. ed. 2 (1762) 82.

Batan Islands, Luzon (Rizal), *B. S. 4041 Fénix*, Merrill *Phil. Pl. 1406*. In open waste places. Temperate and tropical regions generally.

SETARIA VIRIDIS (Linn.) Beauv. Agrost. (1812) 51; F.-Vill. Novis. App. (1882) 312; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 366, Fl. Manila (1912) 95.

Luzon (Cagayan, Bontoc, Rizal). A casual introduction, occurring rarely in open waste places. A temperate and subtropical weed.

Local names: Dapdapígan (Bon.); upoan (Bon.).

EXCLUDED SPECIES

SETARIA GLOBULARIS Presl Rel. Haenk. 1 (1830) 314; Kunth Enum. 1 (1833) 151; Miq. Fl. Ind. Bat. 3 (1857) 314; F.-Vill. Novis. App. (1882) 312; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 52.

Panicum globulare Steud. Syn. Pl. Glum. 1 (1854) 51.

"Hab. in insulis Philippinis." The species is certainly not Philippine, but the specimens on which it was based were probably of Mexican origin. A sheet of Presl's type material in the herbarium of the Missouri Botanical Garden consists of *Setaria caudata* Lam. and *S. composita* Kunth.

SETARIA MACROSTACHYA HBK.; F.-Vill. Novis. App. (1882) 312.

33. CENCHRUS Linnaeus

CENCHRUS VIRIDIS Spreng. Syst. Veg. 1 (1825) 301; Chase in Contr. U. S. Nat. Herb. 22 (1920) 57, f. 11.

Cenchrus inflexus R. Br. Prodr. (1810) 50, non Poir.

Cenchrus echinatus Presl Rel. Haenk. 1 (1830) 317; Miq. Fl. Ind. Bat. 3 (1857) 472; F.-Vill. Novis. App. (1882) 313; Perk. Frag. Fl. Philip. (1904) 145; Merr. in Philip. Journ. Sci. 1 (1906) 367, Fl. Manila (1912) 95, Sp. Blancoanae (1918) 68, non Linn. .

Cenchrus hexaflorus Blanco Fl. Filip. (1837) 36, ed. 2 (1845) 24, ed. 3, 1 (1877) 46.

A widely distributed weed in the settled areas of the Philippines, occurring generally in open waste places in and about towns. A native of tropical America, now also found in Guam, Siam, Indo-China, New Guinea, and northern Australia. Dr. A. B. Rendle has critically compared our material with the type of *Cenchrus inflexus* R. Br.

The Philippine form has been confused with *C. echinatus* Linn.

Local names: Agíngai (Tag.); sagisí (Ilk.).

34. PENNISETUM Persoon

PENNISETUM ALOPECUROIDEUM (Linn.) Spreng. Syst. 1 (1825) 303.

Panicum alopecuroideum Linn. Sp. Pl. (1753) 55.

Alopecurus hordeiformis Lour. Fl. Cochinch. (1790) 48, non ? Linn.

Pennisetum compressum R. Br. Prodr. (1810) 195; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 368.

Gymnothrix nigricans Presl Rel. Haenk. 1 (1830) 315.

Pennisetum nigricans Trin. ex Steud. Nomencl. ed. 2, 2 (1841) 297;

Miq. Fl. Ind. Bat. 3 (1857) 470; F.-Vill. Novis. App. (1882) 312.

Pennisetum cenchroides F.-Vill. Novis. App. (1882) 313, non Spreng.

Luzon (Ilocos Norte, Bontoc, Benguet), *Merrill 7704, 4435, Williams 1201, Elmer 5756, B. S. 31898 Santos, 5932, 5365, 27422 Ramos, 12708 Fénix, 8763 McGregor*. On open slopes at medium altitudes, chiefly in the pine region, ascending to 2,000 m. India to Indo-China, China, Japan, and Australia.

Local name: Kaibuan (Ig.).

PENNISETUM MACROSTACHYUM Trin. in Mém. Acad. St. Pétersb. VI

3² (1835) 177; Ceron Cat. Pl. Herb. Manila (1892) 121; Merr. in

Philip. Journ. Sci. 1 (1906) Suppl. 368.

Sericura elegans Hassk. in Flora 25 (1842) Beibl. 2: 2.

Luzon (Laguna, Batangas, Rizal, Tayabas, Sorsogon), Mindoro, Masbate, Panay, Cebu, Negros, Mindanao, Basilan, *Merrill 3381, Williams 2622, Elmer 9674, 81324, 15327, B. S. 31306 Ramos & Edaña, 24939 Fénix, 14749, 11079, 5217 Ramos, 15494 Reillo*. In old clearings at low and medium altitudes. Malay Peninsula and Archipelago to Polynesia.

Local names: Daráui (C. Bis.); gulálai (Sub.); handaláui (C. Bis.); lagóli (Bag.).

35. CHAMAERAPHIS R. Brown

CHAMAERAPHIS MURICATA (Linn. f.) comb. nov.

Andropogon muricatus Linn. f. Suppl. (1781) 433; Stapf in Kew Bull. (1906) 348 in nota.

Panicum asperum Koen. in Naturf. 23 (1788) 209.

Panicum spinescens R. Br. Prodr. (1810) 193.

Chamaeraphis spinescens Poir. in Lam. Encycl. Suppl. 2 (1811) 189;

F.-Vill. Novis. App. (1882) 313; Hook. f. Fl. Brit. Ind. 7 (1897) 62.

Chamaeraphis aspera Nees in Wall Cat. (1849) No. 8679; Mez in

Perk. Frag. Fl. Philip. (1904) 145; Merr. in Philip. Journ. Sci. 1

(1906) 368, Fl. Manila (1912) 96.

Central Luzon to Palawan and Mindanao, *Merrill 375, 4194, 7060, 9467, B. S. 29500 Ramos & Edaña*. In shallow stagnant water at low altitudes. India to China through Malaya to tropical Australia.

36. THUAREA Persoon

THUAREA INVOLUTA (Forst.) R. & S. Syst. 2 (1817) 782; Merr. in

Philip. Journ. Sci. 1 (1906) Suppl. 369, Fl. Manila (1912) 97.

Ischaemum involutum Forst. Prodr. (1786) 73.

Thuarea sarmentosa Pers. Syn. 1 (1805) 110; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97 (*Thouarea*); F.-Vill. Novis. App. (1882) 313; Usteri Beitr. Ken. Phil. Veg. (1905) 133; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 28.

Luzon (Cagayan, Bataan, Rizal), Mindoro, Palawan, Samar, Mindanao. On sandy beaches and in thickets back of the beach, often common. Ceylon to Madagascar through Malaya to Australia and Polynesia.

37. SPINIFEX Linnaeus

SPINIFEX LITTOREUS (Burm. f.) Merr. in Philip. Journ. Sci. 7 (1912) Bot. 229, Fl. Manila (1912) 97, Sp. Blancoanae (1918) 69.

Stipa littorea Burm. f. Fl. Ind. (1768) 29.

Stipa spinifex Linn. Mant. 1 (1767) 84; Blanco Fl. Filip. (1837) 41; ed. 2 (1845) 29, ed. 3, 1 (1877) 54.

Spinifex squarrosus Linn. Mant. 2 (1771) 163; Blanco Fl. Filip. (1837) 46; ed. 2 (1845) 31; ed. 3, 1 (1877) 57; F.-Vill. Novis. App. (1882) 313; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 369.

On sandy seashores from the Batan Islands to Palawan and Mindanao. India to China and Malaya.

Local names: Baki-báki (Bis.); tartarái (Ilk.); tayung-táyung (C. Bis.).

38. LEPTASPIS R. Brown

LEPTASPIS CUMINGII Steud. Syn. Pl. Glum. 1 (1854) 416; Miq. Fl. Ind. Bat. 3 (1857) 375; F.-Vill. Novis. App. (1882) 318; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 370.

Leptaspis cochleata Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 81, non Thwaites.

Mindanao (Misamis, Davao), Bongao, *Copeland 703*, *Cuming 1627*, *B. S. 36319 Yates*. In forests at low or medium altitudes. Endemic.

LEPTASPIS URCEOLATA (Roxb.) R. Br. in Benn. Pl. Jav. Rar. (1838) 23, *t. 6*; Miq. Fl. Ind. Bat. 3 (1854) 374; F.-Vill. Novis. App. (1882) 318; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 370; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 194.

Pharus urceolatus Roxb. Fl. Ind. ed. 2, 3 (1832) 611.

Leptaspis manillensis Steud. Syn. Pl. Glum. 1 (1854) 8; Miq. Fl. Ind. Bat. 3 (1857) 374; F.-Vill. Novis. App. (1882) 318.

Luzon (Camarines, Sorsogon), Polillo, Palawan, Samar, Leyte, Negros, Mindanao. In thickets and forests at low and medium altitudes; represented by numerous collections. Ceylon, Malay Peninsula, Sumatra, Java, through the Moluccas to New Guinea.

Local names: Alaplap (Mbo.); baliotus (Bag.); budhak-budhak (Sul.); handalokot (Mbo.); madioug (S. L. Bis.); rokot-dókot (S. L. Bis.); sa-lindaget (Mbo.); taliñgagito (Sub.).

39. ORYZA Linnaeus

ORYZA MEYERIANA (Zoll. & Morr.) Benth. ex Pilger in Perk. Frag. Fl. Philip. (1904) 145; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 370.

Padia meyeriana Zoll. & Morr. Verz. Ind. Arch. (1854) 103; Steud. Syn. Pl. Glum. 1 (1854) 3; Miq. Fl. Ind. Bat. 3 (1857) 373.

Oryza granulata Nees & Arn. in Wall. Cat. (1830) No. 2354, *nomen nudum*; Hook. f. Fl. Brit. Ind. 7 (1897) 93; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 7.

Luzon (Isabela, Ifugao, Nueva Vizcaya, Rizal, Tayabas), Palawan, Panay, Mindanao (Zamboanga), Merrill 116, 6697, 8060, 7244, B. S. 19967 McGregor, 20677 Escritor, 343 Bermejós, 8036 Ramos. In forests, bamboo thickets, etc., at low and medium altitudes. India to Java, Borneo, and Celebes.

ORYZA MINUTA Presl Rel. Haenk. 1 (1830) 208; Kunth Enum. 1 (1833) 7; Miq. Fl. Ind. Bat. 3 (1857) 371; F.-Vill. Novis. App. (1882) 319.

Oryza manillensis Merr. in Philip. Journ. Sci. 3 (1908) Bot. 219, Fl. Manila (1912) 98.

Oryza latifolia F.-Vill. Novis. App. (1882) 319, non Desv.

Luzon (Rizal, Laguna, Tayabas, Sorsogon), Mindanao (Zamboanga), Presl! in Herb. Bernhardi, Merrill 8112, 7402, Phil. Pl. 128, Elmer 17407, B. S. 29492 Ramos & Edaño, 2194, 12102, 12044, 13269 Ramos, 36714 Ramos & Edaño. In open wet places at low and medium altitudes. Borneo, Malay Peninsula.

This is I believe the Malay Peninsula and Bornean form currently referred to *Oryza latifolia* Desv., but I believe the type of Desvaux's species was an American plant. Of the specimens cited, Elmer 17407, B. S. 36714 Ramos & Edaño, and Merrill 8112, being more robust than the others, closely match the Malay Peninsula form.

Local names: Palai-máya (Sul.); palai-pálai (Sub.); baili-gaja (Sul.).

ORYZA SATIVA Linn. Sp. Pl. (1753) 333; F.-Vill. Novis. App. (1882) 318; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 370, Fl. Manila (1912) 97, Sp. Blancoanae (1918) 69, 70.

Oryza aristata Blanco Fl. Filip. (1837) 274; ed. 2 (1845) 190; ed. 3, 1 (1877) 339.

Oryza sativa Linn. vars. *binamban*, *glutinosa*, *lamuyo*, *pilosa*, *praecox*, *quinanda*, *rubra*, et *violacea* Blanco l. c.

Oryza glutinosa Lour. Fl. Cochinch. (1790) 215; F.-Vill. Novis. App. (1882) 319.

Oryza montana Lour. l. c. 215; F.-Vill. l. c.

Oryza praecox Lour. l. c. 215; F.-Vill. l. c.

Extensively cultivated throughout the Philippines but not a native of the Archipelago, although of prehistoric introduction. The rice plant is enormously variable and probably several hundred distinct cultural forms occur in the Archipelago. No attempt has been made to compile the very numerous local names currently applied to special varieties. Rice.

Local names: Ammái (Ibn. It.); págai (Il.); págei (Bon.); pálai (Tag.); párai (Bik.); pai (Sul.).

40. LEERSIA Swartz

LEERSIA HEXANDRA Sw. Prodr. (1788) 21; Llanos Frag. Pl. Filip. (1851) 26; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 18; F.-Vill. Novis. App. (1882) 318; Vidal Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Filip. (1886) 288; Pilger in Perk. Frag. Fl. Philip. (1904) 145; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 371, Fl. Manila (1912) 98.

Leersia luzonensis Presl Rel. Haenk. 1 (1830) 207; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 97; Miq. Fl. Ind. Bat. 3 (1857) 367; F.-Vill. Novis. App. (1882) 318.

Leersia glaberrima Trin. in Mém. Acad. St. Pétersb. VI 5 (1839) 173; Miq. Fl. Ind. Bat. 3 (1857) 368; F.-Vill. Novis. App. (1882) 318.

Homalocenchrus hexandrus O. Kuntze Rev. Gen. Pl. (1891) 777.

Northern Luzon to Mindanao. In open wet places at low and medium altitudes, ascending to 1,500 m. In the vicinity of Manila this is very extensively cultivated in paddies for green forage.

Local names: Bárit (Tag.); buñgálon (Bis.); zacáte (Sp.).

41. MICROLAENA R. Brown

MICROLAENA STIPOIDES (Labill.) R. Br. Prodr. (1910) 210; Benth. Fl. Austral. 7 (1878) 552; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 269; Merr. op. cit. 371, 5 (1910) Bot. 328.

Ehrhartia stipoides Labill. Pl. Nov. Holl. 1 (1804) 91, t. 118.

Luzon (Abra, Bontoc, Lepanto, Benguet, Zambales), Merrill 4543, 4696, 4831, 6570, Phil. Pl. 1343, B. S. 31957 Santos, 5128, 7064 Ramos. On open slopes and ridges, along mountain trails, etc., altitude 1,800 to 2,300 m. Australia, New Zealand, Hawaii.

42. PHALARIS Linnaeus

PHALARIS MINOR Retz. Obs. 3 (1783) 8; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 421.

Luzon (Benguet), B. S. 2785 Mearns. Altitude about 1,500 m; probably a mere casual here. Europe, India, South Africa, Australia.

43. ANTHOXANTHUM Linnaeus

ANTHOXANTHUM LUZONIENSE Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 178, 372, 5 Bot. (1910) Bot. 328.

Luzon (Benguet), Merrill 4713, 6614, Phil. Pl. 184, B. S. 8890 McGregor, F. B. 16188 Curran, Merritt, & Zschokke. In open grasslands, altitude 2,000 to 2,700 m. Endemic.

44. ARISTIDA Linnaeus

ARISTIDA CULIONENSIS Pilg. ex Perk. Frag. Fl. Philip. (1904) 145; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 372.

Culion, Merrill 471, 515, B. S. 41334 Ramos. In open grasslands at low altitudes. Endemic.

ARISTIDA CUMINGIANA Trin. & Rupr. in Mém. Acad. St. Pétersb. VI 7 (1849) 141; Steud. Syn. Pl. Glum. 1 (1854) 140; Miq. Fl. Ind. Bat. 3 (1857) 381; Vid. Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 372, 5 (1910) Bot. 328, Fl. Manila (1912) 99.

Aristida capillacea Cav. Ic. 5 (1799) 43, t. 468, f. 1, non Lam.

Chaetaria trichodes Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 101.

Aristida trichodes Walp. Ann. 3 (1853) 753.

Luzon (Ilocos Norte, Lepanto, Ifugao, Bontoc, Benguet, Pangasinan, Rizal), Mindanao (Bukidnon). In open dry grasslands at low and medium altitudes, ascending to 1,500 m. India to tropical Africa and southern China.

ARISTIDA TENUISETULOSA (Pilg.) Mez in Fedde Repert. 17 (1921) 147.

Aristida stipoides R. Br. var. *tenuisetulosa* Pilg. ex Perk. Frag. Fl. Philip. (1904) 146; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 372.

Luzon (Amburayan, Ilocos Norte, Ilocos Sur, Zambales, Pampanga, Bataan), Merrill 329, F. B. 5855, 6361 Curran, 12476, 15657 Merritt & Darling, B. S. 2334 Mearns. In dry open sandy soil at low and medium altitudes. Endemic, but closely allied to the Australian *A. stipoides* R. Br.

EXCLUDED SPECIES

ARISTIDA LAXA Cav. Ic. 5 (1799) 44, t. 470, f. 1; Kunth Enum. 1 (1833) 192; Miq. Fl. Ind. Bat. 3 (1857) 381; F.-Vill. Novis. App. (1882) 319.

"Habitat prope Montevideo, et in insulis Philippicis." This is probably a Mexican species, certainly not Philippine, the latter record being based on a wrongly localized specimen of Née's collection.

ARISTIDA LUZONIENSIS Cav. Ic. 5 (1799) 45, t. 470, f. 2; Spreng. Syst. 1 (1825) 266; Kunth Enum. 1 (1833) 192; Miq. Fl. Ind. Bat. 3 (1857) 381; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 99; Novis. App. (1882) 319.

Chaetaria luzoniensis Beauv. Agrost. (1812) 30.

"Habitat in insula Luzon altera ex Philippicis." Certainly not a Philippine species, but American or Australian.

ARISTIDA MURINA Cav. Ic. 5 (1799) 44, t. 469, f. 1; Kunth Enum. 1 (1833) 192; F.-Vill. Novis. App. (1882) 319.

"Habitat in Mindanao insula prope Samboangan [= Zamboanga]." Undoubtedly a wrongly localized species, and Australian or American rather than Philippine.

ARISTIDA RIGIDA Cav. Ic. 5 (1799) 44, t. 469, f. 2; Kunth Enum. 1 (1833) 192; Miq. Fl. Ind. Bat. 3 (1857) 381; F.-Vill. Novis. App. (1882) 319.

"Habitat in insulis Philippicis." Unquestionably an erroneously localized species, and Australian or American rather than Philippine.

ARISTIDA SORZOGONENSIS Presl Rel. Haenk. 1 (1830) 224; Kunth Enum. 1 (1833) 192; Miq. Fl. Ind. Bat. 3 (1857) 381; F.-Vill. Novis. App. (1882) 319.

"Hab. in Luzonia ad Sorzogon." This record is probably based on an erroneously labeled specimen, the species in all probability being American.

44a. MUEHLENBERGIA Schreber

MUEHLENBERGIA ARISANENSIS Hayata Ic. Pl. Formosa 7 (1918) 87, f. 54.

Luzon (Lepanto, Bontoc, Benguet), Merrill 4716, Clemens 9203, Vano-verbergh 1153, B. S. 40213, 40296, 40360 Ramos & Edaña. Along small streams, in old clearings, and in the mossy forest, altitude 1,300 to 2,400 m. Formosa. Closely allied to *M. japonica* Steud.

45. SPOROBOLUS R. Brown

SPOROBOLUS DIANDRUS (Retz.) Beauv. Agrost. (1812) 26; Miq. Fl. Ind. Bat. 3 (1857) 375; F.-Vill. Novis. App. (1882) 321; Pilger in Perk. Frag. Fl. Philip. (1904) 146; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 373.

Agrostis diandra Retz. Obs. 5 (1786) 19.

Vilfa diandra Steud. Syn. Pl. Glum. 1 (1854) 155.

Luzon (Ilocos Norte, Pampanga, Rizal, Tayabas, Albay, Sorsogon), Mindoro, Palawan, Panay, Mindanao. In lawns and waste places about towns, perhaps introduced. Tropical Asia through Malaya to Australia.

SPOROBOLUS INDICUS (Linn.) R. Br. Prodr. (1810) 170; F.-Vill. Novis. App. (1882) 321; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 373, Fl. Manila (1912) 98, Sp. Blancoanae (1918) 71.

Agrostis indica Linn. Sp. Pl. (1753) 63.

Sporobolus elongatus R. Br. Prodr. (1810) 170; Usteri Beitr. Ken. Philip. Veg. (1905) 133.

Spermachiton involutum Llanos Fragm. Pl. Filip. (1851) 25; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 17.

Luzon (Cagayan, Isabela, Bontoc, Benguet), Mindanao (Lanao). Common at medium altitudes in open grasslands, ascending to 2,000 m. Pan-tropic.

Local names: Bakit (P. Bis.); bangkit (P. Bis.); lusai (Sub.); sañg-saṅgitan (Bon.).

SPOROBOLUS PILIFERUS (Trin.) Kunth Enum. 1 (1833) 611; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 373.

Vilfa pilifera Trin. Diss. (1828) 157, Sp. Gram. Ic. t. 58.

Sporobolus ciliatus Presl Rel. Haenk. 1 (1830) 242; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 53, t. 30; Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 269.

Luzon (Bontoc, Benguet), Merrill 4362, 4718, Phil. Pl. 194, Williams 1957, B. S. 5361, 5905 Ramos. On banks, in open grasslands, etc., at medium altitudes, ascending to 1,800 m. India to the Malay Peninsula.

SPOROBOLUS VIRGINICUS (Linn.) Kunth Rev. Gram. 1 (1829) 67;
Merr. in Philip. Journ. Sci. 5 (1910) Bot. 171, Fl. Manila (1912) 99.
Agrostis virginica Linn. Sp. Pl. (1753) 63.

Sporobolus humilis Presl Rel. Haenk. 1 (1830) 241; Miq. Fl. Ind.
Bat. 3 (1857) 376; F.-Vill. Novis. App. (1882) 321; Scribn. in
Rept. Mo. Bot. Gard. 10 (1899) 53, t. 30; Merr. in Philip. Journ.
Sci. 1 (1906) Suppl. 374.

Vilfa humilis Steud. Syn. Pl. Glum. 1 (1854) 153.

Luzon (Cagayan, Rizal), *Merrill Phil. Pl. 574, B. S. 7881 Ramos*. Along
tidal streams within the influence of salt or brackish water. All or most
warm countries.

EXCLUDED SPECIES

SPOROBOLUS SCOPARIUS Presl Rel. Haenk. 1 (1830) 243; Kunth Enum. 1
(1833) 216; Miq. Fl. Ind. Bat. 3 (1857) 376; F.-Vill. Novis. App.
(1882) 321; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 374.

Vilfa scoparia Steud. Syn. Pl. Glum. 1 (1854) 156.

"Hab. ad portum Sorzogon" Presl. Certainly not a Philippine species,
but probably based on American material.

46. GARNOTIA Brongniart

GARNOTIA STRICTA Brongn. in Bot. Duperr. Voy. (1829) 132, t. 21;
Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 374.

Northern Luzon to Mindanao. On talus slopes, along streams, etc.,
at low and medium altitudes; represented by numerous collections. In-
dia to Polynesia and Hawaii.

Var. **LONGISETA** (Merr.) Hack. in Allgem. Bot. Zeitschr. 12 (1909)
141.

Garnotia longisetata Merr. ex Hack. l. c. in syn.

Luzon (Bontoc, Benguet, Nueva Vizcaya, Bulacan, Bataan, Laguna).
Chiefly in the mossy forest, altitude up to 2,300 m, but occurring also at
low or medium altitudes. Endemic.

Local names: Balahibong-usá (Tag.); bariri (Bag.); pauai (Ig.).

47. AGROSTIS Linnaeus

AGROSTIS ALBA Linn. Sp. Pl. (1753) 63; Hook. f. Fl. Brit. Ind. 7 (1897)
254.

Luzon (Benguet), *B. S. 40496 Ramos & Edaña*. Along trails in open
forest, altitude about 1,500 m; probably introduced. A native of temperate
Europe and Asia, now widely distributed in temperate and subtemperate
countries.

AGROSTIS ELMERI Merr. in Govt. Lab. Publ. (Philip.) 29 (1905) 7,
Philip. Journ. Sci. 1 (1906) Suppl. 375, 5 (1910) Bot. 328.

Luzon (Bontoc, Lepanto, Benguet), Negros. In the mossy forest and
in open places, altitude 1,600 to 2,800 m; represented by numerous col-
lections. Southern China. To be compared with *A. micrantha* Steud.

Local name: Sasamon (Ig.).

48. ANISELYTRON Merrill

ANISELYTRON AGROSTOIDES Merr. in Philip. Journ. Sci. 5 (1910) Bot. 329.

Luzon (Benguet), *Merrill 6483*. In the mossy forest on Mt. Pulo, altitude about 2,400 m. A monotypic endemic genus known only from the original collection.

49. CALAMAGROSTIS Roth

CALAMAGROSTIS ARUNDINACEA (Linn.) Roth var. **NIPPONICA** (Franch. & Sav.) Hack. in Bull. Herb. Boiss. 7 (1899) 652, Philip. Journ. Sci. 1 (1906) Suppl. 269; Merr. op. cit. 375.

Calamagrostis nipponica Franch. & Sav. Enum. Pl. Jap. 3 (1879) 599.

Luzon (Lepanto, Benguet), Mindanao (Davao), *Merrill 4564, 4697, 4701, 7766, B. S. 31740 Santos, 4487 Mearns*. On open grassy slopes, altitude 1,800 to 2,400 m. Europe to India, China and Japan.

Local names: Baríri (Bag.); sangsangitau (Ig.); sasamon (Ig.).

CALAMAGROSTIS FILIFOLIA Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 179, 375, 5 (1910) Bot. 330.

Luzon (Benguet), *Merrill 4839, 6610, 6611, 4537, 4715, B. S. 31979 Santos, 8891 McGregor*. On open slopes, altitude 1,800 to 2,200 m. Endemic.

Local name: Añgangsúg (Ig.).

50. ERIACHNE R. Brown

ERACHNE PALLESCENS R. Br. Prodr. (1810) 184; Benth. Fl. Austral. 7 (1878) 630; F.-Vill. Novis. App. (1882) 321; Pilger in Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 376.

Culion, Palawan, *Merrill 520, 9582*. In open grassy places up to 550 m altitude. India to southern China through Malaya to Australia.

ERACHNE TRISETA Nees ex Steud. Syn. Pl. Glum. 1 (1854) 237; Pilger in Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 376.

Culion, *Merrill*. In open grasslands at low altitudes. India, Ceylon.

EXCLUDED SPECIES

ERACHNE GRACILIS Brongn; F.-Vill. Novis. App. (1882) 321.

51. COELACHNE R. Brown

COELACHNE HACKELII Merr. in Govt. Lab. Publ. (Philip.) 29 (1905) 8, Philip. Journ. Sci. 1 (1906) Suppl. 376.

Luzon (Benguet), *Elmer 5762, Merrill 4338, Phil. Pl. 198, Williams 1196*. Forming dense mats along small cool streams, altitude about 1,500 m. Endemic.

Coelachne pulchella R. Br. and *C. brachiata* Munro were both credited to the Philippines by F.-Villar, Novis. App. (1882) 321, but neither has been found in the Archipelago.

52. **DESCHAMPSIA** Beauvois

DESCHAMPSIA FLEXUOSA (Linn.) Trin. in Bull. Acad. St. Pétersb. 1 (1836) 66; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 330.

Aira flexuosa Linn. Sp. Pl. (1753) 65.

Luzon (Benguet), *Merrill 6354, Phil. Pl. 189, B. S. 8904 McGregor, F. B. 16155 Curran, Merritt, & Zschokke*. In open grasslands at the summit of Mt. Pulog, altitude about 2,600 m. Endemic. North Temperate Zone generally.

53. **AVENA** Linnaeus

AVENA FATUA Linn. Sp. Pl. (1753) 80; F.-Vill. Novis. App. (1882) 318; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 287.

Luzon (Benguet), *B. S. 2812 Mearns*. Probably merely a casual here, altitude about 1,500 m. A widely distributed weed in temperate regions, of European origin.

AVENA SATIVA Linn. Sp. Pl. (1753) 79; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 376.

Occasional in the Mountain Province at and above 1,500 m altitude, here a casual that will probably not persist. Oats.

EXCLUDED SPECIES

AVENA STERILIS Linn.; F.-Vill. Novis. App. (1882) 319.

54. **MICROCHLOA** R. Brown

MICROCHLOA INDICA (Linn. f.) O. Kuntze Rev. Gen. Pl. 3 (1893) 356; Hack. in Fedde Repert. 7 (1909) 373; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 74.

Nardus indica Linn. f. Suppl. (1781) 105.

Rottboellia setacea Roxb. Pl. Coromandel 2 (1798) 18, t. 132.

Microchloa setacea R. Br. Prodr. (1810) 208; F.-Vill. Novis. App. (1882) 319.

Luzon (Ilocos Norte, Bontoc, Lepanto, Rizal), *B. S. 32889 Ramos, Loher 7179, Vanoverbergh 764*. In open dry grasslands at low and medium altitudes. Pantropic.

55. **CYNODON** Persoon

CYNODON ARCUATUS Presl Rel. Haenk. 1 (1830) 290; F.-Vill. Novis. App. (1882) 320; Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 41, t. 40; Merr. in Govt. Lab. Publ. (Philip.) 17 (1904) 9, Philip. Journ. Sci. 1 (1906) Suppl. 28, 377, Fl. Manila (1912) 103.

Luzon (Ilocos Norte, Bontoc, Tarlac, Bataan, Rizal, Laguna), *Merrill 3619, Phil. Pl. 135, Williams 88, B. S. 1392, 32738 Ramos, 23130 Ramos*. In open grasslands at low and medium altitudes. Endemic.

CYNODON DACTYLON (Linn.) Pers. Syn. 1 (1804) 85; Miq. Fl. Ind. Bat. 3 (1857) 382; F.-Vill. Novis. App. (1882) 319; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292; Pilg. ex Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 28, 377, Fl. Manila (1912) 103, Sp. Blancoanae (1918) 71.

Panicum dactylon Linn. Sp. Pl. (1753) 58.

Cynodon linearis Willd. Enum. Hort. Berol. (1809) 90; Presl Rel. Haenk. 1 (1830) 290; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 98.

Panicum glumaepatulum Steud. Syn. Pl. Glum. 1 (1854) 41; F.-Vill. Novis. App. (1882) 312 (*glumaepetalum*).

Digitaria glumaepatula Miq. Fl. Ind. Bat. 3 (1857) 439.

Cynodon rufescens Llanos Frag. Pl. Filip. (1851) 31; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 21, non ? Lagasca.

Capriola dactylon O. Kuntze Rev. Gen. Pl. (1891) 764.

Throughout the Philippines, in open grasslands, fallow rice paddies, waste places, at low and medium altitudes; common. Pantropic.

Local names: Balhálut (Bon.); galud-gálud (Ilk.); gráma (Sp.); kola-tai (Tag.); palot-gálot (Ilk.).

56. CHLORIS Swartz

CHLORIS BARBATA (Linn.) Sw. Fl. Ind. Occ. 1 (1797) 200; Miq. Fl. Ind. Bat. 3 (1857) 387; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; F.-Vill. Novis. App. (1882) 320; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 292; Hook. f. Fl. Brit. Ind. 7 (1897) 292; Pilger in Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 377, Fl. Manila (1912) 103, Sp. Blancoanae (1918) 72.

Andropogon barbatus Linn. Mant. 2 (1771) 302.

Chloris longifolia Steud. Syn. Pl. Glum. 1 (1854) 205; Miq. Fl. Ind. Bat. 3 (1857) 388; F.-Vill. Novis. App. (1882) 320.

Chloris inflata Llanos Frag. Pl. Filip. (1851) 31; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 22, non Link.

A common weed in and about towns in many parts of Luzon, Mindoro, Negros, and Palawan, and probably in other islands. A pantropic weed of American origin.

Local names: Banúko (Ilk.); korokorósan (Tag.).

CHLORIS INCOMPLETA Roth Nov. Sp. Pl. (1821) 60; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 288.

Luzon (Zambales), Palawan, Mindanao (Zamboanga), *Merrill 9275, 9616, 8279, Phil. Pl. 1308, Gates 8145*. In dry open sandy soil at low altitudes, sometimes along the seashore. India to Formosa and Malaya.

CHLORIS MEARNSII Merr. in Philip. Journ. Sci. 3 (1908) Bot. 220.

Luzon (Ilocos Norte), *B. S. 7606, 27201 Ramos, 2294 Mearns*. In dry open places at low altitudes; perhaps but a depauperate form of *C. incompleta* Roth. Endemic.

CHLORIS TENER (Presl) Scribn. in Rept. Mo. Bot. Gard. 10 (1899) 41, t. 40.

Cynodon tener Presl Rel. Haenk. 1 (1830) 291; Kunth Enum. 1 (1833) 260; Miq. Fl. Ind. Bat. 3 (1857) 383; F.-Vill. Novis. App. (1882) 320.

Luzon (Sorsogon), Negros, (*Presl*), *B. S. 9951 Robinson*, as *Cynodon*. In open places at low altitudes. Hainan, Indo-China.

DOUBTFUL AND EXCLUDED SPECIES

CHLORIS CRINITA Lagasca Varied. Cienc. 4 (1805) 143; Kunth Enum. 1 (1833) 268; Miq. Fl. Ind. Bat. 3 (1857) 389; F.-Vill. Novis. App. (1882) 320.

"Insulae Philippinae" Kunth. This was probably based on a Malaspina expedition collection erroneously localized. Judging from the description it is not a *Chloris*.

CHLORIS TRUNCATA R. Br. Prodr. (1910) 186; Kunth Enum. 1 (1833) 266; Miq. Fl. Ind. Bat. 3 (1857) 387; F.-Vill. Novis. App. (1882) 320.

Chloris dolichostachya Lagasca Gen. Sp. Nov. (1816) 5.

Lagasca's species was credited to the Philippines by him; Link has reduced it to *Chloris truncata* R. Br., an Australian species. Nothing conforming to it is known from the Philippines.

CHLORIS INFLATA Link Hort. Berol. 1 (1827) 105.

"Calif.; Ins. Philipp." Link. The species was probably based on a Haenke collection, probably erroneously localized, at least as to the Philippine reference.

CHLORIS RADIATA Sw.; F.-Vill. Novis App. (1882) 320.

CHLORIS RUFESCENS Lagasca Varied. Cienc. 4 (1805) 143; Kunth Enum. 3 (1833) 268; Miq. Fl. Ind. Bat. 3 (1857) 388; F.-Vill. Novis. App. (1882) 320.

"Insulae Philippinae" Kunth. Probably not a Philippine species but an erroneously localized Malaspina Expedition collection.

CHLORIS RHACHITRICA Steud. Syn. Pl. Glum. 1 (1854) 205; Miq. Fl. Ind. Bat. 3 (1857) 388; F.-Vill. Novis. App. (1882) 320.

"Cuming no. 111. Ins. Philip." This number of Cuming's Philippine collection is a fern, and none of his Philippine grasses conform to Steudel's description. The reference may be to Cuming's previous South American collection.

57. **TRIOGON** Roth

TRIOGON CHINENSIS (Franch.) Hack. in Bull. Herb. Boiss. II 3 (1903) 503; Merr. in Philip. Journ. Sci. 14 (1919) 369.

Nardurus filiformis Steud. var. *chinensis* Franch. in Nuov. Arch. Mus. Paris II 7 (1894) 149.

Luzon (Ilocos Norte), *B. S. 33237* Ramos. On dry open rocky slopes at low altitudes. Korea, China, Formosa.

58. **ELEUSINE** Gaertner

ELEUSINE INDICA (Linn.) Gaertn. Fruct. 1 (1788) 8; Presl Rel. Haenk. 1 (1830) 286; Llanos Frag. Pl. Filip. (1851) 45; Miq. Fl. Ind. Bat. 3 (1857) 386; F.-Vill. Novis. App. (1882) 320; Vidal Phan. Cuming. Philip. (1885) 159; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 378, Fl. Manila (1912) 104, Sp. Blancoanae (1918) 72.

Cynosurus indicus Linn. Sp. Pl. (1753) 72.

Eleusine barbata Vidal Rev. Pl. Vasc. Filip. (1886) 292.

Eleusine polydactyla Steud. Syn. Pl. Glum. 1 (1854) 211; Miq. Fl. Ind. Bat. 3 (1857) 386; F.-Vill. Novis. App. (1882) 320, ex descr.

An abundant weed in the settled areas throughout the Philippines. All warm countries.

Local names: Apidan (If.); bakis-bakisan (Tag.); baraṅgan (Bik.); bikad-bikad (Sul.); bila-bíla (P. Bis.); dinapaiuk (If.); gagabútan (Tag.); kabit-kabit (Tag.); palagtiki (Bis.); paraṅgis (Ilk.); paraṅgis-sabúṅgan (Pamp.); sabuṅg-sabuṅgan (Tag., Pamp.); sambáli (Tag.).

EXCLUDED SPECIES

ELEUSINE COROCANA (Linn.) Gaertn.; F.-Vill. Novis. App. (1882) 320.

ELEUSINE VERTICILLATA Roxb.; F.-Vill. Novis. App. (1882) 320.

59. DACTYLOCTENIUM Willdenow

DACTYLOCTENIUM AEGYPTIUM (Linn.) Richt. Pl. Europ. 1 (1889) 68; Merr. Fl. Manila (1912) 104, Sp. Blancoanae (1918) 72.

Cynosurus aegyptius Linn. Sp. Pl. (1753) 72.

Dactyloctenium aegyptiacum Willd. Enum. Hort. Berol. (1809) 1029; Pilg. ex Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 379.

Eleusine aegyptiaca Desf. Fl. Atl. 1 (1798-1800) 85; F.-Vill. Novis. App. (1882) 320; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 293.

Eleusine mucronata Llanos Frag. Pl. Filip. (1851) 46, non Spreng.

Dactyloctenium mucronatum Willd. var. *aegyptiacum* Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102.

Throughout the Philippines. A common weed in the settled areas at low and medium altitudes. Pantropic.

Local names: Alam (Tag.); damúṅg-bálang (Tag.).

60. LEPTOCHLOA Beauvois

LEPTOCHLOA CHINENSIS (Linn.) Nees Syll. Ratisb. 1 (1824) 4; Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; Steud. Syn. Pl. Glum. 1 (1854) 209; Miq. Fl. Ind. Bat. 3 (1857) 389; F.-Vill. Novis. App. (1882) 320; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 293; Pilger in Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 379, Fl. Manila (1912) 105.

Poa chinensis Linn. Sp. Pl. (1753) 69.

Leptochloa tetraquetra Presl Rel. Haenk. 1 (1830) 288; Miq. Fl. Ind. Bat. 3 (1857) 389; F.-Vill. Novis. App. (1882) 320.

Luzon (Cagayan to Camarines, in most provinces), Panay. In open usually wet places along or near streams. India to Japan southward through Malaya to Australia.

Local name: Kurukauáyan (Bik.).

LEPTOCHLOA FILIFORMIS (Pers.) R. & S. Syst. 2 (1817) 580; Presl Rel. Haenk. 1 (1830) 288; Kunth Enum. 1 (1833) 270; Miq. Fl. Ind. Bat. 3 (1857) 389; F.-Vill. Novis. App. (1882) 320; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 380.

Eleusine filiformis Pers. Syn. (1805) 87.

Luzon (Abra, Union), Palawan. In open low lands, along streams, in disturbed soil, etc., at low altitudes. Pantropic.

61. PHRAGMITES Trinius

PHRAGMITES KARKA (Retz.) Trin. ex Steud. Nomencl. ed. 2, 2 (1840) 324; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 381.

Arundo karka Retz. Obs. 4 (1786) 21.

Phragmites roxburghii Steud. l. c.; Nees in Noy. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 173, Hook. Journ. Bot. Kew Miscel. 2 (1850) 101; Miq. Fl. Ind. Bat. 3 (1857) 412; F.-Vill. Novis. App. (1882) 321.

Trichoon roxburghii W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 383, 391.

Luzon (Ilocos Norte, Bontoc, Bataan, Laguna, Camarines), Negros, Mindanao, Merrill 3178, Phil. Pl. 581, Elmer 6851, Williams 221, Clemens 728, B. S. 2213 Mearns, 15910 Fénix. In thickets and along streams at low and medium altitudes. India to Malaya.

Local names: Kanúbong (Bon.); lachitlít (Bag.); lupí (Bik.); sabúnog (P. Bis.); tabúnak (P. Bis.); tabúnog (Bis.); talúbung (Bon.); tantanóbong (Ilk.); tanúbung (Bon.).

PHRAGMITES VULGARIS (Lam.) Trin. Fund. Agrost. (1820) 134; Pilger in Perk. Frag. Fl. Philip. (1904) 147; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 381, Fl. Manila (1912) 100, Sp. Blancoanae (1918) 72.

Arundo vulgaris Lam. Fl. Franc. 3 (1778) 615.

Arundo phragmites Linn. Sp. Pl. (1753) 81.

Arundo tecta Blanco Fl. Filip. (1837) 48; ed. 2 (1845) 33; ed. 3, 1 (1877) 60, non Walt.

Arundo donax Llanos in Mem. Acad. Cienc. Madr. 4 (1858) 507; F.-Vill. & Llanos in Blanco Fl. Filip. ed. 3, 4¹ (1880) 105, non Linn.

Phragmites communis Trin. Fund. Agrost. (1820) 134; Usteri Beitr. Ken. Veg. Philip. (1905) 133.

Arundo madagascariensis F.-Vill. Novis. App. (1882) 321, Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 293, non Kunth.

Neyraudia madagascariensis Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 381, non Hook. f.

Northern Luzon to Palawan and Mindanao, often forming dense thickets along the margins of lakes and slow streams. All warm countries.

Local names: Bagang (Bis.); bugang (C. Bis.); lupí (Bik.); tabúnak (Bis., P. Bis.); tagísi (Ibn.); tambó (Tag.); uba-uba (Sul.); tangbó (C. Bis.); tanóbong (Pang., Ilk.); tantanúbong (Ibn.); tanúbong (Ilk.).

62. DIPLACHNE Beauvois

DIPLACHNE POLYSTACHYA (Forsk.) Backer in Bull. Jard. Bot. Buitenz. III 2 (1920) 325.

Bromus polystachyus Forsk. Fl. Aegypt.-Arab. (1775) 23.

Diplachne fusca Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 381, Fl. Manila (1912) 99, non Beauv.

Luzon (Laguna, Rizal), Mindoro, Mindanao (Cotabato), *Merrill 5104*, *Phil. Pl. 183*, *Kneucker Gram. Exsic. 554a*, *Elmer 18131*, *Williams 2027*. In swampy often brackish places at low altitudes. Tropical Asia and Africa to Australia.

63. ERAGROSTIS Beauvois

ERAGROSTIS AMABILIS (Linn.) W. & A. in Hook. & Arn. Bot. Beechy Voy. (1841) 251; Merr. Interpret. Herb. Amb. (1917) 95.

Poa amabilis Linn. Sp. Pl. (1753) 68.

Poa tenella Linn. Sp. Pl. (1753) 69.

Poa plumosa Link. Hort. Berol. 1 (1827) 192; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102 (f. *laxa*); Miq. Fl. Ind. Bat. 3 (1857) 394; F.-Vill. Novis. App. (1882) 322; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 293; Usteri Beitr. Ken. Philip. Veg. (1905) 132 (f. *breviciliata* Hack.).

Eragrostis tenella R. & S. Syst. 2 (1817) 576; Presl Rel. Haenk. 1 (1830) 274; F.-Vill. Novis. App. (1882) 322; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 383, 3 (1908) Bot. 397, Fl. Manila (1912) 101.

In and about towns throughout the Philippines, a very common weed in open waste places. Old World Tropics generally.

Local names: Bagibuk (Tag.); balisibis (Iv.); kaliraúrau (Tag.); págai-billít (Ilk.); pulpulút (Bon.); sigsigid (Bon.).

ERAGROSTIS CILIANENSIS (All.) Vig.-Lut. in Malpighia 18 (1904) 386; Hubbard in Philip. Journ. Sci. 8 (1913) Bot. 159.

Briza eragrostis Linn. Sp. Pl. (1753) 70.

Poa cilianensis All. Fl. Pedem. 2 (1785) 246, t. 91, f. 2.

Eragrostis major Host. Gram. Austr. 4 (1809) 14, t. 24.

Eragrostis megastachya Link Hort. Berol. 1 (1827) 187; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 293.

Cebu, Negros, *B. S. 11122* Ramos, *Eskridge*. In open waste places at low altitudes. All warm countries.

ERAGROSTIS DISTANS Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 81; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 384, 5 (1910) Bot. 330, Fl. Manila (1912) 102, Sp. Blancoanae (1918) 74.

Uniola paniculata Llanos Frag. Pl. Filip. (1851) 32; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 23, non Linn.

Eragrostis pilosa F.-Vill. Novis. App. (1882) 322, non Beauv.

Luzon (Ilocos Norte, Lepanto, Benguet, Pangasinan, Rizal), Palawan, *Elmer 6608*, *Kneucker Gram. Exsic. 761*, *Merrill 4472*, *B. S. 4821*, *5314* Ramos. In dry open grasslands at low and medium altitudes. Endemic.

ERAGROSTIS ELONGATA (Willd.) Jacq. Eclog. Gram. (1818) 3, t. 3; Presl Rel. Haenk. 1 (1830) 275; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 384.

Poa elongata Willd. Enum. Hort. Berol. 1 (1809) 108.

Eragrostis elegantula Pilger in Perk. Frag. Fl. Philip. (1904) 148; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 384, Fl. Manila (1912) 102, non Steud.

Eragrostis nigra Usteri Beitr. Ken. Philip. Veg. (1905) 132, non Nees.

Eragrostis luzoniensis Steud. Syn. Pl. Glum. 1 (1854) 266; Miq. Fl.

Ind. Bat. 3 (1857) 393; F.-Vill. Novis. App. (1882) 322.

Luzon (Isabela, Ilocos Norte, Ifugao, Bontoc, Benguet, Nueva Vizcaya, Nueva Ecija, Pangasinan, Rizal), Culion, Palawan, Mindanao (Bukidnon). Common in open grasslands at low and medium altitudes. Pantropic.

Hackel has identified this as *Eragrostis atrovirens* Trin. Typical specimens are Merrill 4155, 9321 and Phil. Pl. 1460, 130. Cuming 1416, cotype of *E. luzoniensis* Steud., appears to me to be merely a form with pale spikelets.

ERAGROSTIS INTERRUPTA (R. Br.) Beauv. Agrost. (1812) 71 (non Doell).

Poa interrupta R. Br. Prodr. (1810) 180.

Eragrostis brounei Nees ex Steud. Nomencl. ed. 2, 1 (1841) 562;

Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; F.-Vill. Novis. App.

(1882) 322; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl.

Vasc. Filip. (1886) 293.

Eragrostis cumingii Steud. Syn. Pl. Glum. 1 (1854) 266; Miq. Fl.

Ind. Bat. 3 (1857) 394; F.-Vill. Novis. App. (1882) 322.

Eragrostis zeylanica Nees in Nov. Act. Acad. Nat. Cur. 19 (1843)

Suppl. 1: 204; Hook. Journ. Bot. Kew Miscel. 2 (1850) 102 (var.

glomerata et *minor*); Rolfe in Journ. Bot. 23 (1885) 216; Vidal

Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886)

293; Pilger in Perk. Frag. Fl. Philip. (1904) 148.

Luzon (Abra, Bontoc, Benguet, Union, Nueva Vizcaya, Bataan, Laguna, Zambales), Culion, Cuming 1104, 672, Merrill 686, 4331, Elmer 5707, Williams 1181, 92. In open grasslands at low and medium altitudes. Pantropic.

ERAGROSTIS JAPONICA (Thunb.) Trin. in Mém. Acad. St. Pétersb. VI

1 (1830) 450; Hack. in Bull. Herb. Boiss. 7 (1899) 705; Merr.

in Philip. Journ. Sci. 1 (1906) Suppl. 384, Sp. Blancoanae (1918)

73.

Poa japonica Thunb. Fl. Jap. (1784) 31; Llanos Frag. Pl. Filip.

(1851) 47; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880)

34.

Poa interrupta Lam. Tabl. Encycl. 1 (1791) 185.

Eragrostis aurea Steud. Syn. Pl. Glum. 1 (1854) 267; Miq. Fl. Ind.

Bat. 3 (1857) 394; F.-Vill. Novis. App. (1882) 322.

Panicum leptanthum Steud. op. cit. 79.

Vilfa verticillata Steud. op. cit. 158.

Sporobolus verticillatus Nees in Hook. Journ. Bot. Kew Miscel. 2

(1850) 101; Miq. Fl. Ind. Bat. 3 (1857) 375; F.-Vill. Novis. App.

(1882) 321.

Eragrostis tenuissima Schrad. ex Nees in Fl. Afr. Austral. (1841)

409; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip.

(1886) 293.

Eragrostis interrupta (non Beauv.!) var. *tenuissima* Stapf in Hook. f. Fl. Brit. Ind. 7 (1897) 316.

Poa amboinensis F.-Vill. Novis. App. (1882) 322, ex syn. Llanos, non

Linn.

Eragrostis hapalantha Trin.; Presl Repert. 1 (1833) 252.

Eragrostis minutiflora Presl Rel. Haenk. 1 (1830) 274 (excl. syn. Retz.?).

Eragrostis interrupta Doell in Mart. Fl. Bras. 3: 157; Usteri Beitr. Ken. Philip. Veg. (1905) 132; Pilger in Perk. Frag. Fl. Philip. (1904) 148; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 383, Fl. Manila (1912) 101, non Beauv.

Eragrostis interrupta (non Beauv.!) var. *koenigii* Stapf in Hook. f. Fl. Brit. Ind. 7 (1897) 316.

Northern Luzon (Cagayan) to Palawan and Mindanao. In open wet grasslands, along streams, etc., at low and medium altitudes. Tropical Asia and Africa to Japan and Malaya.

Thunberg's specific name is accepted for this species as a collective one. Merrill 7322, *Phil. Pl.* 165, *Sp. Blancoanae* 709 represent the species as I here interpret it; numerous specimens have been distributed as *E. interrupta* Doell.

ERAGROSTIS LASIOCLADA Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 382.

Culion, Merrill 461. In fallow rice paddies at low altitudes. Endemic.

ERAGROSTIS MANGALORICA Hochst. ex Steud. Syn. Pl. Glum. 1 (1854) 265; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 299, Fl. Manila (1912) 101.

Luzon (Cagayan, Rizal), Merrill 371, *B. S.* 7452, 14569 Ramos. In open waste places at low altitudes. India.

ERAGROSTIS PILOSA (Linn.) Beauv. Agrost. (1812) 71; Merr. Fl. Manila (1912) 102.

Poa pilosa Linn. Sp. Pl. (1753) 68.

Luzon (Benguet, Rizal), Merrill 9657, *Phil. Pl.* 573. A weed in open waste places, of recent introduction, but now locally abundant. All warm countries.

ERAGROSTIS REFLEXA Hack. in Philip. Journ. Sci. 3 (1908) Bot. 168.

Luzon (Rizal), *B. S.* 2067 Ramos. In fallow lands at low or medium altitudes. Endemic.

ERAGROSTIS SPARTINOIDES Steud. Syn. Pl. Glum. 1 (1854) 265; Pilger in Perk. Frag. Fl. Philip. (1904) 148; Merr. Fl. Manila (1912) 102, *Sp. Blancoanae* (1918) 74.

Uniola spicata Llanos Frag. Pl. Filip. (1851) 33; F.-Vill. & Naves in Blanco Fl. Philip. ed. 3, 4¹ (1880) 23, non Linn.

Luzon (Cagayan, Ilocos Norte, Ifugao, Bontoc, Nueva Vizcaya, Rizal, Laguna), Mindoro, Cuming 668, Merrill 7055, *Phil. Pl.* 107, *Sp. Blancoanae* 170, 422, etc. In open grasslands at low altitudes. Endemic.

ERAGROSTIS UNIOLOIDES (Retz.) Nees Steud. Syn. Pl. Glum. 1 (1854) 264; F.-Vill. Novis. App. (1882) 322; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 384, Fl. Manila (1912) 102.

Eragrostis amabilis Auct., non (Linn.) W. & A.

Eragrostis rubens F.-Vill. Novis. App. (1882) 322, non Hochst.?

Poa annua Llanos Frag. Pl. Filip. (1851) 47; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 34, non Lam.

Eragrostis polymorpha R. Br. Prodr. (1810) 180; F.-Vill. Novis. App. (1882) 322.

Luzon (Zambales, Rizal, Sorsogon), *Elmer 14531*, *Merrill Phil. Pl. 160*, *F. B. 5886 Curran*, *B. S. 19218 Reillo*. In open wet places at low altitudes. Old World Tropics, introduced into the New World.

ERAGROSTIS VISCOSA (Retz.) Trin. in Mém. Acad. St. Pétersb. VI 1 (1831) 397; F.-Vill. Novis. App. (1882) 322; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 383, Fl. Manila (1912) 101, Sp. Blancoanae (1918) 73.

Poa viscosa Retz. Obs. 2 (1781) 20.

Eragrostis tenella R. & S. var. *viscosa* Stapf in Hook. f. Fl. Brit. Ind. 7 (1897) 315; Pilger in Perk. Frag. Fl. Philip. (1904) 148.

Cyperus paniculatus Blanco Fl. Filip. (1837) 32, ed. 2 (1845) 22, ed. 3, 1 (1877) 42, non aliorum.

Luzon (Zambales, Rizal), *Merrill Sp. Blancoanae 229*, *Phil. Pl. 148*. In open waste places in and near towns. India to Malaya.

Local name: Kaliraúrau (Tag.).

EXCLUDED SPECIES

ERAGROSTIS CILIATA Nees; F.-Vill. Novis. App. (1882) 322.

64. CENTOTHECA Desvaux

CENTOTHECA LATIFOLIA (Osbeck) Trin. Fund. Agrost. (1820) 141; Merr. Fl. Manila (1912) 100, Sp. Blancoanae (1918) 74.

Holcus latifolius Osbeck Dagbok Ostind. Resa (1757) 247; Linn. Syst. ed. 10 (1759) 1305.

Cenchrus lappaceus Linn. Sp. Pl. ed. 2 (1763) 1488.

Centotheca lappacea Desv. in Nuov. Bull. Soc. Philom. 2 (1810) 189; Presl Rel. Haenk. 1 (1830) 258; Miq. Fl. Ind. Bat. 3 (1857) 398; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 98; F.-Vill. Novis. App. (1882) 322; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 293; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 29, 3 (1908) Bot. 397; Perk. ex Frag. Fl. Philip. (1904) 148.

Melica philippinensis Llanos Frag. Pl. Filip. (1851) 44; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 32.

Centotheca malabarica Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 385, non *Poa malabarica* Linn.

Throughout the Philippines in thickets, forests, and clearings at low and medium altitudes; common. Usually the only grass in some types of forest. Tropical Asia and Africa through Malaya to Polynesia.

Local names: Baili-patung (Sul.); balikbátang (Sub.); kauakauayánan (Tag.); lansa-lansa (Sul.); maríga (Sul.); sikúan (Ig.).

65. LOPHATHERUM Brongniart

LOPHATHERUM GRACILE Brongn. in Bot. Duperr. Voy. (1829) 50, t. 8; F.-Vill. Novis. App. (1882) 323; Pilger in Perk. Frag. Fl. Philip. (1904) 148; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 386, 2 (1907) Bot. 261; C. B. Rob. op. cit. 6 (1911) Bot. 194.

Luzon (Nueva Ecija, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Mindoro, Palawan, Leyte, Mindanao, Basilan, Jolo. In forests, along trails, and in old clearings, chiefly at medium altitudes. India to Japan, southward through Malaya to New Guinea.

The Philippine form is referable to var. *pilosum* Camus in Bull. Mus. Hist. Nat. Paris 25 (1919) 495.

66. POA Linnaeus

POA ANNUA Linn. Sp. Pl. (1753) 68; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 181, 386.

Luzon (Benguet), *Merrill 4288, B. S. 32041 Santos*. Naturalized in open waste places, altitude 1,500 to 2,400 m. Temperate and subtemperate regions generally.

The form reported from the Philippines as *Poa annua* by Llanos, Frag. Pl. Filip. (1851) 47, and by F.-Vill., Novis. App. (1882) 322, was probably *Eragrostis unioides* Nees; there is no description.

POA LUZONIENSIS Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 180, 386.

Luzon (Benguet), *Merrill 4712, B. S. 31911 Santos*. Borders of cold marshes and streams, altitude about 2,400 m. Reported by Ridley from New Guinea, but the specimens should be compared.

67. FESTUCA Linnaeus

FESTUCA MEGALURA Nutt. in Journ. Acad. Sci. Phila. II 1 (1847) 188; Piper in Contr. U. S. Nat. Herb. 10 (1906) 17, t. 5.

Festuca myuros Merr. in Philip. Journ. Sci. 14 (1919) 369, non Linn.

Luzon (Benguet), *B. S. 31913 Santos, Hitchcock*. In open grassy places, altitude 1,400 to 2,300 m. A native of the west coast of North and South America, introduced in Portugal, and a recent introduction in Benguet. It is very similar to *Festuca myuros* Linn. and by some authors is treated as a variety of that species. The identification is by Professor Hitchcock.

68. MONOSTACHYA Merrill

MONOSTACHYA CENTROLEPIDOIDES Merr. in Philip. Journ. Sci. 5 (1910) Bot. 331.

Luzon (Benguet), *F. B. 16088 Curran, Merritt, & Zschokke*. In open grasslands near the summit of Mt. Polog, altitude about 2,800 m. A monotypic endemic genus.

69. RAMOSIA Merrill

RAMOSIA PHILIPPINENSIS Merr. in Philip. Journ. Sci. 11 (1916) Bot. 2.

Luzon (Sorsogon), *B. S. 23607 Ramos*. In swampy places, border of Lake Polog, at medium altitudes. A monotypic endemic genus.

70. BROMUS Linnaeus

BROMUS PAUCIFLORUS (Thunb.) Hack. in Bull. Herb. Boiss. 7 (1899) 713; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 386.

Festuca pauciflora Thunb. Fl. Jap. (1784) 52.

Luzon (Bontoc, Benguet), *Merrill 4714, Clemens 9204, 9091, Vanoverbergh 3938*. In wet open grassy places, altitude 1,500 to 2,400 m. Japan, China.

Bromus commutatus Schrad. is represented by *B. S. 31966 Santos* and *Merrill 6627*, cultivated at Pauai, Benguet; introduced as *B. inermis* Leyss.

DOUBTFUL SPECIES

BROMUS PALLENS Cav. Ic. 6 (1801) 66, *t. 591, f. 1*; Kunth. Enum. 1 (1833) 418; Miq. Fl. Ind. Bat. 3 (1857) 398; F.-Vill. Novis. App. (1882) 323.

"Habitat in Manilae vicinis, ibique *Acanthus ilicifolius* Née legit." The figure and description appertain to *Bromus*, but no representative of the genus is known to occur at low altitudes in the Philippines. The description was probably based on an erroneously localized plant.

BROMUS LUZONENSIS Presl Rel. Haenk. 1 (1828) 262.

Triticum luzonense Kunth Enum. 1 (1833) 446; Miq. Fl. Ind. Bat. 3 (1857) 402; F.-Vill. Novis. App. (1882) 323.

"Hab. in Luzonia" Presl. The type was certainly an erroneously localized specimen, probably from tropical America.

71. **BRACHYPODIUM** Beauvois

BRACHYPODIUM SYLVATICUM Beauv. Agrost. (1812) 146, subsp. **LUZONIENSE** Hack. in Philip. Journ. Sci. 1 (1906) Suppl. 269; Merr. op. cit. 387, 5 (1910) Bot. 332.

Luzon (Benguet), *Merrill 4710, 4536, 4830, F. B. 16187 Curran, Merritt, & Zschokke*. On open slopes and in the mossy forest, altitude 1,600 to 2,500 m.

Var. **ASPERUM** Hack. l. c.; Merr. l. c.

Luzon (Benguet), *Merrill 4698*. On steep slopes in damp, shaded ravines, altitude about 1,700 m. The two Philippine forms endemic, the species Europe and northern Africa to India and Ceylon.

72. **MONERMA** Beauvois

MONERMA REPENS (Forst.) Beauv. Agrost. (1812) 117; Hack. in Govt. Lab. Publ. (Philip.) 35 (1906) 81; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 387.

Rottboellia repens Forst. Prodr. (1786) 9.

Lepturus repens R. Br. Prodr. (1910) 207.

Mindoro, Culion, Bancoran, Cajoacan, Palmas, Mindanao, *Merrill 606, 3331, 5215, 5354, 11644, Phil. Pl. 153*. In sandy soil back of the beach, about sea cliffs, etc. Ceylon through Malaya to tropical Australia and Polynesia.

73. **TRITICUM** Linnaeus

TRITICUM VULGARE Vill. Hist. Pl. Dauph. 2 (1779) 153; Nees in Hook. Journ. Bot. Kew Miscel. 2 (1850) 102; F.-Vill. Novis. App. (1882) 323; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 294; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 388.

Formerly cultivated in certain provinces in Luzon, now planted to a limited extent in northern Luzon. Wheat.

74. ARUNDINARIA Michaux

ARUNDINARIA NIITAKAYAMENSIS Hayata in Bot. Mag. Tokyo 21 (1907) 49, Journ. Coll. Sci. Tokyo 25¹⁹ (1908) 240; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 267; Merr. op. cit. 332.

Sasa niitakayamensis Camus var. *microcarpa* Camus Bamb. (1913) 24.

Bambusa aff. *B. pygmaea* Miq.; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 261.

Luzon (Benguet), Mindoro, *Elmer 14279, Merrill 4733, 6489, 6222, Phil. Pl. 177, B. S. 8379, 8893 McGregor, 5846 Ramos, F. B. 10823 Curran*. In mossy thickets, sphagnum swamps, and open grasslands, altitude 2,200 to 2,700 m. Formosa.

Local name: Útod (Ig.).

75. BAMBUSA Schreber

BAMBUSA CORNUTA Munro in Trans. Linn. Soc. 26 (1868) 113; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 269; 8 (1913) Bot. 204.

Luzon (Benguet, Nueva Vizcaya), *Merrill 124, 7711*. In ravines along small streams, altitude 1,000 to 1,600 m. Java.

Local name: Lopa (Isn.).

BAMBUSA MERRILLII Gamble in Philip. Journ. Sci. 5 (1910) Bot. 269; Camus Bamb. (1913) 128.

Luzon (Nueva Vizcaya), *Merrill 229*. In forests, altitude about 600 m. Endemic.

BAMBUSA MULTIPLEX (Lour.) Raeusch. Nomencl. ed. 3 (1797) 108; Schultes f. Syst. 7 (1830) 1350; Steud. Syn. Pl. Glum. 1 (1854) 330; Munro in Trans. Linn. Soc. 26 (1868) 119.

Arundo multiplex Lour. Fl. Cochinch. (1790) 58, ed. Willd. (1793) 119.

Ludolphia glaucescens Willd. in Ges. Naturf. Fr. Berl. Mag. 2 (1808) 320.

Bambusa nana Roxb. Hort. Beng. (1814) 25, *nomen nudum*, Fl. Ind. ed. 2, 2 (1832) 199; F.-Vill. Novis. App. (1882) 323; Gamble in Ann. Bot. Gard. Calc. 7 (1896) 40, t. 38, Philip. Journ. Sci. 5 (1910) Bot. 268; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 389; Camus Bamb. (1913) 121.

Arundinaria glaucescens Beauv. Agrost. (1812) 144, 152.

Bambusa glaucescens Sieb. ex Munro in Trans. Linn. Soc. 26 (1868) 89, in syn.; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 230, Fl. Manila (1912) 105.

Planted here and there for ornamental purposes, sometimes for hedges. Probably a native of China, now widely distributed in cultivation.

Local names: Caña de china (Sp.); kauáyan-china (Bik., Tag.); kauáyan-sína (Bik., Tag.).

BAMBUSA SPINOSA Roxb. Hort. Beng. (1814) 25 [non Fl. Ind. ed. 2, 2 (1832) 305]; Merr. Interp. Herb. Amb. (1917) 97, Sp. Blancoanae (1918) 75.

Bambusa spinosa Blume ex Nees in Flora 8 (1825) 580.

Bambusa blumeana Schultes f. Syst. 7² (1830) 1343; F.-Vill. Novis.

App. (1882) 323; Gamble in Ann. Bot. Gard. Calc. 7 (1896) 50, t. 47; Philip. Journ. Sci. 5 (1910) Bot. 270, 8 (1913) Bot. 204; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 29, 388, Fl. Manila (1912) 106, Am. Journ. Bot. 3 (1916) 60; Camus Bamb. (1913) 130.

Bambus pungens Blanco Fl. Filip. (1837) 270; Miq. Fl. Ind. Bat. 3 (1857) 421; Munro in Trans. Linn. Soc. 26 (1868) 119.

Bambus arundo Blanco Fl. Filip. ed. 2 (1845) 188; ed. 3, 1 (1877) 335, t. 100, non Nees, nec Wight.

Bambusa arundinacea F.-Vill. Novis. App. (1882) 323, non Retz.

Bambusa teba Miq. Fl. Ind. Bat. 3 (1857) 418; Munro in Trans. Linn. Soc. 26 (1868) 122.

Northern Luzon to Palawan and Mindanao. A universally planted bamboo at low and medium altitudes throughout the settled areas; the only spiny bamboo in the Philippines. Southern China to the Malay Peninsula and Archipelago to the Moluccas.

The most common and conspicuous bamboo in the settled areas, but in the Philippines apparently always planted. This indicates its probable prehistoric introduction by the early Malay invaders. The species very rarely flowers and then usually only when clumps have been severely injured by fire or by too extensive cutting.

Local names: Aonóo (Bik.); batákan (Bis.); baugín (Pamp.); dugían (Bik.); caña espina (Sp.); kaaóno (Bis.); kauáyan (P. Bis., Bon., C. Bis., Bik., Ilk., Pamp., Sub., Sul., Tag.); kauáyan-gid (P. Bis.); kauáyan-ñg-bayóg (Ilk.); kauáyan-potog (Sbl.); kauáyan-siitan (Ilk.); kauáyan-tinik (Tag.); kauáyan-totóo (Tag., Bik.); kabugáuan (Bik.); lamnuan (Is.); marurúgi (Bik.); pasíñgan (Ibn., Is.); paua (Bis.); rugían (Bik.).

BAMBUSA VULGARIS Schrad. ex Wendl. Collect. Pl. 2 (1808) 26, t. 47; Gamble in Ann. Bot. Gard. Calcutta 7 (1896) 43, t. 40, Philip. Journ. Sci. 5 (1910) Bot. 268, 8 (1913) Bot. 204; Merr. Fl. Manila (1912) 105, Am. Journ. Bot. 3 (1916) 61, Sp. Blancoanae (1918) 75; Camus Bamb. (1913) 123.

Bambusa monogyna Blanco Fl. Filip. (1837) 286, ed. 2 (1845) 187, ed. 3, 1 (1877) 333; Miq. Fl. Ind. Bat. 3 (1857) 420; Munro in Trans. Linn. Soc. 26 (1868) 119; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 29, 389; Camus Bamb. (1913) 132.

Bambusa mitis Blanco Fl. Filip. (1837) 271, ed. 2 (1845) 188, ed. 3, 1 (1877) 336, non Poir.

Bambusa blancoi Steud. Syn. Pl. Glum. 1 (1854) 331; Miq. Fl. Ind. Bat. 3 (1857) 431; Munro in Trans. Linn. Soc. 26 (1868) 120; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 389; Camus Bamb. (1913) 134.

Dendrocalamus sericeus F.-Vill. Novis. App. (1882) 324, non Munro.

Dendrocalamus strictus F.-Vill. Novis. App. (1882) 324, non Nees.

Northern Luzon to Palawan and Mindanao, in most islands and provinces, in the settled areas at low and medium altitudes. Probably an introduced plant in the Philippines. Pantropic in cultivation.

Local names: Bolínau (P. Bis.); burírau (P. Bis.); bútong (Bik.); kabalóan (Bik.); kauáyan (Ibn.); kauáyan-china (Tag.); kauáyan-kilíng (Ilk., Tag.); kauáyan-kiting (Sbl.); kilíng (Ilk., Tag.); lúnas (P. Bis.);

marobál (Bik.); patong (Tagb.); patung (Sul.); sinámbang (C. Bis.); taiuának (Tag.); taríng (Tagb.); teuának (Tag.); tilíng (P. Bis.).

Var. **STRIATA** (Lodd.) Gamble in Ann. Bot. Gard. Calc. 7 (1896) 44, Philip. Journ. Sci. 5 (1910) Bot. 269; Merr. Fl. Manila (1912) 105.

Bambusa striata Lodd. ex Penny Cyclop. 3 (1835) 357.

Occasionally cultivated for ornamental purposes; probably originating in China.

EXCLUDED SPECIES

BAMBUSA TULDOIDES Munro; F.-Vill. Novis. App. (1882) 323.

76. **GUADUA** Kunth

GUADUA PHILIPPINENSIS Gamble in Philip. Journ. Sci. 8 (1913) Bot. 203.

Mindanao (Davao), *Piper* 475. Probably from thickets or secondary forests; collected at Mati. Endemic. The genus is otherwise American.

77. **GIGANTOCHLOA** Kurz

GIGANTOCHLOA LEVIS (Blanco) Merr. in Am. Journ. Bot. 3 (1916) 61, Sp. Blancoanae (1918) 76.

Bambusa levis Blanco Fl. Filip. (1837) 272, ed. 2 (1845) 189, ed. 3, 1 (1877) 337; Miq. Fl. Ind. Bat. 3 (1857) 421; Munro in Trans. Linn. Soc. 26 (1868) 121; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 93, Philip. Journ. Sci. 1 (1906) Suppl. 388; Camus Bamb. (1913) 134.

Gigantochloa scribneriana Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 390; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 270, 8 (1913) Bot. 204; Camus Bamb. (1913) 143.

Dendrocalamus flagellifer F.-Vill. Novis. App. (1882) 324, non Munro.

Gigantochloa robusta Gamble in Philip. Journ. Sci. 8 (1913) Bot. 204, non ? Kurz.

Gigantochloa atter C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 194, non Kurz.

Dendrocalamus curranii Gamble in Philip. Journ. Sci. 5 (1910) Bot. 271; C. B. Rob. op. cit. 6 (1911) Bot. 194; Camus Bamb. (1913) 158.

Dendrocalamus latiflorus Gamble in Philip. Journ. Sci. 5 (1910) Bot. 291, non Munro.

Luzon (Bulacan, Bataan, Rizal, Laguna, Tayabas), Culion, Polillo, Leyte, Camiguin de Misamis, *Loher* 7208, *Scribner* 14, *Merrill Phil. Pl.* 586, 590, *Sp. Blancoanae* 310, *B. S.* 10414, 10417 *McGregor*, 6313, 11838 *Robinson*, 14696 *Ramos*, *F. B.* 22771 *De Mesa & Rosario*, 10177 *Curran*, 25405 *Labitig*, 25418 *Duldulao*. Usually if not always planted, and probably an introduced species in the Philippines. To be compared critically with the Malayan species.

Local names: Bóko (P. Bis.); bólo (P. Bis.); bonsína (Tag.); bóton (P. Bis.); búhong-china (Tag.); búlung-sína (Tag.); butón (C. Bis.); kabolian (Bik.); kauáyan-china (Tag.); kauáyan-putí (Tag.); kauáyan-sína (Tag.); kayali (C. Bis.); patong (Bik., Bis., Sul.).

78. DENDROCALAMUS Nees

DENDROCALAMUS MERRILLIANUS (Elm.) Elm. Leaf. Philip. Bot. 7 (1915) 2675.

Gigantochloa merrilliana Elm. Leaf. Philip. Bot. 1 (Jan., 1908) 273; Camus Bamb. (1913) 139.

Dendrocalamus parviflorus Hack. in Philip. Journ. Sci. 3 (July, 1908) 168; Gamble op. cit. 5 (1910) Bot. 272; Camus Bamb. (1913) 159.

Luzon (Ilocos Sur, Abra, Nueva Ecija, Rizal, Zambales, Pangasinan, Bulacan), Leyte, Cebu, Bohol, Mindanao (Lanao), *Elmer* 7283, *Clemens*, *Barrow* 30, *F. B.* 27332 *Lopez & Reyes*, 27319, 27329 *Reyes*, 27219 *Duran*, 27177 *Maneja & Bawan*, *B. S.* 22466 *Ramos*, 27706 *Fénix*, 17949 *Otanes*, mostly sterile specimens. In and about towns, chiefly planted. Endemic.

Local names: Bayóg (Ilk., Ibn., Tag., Sbl.); bayúgin (Tag.); botóng (C. Bik., Bis.); bukáui (Tag.); butóng (P. Bis.); kauáyan (Tag., C. Bis., Ilk.); kauáyan-bayóg (Pang.).

EXCLUDED SPECIES

DENDROCALAMUS GIGANTEUS Munro; F.-Vill. Novis. App. (1882) 324.

DENDROCALAMUS LATIFLORUS Munro; F.-Vill. l. c.; Vidal Sinopsis Atlas (1883) 42, t. 96, f. B.

Vidal's figure was copied from Munro's plate, *vide* Vidal l. c.; see *Gigantochloa levis* Merr.

79. CEPHALOSTACHYUM Munro

CEPHALOSTACHYUM MINDORENSE Gamble in Philip. Journ. Sci. 5 (1910) Bot. 272.

Luzon (Camarines), Mindoro, *F. B.* 11421, 8619, 6213 *Merritt*, 10667 *Curran*, 27424 *Ablaza & Ramos*. In thickets and forests at low altitudes, locally abundant. Endemic.

This is presumably the form reported from the Philippines by F.-Villar, Novis. App. (1882) 324, as *C. capitatum* Munro, a species that does not occur in the Archipelago. To be compared here is *Vidal* 4021, reported by Ceron, Cat. Pl. Herb. Manila (1892) 185, as *Oxytenanthera* sp.

Local names: Bagtók (Bik.); baktó (Tag.); bíkal (Bik.).

80. SCHIZOSTACHYUM Nees

SCHIZOSTACHYUM CURRANII Gamble in Philip. Journ. Sci. 5 (1910) Bot. 277; Camus Bamb. (1913) 177.

Luzon (Ifugao, Benguet), *F. B.* 10849 *Curran*, 18014 *Merritt*, *B. S.* 19778 *McGregor*, *Merrill Phil. Pl.* 1217. In borders of the mossy forest, altitude about 2,000 m. Endemic.

SCHIZOSTACHYUM DIELSIIANUM (Pilger) Merr. in Philip. Journ. Sci. 1 Suppl. (1906) 391; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 274, 8 (1913) Bot. 205; Camus Bamb. (1913) 174, t. 96, f. B.

Dinochloa dielsiana Pilger in Perk. Frag. Fl. Philip. (1904) 148.

Palawan, *Merrill* 711. In thickets and forests at low altitudes. Endemic.

By no means certainly distinct from *S. diffusum* Merr. Numerous specimens distributed as *S. dielsianum* Merr. appear to be referable to *S. diffusum*, but many intergrades occur.

SCHIZOSTACHYUM DIFFUSUM (Blanco) Merr. in Am. Journ. Bot. 3 (1916) 62, Sp. Blancoanae (1918) 78.

Bambus diffusa Blanco Fl. Filip. (1837) 269; ed. 2 (1845) 187, ed. 3, 1 (1877) 334; Miq. Fl. Ind. Bat. 3 (1857) 420; Munro in Trans. Linn. Soc. 26 (1868) 118.

Schizostachyum acutiflorum Munro in Trans. Linn. Soc. 26 (1868) 137; F.-Vill. Novis. App. (1882) 324; Vidal Phan. Cuming. Philip. (1885) 159, Rev. Pl. Vasc. Filip. (1886) 294; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 391, 3 (1908) Bot. 397; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 273, 8 (1913) Bot. 205; Camus Bamb. (1913) 174, t. 95, f. A.

Dinochloa diffusa Merr. in Govt. Lab. Publ. (Philip.) 29 (1905) 7.
? *Schizostachyum blumei* F.-Vill. Novis. App. (1882) 324, non Nees.
Ischurochloa floribunda F.-Vill. in Blanco Fl. Filip. ed. 3, 4¹ (1880) 14, non Büse.

Dinochloa major Pilger in Perk. Frag. Fl. Philip. (1904) 149; Camus Bamb. (1913) 170.

Babuyan Islands, Luzon (Cagayan, Isabela, Apayao, Ilocos Norte, Ilocos Sur, Benguet, Nueva Vizcaya, Pangasinan, Zambales, Bulacan, Nueva Ecija, Pampanga, Rizal, Bataan, Laguna, Tayabas, Camarines), Mindoro, Palawan, Leyte. Common in secondary and primary forests at low and medium altitudes, ascending to 1,700 m. Variable. Endemic.

Local names: Baliáro (Ilk.); balíkau (Ilk.); bíkil (Ilk.); bongbong (Bis.); loob (Bis.); indi, inri, hindi (Bik.); butor (Isn.); bábui (Bik.); lilit (Pamp.); bíkal (Ilk., Ibn., Pamp., Pang., Tag.); bíkal-bábi (Pamp.); bíkal-bábui (Sbl.); gimak (Tag.); mangnau (Ilk.); tagisí (Tag.); usíu (Tag.).

SCHIZOSTACHYUM FENIXII Gamble in Philip. Journ. Sci. 6 (1911) Bot. 289, 8 (1913) Bot. 205.

Luzon (Cagayan, Abra, Ilocos Sur, Benguet), Panay, *F. B. 17815 Curran, 23567 Vergara, 25482, 25086 Paraiso, Fénix, McVey*. In thickets at low and medium altitudes. Endemic.

Local names: Bólo (Ilk.); páua (P. Bis.); púser (Ilk.).

SCHIZOSTACHYUM LIMA (Blanco) Merr. in Am. Journ. Bot. 3 (1916) 62, Sp. Blancoanae (1918) 77.

Bambus lima Blanco Fl. Filip. (1837) 271; ed. 2 (1845) 189; ed. 3, 1 (1877) 336; Miq. Fl. Ind. Bat. 3 (1857) 421; Munro in Trans. Linn. Soc. 26 (1868) 121; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 388; Camus Bamb. (1913) 134.

Bambusa longinodis F.-Vill. Novis. App. (1882) 323, non ? Miq.

Schizostachyum hallieri Gamble in Philip. Journ. Sci. 5 (1910) Bot. 274, 8 (1913) Bot. 205; Camus Bamb. (1913) 175, t. 97, f. B.

Schizostachyum brachycladum Gamble in Philip. Journ. Sci. 8 (1913) Bot. 206, non Kurz.

Schizostachyum longispiculatum Gamble in Philip. Journ. Sci. 8 (1913) Bot. 206, non Kurz.

Luzon (Nueva Vizcaya, Bataan, Laguna, Sorsogon), Mindoro, Palawan, Leyte, Mindanao, Basilan, *Hallier, Merrill 581, 9226, Sp. Blancoanae 41, Phil. Pl. 1472, as S. longispiculatum Kurz, Elmer 7145, 10838, 12958, 12599, 16621, Wenzel 1357, Weber 1111, Wester 145, B. S. 20120 McGregor,*

F. B. 11374 Merritt. In thickets and secondary forests at low and medium altitudes, sometimes gregarious; a species having unusually long internodes, these being up to 120 cm in length. Endemic.

Local names: Ános (Tag.); bagákai (C. Bis., P. Bis., Bik.); bitu (Pamp.); bolo (Ilk.); buho (P. Bis.); lakap (Buk.); nap-nap (C. Bis.); sumbiling (Tagb.).

SCHIZOSTACHYUM LUMAMPAO (Blanco) Merr. in Am. Journ. Bot. 3 (1916) 65, Sp. Blancoanae (1918) 77.

Bambus lumampao Blanco Fl. Filip. (1837) 272, ed. 2 (1845) 189; ed. 3, 1 (1877) 338; Miq. Fl. Ind. Bat. 3 (1857) 421; Munro in Trans. Linn. Soc. 26 (1868) 118; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 389; Camus Bamb. (1913) 132.

Dendrocalamus membranaceus F.-Vill. Novis. App. (1882) 324, non Munro.

Schizostachyum mucronatum Hack. in Philip. Journ. Sci. 3 (1908) Bot. 169; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 276; Camus Bamb. (1913) 175.

Schizostachyum hirtiflorum Hack. in Philip. Journ. Sci. 2 (1907) Bot. 420; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 275; Camus Bamb. (1913) 176.

Luzon (Cagayan, Isabela, Ilocos Norte, Ilocos Sur, Benguet, Pangasinan, Pampanga, Rizal, Bataan, Tayabas, Camarines), Panay, *Merrill Sp. Blancoanae* 891, *Elmer* 6173, *B. S.* 7825 *Ramos*, 2310 *Mearns*, 29875 *Fénix*, *F. B.* 21926 *Alejandro*, 26058 *Barros*, 25829 *Paraíso*, 22808 *Díaz & Alambra*, 24113 *Velasco*, 23560 *Vergara*, 20945 *De Leon*, 20870 *Bernardo*, 10414 *Curran*, 3615 *Maule*. In secondary forests, but more frequently gregarious and occupying extensive areas at low and medium altitudes where forests have been destroyed. Endemic.

Local names: Bakákan (Tagb.); bóho (Tag., Bik.); bokáui (Tag.); bólo (Tag., Ilk., Pang., Bik., P. Bis., Ibn.); búlo (Ibn.); caña bóho (Sp.); dasó (Bik.); lumampáu (Tag.); napnáp (Ilk.); oras (Bik.); valú (Ibn.); vúlu (It.).

SCHIZOSTACHYUM LUZONICUM Gamble in Philip. Journ. Sci. 5 (1910) Bot. 277; Camus Bamb. (1913) 177.

Luzon (Zambales), *F. B.* 8411 *Curran & Merritt*, 5926 *Curran*. In forests and in ravines, altitude about 800 m. Endemic.

SCHIZOSTACHYUM PALAWANENSE Gamble in Philip. Journ. Sci. 5 (1910) Bot. 274, 8 (1913) Bot. 205; Camus Bamb. (1913) 175.

Palawan, *B. S.* 277 *Bermejós*, *Elmer* 12618. In forests at low altitudes. Endemic.

Local name: Bugto (Tagb.).

SCHIZOSTACHYUM TEXTORIUM (Blanco) Merr. in Am. Journ. Bot. 3 (1916) 64, Sp. Blancoanae (1918) 76.

Bambus textoria Blanco Fl. Filip. (1837) 270; ed. 2 (1845) 188; ed. 3, 1 (1877) 335; Miq. Fl. Ind. Bat. 3 (1857) 421; Munro in Trans. Linn. Soc. 26 (1868) 122; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 389; Camus Bamb. (1913) 135.

Gigantochloa atter F.-Vill. Novis. App. (1882) 323, non Kurz.

Schizostachyum merrillii Gamble in Philip. Journ. Sci. 5 (1910) Bot. 278; Camus Bamb. (1913) 174.

Luzon (Batangas, Rizal), *Merrill 1744, Sp. Blancoanae 714*. In secondary forests at low altitudes. Endemic.

Local name: Kalbáng (Tag.).

SCHIZOSTACHYUM TOPPINGII Gamble in Philip. Journ. Sci. 5 (1910) Bot. 276; Merr. Fl. Manila (1912) 106; Camus Bamb. (1913) 178.

Luzon (Nueva Vizcaya, Rizal), Mindoro, *B. S. 5222 Topping, F. B. 9500 Merritt, B. S. 39366, 39713 Ramos*. In thickets and secondary forests at low and medium altitudes. Endemic.

81. DINOCHLOA Büse

DINOCHLOA ELMERI Gamble in Philip. Journ. Sci. 5 (1910) Bot. 280; Camus Bamb. (1913) 170.

Luzon (Benguet), *Elmer 6542*. In the mossy forests on Mt. Santo Tomas, altitude about 2,200 m. Endemic.

DINOCHLOA LUÇONIAE (Munro) comb. nov.

Bambusa luçoninae Munro in Trans. Linn. Soc. 26 (1868) 115; F.-Vill.

Novis. App. (1882) 323 (*luçoninae*); Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 389; Camus Bamb. (1913) 134.

Dinochloa aguilarii Gamble in Philip. Journ. Sci. 5 (1910) Bot. 280; Camus Bamb. (1913) 170.

Dinochloa ciliata Kurz in Journ. As. Soc. Beng. 42² (1873) 253, in nota; Camus Bamb. (1913) 170, t. 95, f. D; Merr. in Philip. Journ. Sci. 11 (1916) Bot. 51.

Dinochloa scandens O. Kuntze var. *angustifolia* Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 392; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 279, 8 (1913) Bot. 206; Camus Bamb. (1913) 169.

Dinochloa tjankorreh Büse var. *angustifolia* Hack. ex Merr. l. c. in syn.

Luzon (Cagayan, Ifugao, Ilocos Norte, Nueva Ecija, Bulacan, Zambales, Cavite, Bataan, Laguna, Tayabas), Mindoro, Leyte, Samar, Mindanao, Basilan. In forests at low and medium altitudes. Borneo.

I have examined the type of *Bambusa luçoninae* Munro, which was from Mt. Banahao, and which is preserved in the United States National Herbarium. I fail to see how *D. aguilarii* Gamble can be distinguished from *D. ciliata* Kurz, or either from *D. luçoninae* Merr.

Local names: Bika (Ibn.); bíkal (Ibn.); buloau (Sub.); bolokáui (C. Bis.); malilit (Sbl.); osíu (Tag.); timak (Tag.); usíu (Tag.).

DINOCHLOA PUBIRAMEA Gamble in Philip. Journ. Sci. 5 (1910) Bot. 279; Camus Bamb. (1913) 171.

Dinochloa scandens O. Kuntze var. *pubiramea* Merr. ex Gamble l. c. in syn.

Leyte, Samar, Negros, Mindanao (Surigao), Basilan, *Bolster 319, Wenzel 1310, F. B. 5589 Everett, 26268 Mallonga, B. S. 17639 Ramos*. In forests at low and medium altitudes. Borneo.

Local name: Bukau (Yak.).

DINOCHLOA SCANDENS (Blume) O. Kuntze Rev. Gen. Pl. (1891) 773; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 392; Gamble in Philip. Journ. Sci. 5 (1910) Bot. 278, 8 (1913) Bot. 206; Camus Bamb. (1913) 169.

Bambusa scandens Blume ex Nees in Flora 7 (1824) 291.

Nastus tjangkorreh Schultes Syst. Veg. 7² (1830) 1358.

Dinochloa tjankorreh Büse in Miq. Pl. Jungh. (1854) 388; Miq. Fl. Ind. Bat. 3 (1857) 415; Munro in Trans. Linn. Soc. 26 (1868) 153; F.-Vill. Novis. App. (1882) 324; Vidal Sinopsis Atlas (1883) 42, t. 96, f. C, Phan. Cuming. Philip. (1885) 157, Rev. Pl. Vasc. Philip. (1886) 294; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 29.

Dinochloa macrocarpa Elm. Leaf. Philip. Bot. 7 (1915) 2675.

Luzon (Bulacan, Laguna, Camarines, Sorsogon), Polillo, Mindoro, Palawan, Balabac, Samar, Panay, Sibuyan, Biliran, Leyte, Mindanao, Basilan. Common in primary forests at low and medium altitudes. Malay Peninsula and Archipelago.

Local names: Baká (Tagb.); balikau (Tagb.); biá (Tagb.); biká (Bon.); balilit (Bag.); bayokáui (P. Bis.); bukáu (Yak.); bukáui (Tag.); bulukáui (P. Bis.); burukáui (S. L. Bis.); timak (Mang.); usíu (Bik., Tag.).

EXCLUDED GENERA AND SPECIES

LAPPAGO RACEMOSA Houck.; F.-Vill. Novis. App. (1882) 313.

BEESEA RHEEDII Kunth; Vidal Cat. Pl. Prov. Manila (1880) 48; F.-Vill. Novis. App. (1882) 324.

PHYLLSTACHYS BAMBUSOIDES Sieb.; F.-Vill. Novis. App. (1882) 323.

PHYLLSTACHYS NIGRA Munro; F.-Vill. l. c.

PAPPOPHORUM PHLEOIDES (Cav.) R. & S.

"Arab. vel ins. Philipp.?" Steud. Syn. Pl. Glum. 1 (1854) 199. 'This is an American species erroneously credited to the Philippines by Steudel.

STENOTAPHRUM COMPLANATUM Schrank.; F.-Vill. Novis. App. (1882) 313.

TEINOSTACHYUM sp.; Gamble in Philip. Journ. Sci. 8 (1913) Bot. 205.

This reference was based on a sterile specimen of what will probably prove to be a species of *Schizostachyum*.

Several species of the Chlorideae, chiefly of the genus *Bouteloua*, have been erroneously credited to the Philippines on the basis of American specimens collected by members of the Malaspina Expedition and erroneously localized as Philippine. They are:

BOUTELOUA CURTIPENDULA (Mich.) Torr.; *Eutriana curtispindula* Trin.; Kunth Enum. 1 (1833) 280; Miq. Fl. Ind. Bat. 3 (1857) 383; F.-Vill. Novis. App. (1882) 320=*Bouteloua racemosa* Lagasca.

BOUTELOUA BARBATA Lagasca; *Eutriana barbata* Kunth Rev. Gram. 1 (1829) 96; Enum. 1 (1833) 282; Miq. Fl. Ind. Bat. 3 (1857) 284; F.-Vill. Novis. App. (1882) 321.

BOUTELOUA TENUIS Griseb.; *Chondrosium tenue* Beauv. Agrost. (1812) 41; Kunth Enum. 1 (1833) 76; F.-Vill. Novis. App. (1882) 320.

BOUTELOUA SIMPLEX Lagasca; *Chondrosium simplex* Kunth Enum. 1 276; F.-Vill. Novis. App. (1882) 320.

POLYSCHISTIS PAUPERCULA Presl. Rel. Haenk. 1 (1830) 294, *t.* 41, *f.* 2; Kunth Enum. 1 (1833) 282; Miq. Fl. Ind. Bat. 3 (1859) 384; F.-Vill. Novis. App. (1882) 321.

"Hab. in insula Luzonia" Presl. The specimens were erroneously localized; a species of tropical America = *Pentarrhaphis* sp.

ATHEROPOGON PROCUMBENS Jacq.; R. & S. Syst. 2 (1825) 292.

"Ins. Philipp. Mexico" Roemer and Schultes. This is one of the American species of *Bouteloua*.

CYPERACEAE

1. LIPOCARPHA R. Brown

LIPOCARPHA ARGENTEA R. Br. in Tuckey's Congo Append. (1818) 459; Miq. Fl. Ind. Bat. 3 (1856) 331; F.-Vill. Novis. App. (1882) 308; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 285; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 109.

Lipocarpa laevigata Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28.

Kyllinga albescens Steud. Syn. Pl. Glum. 2 (1855) 68; Miq. Fl. Ind. Bat. 3 (1856) 294; Naves Novis. App. (1882) 300.

Luzon (Cagayan, Lepanto, Bontoc, Benguet, Nueva Vizcaya, Nueva Ecija), Mindanao (Bukidnon), *B. S.* 26007, 26151 *Fénix*, 14252 *McGregor*, 7864 *Ramos*, *Merrill* 215, *Phil. Pl.* 570, *Elmer* 6484, *Williams* 1224. In open wet places chiefly at medium altitudes, ascending to 1,800 m. Warmer parts of the Old World.

Local name: Balábak (Buk.).

LIPOCARPHA MICROCEPHALA (R. Br.) Kunth Enum. 2 (1837) 268; Miq. Fl. Ind. Bat. 3 (1856) 331; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 668, Philip. Journ. Sci. 2 (1907) Bot. 110; F.-Vill. Novis. App. (1882) 308; Merr. Fl. Manila (1912) 119.

Hypaelyptum microcephalum R. Br. Prodr. (1810) 220.

Ascolepis kyllingioides Steud. in Zoll. Verz. Ind. Archip. (1854) 61.

Kyllinga squarrosa Steud. Syn. Pl. Glum. 2 (1855) 68.

Luzon (Apayao, Ifugao, Bulacan, Rizal), *Merrill* 3625, *Phil. Pl.* 547, *B. S.* 21712, 12177 *Ramos*, 28174 *Fénix*. In open wet grasslands at low and medium altitudes. Japan to the Malay Peninsula and tropical Australia.

2. HYPOLYTRUM L. C. Richard

HYPOLYTRUM COMPACTUM Nees in Linnaea 9 (1834) 288, Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 73; Steud. Syn. Pl. Glum. 2 (1855) 132; Miq. Fl. Ind. Bat. 3 (1856) 333; Boeck. in Linnaea 37 (1871) 127; F.-Vill. Novis. App. (1882) 309; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 29; C. B. Clarke op. cit. 2 (1907) Bot. 109.

Luzon (Benguet, Pangasinan, Zambales, Nueva Vizcaya, Bulacan, Bataan), Palawan, *B. S.* 29983 *Fénix*, 29537, 29428 *Ramos & Edaño*, *Elmer* 8889, 13041, *Merrill* 2496, 7597, 5184, *Phil. Pl.* 1434. In forests at low and medium altitudes, often common. Indo-China, Andaman Islands, Borneo.

Local name: Túhog-dalág (Tag.).

HYPOLYTRUM LATIFOLIUM L. C. Rich. in Pers. Syn. 1 (1805) 70; Hook. f. in Curtis's Bot. Mag. III 33 (1877) t. 6282; F.-Vill. Novis. App. (1882) 309; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 108; Merr. op. cit. 264.

Scirpus anomalus Retz. Obs. 5 (1789) 15, non *Hypolytrum anomalum* Steud.

Albikia schoenoides Presl Rel. Haenk. 1 (1828) 185, t. 34.

Hypolytrum schoenoides Nees in Linnaea 9 (1834) 288.

Hypolytrum trinervium Kunth Enum. 2 (1837) 272; Miq. Fl. Ind. Bat. 3 (1856) 332; F.-Vill. Novis. App. (1882) 308.

Luzon (Tayabas, Pangasinan, Camarines), Palawan, Biliran, Panay, Mindanao, Basilan, Merrill 9449, Phil. Pl. 537, F. B. 20186 Aguilar, B. S. 13115 Foxworthy & Ramos, 18561 McGregor, 34841 Ramos & Pascasio, 17787 Otanes. In forests at low and medium altitudes. India to Formosa through Malaya to Polynesia.

Local names: Baliis (Tag.); bubugo (Sub.); marakibkib (Neg.); parai-biau (Sul.); root (Neg.).

HYPOLYTRUM SCIRPOIDES (Presl) comb. nov.

Albikia scirpoides Presl Rel. Haenk. 1 (1828) 185, t. 35, excl. syn.

Hypolytrum philippense C. B. Clarke in Philip. Journ. Sci. 2 (1908) Bot. 109.

Hypolytrum viridinux C. B. Clarke l. c.

Throughout the Philippines, Elmer 6223, 15916, 18198; Merrill 7291, 10430, 744, 5126, 8186, B. S. 29589, 28654, 28462 Ramos & Edaña, 34394 Ramos & Pascasio, 28164 Fénix, 23601, 13929, 20424, 15314, 1853, 7389, 33162 Ramos, 14238, 32473 McGregor. In forests at low and medium altitudes; often common. Borneo.

Local name: Alisáhis (Buk.).

The Philippine form Presl figured and described is unmistakably the same as *H. philippense* C. B. Clarke; Presl erred in referring to it *H. nemorum* Beauv. and *Schoenus nemorum* Vahl.

3. CYPERUS Linnaeus

CYPERUS COMPRESSUS Linn. Sp. Pl. (1753) 46; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 56; Hook. Journ. Bot. Kew Miscel. 6 (1854) 27; Naves Novis. App. (1882) 302; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 282; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 84; Merr. op. cit. 3 (1908) Bot. 398, Fl. Manila (1912) 111, Sp. Blancoanae (1918) 79.

Cyperus humilis Llanos Frag. Pl. Filip. (1851) 13; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 7, non Kunth.

Cyperus compressus Linn. var. *brachiatatus* Nees in Linnaea 35 (1868) 518.

Cyperus meyenii Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 57.

Throughout the Philippines in open grasslands, waste places, etc., at low and medium altitudes; common. Pantropic.

Local names: Ananá (Ibn.); gísai-kalabáu (Tag.); kaptos (Iv.); túhog-dalág (Tag.); sangsaṅga (Bon.).

- CYPERUS DIFFORMIS** Linn. Amoen. Acad. 4 (1759) 302; Presl Rel. Haenk. 1 (1828) 69; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 70, Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; Miq. Fl. Ind. Bat. 3 (1856) 269; Naves Novis. App. (1882) 302; Vidal Phan. Cuming. Philip. (1855) 155, Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 82; Merr. Fl. Manila (1912) 110, Sp. Blancoanae (1918) 78.
- Cyperus subrotundus* Naves Frag. Pl. Filip. (1851) 14; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 8.
- Cyperus goeringii* Steud. Syn. Pl. Glum. 2 (1855) 24; Naves Novis. App. (1882) 303.

Throughout the Philippines in open wet places, ascending to 2,000 m; a characteristic rice-paddy weed. Old World Tropics generally, introduced in Mexico.

Local names: Baki-báki (Bis.); ballayang (Ilk.); bankoan (Bis.); gilamhon (Bis.); pukuugan (Bon.).

- CYPERUS DIFFUSUS** Vahl Enum. 2 (1806) 321; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 58, Hook. Journ. Bot. Kew Miscel. 6 (1854) 28 (var. *longifolius* Nees); Miq. Fl. Ind. Bat. 3 (1856) 264; Naves Novis. App. (1882) 302; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 83; Merr. op. cit. 262, Fl. Manila (1912) 110.

Cyperus scirpoideus Presl Rel. Haenk. 1 (1828) 178, non Vahl.

Cyperus sorzogonensis Presl op. cit. 351.

Cyperus longifolius Dcne. in Nuov. Ann. Mus. Paris 3 (1834) 359; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28.

Throughout the Philippines, in thickets, secondary and primary forests, and in old clearings at low and medium altitudes; common. Pantropic.

Local names: Barsaṅgá-bákir (Ilk.); gumi-gúmi (Tag.); haras (Tag.); nipinpin-di-gúbat (Sbl.); túhog-dalág (Tag.).

Var. **PUBESQUAMA** (Steud.) Kükenth. in herb. comb. nov.

Cyperus pubesquama Steud. in Zoll. Verz. Ind. Archipel. (1854) 62;

Miq. Fl. Ind. Bat. 3 (1856) 266; Naves Novis. App. (1882) 302;

C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 83.

Cyperus lagorensis Steud. Syn. Pl. Glum. 2 (1855) 36; Miq. Fl. Ind. Bat. 3 (1856) 275.

Cyperus bancanus Naves Novis. App. (1882) 302; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 30; C. B. Clarke op. cit. 2 (1907) Bot. 83, non Miq.

Cyperus calacaryensis Steud. Syn. Pl. Glum. 2 (1855) 34; Miq. Fl. Ind. Bat. 3 (1856) 275; Naves Novis. App. (1882) 304.

Luzon (Cagayan, Isabela, Bulacan, Rizal, Bataan, Laguna), Mindoro, Leyte, Mindanao, *F. B. 2581 Meyer, B. S. 13951, 8015 Ramos, Merrill 5054, 5692, Phil. Pl. 523*. Habitat of the species from which it is scarcely distinguishable. India to Malaya.

Var. **CELEBICA** Miq. Fl. Ind. Bat. 3 (1856) 283.

Throughout the Philippines, *Elmer 18286, 6674, 8306, Williams 85, B. S. 28839 Ramos & Edaño, 16166 Reillo, F. B. 20172 Aguilar*. Habitat of the species. Malaya.

CYPERUS DIGITATUS Roxb. Hort. Beng. (1814) 81, *nomen nudum*, Fl. Ind. 1 (1820) 209; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 86.

Cyperus auricomus Naves Novis. App. (1882) 305; Ceron Cat. Pl. Herb. Manila (1892) 177, non Sieber.

Babuyan Islands, Luzon (Cagayan, Isabela, Nueva Vizcaya, Rizal, Pangasinana, Laguna), Leyte, Mindanao, *B. S. 1450, 8075, 8151, 13516 Ramos, Weber 1083, Merrill 2317*. In open wet places at low and medium altitudes; widely scattered. Pantropic.

CYPERUS DISTANS Linn. f. Suppl. (1781) 103; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27; Naves Novis. App. (1882) 303; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 84; Merr. op. cit. 3 (1908) Bot. 398, 5 (1910) Bot. 333, Fl. Manila (1912) 111.

Cyperus kurrui Steud. Syn. Pl. Glum. 2 (1855) 38; Miq. Fl. Ind. Bat. 3 (1856) 279; Naves Novis. App. (1882) 305.

Throughout the Philippines. Common in open rather wet places at low and medium altitudes. Pantropic.

Local names: Amúsan (Tag.); bakabakáhan (Tag.); barisañgá (Ilk.); burabongdai (S. L. Bis.); gagauan (Bon.); hañgañga (If.); langalang (Bik.); muthang-kalabáu (Tag.); pulukgalo (Buk.); talúbak (Buk.).

CYPERUS ELATUS Linn. Amoen. Acad. 4 (1759) 301; Naves Novis. App. (1882) 304; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 86.

Cyperus bispicatus Steud. in Zoll. Verzeich. Ind. Archipel. (1854) 62; Naves Novis. App. (1882) 303.

Luzon (Laguna), Biliran, Leyte, *Wenzel 1279, B. S. 14955 Ramos, 23140, 18897 McGregor*, distributed as *C. radiatus* Vahl, and not certainly distinct from Vahl's species. In swamps and open wet places at low altitudes. India to Malaya.

Local name: Kobong-kóbong (Bis.).

CYPERUS ELEUSINOIDES Kunth Enum. 2 (1837) 39; Naves Novis. App. (1882) 303; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 84.

Luzon (Cagayan, Bontoc, Benguet, Nueva Vizcaya), Mindanao (Bukidnon), *Merrill 226, B. S. 24950 Fénix, 22728 Castillo, 8747, 14264 McGregor*. In open wet places at low and medium altitudes, not common. Tropical Asia and Africa through Malaya to tropical Australia.

Local names: Gauang (Bon.); salinñgañga (Bon.).

CYPERUS FLABELLIFORMIS Retth. Descr. Nov. Pl. (1773) 42; Merr. Fl. Manila (1912) 110.

Cultivated for ornamental purposes, occasional in waste places in Manila but scarcely established. A native of Africa and southwestern Asia.

CYPERUS HASPAN Linn. Sp. Pl. (1753) 45; Miq. Fl. Ind. Bat. 3 (1856) 267; Naves Novis. App. (1882) 302; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 82; Merr. op. cit. 3 (1908) Bot. 398, Fl. Manila (1912) 110, Sp. Blancoanae (1918) 79.

Cyperus caespitosus Llanos Frag. Pl. Filip. (1851) 14 (*caespitosus*); F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 8; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 86, non Poir.

Cyperus dehiscens Naves Novis. App. (1882) 303, non Kunth.

Cyperus flavidus C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 82, non Retz.

In open wet places throughout the Philippines; a common and characteristic rice-paddy weed. Pantropic.

I fail to see how Philippine specimens referred to *C. flavidus* Retz. can be distinguished from *C. haspan* Linn.

Local names: Balabalangutan (Tag.); barsaṅgá (Ilk.); buṅgot-buṅgot (P. Bis.); manik-manikan (Tag.); misai-kalabau (Tag.); zaingal (Sub.).

CYPERUS IRIA Linn. Sp. Pl. (1753) 45; Presl Rel. Haenk. 1 (1828) 174; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 70, Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; Naves Novis. App. (1882) 303; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 83 (incl. var. *paniciformis* C. B. Clarke); Merr. Fl. Manila (1912) 111, Sp. Blancoanae (1918) 79.

Cyperus nuttallii Llanos Frag. Pl. Filip. (1851) 14; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 9, non Torr.

Chlorocyperus iria Rikli in Jahrb. Wissensch. Bot. 27 (1895) 564; Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 6; Kneucker Cyp. Exsic. No. 237.

In open wet places throughout the Philippines; a common and characteristic rice-paddy weed. Old World Tropics generally.

Local names: Alinang (Bik.); okokiang (Bon.); paiung-paiung (Bik.); sud-sud (Bik.); taga-taga (Bik.).

CYPERUS LEUCOCEPHALUS Retz. Obs. 5 (1789) 11; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 82.

Sorostachys kyllingioides Steud. in Flora 33 (1850) 229, *nomen nudum*, Syn. Pl. Glum. 2 (1855) 71; Miq. Fl. Ind. Bat. 3 (1856) 296.

Cyperus sorostachys Boeck. in Linnaea 35 (1868) 588; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 48.

Cyperus pulchellus R. Br. Prodr. (1810) 213; Naves Novis. App. (1882) 301.

Mindanao (Misamis), *Cuming* 1617. This does not seem to have been found in the Philippines since Cuming collected it, between 1836 and 1840. Pantropic.

CYPERUS MALACCENSIS Lam. Tabl. Encycl. 1 (1791) 146; Usteri Beitr. Ken. Philip. Veg. (1905) 131; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 30, Fl. Manila (1912) 111; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 84.

Chlorocyperus malaccensis Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 6; Kneucker Cyp. Exsic. No. 238.

Central Luzon to Palawan and Mindanao, on the muddy margins of tidal streams within the influence of salt or brackish water. Tropical Asia through Malaya to Australia and Polynesia.

Local names: Baga-as (P. Bis.); balangót (Tag., Bik., Bis., Pamp.); balongát (Pamp.); barangót (Bik.); talaid (Bag.); tíkog (Mbo.).

CYPERUS PILOSUS Vahl Enum. 2 (1806) 354; Naves Novis. App. (1882) 303; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 84; Merr. Fl. Manila (1912) 111.

Cyperus marginellus Nees in Wight Contr. (1834) 83, Hook. Journ. Bot. Kew Miscel. 6 (1854) 28.

Cyperus pauciflorus Steud. Syn. Pl. Glum. 2 (1855) 34; Miq. Fl. Ind. Bat. 3 (1856) 275; Naves Novis. App. (1882) 304.

Cyperus obliquus Nees in Wight Contr. (1834) 86; Naves Novis. App. (1882) 305.

Duvaljouvea pilosa Palla in Koch Syn. ed. 3 (1905) 2555; Kneucker Cyp. Exsic. No. 245.

In open wet places throughout the Philippines; common. Tropical Asia and Africa through Malaya to Australia.

Local name: Paragi (Sub.).

CYPERUS RADIATUS Vahl Enum. 2 (1806) 369; Naves Novis. App. (1882) 305; Vidal Phan. Cuming. Philip. (1885) 155 (var. *elongata* Boeck.), Rev. Pl. Vasc. Filip. (1886) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 85; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 398, Fl. Manila (1912) 112, Sp. Blancoanae (1918) 78.

Cyperus racemosus Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27; Naves Novis. App. (1882) 303, ? non Retz.

Cyperus macrosciadion Steud. Syn. Pl. Glum. 2 (1855) 37; Miq. Fl. Ind. Bat. 3 (1856) 277; Naves Novis. App. (1882) 305.

Cyperus anabaptistus Steud. Syn. Pl. Glum. 2 (1855) 37; Miq. Fl. Ind. Bat. 3 (1856) 277; Naves Novis. App. (1882) 305; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 86, ex descr.

Cyperus cumingii Steud. l. c. in syn.

Cyperus imbricatus Llanos Frag. Pl. Filip. (1851) 17; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 11, non Retz.

Cyperus spicatus Presl Rel. Haenk. 1 (1828) 173; Steud. Syn. Pl. Glum. 2 (1855) 52; Miq. Fl. Ind. Bat. 3 (1856) 287; Naves Novis. App. (1882) 306; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 87, ex descr.

Dichostylis radiatus Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 8, Kneucker Cyp. Exsic. No. 247.

Throughout the Philippines in open wet places at low and medium altitudes. Pantropic.

Local names: Alinang (P. Bis.); balabalanġutan (Tag.); balayang (Ilk.); dagko (Bis.); obod-obod (Bis.); upopi (Ibn.).

CYPERUS ROTUNDUS Linn. Sp. Pl. (1753) 45; Presl Rel. Haenk. 1 (1828) 175; Blanco Fl. Filip. (1837) 31, ed. 2 (1845) 21, ed. 3, 1 (1877) 40; Miq. Fl. Ind. Bat. 3 (1856) 274; Naves Novis. App. (1882) 304; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 282; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 85; Merr. Fl. Manila (1912) 112, Sp. Blancoanae (1918) 79.

Cyperus hexastachyus Rottb. Descr. Ic. (1773) 28, t. 14 f. 2; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 60, var. *castaneus* Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27.

Cyperus curvatus Llanos Frag. Pl. Filip. (1851) 15; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 9, non Vahl.

Throughout the Philippines in open areas at low and medium altitudes. Common in lawns, along roads, and in waste places. Pantropic.

Local names: Barsaṅgá (Ilk.); boto-botónes (Bik.); galonálpas (Pamp.); kungsi (Pamp.); malaapúlíd (Pamp.); motá (Pamp.); muthá (Tag.); omadiung (Pamp.); onoran (Pamp.); sur-sur (Pamp.); tarugug (Bik.).

CYPERUS STOLONIFERUS Retz. Obs. 4 (1786) 10; Presl Rel. Haenk. 1 (1828) 169; Boeck. in Linnaea 35 (1868) 489; Naves Novis. App. (1882) 302; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 85.

Luzon (Rizal), Panay, *McGregor 41, Copeland*. In open places at low altitudes; rare and local. Tropical Asia to Mauritius, Formosa, and through Malaya to tropical Australia.

CYPERUS UNCINATUS Poir. in Lam. Encycl. 7 (1806) 247; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 82.

Cyperus cuspidatus HBK. Nov. Gen. Sp. Pl. 1 (1815) 204; Naves Novis. App. (1882) 301; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 282.

Cyperus solutus Steud. Syn. Pl. Glum. 2 (1855) 14; Miq. Fl. Ind. Bat. 3 (1856) 263; Naves Novis. App. (1882) 302.

Cyperus pusillus Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27, non ? Vahl.

Luzon (Bontoc, Benguet, Bulacan, Rizal, Laguna), Mindanao (Lanao), *Merrill Phil. Pl. 1488, Williams 1975bis, 1969bis, B. S. 19213 Ramos*. In open lands, old clearings, etc., at low and medium altitudes. Pantropic.

Local name: Salasa (Tag.).

CYPERUS ZOLLINGERI Steud. in Zoll. Verz. Ind. Archipel. (1854) 62; Syn. Pl. Glum. 2 (1855) 17; Miq. Fl. Ind. Bat. 3 (1856) 264; Naves Novis. App. (1882) 302; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 85; Merr. Fl. Manila (1912) 112.

Cyperus tenuiculmis Boeck. in Linnaea 36 (1869) 286; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 50; Naves Novis. App. (1882) 306.

Luzon (Cagayan, Ilocos Norte, Benguet, Rizal, Sorsogon), Palawan, Panay, Mindanao, *Elmer 16488, 6473, Merrill 7667, 9310, Phil. Pl. 530, B. S. 32583 McGregor, 29513 Ramos & Edaño, 1106, 4636, 7672, 7820 Ramos*. In open grasslands at low and medium altitudes, ascending to 1,600 m. Tropical Asia and Africa through Malaya to tropical Australia.

Local names: Dáat-láua (Tag.); pakama (Bag.).

DOUBTFUL AND EXCLUDED SPECIES

CYPERUS BULBOSUS Vahl; Naves Novis. App. (1882) 304.

CYPERUS CASTANEUS Willd.; Naves Novis. App. (1882) 301.

CYPERUS CEPHALOTES Vahl; Naves Novis. App. (1882) 301.

CYPERUS CORYMBOSUS Rotth.; Naves Novis. App. (1882) 303.

CYPERUS ESCULENTUS Linn.; Naves Novis. App. (1882) 304.

CYPERUS EUMORPHUS Steud.; Naves Novis. App. (1882) 303.

CYPERUS LAEVIGATUS Linn.; Naves Novis. App. (1882) 301.

CYPERUS LONGUS Linn.; Naves Novis. App. (1882) 304.

CYPERUS ORNATUS R. Br.; Naves Novis. App. (1882) 303.

CYPERUS PANGOREI Rottb.; Naves Novis. App. (1882) 304.

CYPERUS RADIANS Nees; Naves Novis. App. (1882) 305.

CYPERUS SEXFLORUS R. Br.; Naves Novis. App. (1882) 304.

CYPERUS TEGETIFORMIS Roxb.; Naves Novis. App. (1882) 303.

CYPERUS CLAVATUS Lam.; Usteri Beitr. Ken. Philip. Veg. (1905) 131.

A lapsus calami; no such name occurs in botanical literature.

CYPERUS CRUENTUS Rottb.; Boeck. in Linnaea 36 (1869) 338; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 51; Naves Novis. App. (1882) 305.

The Philippine record was based on *Cuming* 2372, which was from Malacca, not from the Philippines.

CYPERUS HAEMATODES Endl.; Steud. Syn. Pl. Glum. 2 (1855) 44; Miq. Fl. Ind. Bat. 3 (1856) 282; Naves Novis. App. (1882) 305.

The type was from Norfolk Island; the Philippine record is due to Steudel's error in transcribing the type locality as "Ins. Norf. Philipp."

CYPERUS LAMPROCARPUS Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27; Boeck. in Linnaea 35 (1868) 490; Naves Novis. App. (1882) 306.

The type is *Cuming* 2437, which was from Sumatra, not from the Philippines. It is supposed to be a synonym of *Cyperus stoloniferus* Retz.

CYPERUS TUBEROSUS Rottb.; Miq. Fl. Ind. Bat. 3 (1856) 265.

"Luzon ?" *fide* Miquel l. c. The record must be an erroneous one as the species is not known from the Philippines.

CYPERUS MINUTIFLORUS Presl Rel. Haenk. 1 (1830) 351; Steud. Syn. Pl. Glum. 2 (1855) 52; Miq. Fl. Ind. Bat. 3 (1856) 287; Naves Novis. App. (1882) 306; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 87.

Cyperus micranthus Presl Rel. Haenk. 1 (1828) 178, non Nees.

Cyperus breviflorus Dietr. Sp. Pl. 2 (1833) 316.

Cyperus multiflorus Kunth Enum. 2 (1837) 562 err. typ.

"Hab. in insula Luzon."

CYPERUS PHILIPPENSIS Presl Rel. Haenk. 1 (1828) 174; Steud. Syn. Pl. Glum. 2 (1855) 52; Miq. Fl. Ind. Bat. 3 (1856) 287; Naves Novis. App. (1882) 306; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 87.

"Hab. in insula Luzon." It is desirable that the types of the last two be examined as from the description I cannot place either of these satisfactorily. Both are probably referable to some of the admitted species listed above.

CYPERUS EXALTATUS Retz.; Naves Novis. App. (1882) 304; Ceron Cat. Pl. Herb. Manila (1892) 178; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 86.

Vidal's specimen on which Ceron's record is based is *C. elatus* Linn. *fide* C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 86. I suspect that Llanos's plant cited by Clarke as *C. exaltatus* Retz. is either *C. radiatus* Vahl or *C. elatus* Linn.

4. PYCREUS Beauvois

PYCREUS ERAGROSTIS (Vahl) Merr. Fl. Manila (1912) 108.

Cyperus eragrostis Vahl Enum. 2 (1806) 322; Naves Novis. App. (1882) 300; Suringar Geslacht. Cyp. (1898) 65.

Pycreus sanguinolentus Nees in Linnaea 9 (1834) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 79.

Cyperus sanguinolentus Vahl Enum. 2 (1806) 351.

Cyperus atratus Steud. in Zoll. Verz. Ind. Archip. (1854) 62; Naves Novis. App. (1882) 301.

Luzon (Ilocos Norte, Bontoc, Benguet, Pangasinan, Rizal, Laguna, Sorogon), Mindanao, *Elmer* 14336, 14323, *Merrill* 7354, 9792, *Phil. Pl.* 514, *Williams* 1972, *B. S.* 27620, 4893, 12039 *Ramos*. In open wet places at low and medium altitudes, ascending to 1,600 m. Widely distributed in the warmer parts of the Old World.

PYCREUS GLOBOSUS (All.) Reichb. Fl. Germ. Exkurs. (1830-32) 140¹⁰;

C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 81 (incl. var. *nilagiricus* C. B. Clarke).

Cyperus globosus All. Fl. Pedem. Auctuar. (1789) 49; Naves Novis. App. (1882) 301; Suringar Geslacht. Cyp. (1898) 58.

Cyperus vulgaris Kunth Enum. 2 (1837) 4; Miq. Fl. Ind. Bat. 3 (1856) 256.

Cyperus lanceolatus Presl Rel. Haenk. 1 (1828) 167, non ? Poir.

Luzon (Bontoc, Lepanto, Benguet, Nueva Vizcaya), Leyte, Mindanao (Lanao), *Merrill* 237, 4630, *Phil. Pl.* 556, *Elmer* 6483, *Williams* 1242, *B. S.* 31810 *Santos*, 14261, 14257 *McGregor*, 5972, 8238 *Ramos*. In open usually wet places at medium altitudes, ascending to 2,000 m. Warmer parts of the Old World.

Local name: Tantanud (Bon.).

PYCREUS NITENS (Retz.) Nees in Nov. Act. Acad. Nat. Cur. 19 (1843)

Suppl. 1: 53; Merr. Fl. Manila (1912) 108.

Cyperus nitens Retz. Obs. 5 (1789) 13; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283; Suringar Geslacht. Cyp. (1898) 55.

Cyperus pumilus Linn. Amoen. Acad. 4 (1759) 302; Naves Novis. App. (1882) 300, non *Pycreus pumilus* Nees.

Pycreus pulvinatus Nees in Linnaea 9 (1834) 283; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 79.

Cyperus gymnoleptus Steud. Syn. Pl. Glum. 2 (1855) 3; Miq. Fl. Ind. Bat. 3 (1856) 255.

Northern Luzon to Mindanao, in most or all islands and provinces. In open usually damp places at low and medium altitudes. Common in the warmer parts of the Old World; Florida (probably introduced).

PYCREUS ODORATUS (Linn.) Urban Symb. Antil. 2 (1900) 164; Merr. Fl. Manila (1912) 108, Interpret. Herb. Amb. (1917) 104, Sp. Blancoanae (1918) 80.

Cyperus odoratus Linn. Sp. Pl. (1753) 46, excl. syn. Sloane.

Pycreus polystachyus Beauv. Pl. Oware et Benin 2 (1807) 48, t. 86, f. 2; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 80; Merr. op. cit. 3 (1908) Bot. 397.

Cyperus polystachyus R. Br. Prodr. (1810) 214; Miq. Fl. Ind. Bat. 3 (1856) 258; Naves Novis. App. (1882) 301.

Cyperus strigosus Llanos Frag. Pl. Filip. (1851) 16; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 10, non Linn.

Throughout the Philippines. Common in open usually damp grasslands, fallow rice paddies, etc., at low altitudes. Pantropic.

Local names: Alúsang-pasígan (Tag.); haná (Iv.).

Var. **PANICULATUS** (Rottb.) Merr. Fl. Manila (1912) 109.

Cyperus paniculatus Rottb. Descr. Ic. (1773) 40.

Pycrus polystachyus Beauv. var. *laxiflorus* Benth. Fl. Austral. 7 (1878) 261; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 80.

Luzon (Isabela, Pampanga, Rizal), Negros. Occasional; habitat of the species.

Var. **HOLOSERICEUS** (Link) Merr. in Journ. Str. Branch Roy. As. Soc. 76 (1917) 79.

Cyperus holosericeus Link Hort. Berol. 1 (1827) 317.

Luzon (Zambales, Laguna), Palmas, Merrill 5343, Bolster 11, Baker 983, B. S. 22986 McGregor. In open wet places; rare and local. Borneo, its other range uncertain.

PYCREUS SULCINUX C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 593; Philip. Journ. Sci. 2 (1907) Bot. 80.

Cyperus sulcinux C. B. Clarke in Journ. Linn. Soc. Bot. 21 (1884) 593; Suringar Geslacht. Cyp. (1898) 63.

Luzon (Benguet, Bulacan), Palawan, Mindanao (Lanao), Merrill 4289, 9278, Phil. Pl. 548, Williams 1969, Elmer 6579, B. S. 21742 Ramos. In open places, in dry or damp soil, sea level to 1,200 m altitude; local. Warmer parts of the Old World.

PYCREUS UNIOLOIDES (R. Br.) comb. nov.

Cyperus unioides R. Br. Prodr. (1810) 216.

Cyperus angulatus Nees in Wight Contrib. (1834) 73.

Pycrus angulatus Nees in Linnaea 9 (1834) 283.

Luzon (Bontoc, Benguet, Nueva Vizcaya), Mindanao (Lanao), Williams 1971, B. S. 14228 McGregor, 8262 Ramos. In boggy places at medium altitudes. Pantropic.

DOUBTFUL AND EXCLUDED SPECIES

CYPERUS ALBUS Presl Rel. Haenk. 1 (1828) 175; Miq. Fl. Ind. Bat. 3 (1856) 287; Steud. Syn. Pl. Glum. 2 (1855) 52; Naves Novis. App. (1882) 306; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 81.

"Hab. in insula Luzon" Presl. This is a *Pycrus* judging from Presl's description of the styles as 2-fid. It is necessary to examine the type to determine its status.

CYPERUS LUZONENSIS Presl Rel. Haenk. 1 (1828) 174; Steud. Syn. Pl. Glum. 2 (1855) 52; Miq. Fl. Ind. Bat. 3 (1856) 287; Naves Novis. App. (1882) 306; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 81.

"Hab. in insula Luzon" Presl. The type should be examined as the status of the species is hardly determinable from the description alone.

CYPERUS FLAVESCENS Linn.; Naves Novis. App. (1882) 301.

CYPERUS FLAVICOMUS Michx.; Naves l. c.

5. MARISCUS Gaertner

MARISCUS CYPERINUS (Retz.) Vahl Enum. 2 (1806) 377; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 87; Merr. op. cit. 3 (1908) Bot. 398, 5 (1910) Bot. 333, Fl. Manila (1912) 113.

Kyllinga cyperinus Retz. Obs. 6 (1791) 21.

Mariscus sundaicus Miq. Fl. Ind. Bat. 3 (1856) 289.

Cyperus sundaicus Naves Novis. App. (1882) 306.

Mariscus umbellatus Moritz Verz. Zoll. Pflanz. (1845) 98, vix Vahl.

Cyperus umbellatus Naves Novis. App. (1882) 305, vix Benth.

Cyperus manilensis Boeck. in Engl. Bot. Jahrb. 5 (1884) 501.

Mariscus philippensis Steud. Syn. Pl. Glum. 2 (1855) 66; Miq. Fl. Ind. Bat. 3 (1856) 290; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 88.

Cyperus steudelianus Naves Novis. App. (1882) 306; Boeck. in Engl. Bot. Jahrb. 5 (1884) 91.

Mariscus umbellatus Presl Rel. Haenk. 1 (1828) 191, non Vahl.

Cyperus cylindrostachys Boeck. in Linnaea 36 (1869) 383, p. p.; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283.

Throughout the Philippines at low and medium altitudes, in open waste places, old clearings, etc. India to Malaya and Polynesia.

Local names: Alúsang (Tag.); alúsang-párang (Tag.); bubuyangkat (Sub.); busikad-dako (C. Bis.); haná (Iv.); saka-án (Ig.); silal (Buk.); ubud-úbud (Tag.).

MARISCUS DILUTUS (Vahl) Nees in Wight Contrib. (1834) 90; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 231, Fl. Manila (1912) 113, Sp. Blancoanae (1918) 80, 81.

Cyperus dilutus Vahl Enum. 2 (1806) 357; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 65; Miq. Fl. Ind. Bat. 3 (1856) 285; Naves Novis. App. (1882) 305; Vidal Phan. Cuming. Philip. (1885) 155; Vidal Rev. Pl. Vasc. Filip. (1886) 283.

Mariscus microcephalus Presl Rel. Haenk. 1 (1828) 182; Steud. Syn. Pl. Glum. 2 (1855) 66; Miq. Fl. Ind. Bat. 3 (1856) 290; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 88.

Cyperus haenkeanus Kunth Enum. 2 (1837) 93.

Cyperus quadriflorus Naves Frag. Pl. Filip. (1851) 17 (*cuadriflorus*); F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 3¹ (1880) 12.

Cyperus luzoniensis Llanos l. c.; F.-Vill. & Naves op. cit. 11.

Cyperus septatus Steud. Syn. Pl. Glum. 2 (1855) 46; Naves Novis. App. (1882) 305.

Cyperus microcephalus Naves Novis. App. (1882) 304, non R. Br.

Sphaeromariscus microcephalus Camus in Not. Syst. 1 (1910) 239.

Duvaljouvea diluta Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 8; Kneucker Cyp. Exsic. No. 246.

Throughout the Philippines, in all or most islands and provinces. Common in open wet places at low and medium altitudes. Tropical Asia to Malaya and the Mascarene Islands.

Local names: Baki-báking-pulá (S. L. Bis.); durugi (Sub.); giron (Bag.) kadang-kadang (Bik.); tíkai (Tag.).

MARISCUS MERRILLII C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 87.
Luzon (Cavite), *Merrill 4170*. On wet banks in ravines, altitude about 300 m. Endemic.

MARISCUS NIVEUS (Murr.) Merr. in Philip. Journ. Sci. 14 (1919) 369.
Schoenus niveus Murr. in Linn. Syst. Veg. ed. 13 (1774) 81.
Scirpus glomeratus Linn. Sp. Pl. (1753) 52, excl. syn. Gronov., non *Mariscus glomeratus* Barton.
Schoenus coloratus Linn. var. β Linn. Sp. Pl. ed. 2 (1762) 64.
Cyperus kyllingiaeoides Vahl Enum. 2 (1806) 312.
Mariscus dregeanus Kunth Enum. 2 (1837) 120; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 620.
Cyperus irroratus Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28.
Luzon (Ilocos Norte), *32826 Ramos*. On dry open rocky slopes at low altitudes. India to tropical Africa, the Malay Peninsula, Singapore, and Borneo.

MARISCUS PENNATUS (Lam.) comb. nov.

Cyperus pennatus Lam. Tabl. Encycl. 1 (1791) 144; Miq. Fl. Ind. Bat. 3 (1856) 281; Naves Novis. App. (1882) 304; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283.
Cyperus stuppeus Forst. f. Prodr. (1786) 89, *nomen nudum*.
Mariscus stuppeus Merr. in Philip. Journ. Sci. 3 (1908) Bot. 398, Fl. Manila (1912) 113, Sp. Blancoanae (1918) 80.
Cyperus canescens Vahl Enum. 2 (1806) 355; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27; Boeck. in Linnaea 36 (1869) 340; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 51.
Mariscus albescens Gaudich. Bot. Freyc. Voy. (1826) 415; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 88.
Cyperus holciflorus Presl Rel. Haenk. 1 (1828) 173; Steud. Syn. Pl. Glum. 2 (1855) 43; Miq. Fl. Ind. Bat. 3 (1856) 282; Naves Novis. App. (1882) 305.
Cyperus bracteolatus Steud. Syn. Pl. Glum. 2 (1855) 49; Miq. Fl. Ind. Bat. 3 (1856) 286; Naves Novis. App. (1882) 305.
Cyperus bracteatus Vidal Phan. Cuming. Philip. (1885) 155.
Cyperus ovatus Naves Frag. Pl. Filip. (1851) 15; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 3 (1880) 10.
Cyperus nitidulus Boeck. in Linnaea 36 (1869) 363; Naves Novis. App. (1882) 305; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 51; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283.
Cyperus anomalus Steud. Syn. Pl. Glum. 2 (1855) 37; Miq. Fl. Ind. Bat. 3 (1856) 277; Naves Novis. App. (1882) 305.
Cyperus stigmatosus Steud. in Zoll. Verz. Ind. Archip. (1854) 96, Syn. Pl. Glum. 2 (1855) 38; Naves Novis. App. (1882) 305.

Along shores of tidal streams, inner borders of mangrove swamps, etc., within the influence of salt or brackish water throughout the Philippines. Tropical Africa to southeastern Asia, Malaya, and Polynesia.

Local names: Kai-kai (Mbo.); obod-óbod (Ibn.); tuhog-dalág (Tag.).

MARISCUS SIEBERIANUS Nees in Linnaea 9 (1834) 286; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 88.

Northern Luzon to Palawan and Mindanao, in most or all islands and provinces. In old clearings, open grasslands, etc., at low and medium altitudes. Pantropic; considered by Kükenthal to be a variety of *M. umbellatus* Vahl.

Local names: Kupaíupi (Sub.); okokiáng (Bon.); mañgilang (Sub.).

MARISCUS TENUIFOLIUS Nees in Mart. Fl. Brazil 2¹ (1843) 46; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 87.

Mariscus flabelliformis C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 89; Merr. Fl. Manila (1912) 113, non ? HBK.

Northern Luzon to Palawan and Mindanao. Scattered in thickets and secondary forests at low and medium altitudes. India to the Malay Peninsula.

Local name: Bakis-bakísan (Tag.).

6. JUNCCELLUS C. B. Clarke

JUNCCELLUS PYGMAEUS (Rottb.) C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 596, Philip. Journ. Sci. 2 (1907) Bot. 81; Merr. Fl. Manila (1912) 109.

Cyperus pygmaeus Rottb. Descr. Ic. (1773) 20, t. 14, f. 4, 5; Miq. Fl. Ind. Bat. 3 (1856) 261; Naves Novis. App. (1882) 301.

Dichelostylis pygmaea Nees in Linnaea 9 (1834) 289; Kneucker Cyp. Exsic. No. 248.

Luzon (Cagayan, Rizal, Laguna), Mindanao (Lanao), Merrill Phil. Pl. 509, B S. 22984, 2684 Ramos, 22714 Castillo. In open wet places, river banks, etc., at low and medium altitudes. Africa, India to Korea southward to Australia.

7. TORULINIUM Desvaux

TORULINIUM FERAX (L. C. Rich.) Ham. Prodr. Pl. Ind. Occ. (1825) 15 (*ferox*); Merr. Fl. Manila (1912) 114.

Cyperus ferax L. C. Rich. in Act. Soc. Hist. Nat. Paris 1 (1792) 106; Naves Novis. App. (1882) 305; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283.

Torulinium confertum Desv. ex Ham. Prodr. Pl. Ind. Occ. (1825) 15; Usteri Beitr. Ken. Philip. Veg. (1905) 131; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 89.

Diclidium elatum Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 27.

Cyperus calopecterus Miq. Fl. Ind. Bat. 3 (1856) 282; Naves Novis. App. (1882) 305.

Cyperus holophyllus Miq. op. cit. 283; Naves l. c.

Mariscus ferax C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 624.

Cyperus haenkei Presl Rel. Haenk. 1 (1828) 172.

Throughout the Philippines. In open wet places at low and medium altitudes. Pantropic.

Local names: Bilíran (S. L. Bis.); pulakgalau (Sub.).

8. KYLLINGA Rottboell

KYLLINGA BREVIFOLIA Rottb. Descr. Ic. (1773) 13, t. 4, f. 3; Naves Novis. App. (1882) 300; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 78; Merr. Fl. Manila (1912) 107.

Kyllinga rigidula Steud. Syn. Pl. Glum. 2 (1855) 71; Miq. Fl. Ind. Bat. 3 (1856) 294; Naves Novis. App. (1882) 300, quoad pl. Philip.

Kyllinga caespitosa Nees var. *robusta* Boeck. in *Linnaea* 35 (1868) 413; Scheff. in *Nat. Tijdschr. Nederl. Ind.* 34 (1874) 45; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 283. *Kyllinga longiculmis* Miq. Fl. Ind. Bat. 3 (1856) 292; Naves Novis. App. (1882) 300.

Kyllinga fuscata Miq. l. c. 294; Naves l. c.

Kyllinga squarrosa Steud. Syn. Pl. Glum. 2 (1855) 68; Naves l. c.

Kyllinga sororia Kunth Enum. 2 (1837) 131; Naves l. c.

Throughout the Philippines in open grasslands, waste places, etc., at low and medium altitudes; common. Pantropic.

Local names: Bibi-inok (Bon.); kadmakat (Ig.); pugo-pugo (C. Bis.).

KYLLINGA CYLINDRICA Nees in Wight Contrib. (1834) 91; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 79.

Luzon (Benguet, Laguna), Mindanao (Bukidnon, Lanao); *Elmer* 6500, *Merrill Phil. Pl.* 571, *Williams* 1225, *B. S.* 26039 *Fénix*, 14033 *Robinson*, 10980 *Ramos*. In open grasslands at medium altitudes, ascending to 1,600 m. Warmer parts of Africa and Asia, through Malaya to Australia. Considered by Kükenthal to be a variety of *K. odorata* Vahl.

KYLLINGA INTERMEDIA R. Br. Prodr. (1810) 219; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 78; Merr. op. cit. 262, 5 (1910) Bot. 333.

Kyllinga oligostachya Boeck. in *Linnaea* 35 (1868) 407.

Luzon (Ifugao, Bontoc, Lepanto, Benguet, Nueva Vizcaya, Laguna), Mindoro, Negros, Mindanao, *B. S.* 14249 *McGregor*, 14125 *Robinson*, *Elmer* 6494, 10021, 9675, *Merrill* 6539, 4618, 4624, 4371, 4421, 4695, *Phil. Pl.* 539; *Williams* 1228. In open wet places, on slopes, about cliffs, etc., at medium altitudes, ascending to 2,200 m. Formosa, Australia, Samoa. Considered by Kükenthal to be a variety of *K. brevifolia* Rottb.

Local names: Busikad (C. Bis.); tabtabúlog (Buk.).

KYLLINGA MELANOSPERMA Nees in Wight Contrib. (1834) 91; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 588.

Kyllinga pungens C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 78, non Link.

Kyllinga bifolia Miq. Fl. Ind. Bat. 3 (1856) 293; Naves Novis. App. (1882) 300.

Luzon (Ilocos Norte, Ifugao, Kalinga, Benguet, Nueva Vizcaya), Mindanao (Lanao), *B. S.* 27159 *Ramos*, 14236, 19995 *McGregor*, *Merrill* 216, 4649, *Phil. Pl.* 1761, 572, *Williams* 1235, *Elmer* 6292, 6495. In open wet places at medium altitudes, ascending to 2,000 m. India to Madagascar and Malaya.

KYLLINGA MONOCEPHALA Rottb. Descr. Ic. (1773) 13, t. 4, f. 4; Miq. Fl. Ind. Bat. 3 (1856) 291; Naves Novis. App. (1882) 199; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 77; Merr. op. cit. 3 (1908) Bot. 397, Fl. Manila (1912) 108, Sp. Blancoanae (1918) 81. *Kyllinga triceps* Blanco Fl. Filip. (1837) 34, ed. 2 (1845) 23, ed. 3, 1 (1877) 44; Steud. Syn. Pl. Glum. 2 (1855) 72; Naves Novis. App. (1882) 300 (Linn. f., p. p.).

- Kyllinga mindorensis* Steud. Syn. Pl. Glum. 2 (1855) 67; Miq. Fl. Ind. Bat. 3 (1856) 292; Naves Novis. App. (1882) 300.
Kyllinga monocephala Rottb. var. *mindorensis* Boeck. in Linnaea 35 (1868) 424; Vidal Phan. Cuming. Philip. (1885) 155, Rev. Pl. Vasc. Filip. (1886) 284.
Kyllinga gracilis Kunth Enum. 2 (1837) 134; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28; Naves Novis. App. (1882) 300.

Throughout the Philippines. Common in waste places, open grasslands, etc., at low and medium altitudes. Pantropic.

Local names: Anúang (Tag.); barubotónes (Bis.); bolobotónes (Bis.); borobotónes (Bis.); botoncillo (Sp.); bosobotónes (Bis.); borsa-nga-dadakkel (Ilk.); baki-báki (S. L. Bis.); bosikad (C. Bis.); boskad (Bis.); busikad (P. Bis.); katutu (Mag.); kurukamóting-orig (Bik.); malabotónes (Bis.); muthá (Tag.); mustra (Tag.); puñgós (S. L. Bis.); sangsangítan (Bon.); sud-sud (Bis.); uli-uli (Bag.).

EXCLUDED SPECIES

KYLLINGA PUMILA Steud.; Naves Novis. App. (1882) 300.

9. FUIRENA Rottboell

FUIRENA CILIARIS (Linn.) Roxb. Hort. Beng. (1814) 81, Fl. Ind. 1 (1820) 184; Merr. Fl. Manila (1912) 119, Sp. Blancoanae (1918) 81.

Scirpus ciliaris Linn. Mant. 2 (1771) 182.

Fuirena glomerata Lam. Tabl. Encycl. 1 (1791) 150; F.-Vill. Novis. App. (1882) 309; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 101.

Fuirena striata Llanos Frag. Pl. Filip. (1851) 21; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 14.

Throughout the Philippines, in rice paddies, open wet lands, etc., at low and medium altitudes, often abundant. Old World Tropics generally.

Local name: Pugápuk (Tag.).

FUIRENA UMBELLATA Rottb. Deser. Ic. (1773) 70, t. 19, f. 3; F.-Vill. Novis. App. (1882) 308; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 101.

Fuirena paniculata Lam. Tabl. Encycl. 1 (1791) 150, t. 39.

Fuirena pentagona Nees in Wight Contrib. (1834) 93, Hook. Journ. Bot. Kew Miscel. 6 (1854) 28.

Throughout the Philippines in open wet lands at low and medium altitudes. Pantropic.

10. SCIRPUS Linnaeus

SCIRPUS ARTICULATUS Linn. Sp. Pl. (1753) 47; Rolfe in Journ. Bot. 23 (1885) 216; Ceron Cat. Pl. Herb. Manila (1892) 178; Clarke in Philip. Journ. Sci. 2 (1907) Bot. 99; Merr. Fl. Manila (1912) 118, Sp. Blancoanae (1918) 81.

Carex glomerata Blanco Fl. Filip. ed. 2 (1845) 24, ed. 3, 1 (1877) 45; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 90, non Thunb.

Scirpus incurvatus Roxb. Fl. Ind. 1 (1820) 216; Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 105.

Luzon (Cagayan, Ilocos Norte, Pangasinan, Nueva Ecija, Rizal, Laguna), *Merrill Sp. Blancoanae* 268, *Phil. Pl.* 513, *B. S.* 2295, 2298 Mearns, 4949, 7447 Ramos. In open wet places in settled areas at low altitudes. Old World Tropics generally.

Local name: Apúrau (Pang.).

SCIRPUS ERECTUS Poir. in *Lam. Encycl.* 6 (1804) 761; Usteri Beitr. Ken. Philip. Veg. (1905) 132; C. B. Clarke in *Philip. Journ. Sci.* 2 (1907) Bot. 99; Merr. Fl. Manila (1912) 118.

Scirpus juncoides Roxb. Fl. Ind. 1 (1820) 216; Ceron Cat. Pl. Herb. Manila (1892) 178.

Scirpus luzonensis Presl Rel. Haenk. 1 (1828) 193; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 93; Steud. Syn. Pl. Glum. 2 (1855) 84; Miq. Fl. Ind. Bat. 3 (1856) 304.

Luzon (Ifugao, Bontoc, Benguet, Nueva Vizcaya, Rizal, Bulacan, Sorso-gon), Polillo, Leyte, Biliran, Mindanao. In open wet places, rice paddies, etc., at low and medium altitudes. Warmer parts of both hemispheres.

Local names: Bitubituínan (Tag.); gúmi (Tag.); paratupit (Ilk.).

SCIRPUS GROSSUS Linn. f. Suppl. (1781) 104; Presl Rel. Haenk. 1 (1828) 195; Miq. Fl. Ind. Bat. 3 (1856) 307; F.-Vill. Novis. App. (1882) 308; Merr. Fl. Manila (1912) 117, *Sp. Blancoanae* (1918) 82.

Cyperus difformis Blanco Fl. Filip. (1837) 32, ed. 2 (1845) 22, ed. 3, 1 (1877) 41; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 90, non Linn.

Scirpus kysoor Roxb. Hort. Beng. (1814) 6; Llanos Frag. Pl. Filip. (1851) 20 (*kisoor*); F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 14 (*kisoor*).

Scirpus grossus Linn. f. var. *kysoor* (Roxb.) C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 660, *Philip. Journ. Sci.* 2 (1907) Bot. 100.

Schoenoplectus grossus Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 3.

Central Luzon to Mindanao, in most islands and provinces, *Merrill Sp. Blancoanae* 564, 692, *Phil. Pl.* 515, *Kneucker Cyp. Exsic.* 220. In fresh-water swamps at low altitudes. India to Indo-China and Malaya.

Local names: Agas (Bik.); бага-ás (P. Bis.); baki-báki (P. Bis.); balakbák (Pang.); balañgót (P. Bis.); bangkúang (Bik.); ragiudíu (Bik.); tikúg (Mbo.); tíkiu (Tag.); títiu (Tag.).

SCIRPUS LACUSTRIS Linn. Sp. Pl. (1753) 48; Merr. in *Philip. Journ. Sci.* 5 (1910) Bot. 172.

Luzon (Cagayan, Zambales, Benguet), *F. B.* 17290 Curran, *Merrill Phil. Pl.* 1792. In fresh-water swamps, sea level to 1,300 m altitude. In most warm countries except Malaya and South America.

Local name: Tiker (Ilk.).

SCIRPUS MERRILLII (Palla) Kükenth. in herb. comb. nov.

Schoenoplectus merrillii Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 3.

Cyperus tenellus C. B. Clarke in *Philip. Journ. Sci.* 2 (1907) Bot. 81, non Linn. f.

Scirpus inundatus C. B. Clarke op. cit. 99, non Poir.

Luzon (Bontoc, Benguet), Negros, Mindanao (Davao), *Merrill 4740, 6639, 7680, Phil. Pl. 511, Kneucker Cyp. Exsic. 223, Elmer 11379a, B. S. 31748 Santos, 3331 McGregor*. In open wet places on the higher mountains, altitude 1,600 to 2,700 m. Endemic.

Local name: Salaisoi (Bag.).

SCIRPUS MUCRONATUS Linn. Sp. Pl. (1753) 50; F.-Vill. Novis. App. (1882) 308; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 100; Merr. Fl. Manila (1912) 118.

Scirpus acutus Presl Rel. Haenk. 1 (1828) 192, non Muhl.

Scirpus preslii Dietr. Sp. Pl. 2 (1833) 175 (*presslii*); Steud. Syn. Pl. Glum. 2 (1855) 84; Miq. Fl. Ind. Bat. 3 (1856) 305; F.-Vill. Novis. App. (1882) 308 (*preslei*).

Northern Luzon to Mindanao, in most islands and provinces. In open wet places at low and medium altitudes, ascending to 1,500 m. Warmer parts of the Old World.

Local names: Kanubsúban (Tag.); libíran (Tag.); parapipit (Ilk.); pulutapit (Bon.); pupuegan (Bon.).

SCIRPUS PULOGENSIS Merr. in Philip. Journ. Sci. 5 (1910) Bot. 333.

Luzon (Benguet), *Merrill 7789, 6616, 6550, B. S. 31750 Santos, F. B. 16134 Curran, Merritt, & Zschokke*. In very damp ravines and on open grassy slopes on the highest mountains, altitude 2,400 to 2,700 m. Endemic.

SCIRPUS SUPINUS Linn. Sp. Pl. (1753) 49; F.-Vill. Novis. App. (1882) 308; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 99 (incl. var. *uninodis* C. B. Clarke); Merr. Fl. Manila (1912) 118.

Luzon (Ilocos Norte, Rizal), *Merrill 3654, Phil. Pl. 504, B. S. 7654, 21970 Ramos*. In open wet places, rice paddies, etc., at low altitudes. Europe to southern Africa and Australia.

SCIRPUS TERNATANUS Reinw. ex Miq. Fl. Ind. Bat. 3 (1856) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 100 (*ternatensis*).

Scirpus chinensis Munro in Seem Bot. Voy. Herald (1857) 423.

Luzon (Ifugao, Kalinga, Benguet, Nueva Vizcaya, Rizal, Tayabas), Mindanao (Davao, Bukidnon, Zamboanga), *B. S. 31934 Santos, 3173, 26750 Ramos, 29113, 37489, 39192 Ramos & Edaño, 19737 McGregor, Merrill 2315, Phil. Pl. 559, Williams 1249, Elmer 6287, 5790*. In open wet places and also in thickets chiefly at medium altitudes, ascending to 2,200 m. India to China and the Riu Kiu Islands, Celebes, and the Moluccas.

Local names: Daátan (Tag.); sagari (Bag.); sañgañga (Klg.); siilak (Ig.).

SCIRPUS TRIQUETER Linn. Mant. 1 (1767) 29; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 100 (var. *segregata* C. B. Clarke).

Luzon, (*Lohr 802*), fide C. B. Clarke. I have seen no specimens of this species. Europe to Japan, also in South Africa and New Guinea.

EXCLUDED SPECIES

SCIRPUS FLUCTANS Linn.; F.-Vill. Novis. App. (1882) 308.

SCIRPUS MARITIMUS Linn.; F.-Vill. l. c.

SCIRPUS MAXIMUS Linn.; F.-Vill. l. c.

11. ELEOCHARIS R. Brown

(Heleocharis auct.)

ELEOCHARIS ACICULARIS (Linn.) R. & S. Syst. 2 (1817) 154; Merr. in Philip. Journ. Sci. 9 (1914) Bot. 264.

Scirpus acicularis Linn. Sp. Pl. (1753) 48.

Luzon (Bontoc, Benguet), Merrill 7665, Vanoverbergh 293. About rice paddies, altitude 1,200 to 1,500 m. North Temperate Zone generally, extending southward to southern China.

ELEOCHARIS AFFLATA Steud. in Zoll. Verz. Ind. Archip. (1854) 62; Syn. Pl. Glum. 2 (1855) 76; Naves Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 90; Merr. op. cit. 3 (1908) Bot. 398.

Eleocharis microcarpa C. B. Clarke in Philip. Journ. Sci. 2 (1908) Bot. 91, non Torr.

? *Eleocharis pellucida* Presl Rel. Haenk. 1 (1828) 196; Steud. Syn. Pl. Glum. 2 (1855) 80; Miq. Fl. Ind. Bat. 3 (1856) 301; Naves Novis. App. (1882) 307 (*Heleocharis*).

Eleocharis subprolifera Steud. in Zoll. Verz. Ind. Archip. (1854) 62; Naves Novis. App. (1882) 307 (*Heleocharis*).

Babuyan Islands, Luzon (Bontoc, Lepanto, Benguet), Elmer 6299, 5751, Williams 1230, 1243, 1247, Merrill 4621, 7703, Phil. Pl. 555. In shallow water, swampy places, rice paddies, etc., altitude 1,000 to 1,600 m. Japan to China and India, southward to Borneo.

ELEOCHARIS ATROPURPUREA (Retz.) Presl Rel. Haenk. 1 (1828) 196; Kunth Enum. 2 (1837) 151; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 90.

Scirpus atropurpureus Retz. Obs. 5 (1789) 14.

Admitted on the authority of C. B. Clarke. I have seen no Philippine material representing the species. Warmer parts of the world.

ELEOCHARIS CARIBAEA (Rottb.) Blake in Rhodora 20 (1918) 24; Merr. Sp. Blancoanae (1918) 83.

Scirpus caribaeus Rottb. Descr. Pl. Rar. Progr. (1772) 24, Descr. Ic. Nov. Pl. (1772) 46, t. 24.

Eleocharis capitata Miq. Fl. Ind. Bat. 3 (1856) 299; Naves Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 90; Merr. Fl. Manila (1912) 114, non R. Br.

Eleocharis retroflexus Llanos Frag. Pl. Filip. (1851) 19; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 13, non Poir.

Luzon (Cagayan, Ilocos Norte, Bataan, Rizal, Laguna), Cebu. Common in open wet lands at low altitudes. Pantropic.

Blake has shown that the type of *Scirpus capitatus* Linn. is identical with the plant commonly known as *Eleocharis tenuis* (Schultes) Willd., which should henceforth bear the name *Eleocharis capitata* (Linn.) R. Br.

ELEOCHARIS DULCIS (Burm. f.) Trin. ex Henschel Vita Rumph. (1833) 186; Merr. Interpret. Herb. Amb. (1917) 104, Sp. Blancoanae (1918) 82.

Andropogon dulcis Burm. f. Fl. Ind. (1768) 219.

Hippuris indica Lour. Fl. Cochinch. (1790) 16.

Eleocharis plantaginea R. Br. Prodr. (1810) 224; Naves Novis. App. (1882) 306.

Carex tuberosa Blanco Fl. Filip. (1837) 35, ed. 2 (1845) 24, ed. 3, 1 (1877) 45, t. 15, non Degl.

Scirpus plantaginoides Rottb. Descr. Nov. Pl. (1773) 45.

Scirpus plantagineus Retz. Obs. 5 (1789) 14.

Eleocharis plantaginoides W. F. Wight in Contrib. U. S. Nat. Herb. 9 (1905) 268.

Eleocharis tuberosa Naves Novis. App. (1882) 306, non ? Schultes.

Eleocharis tumida R. & S. Syst. Mant. 2 (1824) 86; Naves Novis. App. (1882) 306.

Luzon (Cagayan, Bulacan, Laguna, Camarines), Negros, *B. S.* 7448 Ramos, 12332 McGregor, Merrill *Sp. Blancoanae* 395, *Phil. Pl.* 521. In open wet lands, shallow swamps, etc., at low altitudes. India to Malaya.

Local names: Apulid (Tag., Bik.); cabezas de negrito (Sp.); kalaṅgub (C. Bis); potok (Tag.).

ELEOCHARIS EQUISETINA Presl Rel. Haenk. 1 (1828) 195; Steud. Syn. Pl. Glum. 2 (1855) 82; Miq. Fl. Ind. Bat. 3 (1856) 302; Naves Novis. App. (1882) 306; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 89; Merr. Fl. Manila (1912) 114.

Heleocharis plantaginea Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284, non R. Br.

Luzon (Rizal, Sorsogon), Mindanao (Lanao), *Elmer* 14341, *Kneucker Cyp. Exsic.* 224, Merrill *Phil. Pl.* 531. In open wet lands at low altitudes. Ceylon to Madagascar, Malaya, and New Caledonia.

ELEOCHARIS FISTULOSA (Poir.) Schultes f. in Roem. & Schultes Syst. Mant. 2 (1824) 89; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 626; Naves Novis. App. (1882) 306; Camus in Lecomte Fl. Gén. Indo-Chine 7 (1912) 84, f. 13, 3, 4.

Scirpus fistulosus Poir. in Lam. Encycl. 6 (1804) 749.

Eleocharis planiculmis Steud. Syn. Pl. Glum. 2 (1855) 80; Naves Novis. App. (1882) 306.

Luzon (Rizal), Mindanao (Davao, Lanao), Merrill 9790, Copeland 441. In open wet places at low altitudes. Pantropic.

ELEOCHARIS OCHREATA Nees in Linnaea 9 (1834) 294; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 90.

Luzon, (*Loher* 742). Admitted on the authority of C. B. Clarke; I have seen no specimens. Pantropic.

ELEOCHARIS RETROFLEXA (Poir.) Urban Symb. Antil. 2 (1900) 165 (*Heleocharis*).

Scirpus retroflexus Poir. in Lam. Encycl. 4 (1804) 753.

Eleocharis chaetaria R. & S. Syst. 2 (1817) 154; Naves Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 90.

Luzon (Laguna, Sorsogon), Polillo, Negros, Mindanao, *Elmer* 14558, *Clemens* 845, *F. B.* 19276 *Curran*, *B. S.* 9029 *Robinson*, 10049 *Ramos*. Along streams and in rice paddies at low and medium altitudes. Pantropic.

ELEOCHARIS SPIRALIS (Rottb.) Steud. Syn. Pl. Glum. 2 (1855) 81; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 627; Merr. in Philip. Journ. Sci. 14 (1919) 370.

Scirpus spiralis Rottb. Descr. Ic. (1772) 45, t. 15, f. 1.

Luzon (Rizal), *Merrill 9788*. In open wet lands at low altitudes. India to Ceylon, Burma, and Indo-China.

ELEOCHARIS TETRAQUETRA Nees in Wight Contrib. (1834) 113; Naves Novis. App. (1882) 307.

Luzon (Bontoc, Benguet), Mindanao (Lanao), *Merrill Phil. Pl. 557, 1762, Clemens 1105, Vanoverbergh 461*. In swamps, altitude 650 to 1,800 m. India to China and tropical Australia.

ELEOCHARIS VARIEGATA Presl Oken Isis 21 (1828) 269, var. **LAXIFLORA** (Thw.) C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 626, Philip. Journ. Sci. 2 (1907) Bot. 90; Merr. Fl. Manila (1912) 114.

Luzon (Bulacan, Rizal, Camarines), Biliran, Mindanao, *Merrill Phil. Pl. 520, 1461, B. S. 33928 Ramos & Edaño, 18515 McGregor, 26129 Fénix, 1112 Ramos*. In open wet places at low altitudes. India to Madagascar, China, Malaya, and Polynesia.

12. FIMBRISTYLIS Vahl

FIMBRISTYLIS ACUMINATA Vahl Enum. 2 (1806) 285; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29; Naves Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 91; Merr. Fl. Manila (1912) 115.

Luzon (Bulacan, Rizal, Zambales, Albay, Sorsogon), *Merrill 9798, 7355, Phil. Pl. 546, Kneucker Cyp. Exsic. 234, Elmer 14322*. In open wet places, old rice paddies, etc., at low and medium altitudes. Tropical Asia, Malaya, and tropical Australia.

Local names: Puyóng-usá (Tag.); surusibúyas (Bik.).

FIMBRISTYLIS AESTIVALIS (Retz.) Vahl Enum. 2 (1806) 288; F.-Vill. Novis. App. (1882) 308; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 94.

Scirpus aestivalis Retz. Obs. 4 (1786) 12.

Fimbristylis dichotoma Presl Rel. Haenk. 1 (1828) 191, non Vahl.

Luzon (Cagayan, Isabela, Bontoc, Benguet, Pangasinan, Pampanga, Rizal), Mindanao (Lanao), *Merrill 1465, 4292, Phil. Pl. 554, Elmer 6070, B. S. 17847 Otañes, 7473, 8066, 3358, 5971 Ramos*. In open damp places, rice paddies, etc., at low and medium altitudes.

Var. **MACROSTACHYA** Benth. Fl. Austral. 7 (1878) 310; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 94.

Leyte, (*Jagor 1008*). The species India to Amurland southward to Australia, the variety in Australia.

Local names: Boboók (Bon.) lamlamsit (Bon.); sirau-sirau (Ilk.).

FIMBRISTYLIS ANNUA (All.) R. & S. Syst. 2 (1817) 95.

Scirpus annuus All. Fl. Pedem. 2 (1785) 277.

Scirpus diphyllus Retz. Obs. 5 (1789) 15.

Fimbristylis diphylla Vahl Enum. 2 (1806) 289; F.-Vill. Novis. App. (1882) 307, Vidal Rev. Pl. Vasc. Filip. (1886) 284; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 30, 3 (1908) Bot. 398, Fl. Manila (1912) 116; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 93.

- Fimbristylis tomentosa* Vahl Enum. 2 (1806) 290; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 81.
- Fimbristylis brizoides* Sm. var., Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 28.
- Fimbristylis philippica* Steud. Syn. Pl. Glum. 2 (1855) 116; Miq. Fl. Ind. Bat. 3 (1856) 324; F.-Vill. Novis. App. (1882) 308.
- Fimbristylis communis* Kunth Enum. 2 (1837) 234; Miq. Fl. Ind. Bat. 3 (1856) 323.
- Fimbristylis affinis* Presl Rel. Haenk. 1 (1828) 191; Kunth Enum. 2 (1837) 234; F.-Vill. Novis. App. (1882) 308.
- ? *Scirpus falcatus* Llanos Frag. Pl. Filip. (1851) 20; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 13; Merr. Sp. Blancoanae (1918) 83.
- Fimbristylis dichotoma* C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 93, non Vahl.

Throughout the Philippines in open waste places at low and medium altitudes, abundant and variable; in the Philippines various forms or varieties occur that have been characterized as species, such as *F. podocarpa* Nees, *F. tomentosa* Vahl, *F. diphylla* Vahl, *F. calocarpa* Steud., and *F. royeniana* Nees. Pantropic.

Local names: Baliótas (Bag.); bubáging (Sub.); gilal (Sub.); tabtábin (Sbl.); talágig (Bag.); tayok-táyok (Sbl.).

FIMBRISTYLIS CAPITULIFERA Merr. in Philip. Journ. Sci. 9 (1914) Bot. 265.

Batan and Babuyan Islands, *B. S.* 4042, 3575, 3926 *Fénix*, 3172, 3171, 3173 *Mearns*, 10203 *McGregor*. In open places, usually near the sea. Endemic.

FIMBRISTYLIS COMPLANATA (Retz.) Link. Hort. Berol. 1 (1827) 292; Usteri Beitr. Ken. Philip. Veg. (1905) 131 (var. *kraussiana* Hochst.); C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 96.

Scirpus complanatus Retz. Obs. 5 (1789) 14.

Trichelostylis complanata Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29.

Fimbristylis amblyphylla Steud. Syn. Pl. Glum. 2 (1855) 117; Miq. Fl. Ind. Bat. 3 (1856) 324; F.-Vill. Novis. App. (1882) 308.

Fimbristylis autumnalis Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284, non R. & S.

Isolepis willdenowii R. & S. Syst. 2 (1817) 320; Presl Rel. Haenk. 1 (1828) 189.

Luzon (Ilocos Norte, Bontoc, Pampanga, Rizal, Laguna, Camarines, Sorsogon), Mindoro, Palawan, Negros, Panay, Mindanao, *Merrill Phil. Pl.* 1425, *Elmer* 18096, 14324, *B. S.* 31445, 30941, 33500 *Ramos & Edaño*, 27619, 32956 *Ramos*. In open wet places at low and medium altitudes. All warm countries.

Local name: Baki-báki (P. Bis.).

FIMBRISTYLIS CORNICULATA Merr. in Philip. Journ. Sci. 7 (1912) Bot. 231, Fl. Manila (1912) 116.

Luzon (Rizal), *Merrill* 7359. In open sterile grasslands at low altitudes. Endemic.

FIMBRISTYLIS DIPSACEA (Rottb.) Benth. ex C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 635, Philip. Journ. Sci. 2 (1907) Bot. 93; Merr. Fl. Manila (1912) 116.

Scirpus dipsaceus Rottb. Descr. Ic. (1773) 56, t. 12, f. 1.

Echinolytrum dipsaceum Desv. Journ. Bot. 1 (1808) 21, t. 1.

Luzon (Rizal, Laguna), Mindanao (Lanao), *B. S. 3321 Ramos, 22985 McGregor, Merrill 5105*. On muddy lake and stream margins. Tropical Africa and Asia north to the Amur region.

FIMBRISTYLIS FERRUGINEA (Linn.) Vahl Enum. 2 (1806) 291; F.-Vill. Novis. App. (1882) 307; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 94; Merr. op. cit. 1 (1906) Suppl. 30, Fl. Manila (1912) 94.

Scirpus ferrugineus Linn. Sp. Pl. (1753) 50.

Widely distributed in the Philippines in open wet lands, especially those subject to the influence of brackish water, *Kneucker Cyp. Exsic. 196a, Merrill 546, 4252, Phil. Pl. 549*. Pantropic.

FIMBRISTYLIS FUSCA (Nees) Benth. ex C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 649, Philip. Journ. Sci. 2 (1907) Bot. 98.

Abildgaardia fusca Nees in Wight Contrib. (1834) 95.

Gussonea pauciflora Brongn. in Bot. Duperr. Voy. 2 (1829) 171, t. 34, B, non *Fimbristylis pauciflora* R. Br.

Abildgaardia pauciflora Kunth Enum. 2 (1837) 298.

Cladium cyperoides Merr. in Philip. Journ. Sci. 7 (1912) Bot. 74.

Luzon (Cagayan, Bontoc, Ilocos Norte, Rizal), Palawan, *Merrill 2785, 9389, Vanoverbergh 273, B. S. 32809, 32742, 7859, 7836 Ramos*. In open, rather dry grasslands at low and medium altitudes, ascending to 1,300 m. India to Malaya.

FIMBRISTYLIS GLOBULOSA (Retz.) Kunth Enum. 2 (1837) 231; F.-Vill. Novis. App. (1882) 308; Usteri Beitr. Ken. Philip. Veg. (1905) 132; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 96.

Scirpus globulosus Retz. Obs. 6 (1791) 19.

Fimbristylis torresiana Gaudich. Bot. Freyc. Voy. (1826) 413; F.-Vill. Novis. App. (1882) 308.

Fimbristylis utilis Elm. Leaf. Philip. Bot. 3 (1910) 855.

Luzon (Ilocos Norte, Lepanto, Bontoc, Tayabas, Sorsogon), Polillo, Panay, Leyte, Samar, Negros, Bohol, Mindanao, *Elmer 12571, 15584, Wenzel 1532, Merrill Phil. Pl. 1446, B. S. 34777 Ramos & Pascasio, 30672 Ramos & Edaño*. In open wet places at low altitudes. India to Malaya and Polynesia.

Local names: Anahiúan (Mbo., Sub., C. Bis.); badang-badang (Ilk.); pilokong-kabo (Mbo.); sud-sud (Buk.); tayok-táyok (P. Bis.); tíkog (C. Bis., P. Bis.).

FIMBRISTYLIS JUNCIFORMIS (Retz.) Kunth Enum. 2 (1837) 239; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 97.

Scirpus junciformis Retz. Obs. 6 (1791) 19.

Fimbristylis brevifolia Presl Rel. Haenk. 1 (1828) 193, non R. Br.

Fimbristylis brachyphylla Presl op. cit. 351, non Schultes.

Fimbristylis haenkei Dietr. Sp. Pl. 2 (1833) 161.

Fimbristylis falcata Kunth Enum. 2 (1837) 239; F.-Vill. Novis. App. (1882) 308.

Admitted on the authority of C. B. Clarke who apparently examined Presl's specimen, which was not from Monte Rey, California, but possibly from the Philippines, and also a specimen collected by Callery. India to Indo-China.

FIMBRISTYLIS LEPTOCLADA Benth. Fl. Hongk. (1861) 647; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 647.

Luzon (Bulacan), Culion, Palawan, *Merrill 9285, Phil. Pl. 1426, B. S. 41314 Ramos*. In dry open sandy soil at low altitudes. Ceylon, Malay Peninsula, Borneo, southern China.

FIMBRISTYLIS MARIANA Gaudich. in Bot. Freyc. Voy. (1826) 413, var. **FOENEA** Kükenth. in Fedde Repert. 16 (1920) 432.

Luzon (Cagayan, Isabela, Ilocos Norte), *B. S. 7894, 8074, 27625 Ramos, F. B. 16756 Curran*. In open wet places. Marianne and Caroline Islands, the variety endemic.

The specimens look very much like *F. schoenoides* Vahl, and three of those cited were originally referred by Kükenthal to *F. mariana* Gaudich. var. *maxima* (K. Schum.) Kükenth.

FIMBRISTYLIS MILIACEA (Linn.) Vahl Enum. 2 (1806) 287; F.-Vill. Novis. App. (1882) 308; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 96; Merr. op. cit. 1 (1906) Suppl. 30, 3 (1908) Bot. 398, Fl. Manila (1912) 117, Sp. Blancoanae (1918) 83.

Scirpus miliaceus Linn. Syst. ed. 10 (1759) 868.

Scirpus niloticus Blanco Fl. Filip. (1837) 33, ed. 2 (1845) 23, ed. 3, 1 (1877) 43, non Gmel.

Trichelostylis miliacea Nees & Arn. in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 84, Hook. Journ. Bot. Kew Miscel. 6 (1854) 29.

Isolepis miliacea Presl Rel. Haenk. 1 (1828) 189.

Throughout the Philippines in open wet places; a characteristic rice-paddy weed. Pantropic.

Local names: Agor (Tag.); gúmi (Pang.); sirau-sirau (Ilk.); sirisibúyas (Bik.); taulat (Tag.); ubod-úbod (Tag.).

FIMBRISTYLIS MONOSTACHYA (Linn.) Hassk. Pl. Jav. Rar. (1848) 61; F.-Vill. Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 97; Merr. Fl. Manila (1912) 117.

Cyperus monostachyus Linn. Mant. 2 (1771) 180.

Abildgaardia monostachya Vahl Enum. 2 (1806) 296.

Abildgaardia compressa Presl Rel. Haenk. 1 (1828) 179; Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 74; Miq. Fl. Ind. Bat. 3 (1856) 297; Steud. Syn. Pl. Glum. 2 (1855) 72.

Fimbristylis compressa F.-Vill. Novis. App. (1882) 307.

Throughout the Philippines in open grasslands at low and medium altitudes. Pantropic.

FIMBRISTYLIS PALUDOSA Merr. in Philip. Journ. Sci. 9 (1914) Bot. 265.

Luzon (Bontoc, Benguet), *Williams 1239, Merrill Phil. Pl. 551, Elmer 6497*. In cold open swamps, altitude 1,300 to 1,600 m. Endemic, unless referable to *F. salbundia* Kunth.

FIMBRISTYLIS PINETORUM Merr. in Philip. Journ. Sci. 9 (1914) Bot. 266.

Luzon (Benguet), *Merrill 7664, Phil. Pl. 558*. On open grassy slopes in thin pine forests, altitude about 1,500 m. Endemic.

FIMBRISTYLIS PODOCARPA Nees in Wight Contrib. (1834) 98, Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 77; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 94.

Luzon, (*Loher 766*). Admitted on the authority of C. B. Clarke; I have seen no specimens. India to Malaya and the Marianne Islands.

FIMBRISTYLIS POLYTRICHOIDES (Retz.) Vahl Enum. 2 (1806) 248 (*polythricoides*); F.-Vill. Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 92; Merr. Fl. Manila (1912) 115. *Scirpus polytrichoides* Retz. Obs. 4 (1786) 11.

Luzon (Pampanga, Rizal), *Kneucker Cyp. Exsic. 233, Merrill 9803, 4249, Phil. Pl. 522*. In open wet places, especially near the sea or brackish streams. Tropical Africa and Asia to Japan, through Malaya to Australia.

FIMBRISTYLIS QUINQUANGULARIS Kunth Enum. 2 (1837) 229; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 96.

Luzon (Cagayan, Ilocos Norte, Ilocos Sur, Bulacan, Sorsogon), Negros, Panay, Mindanao, *Merrill Phil. Pl. 1423, B. S. 14558, 7625, 7673, 7684 Ramos, 15656 Merritt & Darling*. In open wet lands, cultivated areas, etc., at low and medium altitudes. India to the Riu Kiu and Marianne Islands.

FIMBRISTYLIS RIGIDULA Nees in Wight Contrib. (1834) 99; Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 95.

Luzon (Cagayan, Nueva Vizcaya), Panay, Negros, *B. S. 14226, 32233 McGregor, 7856, 32811 Ramos*. In open grasslands at low and medium altitudes. India to China.

FIMBRISTYLIS SCHOENOIDES (Retz.) Vahl Enum. 2 (1806) 286; Naves Novis. App. (1882) 307; Ceron Cat. Pl. Herb. Manila (1892) 178; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 92; Merr. Fl. Manila (1912) 116.

Fimbristylis bispicata Nees in Linnaea 9 (1834) 290, var. *monostachya* Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29, F.-Vill. Novis. App. (1882) 307.

Abildgaardia nervosa Presl Rel. Haenk. 1 (1828) 180; Steud. Syn. Pl. Glum. 2 (1855) 73.

Fimbristylis nutans Naves Novis. App. (1882) 307; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284, non Vahl.

Luzon (Ilocos Norte, Nueva Ecija, Bulacan, Rizal, Albay), Palawan, Merrill 9322, 9796, 7110, Phil. Pl. 1447, 507, B. S. 32690, 32954 Ramos. In open grasslands, fallow rice paddies, etc., at low altitudes. India through Malaya to Australia.

Local name: Gumi-gúmi (Tag.).

FIMBRISTYLIS SETACEA Benth. in Hook. Lond. Journ. Bot. 2 (1843) 239; C. B. Clarke in Philip. Journ. Sci. 2 (1907) 91.

Fimbristylis acuminata Vahl var. *minor* Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29; Miq. Fl. Ind. Bat. 3 (1856) 314; Boeck. in Linnaea 37 (1871) 4; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284.

Fimbristylis bursifolia Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284, sphalm.

Abildgaardia brevifolia Steud. Syn. Pl. Glum. 2 (1855) 72; Miq. Fl. Ind. Bat. 3 (1856) 297.

Luzon (Bulacan, Zambales, Rizal, Laguna, Sorsogon), B. S. 21740 Ramos, F. B. 3390 Ahern's collector, 3156 Curran & Merritt, 5873, 12388 Curran. In open wet lands at low altitudes. Burma to Tropical Australia.

FIMBRISTYLIS SPATHACEA Roth. Nov. Sp. Pl. (1821) 24; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 95; Merr. op. cit. 3 (1908) Bot. 398.

Fimbristylis glomerata Nees in Linnaea 9 (1834) 290; Boeck. in Linnaea 37 (1871) 47.

Scirpus glomeratus Retz. Obs. 4 (1786) 11, non Linn.

Fimbristylus rigida Kunth Enum. 2 (1837) 246; F.-Vill. Novis. App. (1882) 308.

Isolepis haenkei Presl Rel. Haenk. 1 (1828) 187.

Luzon (Cagayan, Ilocos Norte, Union, Zambales, Batangas, Laguna, Tayabas), Panay, Negros, B. S. 32274 McGregor, 33444, 32737, 32795, 29245, 7469 Ramos, 29123 Ramos & Edaño. In open wet or dry places, usually near the sea. Pantropic.

FIMBRISTYLIS SQUARROSA Vahl Enum. 2 (1806) 289; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 635.

Palawan, Mindanao (Bukidnon), Merrill 9566, Phil. Pl. 2093, B. S. 26093 Féñic. Shores of lakes and in wet soil at low altitudes. Pantropic.

FIMBRISTYLIS SUBBISPICATA Nees in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 75; F.-Vill. Novis. App. (1882) 307; Miq. Fl. Ind. Bat. 3 (1856) 315 (var. *caesia* Miq.); C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 92.

Luzon, (Loher 747). Admitted on the authority of C. B. Clarke and, as he suggests, perhaps best referable to *F. schoenoides* Vahl. India to China and Japan.

FIMBRISTYLIS TENERA R. & S. Syst. Mant. 2 (1824) 57; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 95.

Luzon, (Loher 740). Admitted on the authority of C. B. Clarke. Tropical Asia, Africa, and Malaya.

FIMBRISTYLIS TETRAGONA R. Br. Prodr. (1810) 226; F.-Vill. Novis. App. (1882) 307; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 91; Merr. Fl. Manila (1912) 115.

Scirpus tetragonus Poir. in Lam. Encycl. Suppl. 5 (1804) 98.

Luzon (Pangasinan, Rizal), *Merrill 7360, 9793, Loher 749, B. S. 4892 Ramos*. In open rather wet places at low altitudes. India through Malaya to Australia.

FIMBRISTYLIS THOMSONII Boeck. in Linnaea 37 (1871) 37; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 646.

Palawan, *Merrill 9350*. On dry open slopes bordering thickets at low altitudes. India to Indo-China.

FIMBRISTYLIS THOUARSII (Kunth) comb. nov.

Arthrostylis thouarsii Kunth Enum. 2 (1837) 284.

Arthrostylis chinensis Benth. Fl. Hongk. (1861) 397.

Actinoschoenus filiformis Benth. in Hook. Ic. III 4 (1881) 33, *t. 1346*.

Fimbristylis actinoschoenus C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 650, Philip. Journ. Sci. 2 (1907) Bot. 98.

Actinoschoenus thouarsii Benth. in Hook. Ic. III 4 (1881) 33.

Culion, Busuanga, *Merrill 553, B. S. 41184 Ramos*, back of the mangrove swamp. Ceylon to Madagascar, Malay Peninsula, southern China, and the Riu Kiu Islands.

DOUBTFUL AND EXCLUDED SPECIES

FIMBRISTYLIS OBTUSIFOLIA Kunth; F.-Vill. Novis. App. (1882) 308.

FIMBRISTYLIS PAUCIFLORA R. Br.; F.-Vill. Novis. App. (1882) 307.

FIMBRISTYLIS PILOSA Vahl; Presl Rel. Haenk. 1 (1828) 191; Steud. Syn. Pl. Glum. 2 (1855) 117.

"Hab. in insula Luzon ad portum Sorsogon." Vahl's species is from Africa; the Luzon specimens referred here by Presl possibly represent a form of *F. annua* R. & S.

FIMBRISTYLIS PRESLI Kunth Enum. 2 (1837) 228; Steud. Syn. Pl. Glum. 2 (1855) 108; F.-Vill. Novis. App. (1882) 308.

Abildgaardia pubescens Presl Rel. Haenk. 1 (1828) 180.

"Hab. in Mexico et in Luzon" Presl. Probably not Philippine.

FIMBRISTYLIS CYMOSA R. Br. Prodr. (1810) 228; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 97.

"'Toubonia' (1433 Cuming) in hb. Kew." This was from Cuming's earlier collections, probably from Polynesia. The printed label bears the date 1831, so that it cannot possibly be from Cuming's Philippine collections.

13. BULBOSTYLIS Kunth

BULBOSTYLIS BARBATA (Rottb.) Kunth Enum. 2 (1837) 208; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 98; Merr. op. cit. 3 (1908) Bot. 398.

Scirpus barbatus Rottb. Descr. Ic. (1773) 52, *t. 17, f. 4*; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 284.

Isolepis barbata R. Br. Prodr. (1810) 212; Presl Rel. Haenk. 1 (1828) 187.

Oncostylis barbata Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29.
Fimbristylis barbata Benth. Fl. Austral. 7 (1878) 32; F.-Vill. Novis.
 App. (1882) 308; Merr. Fl. Manila (1912) 116.

Isolepis cumingii Steud. Syn. Pl. Glum. 2 (1855) 101; Miq. Fl. Ind.
 Bat. 3 (1856) 310.

Fimbristylis cumingii F.-Vill. Novis. App. (1882) 308.

Batan Islands, Luzon (Cagayan to Sorsogon, in most or all provinces),
 Negros. In open dry sandy places at low and medium altitudes, locally
 abundant, ascending to 1,500 m. All or most warm countries.

Local names: Humot (Iv.); kulilis (Tag.); tirtiris (Ilk.).

BULBOSTYLIS CAPILLARIS (Linn.) Kunth Enum. 2 (1837) 211; Usteri
 Beitr. Ken. Philip. Veg. (1905) 131; C. B. Clarke in Philip. Journ.
 Sci. 2 (1907) Bot. 98 (var. *trifida* C. B. Clarke); Merr. op. cit. 5
 (1910) Bot. 334.

Scirpus capillaris Linn. Sp. Pl. (1753) 49.

Isolepis capillaris R. & S. Syst. 2 (1817) 118; Miq. Fl. Ind. Bat. 3
 (1856) 311.

Luzon (Cagayan, Lepanto, Bontoc, Benguet). In open places, chiefly
 at medium altitudes, ascending to 2,000 m. Pantropic.

14. **SCHOENUS** Linnaeus

SCHOENUS AXILLARIS (R. Br.) Poir. in Lam. Encycl. Suppl. 2 (1811)
 251; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 334.

Chaetospora axillaris R. Br. Prodr. (1810) 233.

Luzon (Benguet), *F. B. 16141 Curran, Merritt, & Zschokke*. On Mt.
 Pulog, probably from the summit grasslands, altitude about 2,800 m.
 Australia, New Zealand.

SCHOENUS FALCATUS R. Br. Prodr. (1810) 232; Merr. in Philip. Journ.
 Sci. 9 (1914) Bot. 268.

Cladium filiforme Merr. in Philip. Journ. Sci. 5 (1910) Bot. 172.

Luzon (Ilocos Norte, Zambales), Palawan, *B. S. 5086, 32717 Ramos & Pas-
 casio, 508, 717 Foxworthy*. In meadows and along streams at low altitudes
 ascending to 600 m. Formosa, Borneo, northeastern Australia.

SCHOENUS MELANOSTACHYS R. Br. Prodr. (1810) 231; Merr. in Philip.
 Journ. Sci. 2 (1907) Bot. 263.

Mindoro, *Merrill 6173*. In an open heath on Mt. Halcon, altitude about
 2,400 m. Borneo, Queensland, New South Wales.

SCHOENUS PHILIPPINENSIS (Palla) Kükenth. in herb. comb. nov.

Helothrix philippinensis Palla in Allgem. Bot. Zeitschr. 17 (1911)
 Beil. 1.

Schoenus apogon C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot.
 102; Merr. op. cit. 5 (1910) 334, non R. & S.

Luzon (Benguet, Tayabas, Laguna), Negros, Mindanao (Davao), *Kneuc-
 ker Cyp. Exsic. 211, Merrill 6508, 6973, Phil. Pl. 540, Elmer 11370*. On the
 highest mountains, in the mossy forest and in open grassy places, altitude
 1,800 to 2,700 m. Endemic.

Local name: Kalesi (Bag.).

15. CLADIUM P. Browne

CLADIUM CRINITUM (Stapf) comb. nov.

Vincentia crinita Stapf in Philip. Journ. Sci. 19 (1921) 65.

Luzon (Albay), *B. S. 2935 Mearns*. On slopes at medium altitudes on Mt. Mayon. Endemic.

CLADIUM DISTICHUM C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 102.

Luzon (Tayabas, east coast), *Merrill 1123*. Probably along streams at low altitudes. Endemic.

Local name: Barokíbok (Tag.).

CLADIUM FALCATUM (Nees) C. B. Clarke in Kew Bull. Add. Series 8 (1908) 46.

Baumea falcata Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29.

Vincentia falcata Stapf in Journ. Linn. Soc. Bot. 42 (1914) 179.

Luzon (Albay), *Cuming 932*. I have seen merely a fragment of the type, and Kloss's New Guinea specimen. New Guinea, a variety in Sumatra.

CLADIUM JUNCOIDES Elm. Leaf. Philip. Bot. 3 (1910) 854.

Luzon (Isabela, Ilocos Norte, Tayabas), Palawan, Sibuyan, Panay, Dinagat, Mindanao (Surigao), *Elmer 12150, 13145, B. S. 34593, 35209, 35446, 34605 Ramos & Pascasio, 32846 Ramos, 25428 Yates, 21172 Escritor, 13209 Foxworthy & Ramos, Merrill Phil. Pl. 1258*. In thickets and forests at low and medium altitudes, along streams, etc. Endemic.

Local name: Magodtañgiád (P. Bis.).

CLADIUM LATIFOLIUM Merr. in Philip. Journ. Sci. 2 (1907) Bot. 262, 3 (1908) Bot. 399.

Vincentia malesica Stapf. in Journ. Linn. Soc. Bot. 42 (1914) 178.

Batan Islands, Luzon (Laguna, Tayabas, Sorsogon, Albay), Mindoro, Negros, Mindanao (Davao), *Elmer 17638, Williams 2614, B. S. 23499 Ramos, 6497 Robinson, 3822 Fénix, 2386 Foxworthy*. On ridges in the mossy forest, and in open places, altitude 1,500 to 2,400 m. Sumatra, Borneo.

Local name: Samalang (Bag.).

CLADIUM PHILIPPINENSE Merr. in Philip. Journ. Sci. 5 (1910) Bot. 171.

Luzon (Zambales, Nueva Ecija, Tayabas, Camarines), Mindoro, Palawan, Sibuyan, Panay, Bucas Grande, Mindanao (Surigao), *F. B. 6724 Merritt, B. S. 34619, 35097 Ramos & Pascasio, 35429 Martelino & Edaño, 33476, 26508 Ramos & Edaño, 20678 Escritor, 13161 Foxworthy & Ramos, Elmer 12832, 12401, Weber 1053, Merrill 758*. On ledges and boulders along streams at low and medium altitudes. Endemic.

Local name: Kumpai (Bag.).

CLADIUM RIPARIUM (Nees) Benth. Fl. Austral. 7 (1878) 405; C. B. Clarke Ill. Cyp. (1909) t. 85, f. 5-8; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 288.

Chapelliera riparia Nees in Lehm. Pl. Preiss. 2 (1846-48) 76.

Baumea riparia Boeck. in Linnaea 38 (1874) 246.

Mindanao (Lanao), *Clemens 916*. In swamps, altitude about 650 m. Australia, with a variety in British India.

EXCLUDED SPECIES

CLADIUM MARISCOIDES F.-Vill. Novis. App. (1882) 309.

This was based on *Baumea mariscoides* Gaudich. non Torr. and is the Marianne Islands *Cladium gaudichaudii* W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 230. It is allied to the Philippine *Cladium juncoides* Elm.

CLADIUM GLOMERATUM F.-Vill. Novis. App. (1882) 309, non R. Br.

This was based on *Baumea glomerata* Gaudich. It is a Moluccan species.

CLADIUM MARISCUS R. Br.; F.-Vill. Novis. App. (1882) 309.

16. REMIREA Aublet

REMIREA MARITIMA Aubl. Pl. Guin. (1775) 45, t. 16; Naves Novis.

App. (1882) 309; Vidal Phan. Cuming. Philip. (1885) 156, Rev.

Pl. Vasc. Filip. (1886) 285; C. B. Clarke in Philip. Journ. Sci. 2

(1907) Bot. 103; C. B. Rob. op. cit. 6 (1911) Bot. 195.

Mariscus pungens Steud. Syn. Pl. Glum. 2 (1855) 60; Miq. Fl. Ind.

Bat. 3 (1856) 288.

Mariscus capitatus Steud. in Zoll. Verz. Ind. Archip. (1854) 63.

Cyperus capitatus Naves Novis. App. (1882) 306, non Retz.

Remirea wightiana Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29.

Luzon (Cagayan, Camarines), Polillo, Mindoro, Panay, Mindanao, B. S. 32220 McGregor, 6305, 9133 Robinson, F. B. 16680 Bacani, 5534 Merritt, Merrill 881, 11542, Copeland 857. On sandy seashores. Pantropic.

17. RYNCHOSPORA Vahl

RYNCHOSPORA CORYMBOSA (Linn.) Britton in Trans. N. Y. Acad. Sci.

11 (1892) 84; W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 366.

Scirpus corymbosus Linn. Cent. Pl. 2 (1756) 7.

Rynchospora aurea Vahl Enum. 2 (1806) 229; Presl Rel. Haenk. 1

(1828) 200; Miq. Fl. Ind. Bat. 3 (1856) 336; F.-Vill. Novis. App.

(1882) 309; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl.

Vasc. Filip. (1886) 285; C. B. Clarke in Philip. Journ. Sci. 2 (1907)

Bot. 102.

Luzon (Laguna, Tayabas, Camarines, Sorsogon), Mindoro, Palawan, Leyte, Biliran, Samar, Basilan. In open swampy places at low altitudes. Pantropic.

Local names: Agas (Bik., Bis.); bariu-báriu (Bik.); ragiudíu (Bik.); rakido (Bik.); piso-piso (Bik.).

RYNCHOSPORA GLAUCA Vahl Enum. 2 (1806) 233, var. **CHINENSIS**

(Boeck.) C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 672,

Philip. Journ. Sci. 2 (1907) Bot. 102; F.-Vill. Novis. App. (1882)

309 (as *R. glauca* Vahl).

Luzon (Bontoc, Lepanto, Benguet), Mindanao (Lanao), Clemens 1203, Merrill 4623, Phil. Pl. 569, Williams 1245, Elmer 5757, B. S. 2493, 2781. Mearns. On open seepage slopes, along small streams, etc., at medium altitudes, ascending to 1,800 m. Pantropic.

RYNCHOSPORA RUBRA (Lour.) Makino in Bot. Mag. Tokyo 17 (1903)

180; Merr. Fl. Manila (1912) 119.

Schoenus ruber Lour. Fl. Cochinch. (1790) 41.

Rhynchospora haenkei Presl Rel. Haenk. 1 (1828) 199; Miq. Fl. Ind.

Bat. 3 (1856) 336.

Rhynchospora wallichiana Kunth Enum. 2 (1837) 289; F.-Vill. Novis. App. (1882) 309.

Rhynchospora wallichiana C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 668; Philip. Journ. Sci. 2 (1907) Bot. 101.

Northern Luzon to Palawan and Mindanao, in most islands and provinces. In open grasslands at low and medium altitudes. Tropical Africa and Asia to Japan, southward through Malaya to Australia.

RYNCHOSPORA WIGHTIANA C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1893) 669; Philip. Journ. Sci. 2 (1907) Bot. 101.

Rhynchospora wightiana Steud. Syn. Pl. Glum. 2 (1855) 148.

Luzon (Pangasinan, Bulacan, Rizal), Merrill *Phil. Pl.* 2099, *Loher* 793, *B. S.* 4822, 21743 *Ramos*. In open grasslands at low and medium altitudes; local. India to Indo-China and southern China.

18. GAHNIA Forster

GAHNIA JAVANICA Moritzi Verz. Zoll. Pflanz. (1845-46) 98; Miq. Fl. Ind. Bat. 3 (1857) 340; F.-Vill. Novis. App. (1882) 309; Vidal Rev. Pl. Vasc. Filip. (1886) 285; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 103; Merr. op. cit. 263, 5 (1910) Bot. 334.

Luzon (Benguet, Laguna, Tayabas), Mindoro, Negros, Mindanao, *Williams* 2621, *Merrill* 6162, 6596, *Phil. Pl.* 538, *Elmer* 11589, *B. S.* 19571 *Ramos*, 38538 *Ramos & Edaño*. In the mossy forest, in open heaths, etc., altitude 1,700 to 2,700 m. Malay Peninsula to western China, south to New Guinea and Polynesia.

Local name: Tamauan (Buk.).

EXCLUDED SPECIES

GAHNIA RAWACENSIS Miq.; F.-Vill. Novis. App. (1882) 309.

19. SCIRPODENDRON Zippel

SCIRPODENDRON GHAERI (Gaertn.) Merr. in Philip. Journ. Sci. 9 (1914) Bot. 268, Interpret. Herb. Amb. (1917) 106.

Chionanthus ghaeri Gaertn. Fruct. 1 (1781) 190, t. 29, f. a-c; Boerl. in Journ. Linn. Soc. Bot. 31 (1896) 246.

Scirpodendron costatum Kurz in Journ. As. Soc. Beng. 38² (1869) 85; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 422; C. B. Rob. op. cit. 6 (1911) Bot. 195.

Scirpodendron pandaniforme Zipp. ex Kurz l. c. in syn.

Scirpodendron sulcatum Miq. Illustr. Fl. Archip. Ind. (1871) 65, t. 28.

Pandanus caricosus Spreng. Syst. 3 (1826) 897.

Luzon (Cagayan, Tayabas), Polillo, Palawan, Biliran, Mindanao (Zamboanga), *Merrill* 5257, 8114, 11218, *Phil. Pl.* 1295, *B. S.* 10249, 18592 *McGregor*, *F. B.* 17828, 19612 *Curran*. In wet places near the seashore and along streams at low altitudes. Ceylon through Malaya to Australia and Polynesia.

Local names: Bilis (Sub.); gáas (Bik.).

20. MAPANIA Aublet

MAPANIA AFFINIS Merr. in Journ. Str. Branch Roy. As. Soc. 85 (1922) 157.

Jolo, *Clemens* 9330. On forested slopes, altitude about 800 m. Borneo.

MAPANIA GRACILLIMA Kükenth. & Merr. in Philip. Journ. Sci. 9 (1914) Bot. 267.

Luzon (Tayabas), Mindanao (Zamboanga), *Merrill 8215, B. S. 26519 Ramos & Edaño*. On forested ridges, altitude about 1,100 m. Endemic.

MAPANIA HUMILIS (Hassk.) F.-Vill. Novis. App. (1882) 309; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 109; C. B. Rob. op. cit. 6 (1911) Bot. 194.

Pandanophyllum humile Hassk. in Tijdschr. Nat. Vereen. 10 (1843) 119; Miq. Fl. Ind. Bat. 3 (1856) 334.

Lipironia humilis Miq. Illustr. Fl. Archip. Ind. (1871) 61, t. 23.

Luzon (Apayao, Rizal, Nueva Vizcaya, Laguna, Tayabas, Sorsogon), Polillo, Mindoro, Palawan, Sibuyan, Samar, Negros, Panay, Mindanao, incl. *Elmer 14864* as *M. alpina* Elm. In damp primary forests at low and medium altitudes. Malay Peninsula and Archipelago.

Local names: Babakal (Sub.); lubigan (P. Bis.); maribari (S. L. Bis.).

MAPANIA KURZII C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1894) 681; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 422; C. B. Rob. op. cit. 6 (1911) Bot. 194.

Mapania banahaensis Elm. Leaf. Philip. Bot. 2 (1909) 574.

Luzon (Laguna, Tayabas, Camarines, Sorsogon), Alabat, Polillo, Catanduanes, Palawan, Panay, Negros, *Elmer 7639, 16222, 15519, Merrill 10505, 7220, 4001, Whitford 1569, B. S. 31414 Ramos & Edaño, 30318, 13347 Ramos, 10248 McGregor*. In primary forests at low and medium altitudes. Malay Peninsula.

MAPANIA MACROCEPHALA (Gaudich.) K. Sch. ex Warb. in Engl. Bot. Jahrb. 13 (1891) 265; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 422.

Hypolytrum macrocephalum Gaudich. Bot. Freyc. Voy. (1826) 414.

Lepironia macrocephala Miq. Illustr. Fl. Archip. Ind. (1871) 64, t. 27.

Mindanao (Davao), Balut, *Merrill 5409, Elmer 11978*. In primary forests, altitude about 650 m. Moluccas to the Bismarck Archipelago.

Local name: Baroñgis (Bag.).

MAPANIA PALUSTRIS (Hassk.) F.-Vill. Novis. App. (1882) 309; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1894) 681; Merr. in Philip. Journ. Sci. 9 (1914) Bot. 267.

Pandanophyllum palustre Hassk. in Tijdschr. Nat. Vereen. 10 (1843) 119.

Lepironia palustris Miq. Illustr. Fl. Archip. Ind. (1871) 63, t. 25.

Mindanao (Zamboanga, Agusan), *Merrill 7298, B. S. 37392 Ramos & Edaño*. In muddy places along small streams in forests at low altitudes. Malay Peninsula and Archipelago.

Local name: Kulibang (Sub.).

21. THORACOSTACHYUM Kurz

THORACOSTACHYUM LUCBANENSE (Elm.) Kükenth. ex Merr. in Philip. Journ. Sci. 11 (1916) Bot. 258.

Mapania lucbanensis Elm. Leaf. Philip. Bot. 2 (1909) 573.

Luzon (Laguna, Tayabas), Panay, Mindanao (Zamboanga), *Elmer B. S. 28469, 29177, 31136 Ramos & Edaño, 22929 McGregor, 20493, 10029 Ramos,*

36625, 37414 *Ramos & Edaño, Elmer 9116*. In primary forests at low and medium altitudes. Amboina.

Local names: Payong-payong (Sub.); tiñgol (Sub.).

22. *SCLERIA* Bergius

SCLERIA ANNULARIS (Nees) Kunth Enum. 2 (1837) 359; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 104.

Hypopyrum annulare Nees in Linnaea 9 (1834) 303.

Luzon, (*Loher 807*), fide C. B. Clarke. Not seen. India and China.

SCLERIA BANCANA Miq. Fl. Ind. Bat. Suppl. (1862) 693; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1894) 693, excl. syn. Presl.

Scleria malaccensis Boeck. in Linnaea 38 (1874) 507.

Busuanga, Palawan, *Merrill 9202, B. S. 41175 Ramos*. In old clearings, borders of thickets, etc., at low altitudes. Tenasserim through Malaya to New Guinea.

SCLERIA CORYMBOSA Roxb. Hort. Beng. (1814) 103, *nomen nudum*, Fl. Ind. ed. 2, 3 (1832) 574; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 104.

Culion, Palawan, *Merrill 656, 9402*. In wet places at low altitudes. India to the Malay Peninsula.

SCLERIA HEBECARPA Nees in Wight Contrib. (1834) 117; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 105.

Scleria wichurai Boeck. in Engl. Bot. Jahrb. 5 (1884) 510.

Luzon (Rizal, Laguna); Palawan, Mindanao (Bukidnon, Zamboanga, Cotabato), *B. S. 23136 McGregor, 12042 Ramos, 29679 Ramos & Edaño, 11703 Robinson*. In thickets and open places at low and medium altitudes. India to Japan and Polynesia.

Var. *PUBESCENS* (Steud.) C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1894) 689.

Scleria pubescens Steud. Syn. Pl. Glum. 2 (1855) 168.

Luzon (Laguna), Panay, Mindanao (Lanao), *Clemens, B. S. 23796 Ramos, 30957 Ramos & Edaño*. In dry open places at low altitudes. India to China and Malaya.

Local name: Dát (Tag.).

SCLERIA LEVIS Retz. Obs. 4 (1786) 13; C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1894) 694.

Mindoro, Palawan, Negros, *Merrill 9244, 8380, F. B. 5529 Merritt*. In open wet places, ravines, etc., sea level to 1,200 m altitude. India to southern China, southward to Java.

SCLERIA LITHOSPERMA (Linn.) Sw. Prodr. (1788) 18; F.-Vill. Novis. App. (1882) 310; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 285; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 103; Merr. Fl. Manila (1912) 120.

Scirpus lithospermus Linn. Sp. Pl. (1753) 51.

Scleria glaucescens Presl Rel. Haenk. 1 (1828) 202.

Northern Luzon to Mindanao, in most islands and provinces. In dry thickets at low and medium altitudes. Pantropic.

Var. *ROXBURGHII* Thw. Enum. Pl. Zeyl. (1859-64) 354; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 104.

Luzon, Mindanao, *B. S.* 16409 Reillo, Merrill 11543, (*Loher* 805, 806). In thickets at low altitudes. India and Ceylon.

Local names: Dáat (Tag.); katábad (Tag.); taláid (Bag.).

SCLERIA LUZONENSIS Palla in Allgem. Bot. Zeitschr. 13 (1907) 49.

Scleria chinensis C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 105; Merr. op. cit. 1 (1906) Suppl. 31, non Nees.

Luzon (Bontoc, Benguet, Nueva Vizcaya, Pampanga, Rizal, Laguna, Bataan, Tayabas), Mindoro, Palawan, Negros, Panay, Mindanao, Merrill 4834, 3958, 3964, 3921, 9523, *Phil. Pl.* 560, *Kneucker Cyp. Exsic.* 167, *Elmer* 8401, *Whitford* 1517, 1518, *Williams* 1227, 730. In primary forests, on ridges, etc., altitude 800 to 2,200 m. Endemic.

Thought by Kükenthal to be but a variety of *S. multifoliata* Boeck.

Local names: Agagedán (Bon.); egededán (Bon.); mankot (Ig.); papan (Bon.).

SCLERIA MERRILLII Palla in Allgem. Bot. Zeitschr. 17 (1911) Beil. 8.

Luzon (Rizal), *Kneucker Cyp. Exsic.* 249, *Merrill Phil. Pl.* 525 (as *S. zeylanica* Poir.), *B. S.* 2029 Ramos. In fallow rice paddies at low altitudes. Endemic.

This is perhaps the same as the form referred by C. B. Clarke to *S. annularis* Kunth.

SCLERIA MOTLEYI C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 104.

Scleria trigona Merr. in Philip. Journ. Sci. 8 (1913) Bot. 363.

Scleria trigonocarpa Ridl. in Journ. Str. Branch Roy. As. Soc. 46 (1906) 228, *nomen*, *Mater. Fl. Malay. Pen.* (Monocot.) 1 (1907) 110.

Luzon (Pangasinan, Bulacan, Rizal, Tayabas, Camarines, Sorsogon), Catanduanes, Busuanga, Palawan, Panay, Leyte, *B. S.* 1854, 30426 Ramos, 20681 *Escritor*, 29706, 33583, 34110 Ramos & Edaño, 35496 *Martelino & Edaño*, 25500 *Yates*, 29896 *Fénix*, *Elmer* 15411 as *S. sorsogonensis* Elm., *Wenzel* 158. In forests at low and medium altitudes. Borneo, Malay Peninsula.

Var. **DENSISPICATA** C. B. Clarke l. c.

Luzon (Ilocos Norte, Nueva Vizcaya, Pampanga), *B. S.* 20997 Santos, 22441, 32845 Ramos. In forests at low and medium altitudes, ascending to 800 m. Endemic.

SCLERIA ORYZOIDES Presl Rel. Haenk. 1 (1828) 201; Miq. *Fl. Ind.*

Bat. 3 (1857) 342; F.-Vill. *Novis. App.* (1882) 310; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 105.

Luzon (Cagayan, Isabela, Camarines), Palawan, Merrill 144, *Phil. Pl.* 1294, *F. B.* 11621 *Curran*, *B. S.* 7777, 8063 Ramos. In fresh-water swamps, old wet rice paddies, etc., at low altitudes. India and tropical East Africa to tropical Australia.

Local name: Agáas (Bik.).

SCLERIA PERGRACILIS (Nees) Kunth Enum. 2 (1837) 354; C. B. Clarke in Hook. f. *Fl. Brit. Ind.* 6 (1894) 685; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 75.

Hypophorum pergracile Nees in Edinb. Philos. Journ. 17 (1834) 267.

Luzon (Bontoc), Mindanao (Cotabato), *Vanoverbergh* 889, *Copeland*. On open slopes, altitude 300 to 1,400 m. Tropical Africa through India to Yunnan and Indo-China.

Local name: Bangbangló (Bon.).

SCLERIA PURPUREO-VAGINATA Boeck. in Engl. Bot. Jahrb. 5 (1884) 513; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 105.

Scleria sumatrensis Vidal Rev. Pl. Vasc. Filip. (1886) 285, non Retz.

Batan Islands, Luzon (Cagayan, Abra, Pangasinan, Nueva Vizcaya, Batan, Rizal, Laguna, Sorsogon), Polillo, Mindoro, Palawan, Leyte, Panay, Negros, Mindanao, *F. B. 1929 Borden, B. S. 31169 Ramos & Edaña, 13932, 26971 Ramos, 29881 Fénix, 14248 McGregor, Wenzel 1691, Elmer 16063, Copeland 95*. In thickets and in open places along streams at low and medium altitudes; scarcely distinguishable from *S. scrobiculata* Nees except by its winged sheaths. Ceram Laut.

Local name: Tañgilad (Tag.).

SCLERIA SCROBICULATA Nees in Wight Contrib. (1834) 117, Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 119; Miq. Fl. Ind. Bat. 3 (1856) 342; F.-Vill. Novis. App. (1882) 310; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 106; Merr. op. cit. 1 (1906) Suppl. 31, 3 (1908) Bot. 399, Fl. Manila (1912) 121, Sp. Blancoanae (1918) 83. *Scleria foveolata* Llanos Frag. Pl. Filip. (1851) 103; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 79, non Cav.

Scleria multifoliata C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 106, p. p., non Boeck.

Throughout the Philippines, in thickets, open places, old clearings, etc., at low and medium altitudes; usually common. Andaman Islands to the Riu Kiu Islands, southward to New Guinea.

Of two of the three specimens referred by Clarke to *S. multifoliata* Boeck. I consider *Topping 460* to be *S. scrobiculata* Nees and *Merrill 4834* to be *S. merrillii* Palla.

Local names: Agagidán (Bon.); aladán (Ilk.); amamgid (Ilk.); amgid (Tag.); árat (Pamp., Tag.); dáat (Tag.); dáhat (Bik.); dat (Pamp., P. Bis.); daut (Tag.); gáat (Iv.); haras (P. Bis.); katábad (Tag.); mangked (Iv.); tangra (Ilk.); ulat (Pang., Tag.).

SCLERIA SUMATRENSIS Retz. Obs. 5 (1789) 19, t. 2; Rolfe in Journ. Bot. 23 (1885) 215; Usteri Beitr. Ken. Philip. Veg. (1905) 132; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 106.

Scleria purpurascens Steud. Syn. Pl. Glum. 2 (1855) 169; F.-Vill. Novis. App. (1882) 310.

Palawan, Mindanao, Basilan, *B. S. 16349 Reillo, Merrill 9750, De Vore & Hoover 7*. In open places and in forests at low and medium altitudes. India to Sumatra, Java, and Borneo.

Local names: Balbalili (Bon.); bulálo (Sub.); pangpáyung (C. Bis.).

SCLERIA TESSELLATA Willd. Sp. Pl. 4 (1805) 315, excl. syn. Rumph.; F.-Vill. Novis. App. (1882) 310; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 104; Merr. Fl. Manila (1912) 120.

Luzon (Lepanto, Bontoc, Benguet, Pangasinan, Bulacan, Rizal), Panay, *B. S. 30025 Fénix, 12221 Ramos, Merrill 4370, 4617, Phil. Pl. 1444, Williams 1970*. In open grasslands at low and medium altitudes, ascending to 1,500 m. India to Japan southward to tropical Australia.

Local name: Katábad (Tag.).

SCLERIA ZEYLANICA Poir. in Lam. Encycl. 7 (1806) 3 p. p.; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 104; Merr. Fl. Manila (1912) 120.

Luzon (Ilocos Norte, Bulacan, Rizal), *Merrill 3665, Phil. Pl. 1451, B. S. 27626, 10905, 21688, 22001 Ramos*. In open rather wet places at low altitudes. India through Malaya to New Caledonia.

Local names: Dáat-babáe (Tag.); dáat-párang (Tag.); pugad-púgad (Tag.).

DOUBTFUL SPECIES

Scleria macrophylla Presl Rel. Haenk. 1 (1828) 200; Miq. Fl. Ind. Bat. 3 (1857) 343; F.-Vill. Novis. App. (1882) 310.

"Hab. in insula Luzon" Presl. This has been reduced to *Scleria bancana* Miq. with which Presl's description does not remotely agree. In its leaves being 16 lines broad, its panicle a foot long and much branched, and in its fruits being the size of a pea *Scleria macrophylla* Presl does not agree with any of the known Philippine or Indo-Malayan species, but apparently belongs in the group of *Scleria macrocarpa* Nees. Presl's type was doubtless from South America or Mexico, not from the Philippines.

23. DIPLACRUM R. Brown

DIPLACRUM CARICINUM R. Br. Prodr. (1910) 241; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 106; Merr. Fl. Manila (1912) 121.

Scleria caricina Benth. Fl. Austral. 7 (1878) 426; F.-Vill. Novis. App. (1882) 309.

Luzon (Abra, Tarlac, Rizal, Camarines), Panay, Bucas Grande, Mindanao, *Merrill 3626, 8236, Loher 7157, B. S. 35034 Ramos & Pascasio, 30954, 33507 Ramos & Edaño*. In open grasslands at low and medium altitudes. India to southern China through Malaya to tropical Australia.

24. UNCINIA Persoon

UNCINIA RUPESTRIS Raoul in Ann. Sci. Nat. III 2 (1844) 117 var.

CAPILLACEA Kükenth. in Engl. Pflanzenreich 38 (1909) 64, Philip. Journ. Sci. 6 (1911) Bot. 58; Merr. op. cit. 5 (1910) Bot. 334.

Luzon (Benguet, Laguna), Mindanao (Davao), *B. S. 19560 Ramos, F. B. 16140 Curran, Merritt, & Zschokke*. In the mossy forest on Mts. Pulog, Banahao, and Apo, altitude 2,200 to 2,500 m. New Zealand, Tasmania.

Local name: Loniai (Bag.).

25. CAREX Linnaeus

CAREX BACCANS Nees in Wight Contrib. (1834) 122; C. B. Clarke in

Curtis's Bot. Mag. III 49 (1893) t. 7288, Journ. Linn. Soc. Bot. 37 (1904) 14, Philip. Journ. Sci. 2 (1907) Bot. 108; Merr. op. cit. 5 (1910) Bot. 334; Kükenth. in Engl. Pflanzenreich 38 (1909) 258, f. 39 E-H, Philip. Journ. Sci. 6 (1911) Bot. 59.

Luzon (Bontoc, Lepanto, Benguet). In damp thickets and in the mossy forest on the higher mountains, altitude 1,600 to 2,400 m. Himalayan region to southern China, Formosa, Sumatra, and Java.

Local names: Gihidsan (Bon.); silak (Ig.); mankat (Ig.).

CAREX BREVICULMIS R. Br. Prodr. (1810) 242, subsp. **ROYLEANA** Nees var. **KINGIANA** (Lév. & Van.) Kükenth. in Engl. Pflanzenreich 38 (1909) 469, Philip. Journ. Sci. 6 (1911) Bot. 62.

Carex kingiana Lév. & Van. in Bull. Acad. Internat. Geog. Bot. 10 (1901) 272.

Luzon (Benguet), *Merrill 6609, 6628, Phil. Pl. 561, B. S. 31771 Santos, 8853 McGregor*. In open grassy meadows and on damp open slopes, altitude 2,200 to 2,700 m. The species in Australia, the subspecies India to Japan, the variety in Japan.

CAREX BREVISCAPA C. B. Clarke in Hook. f. Fl. Brit. Ind. 6 (1894) 736; Kükenth. in Engl. Pflanzenreich 38 (1909) 474.

Luzon (Sorsogon), Palawan, *Merrill 9521, Elmer 15353*. On forested ridges, altitude about 800 m. Ceylon.

CAREX BRUNNEA Thunb. Fl. Jap. (1784) 38; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 107; Merr. op. cit. 1 (1906) Suppl. 31, 5 (1910) Bot. 335; Kükenth. in Engl. Pflanzenreich 38 (1909) 599, Philip. Journ. Sci. 6 (1911) Bot. 63.

Luzon (Ilocos Norte, Abra, Bontoc, Benguet, Lepanto, Kalinga, Zambales, Pampanga, Rizal, Laguna, Tayabas, Bataan), Negros, Mindanao. In the mossy forest, on forested ridges and slopes, altitude 800 to 2,400 m.

Var. **SUBTEIOGYNA** Kükenth. in Fedde Repert. 8 (1910) 8, Philip. Journ. Sci. 6 (1911) Bot. 63.

Luzon (Benguet), *Merrill 4731, 6505, B. S. 8866 McGregor*. Borders of the mossy forest, altitude about 2,200 m. The species Japan to southern Asia, Malaya, Australia, and the Mascarene Islands, the variety endemic.

Local names: Silak (Ig.); tayalid (Bag.).

CAREX CRYPTOSTACHYS Brongn. in Bot. Duperr. Voy. (1828) 152, t. 25; F-Vill., Novis. App. (1882) 310; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 220; Kükenth. in Engl. Pflanzenreich 38 (1909) 471, Philip. Journ. Sci. 6 (1911) Bot. 62.

Luzon (Nueva Vizcaya, Rizal, Laguna, Tayabas, Sorsogon), Mindanao (Zamboanga), *Merrill 8218, Elmer 7306, 14595, 16110, B. S. 1760 Ramos, 29334 Ramos & Edaño*. In forests at medium altitudes, ascending to 1,000 m. Formosa and southern China to the Malay Peninsula, Java, and Waigiu.

CAREX ELMERI Kükenth. in Fedde Repert. 8 (1910) 326, Leaff. Philip. Bot. 3 (1910) 853, Philip. Journ. Sci. 6 (1911) Bot. 64.

Luzon (Bontoc, Benguet, Laguna), Mindoro, Negros, *Vanoverbergh 868, Merrill 6972, 8025*. In forests, altitude 1,500 to 2,200 m. Endemic.

Doubtfully distinct from *C. loheri* C. B. Clarke.

CAREX FILICINA Nees in Wight Contrib. (1834) 123, var. **CEYLANICA** (Boeck.) Kükenth. in Philip. Journ. Sci. 6 (1911) Bot. 59; C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 11 (*C. filicina*), Philip. Journ. Sci. 2 (1907) Bot. 107; Merr. op. cit. 2 (1907) Bot. 263, 5 (1910) Bot. 335; Kükenth. in Engl. Pflanzenreich 38 (1909) 274. *Carex ceylanica* Boeck. in Linnaea 40 (1876) 341; Kükenth. in Engl. Pflanzenreich 38 (1909) 279.

Luzon (Cagayan, Ifugao, Abra, Lepanto, Bontoc, Benguet, Laguna, Tayabas, Rizal), Negros, Mindanao (Bukidnon, Davao), *B. S.* 19559, 7213, 13986 *Ramos*, 19732, 19695, *McGregor*, *Kneucker Caric. Etsic.* 372, *Elmer* 8587, 10728, *Merrill* 7496, *Phil. Pl.* 544, *Loher* 7155. In the mossy forest, altitude 1,400 to 2,500 m. India to China, southward to Java.

Local names: Silak (Ig.); táan (Ig.); taláyig (Bag).

Forma **SATURATA** (C. B. Clarke) Kükenth. in Engl. Pflanzenreich 38 (1909) 275 (as var. of *filicina*), Philip. Journ. Sci. 6 (1911) Bot. 59.

Carex saturata C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 12.

Luzon (Laguna, Tayabas), *Whitford* 949, *F. B.* 866 *Klemme*. In the mossy forest on Mt. Banahao, altitude about 1,800 m. Java, Sumatra, Borneo.

Forma **DEPAUPERATA** Kükenth. l. cc.

Luzon (Lepanto, Bontoc), Mindoro, *Merrill* 4513, 4743, 6200. In the mossy forest, altitude 2,200 to 2,400 m. India to China, Java, and Sumatra.

CAREX FUIRENOIDES Gaudich. Bot. Freyc. Voy. (1826) 412, var. **CIRRHULOSA** (Nees) Kükenth. in Engl. Pflanzenreich 38 (1909) 287, Philip. Journ. Sci. 6 (1911) Bot. 61.

Carex fuirenoides F.-Vill. Novis. App. (1882) 310; C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 107, Journ. Linn. Soc. Bot. 37 (1904) 11, p. p., vix Gaudich.

Carex cirrhulosa Nees in Hook. Journ. Bot. Kew Miscel. 6 (1854) 29.

Carex fibrata Boott ex Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 286, *nomen nudum*.

? *Carex densiflora* Presl Rel. Haenk. 1 (1828) 204; F.-Vill. Novis. App. (1882) 310.

Luzon (Nueva Vizcaya, Rizal, Laguna), *Merrill* 109, *B. S.* 26203 *Ramos*, 29388 *Ramos & Edaño*. In forests at medium altitudes. The variety endemic, the species in the Marianne Islands.

CAREX GRAEFFEANA Boeck. in Flora 58 (1875) 22; C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 5, Philip. Journ. Sci. 2 (1907) Bot. 107; Kükenth. in Engl. Pflanzenreich 38 (1909) 403, Philip. Journ. Sci. 6 (1911) Bot. 62; Merr. op. cit. 5 (1910) Bot. 335.

Luzon (Bontoc, Benguet, Nueva Vizcaya), Negros, Leyte, Mindanao (Bukidnon, Davao), *B. S.* 8177, 8174 *Ramos*, 31742, 32076 *Santos*, 3370, 14401 *McGregor*, 37710, 38642 *Ramos & Edaño*, *Merrill* 6622, 9652, *Phil. Pl.* 566, *Elmer* 9842, 8532, *Wenzel* 625. On slopes in open places, along streams, and in the mossy forest, altitude 1,600 to 2,200 m. Samoa.

Local names: Alasas (Buk.); bagibi (Bag.); giron (Bag.); kigid (Bon.); sidak (Ig.); silak (Ig.).

CAREX INDICA Linn. Mant. 2 (1771) 574, var. **FISSILIS** (Boott) Kükenth. in Engl. Pflanzenreich 38 (1909) 264, Philip. Journ. Sci. 6 (1911) Bot. 59; F.-Vill. Novis. App. (1882) 310 (*indica*); Usteri Beitr. Ken. Philip. Veg. (1905) 132 (*indica*).

Palawan, *B. S.* 844 *Foxworthy*. On river banks, probably at low or medium altitudes. The species India to Java, the variety Borneo to Australia and Samoa.

CAREX JAPONICA Thunb. var. **CHLOROSTACHYS** (D. Don) Kükenth. in Engl. Pflanzenreich 38 (1909) 620.

Carex chlorostachys D. Don in Trans. Linn. Soc. 14 (1825) 330.

Carex doniana Spreng. Syst. 3 (1826) 825.

Luzon (Benguet), *B. S. 40359 Ramos & Edaño*. In openings in the mossy forest on Mt. Simacoco, altitude about 1,900 m. India to Korea, Japan, and Formosa.

CAREX LIGATA Boott in Benth. Fl. Hongk. (1861) 402, var. **NEXA** (Boott) Kükenth. in Engl. Pflanzenreich 38 (1909) 474, Philip. Journ. Sci. 6 (1911) Bot. 63.

Carex nexa Boott. l. c.

Luzon (Kalinga, Bontoc), *Vanoverbergh 496, 1139, B. S. 37491, 37516 Ramos & Edaño, 40765 Ramos*. In forests, altitude about 1,600 m. Central and southern China to the Riu Kiu Islands and Formosa.

Var. **FORMOSENSIS** (Lév. & Van.) Kükenth. in Engl. Pflanzenreich 38 (1909) 474.

Carex formosensis Lév. & Van. in Mém. Soc. Sci. Nat. Cherbourg 25 (1906) 216.

Luzon (Cagayan, Bontoc, Benguet), *Merrill Phil. Pl. 562, 1763 as C. subtransversa* Clarke, *B. S. 31958, 31684 Santos, 13987 Ramos*. Along streams and trails, on cliffs and steep banks, etc., in and near the mossy forest, altitude 1,300 to 2,400 m. Formosa.

Local names: Egegedán (Bon.); silak (Ig.).

CAREX LOHERI C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 14, Philip. Journ. Sci. 2 (1907) Bot. 108; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 335; Kükenth. in Engl. Pflanzenreich 38 (1909) 487, Philip. Journ. Sci. 6 (1911) Bot. 64.

Luzon (Lepanto, Benguet, Zambales, Laguna, Tayabas), *Merrill 4488, 6506, 6607, 6605, Phil. Pl. 563, Elmer 8582, 8444, B. S. 5133, 19565 Ramos*. In the mossy forest, altitude 1,300 to 2,400 m. Endemic.

Forma **GRANDIMASCULA** Kükenth. in Philip. Journ. Sci. 6 (1911) Bot. 64.

Luzon (Benguet), *Merrill 4729*. In clearings near the mossy forest, altitude about 2,200 m. Endemic. Scarcely distinguishable from the species.

Local name: Silak (Ig.).

CAREX MERRILLII Kükenth. in Fedde Repert. 8 (1910) 7, Philip. Journ. Sci. 6 (1911) Bot. 63.

Luzon (Ifugao, Benguet), *Merrill 6623, 7806, Phil. Pl. 568, Yates*. In the mossy forest and in open places along trails, altitude about 2,250 m. Endemic.

CAREX MYOSURUS Nees in Wight Contrib. Bot. Ind. (1834) 122; Kükenth. in Engl. Pflanzenreich 38 (1909) 258, var. **EMINENS** (Nees) Boeck. in Linnaea 39 (1875) 335; Kükenth. op. cit. 259.

Carex eminens Nees in Wight Contrib. Bot. Ind. (1834) 122.

Luzon (Bontoc), *Vanoverbergh 992*. On slopes, altitude about 1,700 m. India.

CAREX NIKKOENSIS Franch. & Savat. Enum. Pl. Japon. 2 (1879) 132, 558; Kükenth. in Engl. Pflanzenreich 38 (1909) 252, f. A-E, Philip. Journ. Sci. 6 (1911) Bot. 59.

Luzon (Benguet), *Merrill 6631*. Along trails in the mossy forest, altitude about 2,300 m. Japan.

CAREX NODIFLORA Boeck. in Engl. Bot. Jahrb. 5 (1884) 516; Kükenth. in Engl. Pflanzenreich 38 (1909) 288, Philip. Journ. Sci. 6 (1911) Bot. 61.

Carex cumingii Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 286; C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 11, Philip. Journ. Sci. 2 (1907) Bot. 107.

Luzon (Isabela, Ilocos Norte, Rizal, Batangas, Laguna, Sorsogon), Alabat, Mindanao (Agusan), *Elmer 6449, 15328, 17781, 8304, Merrill 10470, Phil. Pl. 1481, B. S. 33013, 22411, 3350, 8006 Ramos*. In primary forests at low and medium altitudes. Endemic.

CAREX PALAWANENSIS Kükenth. in Leaf. Philip. Bot. 4 (1911) 1169.

Palawan, *Elmer 13146*. In damp soil in thickets along streams, altitude about 150 m. Endemic.

CAREX PHACOTA Spreng. Syst. 3 (1826) 826; Kükenth. in Engl. Pflanzenreich 38 (1909) 350, f. 56 A-C, Philip. Journ. Sci. 6 (1911) Bot. 62.

Luzon (Bontoc, Benguet), *Merrill Phil. Pl. 565, Williams 1246, B. S. 2505 Mearns*. In open damp places, altitude about 1,500 m. Himalayan region to China, southern Japan, and Java.

CAREX PYCNOTHYRSOS Kükenth. in Philip. Journ. Sci. 6 (1911) Bot. 60.

Luzon (Bontoc, Nueva Ecija), Negros, Mindanao (Agusan), *Merrill Phil. Pl. 543, Weber 1113, B. S. 26309, 37881 Ramos & Edaña*. In primary forests, altitude 250 to 1,200 m. Endemic.

Local name: Harakhaches (Neg.).

CAREX RAFFLESIANA Boott in Trans. Linn. Soc. 20 (1846) 132, Illustr. 1 (1858) 12, f. 33; Kükenth. in Engl. Pflanzenreich 38 (1909) 282, Philip. Journ. Sci. 6 (1911) Bot. 59.

Luzon (Benguet, Pampanga, Bataan, Laguna, Sorsogon), Palawan, Mindanao, *B. S. 8378 McGregor, 665, 679, 554 Foxworthy, 6076 Robinson, Whitford 1121, Merrill 4221, 4742, 4669, Elmer 6039*. In and near the mossy forest, altitude 800 to 2,200 m. Sumatra and Java to northeastern Australia.

Var. **SCABERRIMA** (Boeck.) Kükenth. in Engl. Pflanzenreich 38 (1909) 283, Philip. Journ. Sci. 6 (1911) Bot. 60; Merr. op. cit. 5 (1910) Bot. 335.

Carex bengalensis Roxb. var. *scaberrima* Boeck. in Linnaea 40 (1876) 347; Vidal Phan. Cuming. Philip. (1885) 156, Rev. Pl. Vasc. Filip. (1886) 285 (*bengalensis*).

Carex scaberrima C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 10, Philip. Journ. Sci. 2 (1907) Bot. 107, Kew Bull. Add. Series 8 (1908) 72.

Carex cruciata Merr. in Philip. Journ. Sci. 3 (1908) Bot. 399, non Wahl.

Batan Islands, Luzon (Cagayan to Sorsogon, in most provinces), Leyte, Mindanao, *Elmer* 11607, 17692, 15522, 1154, *Merrill* 7599, 4795, *B. S.* 3801, 12704 *Fénix*, 27035 *Ramos*. Common on ridges in and near the mossy forest, altitude 1,000 to 2,000 m. Sumatra to the Moluccas.

Local names: Chidak (Ig.); ikidsan (Klg.); taláyid (Bag.); tamalang (Bag.).

Var. CONTINUA (C. B. Clarke) Kükenth. in *Philip. Journ. Sci.* 6 (1911) Bot. 60.

Carex continua C. B. Clarke in *Hook. f. Fl. Brit. Ind.* 6 (1894) 717, *Journ. Linn. Soc. Bot.* 36 (1903) 281, 37 (1904) 11, *Philip. Journ. Sci.* 2 (1907) Bot. 107; *Merr. op. cit.* 1 (1906) Suppl. 31; Kükenth. in *Engl. Pflanzenreich* 38 (1909) 281.

Luzon (Ilocos Norte, Bontoc, Kalinga, Benguet, Nueva Vizcaya, Pangasinan, Bataan, Laguna, Tayabas), Mindoro, Negros, Mindanao (Zamboanga), *Merrill* 3197, 6971, 8189, 8208, *Phil. Pl.* 564, *Whitford* 189, 1145, *B. S.* 20999 *Santos*, 33143, 8221, 13495, 15141 *Ramos*, 14246 *McGregor*, *F. B.* 4384 *Merritt*. On ridges in and near the mossy forest, altitude 800 to 2,000 m. India to southern China and the Philippines.

CAREX RAMOSII Kükenth. in *Fedde Repert.* 8 (1910) 8, *Philip. Journ. Sci.* 6 (1911) Bot. 63.

Luzon (Kalinga, Pangasinan, Rizal), *B. S.* 1434, 20990, 40738, 40798 *Ramos*, 11859 *Robinson & Ramos*, 17866 *Otanes*, *Merrill Phil. Pl.* 536. In forests at medium altitudes. Endemic.

CAREX RARA Boott in *Proc. Linn. Soc.* 1 (1845) 284, *Trans. Linn. Soc.* 20 (1846) 139 subsp. **CAPILLACEA** (Boott), Kükenth. in *Engl. Pflanzenreich* 38 (1909) 102, *Philip. Journ. Sci.* 6 (1911) Bot. 58; *Merr. op. cit.* 5 (1910) Bot. 335.

Carex capillacea Boott *Illustr.* 1 (1858) 44, *t.* 110; C. B. Clarke in *Journ. Linn. Soc. Bot.* 37 (1904) 7, *Philip. Journ. Sci.* 2 (1907) Bot. 107.

Luzon (Benguet), *Merrill* 7715, 6626, 6632, 6612, 4732, *Phil. Pl.* 505, *B. S.* 31744 *Santos*, 8333 *McGregor*. In open wet borders of small streams, altitude 2,200 to 2,600 m. Japan and northeastern Asia to the Himalayan region and New South Wales.

CAREX RHIZOMATOSA Steud. in *Zoll. Verz. Ind. Archip.* (1854) 60, *Syn. Pl. Glum.* 2 (1855) 206; *Miq. Fl. Ind. Bat.* 3 (1857) 384; C. B. Clarke in *Journ. Linn. Soc. Bot.* 37 (1904) 12, *Philip. Journ. Sci.* 2 (1907) Bot. 108; Kükenth. in *Engl. Pflanzenreich* 38 (1909) 289, *f.* 44, *Philip. Journ. Sci.* 6 (1911) Bot. 61.

Carex cumingiana Steud. *Syn. Pl. Glum.* 2 (1855) 206; *Miq. Fl. Ind. Bat.* 3 (1857) 349; Boott *Illustr.* 3 (1862) 107, *t.* 324, 325; Boeck. in *Linnaea* 40 (1876) 367; *F.-Vill. Novis. App.* (1882) 310; *Vidal Phan. Cuming. Philip.* (1885) 156, *Rev. Pl. Vasc. Filip.* (1886) 286.

Carex oligostachya Nees in *Hook. Journ. Bot. Kew Miscel.* 6 (1854) 29.

Luzon (Bontoc, Rizal), Mindanao (Davao, Lanao), *B. S.* 2700 *Ramos*, *Clemens* 1096, *Williams* 2615, *Vanoverbergh* 323. On open slopes, altitude 800 to 1,400 m. India to Indo-China, Java, and the Moluccas.

CAREX RHYNCHACHAENIUM C. B. Clarke ex Merr. in Govt. Lab. Publ. (Philip.) 35 (1906) 5, Philip. Journ. Sci. 1 (1906) Suppl. 31, 2 (1907) Bot. 108, Kew Bull. Add. Series 8 (1908) 79; Kükenth. in Engl. Pflanzenreich 38 (1909) 480, Philip. Journ. Sci. 6 (1911) Bot. 62.

Luzon (Kalinga, Pampanga, Bataan, Laguna), Mindanao (Bukidnon, Zamboanga), *Elmer 6983, Merrill 8024, 8213, Phil. Pl. 512, B. S. 37575, 39007, 39081 Ramos & Edaño*. On forested ridges in and near the mossy forest, altitude 800 to 2,100 m. Endemic.

CAREX SUBTRANSVERSA C. B. Clarke in Philip. Journ. Sci. 2 (1907) Bot. 108, Kew Bull. Add. Series 8 (1908) 92; Kükenth. in Engl. Pflanzenreich 38 (1909) 614, Philip. Journ. Sci. 6 (1911) Bot. 63.

Luzon (Benguet), *Merrill 4730, 7714*. On dry open slopes of recent clearings, along trails, etc., altitude 1,900 to 2,300 m. Endemic.

CAREX TRISTACHYA Thunb. Fl. Jap. (1784) 38 var. **POCILLIFORMIS** (Boott) Kükenth. in Engl. Pflanzenreich 38 (1909) 471, Philip. Journ. Sci. 6 (1911) Bot. 62; Merr. op. cit. 5 (1910) 335.

Carex pocilliformis Boott Illustr. 4 (1867) 175, t. 593.

Luzon (Benguet), *Merrill 6606, 6630, 6629, B. S. 8856 McGregor*. In open meadows and on grassy slopes, altitude 2,200 to 2,500 m. Japan to Formosa, the species also in eastern and central China.

CAREX TURRITA C. B. Clarke in Journ. Linn. Soc. Bot. 37 (1904) 13, Philip. Journ. Sci. 2 (1907) Bot. 108, Kew Bull. Add. Series 8 (1908) 74; Kükenth. in Philip. Journ. Sci. 6 (1911) Bot. 63.

Carex walkeri Arn. var. *turrita* Kükenth. in Engl. Pflanzenreich 38 (1909) 546.

Luzon (Abra, Bontoc, Ifugao, Benguet), *Loher 700, B. S. 7288 Ramos, 40354 Ramos & Edaño*. Apparently in open places, altitude probably above 1,400 m. Endemic.

Local name: Tangtaño (Bon.).

EXCLUDED SPECIES

CAREX LONGEBRACTEATA Steud.; F.-Vill. Novis. App. (1882) 310.

CAREX HAENKEANA Presl Rel. Haenk. 1 (1828) 205; Steud. Syn. Pl. Glum. 2 (1855) 241; Miq. Fl. Ind. Bat. 3 (1857) 354; F.-Vill. Novis. App. (1882) 310.

"Hab. in insulis Philippinis" Presl. This is the South American *Carex pseudocyperus* Linn. var. *haenkeana* Kükenth., the record as Philippine being based on an erroneously localized Malaspina Expedition specimen.

PALMAE

1. PHOENIX Linnaeus

PHOENIX HANCEANA Naud. var. **PHILIPPINENSIS**. Becc. in Philip. Journ. Sci. 3 (1908) Bot. 339, 14 (1919) 339.

Batan Islands (Sabtan), *B. S. 3744 Fénix*. Along mountain streams at low or medium altitudes. The species in southeastern China, Hainan, and Formosa.

Local name: Voyáyo (Iv.).

EXCLUDED SPECIES

PHOENIX DACTYLIFERA Linn.; Naves Novis. App. (1882) 282.

The date palm is very rarely found in gardens in the larger towns of the Philippines. It is not adapted to Philippine climatic conditions.

PHOENIX FARINIFERA Roxb.; Naves l. c.

PHOENIX SYLVESTRIS Roxb.; Naves l. c.

2. **CORYPHA** Linnaeus

CORYPHA ELATA Roxb. Fl. Ind. ed. 2, 2 (1832) 176; Merr. Fl. Manila (1912) 123, Sp. Blancoanae (1918) 84; Becc. in Philip. Journ. Sci. 14 (1919) 341, Leaf. Philip. Bot. 8 (1919) 3023; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 68, t. 24-28.

Corypha umbraculifera Blanco Fl. Filip. (1837) 228, ed. 2 (1845) 160, ed. 3, 1 (1877) 290; Naves Novis App. (1882) 281; Vidal Sinopsis Atlas (1883) 41, t. 93, f. A; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88, non Linn.

Corypha sylvestris Mart. Hist. Nat. Palm. 3 (1849) 233, saltem quoad. cit. Philip.

Sagus rumphii Perr. in Mém. Soc. Linn. Paris 3 (1824) 143, non Willd.

Livistona vidalii Becc. in Webbia 1 (1905) 343.

Throughout the Philippines in most islands and provinces, in some regions widely scattered, in others subgregarious and abundant at low and medium altitudes. India to Malaya.

Local names: Bagátai (Is.); buri (Bis., Bik., Pamp., Tag.); busí (Pamp., Bis., Tag.); íbus (Tag.); piét (Tag., Pang.); sílad (Bik.); sílag (Ilk., Pang.); sílal (Sub.); sírar (Bag.); takták (Is.).

CORYPHA MICROCLADA Becc. in Webbia 5 (1921) 7.

Biliran, B. S. 18720 *McGregor*. In open places at low altitudes. Endemic.

3. **LICUALA** Wurmbe

LICUALA SPINOSA Wurmbe in Verh. Bat. Genoots. 2 (1780) 469; Miq. Fl. Ind. Bat. 3 (1855) 53; Naves Novis. App. (1882) 281; Becc. in Perk. Frag. Fl. Philip. (1904) 45, Webbia 1 (1905) 338, Philip. Journ. Sci. 14 (1919) 339; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 88, t. 33.

Culion, Palawan, Balabac, *Merrill 543, Phil. Pl. 1463, B. S. 836 Foxworthy, 269 Bermejós, 448 Mangubat, F. B. 3611 Curran*. In thickets at low altitudes, sometimes immediately back of the mangrove in brackish mud; now cultivated in Manila for ornamental purposes. Malay Peninsula and Archipelago to the Moluccas.

Local names: Balatbát (P. Bis.); ugsáng (Sul.).

EXCLUDED SPECIES

LICUALA ELEGANS Mart.; Naves Novis. App. (1882) 281.

LICUALA RUMPHII Blume; Naves l. c.

LICUALA SPECTABILIS Miq.; Naves l. c.

4. LIVISTONA R. Brown

LIVISTONA MERRILLII Becc. in Perk. Frag. Fl. Philip. (1904) 45, Webbia 1 (1905) 339, Philip. Journ. Sci. 14 (1919) 340.

Livistona whitfordii Becc. in Philip. Journ. Sci. 4 (1909) Bot. 615.

Luzon (Cagayan, Zambales, Tayabas, Camarines), *Merrill 2071, Whitford 731, F. B. 17811, 10409, 17278, 10189, 5876 Curran, 18451 Alvarez, B. S. 33759 Ramos & Edaña*. In forests and in thickets at low altitudes. Endemic.

Local names: Anáhau (Ibn., Bik., Tag.); bal-láng (Ibn.); labik (Neg.); magálai (Bik.); tekis (Sbl.).

LIVISTONA ROBINSONIANA Becc. in Philip. Journ. Sci. 6 (1911) Bot. 230, 14 (1919) 341.

Luzon (Laguna), Polillo, *B. S. 9265 Robinson, 10471 McGregor, Loher 7056*. In forests at low altitudes. Endemic.

Local name: Pilíg (Tag.).

LIVISTONA ROTUNDIFOLIA (Lam.) Mart. var. **LUZONENSIS** Becc. in Leaf. Philip. Bot. 8 (1919) 3025, Philip. Journ. Sci. 14 (1919) 340; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 92, t. 36.

Livistona rotundifolia Becc. in Leaf. Philip. Bot. 2 (1919) 647; Merr. Fl. Manila (1912) 122; Naves Novis. App. (1882) 281, vix Mart.

Corypha minor Blanco Fl. Philip. (1837) 229, ed. 2 (1845) 161, ed. 3, 1 (1877) 290, non Jacq.

Livistona blancoi Merr. Sp. Blancoanae (1918) 84.

Luzon (Benguet, Union, Pangasinan, Zambales, Pampanga, Laguna, Tayabas, Camarines, Albay), Negros, Mindanao, *Merrill 3799, Sp. Blancoanae 919, Elmer 18054, 9293, F. B. 5834, 17344, 17689, 10631 Curran, 15322 Aguilar, 13576 Meyer & Foxworthy, 26228 Mabesa*. In forests at low and medium altitudes; sometimes planted for ornamental purposes.

Local names: Abiáng (Pamp., Pang.); aná-au (Ilk.); anáhau (Bik., P. Bis., Tag.); anáu (Pamp., Ibn.); bagsáng (S. L. Bis.); báhi (Bis.); bal-láng (Ibn.); balak (Mag.); bolong-ulong (Pamp.); bulús (Bik.); buláu (Bik.); bulnó (Bik.); iñgaing (Ig.); labíg (Ilk.); lúyong (Sbl.); palma brava (Sp.); pilíg (Tag.); tikis (Sbl.).

Var. **MICROCARPA** Becc. in Leaf. Philip. Bot. 8 (1919) 3024, Philip. Journ. Sci. 14 (1919) 341.

Livistona microcarpa Becc. in Philip. Journ. Sci. 2 (1907) Bot. 231.

Palawan, Mindanao (Davao), *F. B. 3784 Curran, Elmer 11967*. In forests at low altitudes. The species widely distributed in Malaya, the two varieties endemic.

Local name: Bal-la (Bag.).

Var. **MINDORENSIS** Becc. in Philip. Journ. Sci. 14 (1919) 341.

Livistona mindorensis Becc. in Philip. Journ. Sci. 4 (1909) Bot. 615.

Mindoro, *F. B. 4108 Merritt*. Common in forests at low altitudes on Bungabong River. Endemic.

Local name: Panobau (Tag.).

LIVISTONA SARIBAS (Lour.) Merr. ex A. Chev. in Bull. Econ. Indochine 22 (1919) 501.

Corypha saribas Lour. Fl. Cochinch. (1790) 212, ed. Willd. (1793) 263, excl. syn. Rumph.

Livistona cochinchinensis Mart. Hist. Nat. Palm. 3 (1838) 319; Becc. Malesia 3 (1886) 69, Philip. Journ. Sci. 14 (1919) 340; Becc. & Hook. f. in Hook. f. Fl. Brit. Ind. 6 (1892) 434, Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 92, t. 35.

Livistona spectabilis Griff. in Calcutta Journ. Nat. Hist. 5 (1845) 336, Palms Brit. Ind. (1845) 130, t. 266 C.

Livistona inaequisecta Becc. in Philip. Journ. Sci. 4 (1909) Bot. 616.

Luzon (Cagayan, Laguna), *F. B.* 17145, 17302, 17156, 10079 Curran, 14762 Darling. In open, somewhat swampy regions at low altitudes, in Cagayan gregarious. Malay Peninsula, Indo-China.

Local names; Taráu (Ibn.); tīkal (Tag.).

EXCLUDED SPECIES

LIVISTONA PAPUANA Becc.; Naves Novis. App. (1882) 282.

5. METROXYLON Rottboell

METROXYLON SAGU Rottb. in Nye Samml. Dansk. Vidensk. Selsk. Skrift. 2 (1783) 527; Naves Novis. App. (1882) 276; Merr. Interpret. Herb. Amb. (1917) 112; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 96, t. 37.

Metroxylon rumphii Mart. Hist. Nat. Palm. 3 (1838) 214, 313, t. 102, 159; Naves l. c.; Becc. in Leaf. Philip. Bot. 8 (1919) 3026, Phil. Journ. Sci. 14 (1919) 342, Ann. Bot. Gard. Calcutta 12: t. 105 ined.

Cebu, Bohol, Siquijor, Mindanao (Agusan, Surigao, Misamis, Zamboanga, Cotabato, Davao), Basilan, Sulu. Abundant in fresh-water swamps at low altitudes, in some places planted. Widely distributed in Malaya. The sago palm.

Local names: Ambolong (Mbo.); bagsáng (Bis.); langdáng (Bis.); lumbái (Bis.); lumbiá (C. Bis., Bag.); lumbiág (Sul.); sagú (Mbo., Bis.).

EXCLUDED SPECIES

METROXYLON SYLVESTRE Mart.; Naves Novis. App. (1882) 276.

METROXYLON LONGISPINUM Mart.; Naves l. c.

METROXYLON FILARE Mart.; Naves l. c.

6. COELOCOCCLUS H. Wendland

COELOCOCCLUS AMICARUM (Wendl.) W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 244, t. 45, 46.

Sagus amicarum Wendl. in Bot. Zeit. 36 (1878) 115.

Coelococcus carolinensis Dingl. in Bot. Centralbl. 32 (1887) 349.

Rarely planted, Panay, Mindanao (Zamboanga). Introduced from the Caroline Islands. Polynesian ivory palm.

7. ZALACCA Reinwardt

ZALACCA CLEMENSIANA Becc. in Philip. Journ. Sci. 4 (1909) Bot. 618, 14 (1919) 342, Leaf. Philip. Bot. 8 (1919) 3028, Ann. Bot. Gard. Calcutta 12²: t. 49, ined.

Mindanao (Lanao, Zamboanga, Davao), *Clemens 1109, Elmer 11879, B. S. 37427, 37431, 37430 Ramos & Edaño*. In forests, altitude 600 to 900 m. Endemic.

Local names: Dalúbi (Sub.); kaúbi (Mag.); lakaúbi (Bag.); lubo (Sub.).

EXCLUDED SPECIES

ZALACCA EDULIS Reinw.; Naves Novis. App. (1882) 274.

This has not as yet been found in the Philippines except as a recently introduced and cultivated palm.

8. KORTHALSIA Blume

KORTHALSIA LACINIOSA Mart. Hist. Nat. Palm. 3 (1838) 211, 343; Becc. in Philip. Journ. Sci. 14 (1919) 343, Leaf. Philip. Bot. 8 (1919) 343, Malesia 2 (1884) 74.

Luzon (Tayabas, Camarines, Sorsogon), *Elmer 15634, 16672, F. B. 24537 Aguilar, Klemme, (Vidal 4066)*; possibly also *B. S. 10464 McGregor* from Polillo. In primary forests at low and medium altitudes. Burma and Andaman and Nicobar Islands.

Local name: Dánan (Bik.).

KORTHALSIA MERRILLII Becc. in Ann. Bot. Gard. Calcutta 12²: t. 81, ined., Philip. Journ. Sci. 14 (1919) 342.

Palawan, *Merrill 9410*. In primary forests at low altitudes. Endemic.

KORTHALSIA SCAPHIGEROIDES Becc. in Philip. Journ. Sci. 4 (1909) Bot. 619, 14 (1919) 342, Ann. Bot. Gard. Calcutta 12²: t. 67, ined.

Mindanao (Agusan, Surigao, Zamboanga), Basilan, *Merrill 7313, F. B. 6106, 4816 Hutchinson, 23913 Ponce*. In primary forests at low altitudes. Endemic.

Local name: Kapnigid (C. Bis.).

KORTHALSIA SQUARROSA Becc. in Philip. Journ. Sci. 4 (1909) Bot. 620, 14 (1919) 343, Ann. Bot. Gard. Calcutta 12²: t. 103, ined.

Palawan, Balabac, *Merrill 5384, F. B. 4185 Curran*. In primary forests at low altitudes. Endemic.

EXCLUDED SPECIES

KORTHALSIA ANGUSTIFOLIA Blume; Naves Novis. App. (1882) 273.

KORTHALSIA ROBUSTA Blume; Naves l. c.

9. PLECTOCOMIA Martius and Blume

PLECTOCOMIA ELMERI Becc. in Leaf. Philip. Bot. 8 (1919) 3029, Philip. Journ. Sci. 14 (1919) 342, Ann. Bot. Gard. Calcutta 12²: t. 49 ined.

Mindanao (Davao), *Elmer 11887*. In forests, altitude about 900 m. Endemic.

Local name: Uñgáng (Bag.).

10. **CALAMUS** Linnaeus

CALAMUS ARUGDA Becc. in Philip. Journ. Sci. 4 (1909) Bot. 622, 14 (1919) 352.

Luzon (Cagayan), *F. B.* 6649 *Klemme*, 26618 *Velasco*. In primary forests at low altitudes. Endemic.

Local names: Arúgda (Ibn.); uái (Ilk.).

CALAMUS BICOLOR Becc. in Leaf. Philip. Bot. 8 (1919) 3055, Philip. Journ. Sci. 14 (1919) 356, Ann. Bot. Gard. Calcutta 11 Suppl. 126, t. 73 ined.

Mindanao (Davao), *Elmer* 10541, 10618. In primary forests, altitudes 1,200 to 1,800 m. Endemic.

Local names: Lási (Bag.); rási (Bag.); sambonótan (Bag.).

CALAMUS BLANCOI Kunth Enum. 3 (1841) 595; Mart. Hist. Nat. Palm. 3 (1849) 343; Walp. Ann. 3 (1853) 492, 5 (1858) 832; Miq. Fl. Ind. Bat. 3 (1855) 139; Becc. Rec. Bot. Surv. India 2 (1902) 204, Webbia 1 (1905) 346, Philip. Journ. Sci. 4 (1909) Bot. 635, 14 (1919) 349, Ann. Bot. Gard. Calcutta 11 (1908) 82, 216, t. 64, Leaf. Philip. Bot. 8 (1919) 3048; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88, Sp. Blancoanae (1918) 86.

Calamus gracilis Blanco Fl. Filip. (1837) 267, ed. 2 (1845) 186, ed. 3, 1 (1877) 332, non Roxb.

Calamus buroensis Naves Novis. App. (1882) 275, Vidal Cat. Pl. Prov. Manila (1880) 47, non Mart.

Calamus brevifrons Mart. Hist. Nat. Palm. 3 (1849) 338; Walp. Ann. 3 (1853) 488; Miq. Fl. Ind. Bat. 3 (1855) 127; Naves Novis. App. (1882) 275.

Calamus parvifolius Vidal Phan. Cuming. Philip. (1885) 154, non Roxb.

Luzon (Ilocos Norte), *Cuming* 1225, Zambales, Bataan, Bulacan), Leyte, Mindanao (Zamboanga), *Elmer* 7282, (*Loher* 1376), *Merrill Sp. Blancoanae* 791, *F. B.* 4819 *Hutchinson*, 387 *Maule*, 24697 *Lazaro*, *B. S.* 33047 *Ramos*. In forests at low altitudes. Endemic.

Local names: Abit (Sbl.); alip (Suk.); talóla (Tag.); u-uái (Ibn.); yantók (Tag.).

CALAMUS CUMINGIANUS Becc. in Rec. Bot. Surv. Ind. 2 (1902) 210, Webbia 1 (1905) 346, Ann. Bot. Gard. Calcutta 11 (1908) 99, 348, t. 142, Leaf. Philip. Bot. 8 (1919) 3050, Philip. Journ. Sci. 14 (1919) 351.

Luzon (Tayabas), Mindanao (Agusan), *Cuming* 762, *Elmer* 13646. In primary forests at low and medium altitudes. Endemic.

Local name: Douung-douung (Mbo.).

CALAMUS DIEPENHORSTII Miq. Fl. Ind. Bat. Suppl. (1860-61) 594; Becc. in Ann. Bot. Gard. Calcutta 11 (1908) 96, 322, t. 126, var. **EXULANS** Becc. in Philip. Journ. Sci. 4 (1909) Bot. 627, 6 (1911) Bot. 230, 14 (1919) 350.

Polillo, Palawan, *Merrill* 9357, *B. S.* 756 *Foxworthy*, 9111 *Robinson* (*Loher* 7054 from Rizal Province, Luzon, originally referred here by Beccari is *C. filispadix* Becc. fide Beccari). In primary forests at low altitudes. The variety endemic, the species in the Malay Peninsula, Sumatra, and (?) Borneo.

CALAMUS DIMORPHACANTHUS Becc. in Rec. Bot. Surv. India 2 (1902) 214, Webbia 1 (1905) 355, Ann. Bot. Gard. Calcutta 11 (1908) 117, 479, *t.* 219, Philip. Journ. Sci. 4 (1909) Bot. 631, 14 (1919) 355, Leaf. Philip. Bot. 8 (1919) 3055.

Luzon (Benguet, Union, Laguna), Panay, *Elmer* 6238, 18103, 18337, (*Loher* 1370, 1371, *Vidal* 3956). In forests at medium altitudes, ascending to 1,700 m. Endemic.

Var. **MONTALBANICUS** Becc. in Philip. Journ. Sci. 4 (1909) Bot. 631, 14 (1919) 355.

Luzon (Rizal, Bataan), (*Loher* 7085), *B. S.* 29599 *Ramos & Edaña*. In primary forests at medium altitudes. Endemic.

Var. **ZAMBALENSIS** Becc. in Philip. Journ. Sci. 4 (1909) Bot. 632, 14 (1919) 355.

Luzon (Zambales), *F. B.* 8412 *Curran & Merritt*. In mossy forests on exposed ridges, altitude about 2,000 m. Endemic.

CALAMUS DISCOLOR Mart. Hist. Nat. Palm. 3 (1849) 341; Kunth Enum. 3 (1841) 212; Miq. Fl. Ind. Bat. 3 (1855) 136; Naves Novis. App. (1882) 275; Becc. in Webbia 1 (1905) 355, Ann. Bot. Gard. Calcutta 11 (1908) 120, 495, *t.* 228, Philip. Journ. Sci. 4 (1909) Bot. 635, 14 (1919) 350, Leaf. Philip. Bot. 2 (1909) 649, 8 (1919) 3048.

Calamus lindenii Rod. Ill. Hort. 30 (1883) 157, *t.* 489; Ridley in Journ. Str. Branch Roy. As. Soc. 44 (1905) 200.

Cycas ? hypoleuca Presl Epim. (1851) 238; Walp. Ann. 3 (1852-53) 453; F.-Vill. Novis. App. (1880) 212; Foxw. in Philip. Journ. Sci. 6 (1911) Bot. 152, ex descr.

Luzon (Laguna, Tayabas, Sorsogon), *Elmer* 9299, 16871, *B. S.* 12035 *Ramos, F. B.* 25744 *Mabesa*. In primary forests at low or medium altitudes. Endemic.

Young plants are commonly cultivated in Manila for ornamental purposes.

Local names: Hamlís (Tag.); kumabói (Tag.); likbón (C. Bis.); simurán (P. Bis.); ubánon (C. Bis.); ulisí (C. Bis.); yamíng (P. Bis.).

Var. **NEGROSENSIS** Becc. in Philip. Journ. Sci. 4 (1909) Bot. 635.

Negros, Siargao, Mindanao (Surigao), *F. B.* 12432 *Danao*, 20893 *Tamesis*, 24024 *Razon*. In primary forests at low and medium altitudes. Endemic.

CALAMUS ELMERIANUS Becc. in Leaf. Philip. Bot. 2 (1909) 647, 8 (1919) 3051, Philip. Journ. Sci. 14 (1919) 353, Ann. Bot. Gard. Calcutta 11 Suppl. 69, *t.* 37 ined.

Luzon (Tayabas), Dinagat, Mindanao (Agusan, Davao), *Elmer* 14166, 11756, 9298, 9298, *Merrill* 7289, *F. B.* 24021 *Razon, B. S.* 34909 *Ramos & Pascasio*. In primary forests; sometimes at low altitudes, more often in the mossy forest at 1,300 to 1,800 m.

Local names: Sababai (Mbo.); samánid (Bag.).

CALAMUS FILISPADIX Becc. in Philip. Journ. Sci. 6 (1911) Bot. 230, 14 (1919) 350, Leaf. Philip. Bot. 8 (1919) 3048.

Calamus hookerianus Becc. op. cit. 4 (1909) Bot. 621, non Becc. in Ann. Bot. Gard. Calcutta (1908).

Luzon (Rizal, Tayabas, Camarines, Sorsogon), Polillo, Masbate, Samar, Palawan, Mindanao (Agusan), *Elmer 13949, 16846, 12769, Merrill 7251, F. B. 26219 Mariano, 24824 Valencia, 10630 Curran, B. S. 13128 Foxworthy & Ramos, 10467 McGregor, 33761 Ramos & Edaño, 24427 Ramos*. In forests at low and medium altitudes, ascending to 1,200 m. Endemic.

Local names: Kangnoblób (Mbo.); nókut (S. L. Bis.); pañganpañgánan (Tag.).

CALAMUS FOXWORTHYI Becc. in Philip. Journ. Sci. 14 (1919) 352, *nomen nudum*, Ann. Bot. Gard. Calcutta 11 Suppl. 81, t. 45 ined.

Palawan, *B. S. 690 Foxworthy*. On forested rocky slopes along streams, altitude about 1,000 m. Endemic.

CALAMUS GRANDIFOLIUS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 629, 14 (1919) 352.

Luzon (Laguna, Tayabas), *B. S. 9448 Robinson, (Loher 7088)*.

Local name: Uái (Tag., Neg.).

CALAMUS HALCONENSIS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 633, 14 (1919) 356, Ann. Bot. Gard. Calcutta 11 Suppl. 116, t. 65 ined.

Luzon (Laguna), Mindoro, *F. B. 4399 Merritt, 25742 Mabesa, B. S. 39471 Ramos*. In primary forests, altitude 450 to 1,500 m. Endemic.

Local name: Lambútan (Tag.).

CALAMUS JENNINGSIANUS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 623, 14 (1919) 353.

Mindoro, *F. B. 4400 Merritt*. In forests, altitudes about 1,500 m. Endemic.

CALAMUS MANILLENSIS (Mart.) Wendl. in Kerch. Palm. (1878) 237; Becc. in Rec. Bot. Surv. India 2 (1902) 215, Webbia 1 (1905) 349, Ann. Bot. Gard. Calcutta 11 (1908) 108, 413, t. 226, f. 1, Leaf. Philip. Bot. 8 (1919) 3052, Philip. Journ. Sci. 14 (1919) 352.

Daemonorops ? manillensis Mart. Hist. Nat. Palm. 3 (1849) 330, t. 175, f. 1-3; Walp. Ann. 3 (1853) 480, 5 (1858) 829; Naves Novis. App. (1882) 275.

Luzon (Nueva Vizcaya, Tayabas), Dinagat, Mindanao (Agusan, Davao, Surigao), *Elmer 11714, 10560a, 14011, 14133, 14173, 16811, 17077, B. S. 18994 McGregor, 35255, 34693 Ramos & Pascasio, 26605 Ramos & Edaño*. In forests, altitude 600 to 1,000 m. Endemic.

Local names: Bayábong (Mbo.); bunlak (Mbo.); lintókan (Bag.). saráni (Bag.); tumárom (Mbo.).

CALAMUS MAXIMUS Blanco Fl. Filip. (1837) 266, ed. 2 (1845) 185, ed. 3 1 (1877) 331; Mart. Hist. Nat. Palm. 3 (1849) 343; Miq. Fl. Ind. Bat. 3 (1855) 139; Becc. in Perk. Frag. Fl. Philip. (1904) 45; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88, Sp. Blancoanae (1918) 85.

Calamus merrillii Becc. in Webbia 1 (1905) 347, Ann. Bot. Gard. Calcutta 11 (1908) 105, 390, t. 167, Philip. Journ. Sci. 4 (1909) Bot. 629, 14 (1919) 351, Leaf. Philip. Bot. 8 (1919) 3057.

Luzon (Rizal, Laguna, Tayabas), Masbate, Mindanao (Agusan, Lanao, Davao), Basilan, *Merrill* 1893, *Sp. Blancoanae* 879, *Clemens* 1112, 1124, *Elmer* 11885, 13926, *F. B.* 20654, *Villamil*, 25373, 23988 *Mabesa*, 25353 *Bawan*, 23977 *Maneja*, 26225 *Mariano*, 24322 *Valencia*, 24326 *Acedillo*. In primary forests at medium altitudes, ascending to 1,200 m. Endemic.

Local names: Akab-ákab (Bag.); ápas (Tag.); buai-magi (Yak.); kalapí (Bik.); labni (Pamp.); lokuán (Tag.); palásan (Tag., Bik., Mbo.); parásan (S. L. Bis.); şupot (Tag.); tumalím-bábui (Tag.).

Var. **NANGA** (Becc.) comb. nov.

Calamus merrillii Becc. var. *nanga* Becc. in Philip. Journ. Sci. 14 (1919) 351, Leaf. Philip. Bot. 8 (1919) 3057, Ann. Bot. Gard. Calcutta 11 Suppl. 78, t. 42 ined.

Mindanao (Davao), *Elmer* 11110, 11874. In primary forests, altitude 600 to 1,000 m. Endemic.

Local name: Nañga (Bag.).

Var. **MERRITTIANUS** (Becc.) comb. nov.

Calamus merrittianus Becc. in Philip. Journ. Sci. 2 (1907) Bot. 233, 4 (1909) Bot. 629.

Calamus merrillii Becc. var. *merrittianus* Becc. in Philip. Journ. Sci. 14 (1919) 351, Ann. Bot. Gard. Calcutta 11 Suppl. 78, t. 42 ined.

Luzon (Tayabas), Mindoro, *F. B.* 3912 *Merritt*, 26218 *Mariano*. In primary forests at low altitudes. Endemic.

CALAMUS MEGAPHYLLUS Becc. in Philip. Journ. Sci. 14 (1919) 353, Leaf. Philip. Bot. 8 (1919) 3060, Ann. Bot. Gard. Calcutta 11 Suppl. 66, t. 55 ined.

Mindanao (Agusan, Davao), *Elmer* 11878, 13542. In forests up to 900 m altitude. Endemic.

Local names: Banakbo (Mbo.); lintókan (Bag.).

CALAMUS MELANORHYNCHUS Becc. in Philip. Journ. Sci. 14 (1919) 350, Leaf. Philip. Bot. 8 (1919) 3058, Ann. Bot. Gard. Calcutta 11 Suppl. 30, t. 16 ined.

Mindanao (Davao), *Elmer* 11708. In primary forests, altitude about 1,200 m. Endemic.

Local name: Dalimban (Bag.).

CALAMUS MEYENIANUS Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 425; Becc. in Rec. Bot. Surv. India 2 (1902) 217, *Webbia* 1 (1905) 346, Ann Bot. Gard. Calcutta 11 (1908) 82, 215, t. 63, Philip. Journ. Sci. 14 (1919) 349.

Luzon (Pangasinan, Nueva Vizcaya), *B. S.* 17755 *Otanes*, 11442 *McGregor*. In forests at low and medium altitudes. Endemic.

Local name: Bárit (Ilk.).

CALAMUS MICROCARPUS Becc. in Rec. Bot. Surv. India 2 (1902) 213, *Webbia* 1 (1905) 355, Ann. Bot. Gard. Calcutta 11 (1908) 117, 477, t. 218, Philip. Journ. Sci. 4 (1909) Bot. 627, 6 (1911) Bot. 230, 14 (1919) 356, Leaf. Philip. Bot. 8 (1919) 3055.

Luzon (Rizal, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Mindanao (Davao, Agusan, Lanao), *Elmer 10676, 13551, 15573, 16810, 16991, B. S. 9131 Robinson, 10465 McGregor, 12326 Foxworthy, F. B. 13305 Tamesis, 20667, 20668 Villamil, 27031 Villaflor, 24522 Cortez & Valderrama.* In primary forests at low and medium altitudes, ascending to 1,200 m. Endemic.

Local names Agúsi (Mbo.); kalapít (Bik.); lambútan (Tag.); pagipi (Mbo.); paríti (Bag.); sipai (Tag.); tandúlang-gúbat (Tag.).

Var. **DIMINUTUS** Becc. in Philip. Journ. Sci. 14 (1919) 356.

Luzon (Laguna), *B. S. 16599 Ramos.* In primary forests at low altitudes. Endemic.

CALAMUS MICROSPHAERION Becc. in Perk. Frag. Fl. Philip. (1904) 45, *Webbia 1* (1905) 349, *Ann. Bot. Gard. Calcutta 11* (1908) 114, 453, *t. 204*, Philip. Journ. Sci. 4 (1909) Bot. 627, 14 (1919) 353.

Luzon (Bataan), Culion, *Merrill 507, Whitford, F. B. 20917 De Leon.* In forests at low altitudes. Endemic.

Local names: Kulakíng (Tag.); lábit (Tag.).

Var. **SPINOSIOR** Becc. in Philip. Journ. Sci. 14 (1919) 354.

Palawan, *Merrill 9358, 9247.* In forests at low altitudes. Endemic.

CALAMUS MINDORENSIS Becc. in Philip. Journ. Sci. 2 (1907) Bot. 235, 4 (1909) Bot. 625, 14 (1919) 352, *Ann. Bot. Gard. Calcutta 11 Suppl. 88, t. 49* ined.

Luzon (Laguna), Mindoro, *McGregor 309, F. B. 6217 Merritt, F. B. 25745 Mabesa.* In forests at low altitudes. Endemic.

Local name: Tumulín (Tag.).

CALAMUS MITIS Becc. in Philip. Journ. Sci. 3 (1908) Bot. 341, 14 (1919) 353, *Ann. Bot. Gard. Calcutta 11 Suppl. 68, t. 36* ined.

Batan and Babuyan Islands, Luzon (Ilocos Norte), *B. S. 3817, 4075 Fénix, F. B. 24810 Paraíso.* In forests at low and medium altitudes. Endemic.

Local names: Matkóng (Ilk., Ting.); tebdas (Iv.).

CALAMUS MOSELEYANUS Becc. in Bot. Surv. India 2 (1902) 211, *Webbia 1* (1905) 348, *Ann. Bot. Gard. Calcutta 11* (1908) 105, 396, *t. 171*, *Leaf. Philip. Bot. 8* (1919) 3053, *Philip. Journ. Sci. 14* (1919) 352.

Mindanao (Zamboanga, Davao), Malanipa, *Hallier, Elmer 11886.* In forests at low altitudes, ascending to 1,000 m. Endemic.

Local name: Saráni (Bag.).

CALAMUS MULTINERVIS Becc. in Philip. Journ. Sci. 14 (1919) 352, *Leaf. Philip. Bot. 8* (1919) 3060, *Ann. Bot. Gard. Calcutta 11 Suppl. 88, t. 49* ined.

Mindanao (Davao), *Elmer 11791, 11955.* In primary forests, altitude about 1,200 m. Endemic.

Local names: Balála (Bag.); úbli (Bag.).

CALAMUS ORNATUS Blume ex Schultes f. Syst. 7² (1830) 1326, var. **PHILIPPINENSIS** Becc. in Webbia 1 (1905) 346, Ann. Bot. Gard. Calcutta 11 (1908) 102, 370, Philip. Journ. Sci. 6 (1911) Bot. 230 (excl. syn. Blanco), 14 (1919) 350, Leaf. Philip. Bot. 8 (1919) 3054.

Luzon (Cagayan, Bataan, Laguna, Tayabas, Rizal, Sorsogon), Polillo, Mindoro, Negros, Mindanao (Davao), Surigao, Basilan, *Elmer* 7625, 9086, 11236, 15607, 16886, 18425, 17999, *Whitford* 343, 502, *Williams* 337, *F. B.* 212 *Barnes*, 2854 *Meyer*, 2439 *Borden*, 7154, 17252 *Curran*, 23973, 25041 *Maneja*, 3911, 3913 *Merritt*, 6089 *Everett*, 6107 *Hutchinson*, *B. S.* 9266 *Robinson*, 10461 *McGregor*, 16619, 23364 *Ramos*, 947 *Mangubat*.

In primary forests at low and medium altitudes. Endemic, the species in Malay Peninsula, Sumatra, Borneo, and Java.

Local names: Agúbak (Tag.); alimúran (Ibn.); kalapí (S. L. Bis., P. Bis., Bik.); kayapí (C. Bis.); likúto (Tag.); limúran (Tag., Pamp., Sbl.); lukuán (Tag.); mayaṅgyang (C. Bis.); palaklakánin (Tag.); tubo (Bag.); uái (Tag.).

CALAMUS RAMULOSUS Becc. in Perk. Frag. Fl. Philip. (1904) 46, *Webbia* 1 (1905) 349, Ann. Bot. Gard. Calcutta 11 (1908) 114, 454, *t.* 205, Philip. Journ. Sci. 14 (1919) 354.

Luzon (Tayabas), *Merrill* 2070. In forests at low altitudes. Endemic. Local name: Paulís (Tag.).

CALAMUS REYESIANUS Becc. in Philip. Journ. Sci. 2 (1907) Bot. 237, 14 (1919) 353, Ann. Bot. Gard. Calcutta 11 Suppl. 86, *t.* 36 ined.

Luzon (Laguna, Tayabas), *Reyes*, *B. S.* 13312 *Ramos*, *F. B.* 25746 *Mabesa*. In forests at low and medium altitudes. Endemic.

Local names: Ápas (Tag.); lukuán (Tag.); samúlid (Tag.).

CALAMUS SAMIAN Becc. in Philip. Journ. Sci. 14 (1919) 353, Leaf. Philip. Bot. 8 (1919) 3059, Ann. Bot. Gard. Calcutta 11 Suppl. 92, *t.* 52 ined.

Luzon (Sorsogon), Mindanao (Davao), *Elmer* 11336, 17255. In primary forests, altitude about 1,300 m. Endemic.

Local name: Samian (Bag.).

CALAMUS SIPHONOSPATHUS Mart. Hist. Nat. Palm. 3 (1849) 342; Walp. Ann. 3 (1853) 491, 5 (1858) 832; Miq. Fl. Ind. Bat. 3 (1855) 137; Naves Novis. App. (1882) 275; Usteri Beitr. Ken. Philip. Veg. (1905) 130 (*tiphonolpathus*); Ceron Cat. Pl. Herb. Manila (1892) 175; Becc. in Rec. Bot. Surv. India 2 (1902) 213, *Webbia* 1 (1905) 350, Ann. Bot. Gard. Calcutta 11 (1908) 116, 471, Philip. Journ. Sci. 14 (1919) 354.

Calamus inflatus Warb. in Perk. Frag. Fl. Philip. (1904) 45.

Luzon (Cagayan, Isabela, Pampanga, Rizal, Laguna, Tayabas), Mindoro, Mindanao, *Merrill* 1891, 1643, *Elmer* 11652, *F. B.* 18558 *Alvarez*, 20633 *Villamil*, 23975 *Maneja*, 24321 *Acedillo*, 26223 *Mariano*, 11625 *Fischer*, 23825 *Mabesa*. In primary forests at low and medium altitudes, ascending to 1,200 m. Endemic.

Local names: Bejúco (Sp., Tag.); bíri (Tag.); dagdág (Ilk.); hamlís (Tag.); husi (Neg.); lambútan (Tag.); malaúban (Tag.); oban-óban (Bag.); palimanók (Pamp.); papákin (Tag.); sipai (Tag.); súkol (Tag.); susúko (Tag.); talóla (Tag.); valit (Iv.).

Var. **BATANENSIS** Becc. in Philip. Journ. Sci. 3 (1908) Bot. 342, 14 (1919) 355, Ann. Bot. Gard. Calcutta 11 Suppl. 115 ined.

Batan Islands (Batan), *B. S. 3611 Fénix*. In forests along streams. Endemic.

Var. **FARINOSUS** Becc. in Ann. Bot. Gard. Calcutta 11 (1908) 474, Philip. Journ. Sci. 14 (1919) 355.

Luzon (Rizal), (*Loher 7083*), *F. B. 23972 Maneja*. In forests at medium altitudes. Endemic.

Var. **OLIGOLEPIS** Becc. (*minor et major*) in Webbia 1 (1905) 353, Ann. Bot. Gard. Calcutta 11 (1908) 475, 476, Philip. Journ. Sci. 14 (1919) 355.

Luzon (Rizal, Tayabas), (*Loher, Warburg*). In forests at low or medium altitudes. Endemic.

Var. **POLYLEPIS** Becc. in Webbia 1 (1905) 354, Ann. Bot. Gard. Calcutta 11 (1908) 477, *t. 216*.

Luzon (Rizal), (*Vidal 4068*). In forests at low or medium altitudes. Endemic.

Var. **SUBLEVIS** Becc. l. cc. Philip. Journ. Sci. 14 (1919) 355.

Luzon (Ilocos Norte, Zambales, Rizal, Bataan, Laguna, Tayabas), *Whitford 308, F. B. 1577 Borden, 2635 Meyer, 6225, 6514 Curran, 25739 Mabesa, 13985 Merritt & Darling, 25100 Paraiso, B. S. 26643, 29601 Ramos & Edaño, 4713, 33182 Ramos, 10787 McGregor*. In forests at low and medium altitudes. Endemic.

CALAMUS SPINIFOLIUS Becc. in Rec. Bot. Surv. India 2 (1902) 202, Webbia 1 (1905) 348, Ann. Bot. Gard. Calcutta 11 (1908) 107, 410, *t. 178*, Philip. Journ. Sci. 14 (1919) 353.

Luzon (Pampanga, Bataan), Palawan, Panay, *F. B. 17309 Curran, 20947 De Leon, 1454 Ahern's collector*. In forests at low altitudes. Endemic.

Local names: Kuraklíng (Pamp., Tag.); síka (Tagb.); yantók (Tag.).

CALAMUS SYMPHYSIPUS Mart. in Hist. Nat. Palm. 3 (1849) 336; Becc. in Ann. Bot. Gard. Calcutta 11 (1908) 346, *t. 141*, Leaf. Philip. Bot. 8 (1919) 3049, Philip. Journ. Sci. 14 (1919) 351.

Luzon (Sorsogon), Bucas Grande, Mindanao (Agusan), *Elmer 16701, 13902, B. S. 35035 Ramos & Edaño*. In primary forests up to 900 m altitude. Celebes.

Local name: Bolánog (Mbo.).

CALAMUS TRISPERMUS Becc. in Perk. Frag. Fl. Philip. (1904) 46, Webbia 1 (1905) 349, Ann. Bot. Gard. Calcutta 11 (1908) 108, 412, *t. 180*, Philip. Journ. Sci. 4 (1909) Bot. 625, 14 (1919) 352.

Luzon (Rizal), *Merrill 1645, (Loher 7071)*. In forests at medium altitudes. Endemic.

CALAMUS USITATUS Blanco Fl. Filip. (1837) 265, ed. 2 (1845) 185, ed. 3, 1 (1877) 330, *t.* 99 (as *C. mollis*); Mart. Hist. Nat. Palm. 3 (1849) 340; Miq. Fl. Ind. Bat. 3 (1855) 131; Vidal Sinopsis Atlas (1883) 42, *t.* 93, *f.* D; Merr. Sp. Blancoanae (1918) 85; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) *t.* 11, 12.

Calamus mollis Naves in Blanco Fl. Filip. ed. 3 (1877-83) *t.* 99; Vidal Rev. Pl. Vasc. Filip. (1886) 280; Becc. in Rec. Bot. Surv. India 2 (1902) 204, Perk. Frag. Fl. Philip. (1904) 46, Ann. Bot. Gard. Calcutta 11 (1908) 82, 212, *t.* 61, 62, Webbia 1 (1905) 345, Philip. Journ. Sci. 3 (1908) Bot. 342, 14 (1919) 348, Leaf. Philip. Bot. 8 (1919) 3046; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88, Philip. Journ. Sci. 3 (1908) Bot. 399, Fl. Manila (1912) 124, non Blanco.

Calamus haenkeanus Mart. Hist. Nat. Palm. 3 (1849) 337; Kunth Enum. 3 (1841) 211; Walp. Ann. 2 (1852) 488, 5 1858) 831; Miq. Fl. Ind. Bat. 3 (1855) 127; Naves Novis. App. (1882) 275.

Batan and Babuyan Islands, Luzon (Ilocos Norte, Bontoc, Apayao, Nueva Ecija, Nueva Vizcaya, Bulacan, Rizal, Bataan, Zambales, Laguna, Tayabas, Camarines, Sorsogon), Palawan, Biliran, Samar, Mindanao (Surigao), *Elmer* 11969, 17052, 17727, 18201, *Merrill* 7407, 1642, 1743, 1411, 2077, *Williams* 561, *B. S.* 28131 *Fénix*, 11218, 18634 *McGregor*, 9571 *Robinson*, 4032 *Fénix*, 381 *Ramos*, *F. B.* 23981 *Maneja*, 24536 *Aguilar*, 17034, 5957, 6297, 5982, 6338, 17273, 8439 *Curran*, 2480, 2481, 3026 *Borden*, 24195 *Bawan*, 25102 *Paraíso*. In secondary forests and thickets at low and medium altitudes. Endemic.

Local names: Abét (Tag.); abít (Tag.); arít (Bag.); babúyan (Sbl.); bálit (Ilk.); bári (Ibn.); bárit (Ibn., Ilk.); digtan (Tag.); hanápas (Bik.); kakalsán (Tag.); pagípe (C. Bis.); ritoko (Ting.); tindarára (Bik.); tandúlang-páráng (Tag.); tagalóa (S. L. Bis.); tagókan (P. Bis.); tolólang-lutúkan (Tag.); uéi (Pang.); uái (Tag., Ilk.); uhói (Tagb.); válet (Iv.); yantók (Tag., Sbl.).

Var. **MAJOR** (Becc.) comb. nov.

Calamus mollis var. *major* Becc. in Webbia 1 (1905) 345, Philip. Journ. Sci. 14 (1919) 349.

Luzon (Bataan), *F. B.* 2499 *Meyer*, *Whitford* 80. In forests at low altitudes.

Local name: Uái (Tag.).

Var. **PALAWANICUS** (Becc.) comb. nov.

Calamus mollis var. *palawanicus* Becc. in Philip. Journ. Sci. 2 (1907) Bot. 233, 14 (1919) 345, Leaf. Philip. Bot. 8 (1919) 3047.

Palawan, *Elmer* 12607, *F. B.* 3613 *Curran*, *B. S.* 609 *Foxworthy*, 191, 196 *Bermejós*. In forests at low altitudes. Endemic.

CALAMUS VIDALIANUS Becc. in Rec. Bot. Surv. India 2 (1902) 212, Webbia 1 (1905) 350, Ann. Bot. Gard. Calcutta 11 (1908) 115, 464, *t.* 211, Philip. Journ. Sci. 4 (1909) Bot. 634.

Calamus horrens Vidal Rev. Pl. Vasc. Filip. (1886) 280; Ceron Cat. Pl. Herb. Manila (1902) 175, non Blume.

Luzon (Nueva Ecija, Rizal, Bataan, Tayabas, Camarines), *F. B. 20948 De Leon, 22162 Alvarez, B. S. 13214, 22286 Ramos, 33762 Ramos & Edaño*. In primary forests at low and medium altitudes.

Local names: Butarak (Ilk.); yantók (Tag.).

CALAMUS VINOSUS Becc. in Philip. Journ. Sci. 14 (1919) 352, *nomen nudum*, Leaf. Philip. Bot. 8 (1919) 3061.

Mindanao (Agusan), *Elmer 14153*. On forested ridges, altitude about 1,450 m. Endemic.

CALAMUS VIRIDISSIMUS Becc. in Philip. Journ. Sci. 14 (1919) 353, Leaf. Philip. Bot. 8 (1919) 3059, Ann. Bot. Gard. Calcutta 11 Suppl. 84, t. 47 ined.

Mindanao (Davao), *Elmer 11938*. In forests, altitude about 300 m. Endemic.

Local name: Akal (Bag.).

DOUBTFUL AND EXCLUDED SPECIES

CALAMUS ALBUS Pers.; Naves Novis. App. (1882) 276; Vidal Cat. Pl. Prov. Manila (1880) 47.

CALAMUS CONCINNUS Mart.; Naves op. cit. 275.

CALAMUS CURAG Blanco ex Miq. Fl. Ind. Bat. 3 (1855) 113; Becc. in Ann. Bot. Gard. Calcutta 11 (1908) 498.

Blanco published no such species, but casually mentions a form known locally as *kurag*, or *purag*.

CALAMUS EQUESTRIS Willd.; Naves Novis. App. (1882) 276.

CALAMUS JAVENSIS Blume; Naves op. cit. 275.

CALAMUS NITIDUS Mart.; Linden Ill. Hort. 28 (1881) 16.

"Philippines." This is an Indian species, Linden's specimen being either erroneously named or erroneously localized.

CALAMUS PISICARPUS Blume; Naves op. cit. 275.

CALAMUS PHILIPPENSIS Linden Ill. Hort. 28 (1881) 16, *nomen nudum*, "Philippines."

CALAMUS RHOMBOIDEUS Blume; Naves op. cit. 276.

CALAMUS SCIPIONUM Lour.; Naves l. c.

CALAMUS VERUS Blanco Becc. in Ann. Bot. Gard. Calcutta 11 (1908) 503.

This is merely a *lapsus calami*, Blanco being erroneously cited instead of Loureiro. The page reference given conforms to Loureiro, Flora Cochinchinensis, not to Blanco, Flora de Filipinas.

11. DAEMONOROPS Blume

DAEMONOROPS AFFINIS Becc. in Leaf. Philip. Bot. 8 (1919) 3042, Philip. Journ. Sci. 14 (1919) 359.

Mindanao (Agusan), *Elmer 13978*. In the mossy forests on exposed peaks and ridges, altitude about 1,200 m. Endemic.

Local name: Bag-bág (Mbo.).

DAEMONOROPS CLEMENSIANUS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 636, 14 (1919) 359, Ann. Bot. Gard. Calcutta 12¹ (1911) 163, t. 67.

Mindanao (Lanao), *Clemens 1227*. This is possibly a mixture of the fruits of *D. ochrolepis* Becc. and the leaves of some other species of doubtful status. Endemic.

DAEMONOROPS CURRANII Becc. in Philip. Journ. Sci. 2 (1907) Bot. 238, 4 (1909) Bot. 636, 14 (1919) 359, Ann. Bot. Gard. Calcutta 12 (1911) 142, t. 58, Leaf. Philip. Bot. 8 (1919) 3033.

Palawan, *Elmer 12663*, *F. B. 3791 Curran*. In forests at low altitudes. Endemic.

As later noted by Beccari the portion of leaf-sheath and the fragment of the spadix on the left of the original plate do not belong here but appertain to *D. pedicellaris* Becc.

Local names: Pit-pit (Tagb.); saranoi (Tagb.).

DAEMONOROPS GRACILIS Becc. in Leaf. Philip. Bot. 8 (1919) 3014, Philip. Journ. Sci. 14 (1919) 359.

Palawan, (*Elmer 12945*). In forests. Endemic.

DAEMONOROPS LOHERIANUS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 637, 14 (1919) 358, Ann. Bot. Gard. Calcutta 12¹ (1911) 104, t. 41.

Luzon (Rizal), (*Loher 7073*). In forests at low or medium altitudes. Endemic.

DAEMONOROPS MARGARITAE (Hance) Becc. var. **PALAWANICUS** Becc. in Philip. Journ. Sci. 4 (1909) Bot. 636, 14 (1919) 358, Leaf. Philip. Bot. 8 (1919) 3030, Ann. Bot. Gard. Calcutta 12¹ (1911) 57.

Palawan, *Merrill 9359*, *Elmer 12943*, *B. S. 899 Foxworthy*. In forests at low altitudes. The variety endemic, the species in southeastern China.

DAEMONOROPS MOLLIS (Blanco) Merr. Sp. Blancoanae (1918) 86.

Calamus mollis Blanco Fl. Filip. (1837) 264, ed. 2 (1845) 184, ed. 3, 1 (1877) 329; Mart. Hist. Nat. Palm. 3 (1849) 336; Kunth Enum. 3 (1841) 594; Walp. Ann. 3 (1853) 486, 5 (1858) 831; Miq. Fl. Ind. Bat. 3 (1855) 123.

Daemonorops gaudichaudii Mart. Hist. Nat. Palm. 3 (1849) 331; Walp. Ann. 3 (1853) 481; Naves Novis. App. (1882) 275; Becc. in Perk. Frag. Fl. Philip. (1904) 47, Rec. Bot. Surv. India 2 (1902) 226, Webbia 1 (1905) 355, Philip. Journ. Sci. 3 (1908) Bot. 342, 14 (1919) 359, Leaf. Philip. Bot. 8 (1919) 3031, Ann. Bot. Gard. Calcutta 12¹ (1911) 157, t. 64.

Daemonorops fuscus Mart. Hist. Nat. Palm. 3 (1849) 331; Walp. Ann. 3 (1853) 481, Becc. Rec. Bot. Surv. India 2 (1902) 226; Naves Novis. App. (1882) 274.

Calamus usitatus Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88, non Blanco.

Babuyan Islands to Mindanao, in most islands and provinces, *Whitford 289*, *1871*, *Elmer 11880*, *12494*, *Merrill Sp. Blancoanae 685*, *B. S. 1212*, *24394 Ramos*, *948 Mangubat*, *11358*, *14192 McGregor*, *4066 Fénix*, *29368*, *33720 Ramos & Edaño*, *24876*, *24777 Edaño*, *22577 Ramos & Deroy*, *28401*

Fénix, F. B. 21713 *Abellanosa*, 3741, 6218 *Merritt*, 22163 *Alvarez*, 2601 *Meyer*, 17310, 17218, 17252, 10190 *Curran*, 25040, 23974 *Maneja*, 24108 *Bernardo*, 25519 *Cenabre*, 24825 *Valencia*, 20892 *Tamesis*. In primary forests at low and medium altitudes, often common. Endemic.

Local names: Alimúran (Ibn.); alundayág (Ibn.); babúyan (Tag.); bag-bág (Tag.); baritau (Ibn.); batárag (Ilk.); batásan (P. Bis.); digtán (Tag.); diítan (Tag.); ditán (Ibn., Tag.); gatásan (Tag., S. L. Bis., Bik., P. Bis.); hamlis (Tag.); lakaóan (Bik.); labnig (Bik.); lisan (Sub.); mamuktóng (Tag.); nákat (Bag.); núkot (S. L. Bis.); samúlig (Bik.); sumúlíd (Tag.); tabáñga (Tag.); tákol (Bik.); tamarúra (Tagb.); tapnikid (C. Bis.); uái (Tag., Ilk.); uban-úban (P. Bis.); yantók (Tag.).

DAEMONOROPS OCHROLEPIS Becc. in *Perk. Frag. Fl. Philip.* (1904) 47, *Webbia* 1 (1905) 356, *Philip. Journ. Sci.* 6 (1911) Bot. 230, 14 (1919) 358, *Leaf. Philip. Bot.* 8 (1919) 3032, *Ann. Bot. Gard. Calcutta* 12¹ (1911) 160, t. 65, incl. var. *radulosus* Becc. in *Philip. Journ. Sci.* 4 (1909) Bot. 636, *Ann. Bot. Gard. Calcutta* 12¹ (1911) 162, t. 66.

Luzon (Ilocos Norte, Nueva Ecija, Rizal, Laguna, Tayabas, Sorsogon), Polillo, Mindanao, *Merrill* 2069, *Elmer* 11875, 14120, 14838, 16869, B. S. 13101 *Foxworthy & Ramos*, 13235 *Ramos*, 9039 *Robinson*, 26644 *Ramos & Edaña*, 33048 *Ramos*, F. B. 8490, 10190 *Curran*, 16221 *Mariano*, 25740 *Mabesa*. In primary forests at low and medium altitudes, ascending to 1,200 m. Endemic.

Local names: Ditán (Tag.); ligid-ligid (Tag.); mayáñgan (Mbo.); palaklakánin (Tag.); saladíñgan (Bag.); sumúlíd (Tag.).

DAEMONOROPS OLIGOLEPIS Becc. in *Leaf. Philip. Bot.* 8 (1919) 3035, *Philip. Journ. Sci.* 14 (1919) 359.

Mindanao (Davao), *Elmer* 11757. In primary forests, altitude about 1,100 m. Endemic.

Local name: Rogmán (Bag.).

DAEMONOROPS PANNOSUS Becc. in *Leaf. Philip. Bot.* 8 (1919) 3033, *Philip. Journ. Sci.* 14 (1919) 359.

Mindanao (Davao), *Elmer* 11600. In forests along streams, altitude about 1,200 m. Endemic.

Local name: Sabilóg (Bag.).

DAEMONOROPS PEDICELLARIS Becc. in *Leaf. Philip. Bot.* 8 (1919) 3040, *Philip. Journ. Sci.* 14 (1919) 358.

Mindanao (Agusan, Lanao, Davao), *Elmer* 11896, 14132, 13853, *Clemens* 1230. In primary forests, altitude 650 to 1,250 m. Endemic.

Local names: Hiyod (Mbo.); oban-óban (Mbo.); rogmán (Bag.).

DAEMONOROPS URDANETANUS Becc. in *Leaf. Philip. Bot.* 8 (1919) 3038, *Philip. Journ. Sci.* 14 (1919) 358.

Mindanao (Agusan), *Elmer* 14201. In primary forests, altitude about 1,600 m. Endemic.

Local name: Saháan (Mbo.).

DAEMONOROPS VIRESCENS Becc. in *Perk. Frag. Fl. Philip.* (1904) 47, *Webbia* 2 (1905) 357, *Ann. Bot. Gard. Calcutta* 12¹ (1911) 201, t. 91, *Philip. Journ. Sci.* 14 (1919) 358.

Palawan, *Merrill* 868. In forests at low altitudes. Endemic.

EXCLUDED SPECIES

DAEMONOROPS ACCEDENS Blume; Naves Novis. App. (1882) 274.

DAEMONOROPS ANGUSTIFOLIUS Griff.; Naves op. cit. 274.

DAEMONOROPS CALAPPARIUS Blume; Naves op. cit. 275.

DAEMONOROPS DRACO Mart.; Naves op. cit. 274.

DAEMONOROPS MELANOCHAETES Blume; Mart. Hist. Nat. Palm. 3 (1849) 203; Vidal Cat. Pl. Prov. Manila (1880) 47; Naves Novis. App. (1882) 275.

The Haenke specimen cited by Martius as representing Blume's species was doubtless wrongly identified.

DAEMONOROPS NIGER Blume; Naves op. cit. 274.

DAEMONOROPS OBLONGUS Mart; Naves l. c.

DAEMONOROPS RUMPHII Mart.; Naves l. c.

12. CARYOTA Linnaeus

CARYOTA CUMINGII Lodd. ex Mart. Hist. Nat. Palm. 3 (1849) 315; Walp. Ann. 3 (1853) 466; Miq. Fl. Ind. Bat. 3 (1855) 41; Hook. f. in Curtis's Bot. Mag. III 25 (1869) t. 5762; Naves Novis. App. (1882) 279; Vidal Sinopsis Altas (1883) 42, t. 93, f. B; Becc. in Webbia 1 (1905) 331, Perk. Frag. Fl. Philip. (1904) 48, Philip. Journ. Sci. 14 (1919) 338; Merr. Fl. Manila (1912) 123, Sp. Blancoanae (1918) 87.

Caryota urens Blanco Fl. Filip. (1837) 740, ed. 2 (1845) 510, ed. 3, 3 (1879) 142, t. 349; Naves Novis. App. (1882) 279, Vidal Cat. Pl. Prov. Manila (1880) 47, non Linn.

Caryota merrillii Becc. in Webbia 1 (1905) 333, Philip. Journ. Sci. 14 (1919) 338.

Luzon (Apayao, Union, Rizal, Nueva Ecija, Pangasinan, Nueva Vizcaya, Laguna, Tayabas), Mindoro, Palawan, Guimaras, Panay, Mindanao, Merrill 2380, 8036, 9360, Sp. Blancoanae 736, Elmer 5657, 18286, F. B. 10348, 8490 Curran, 6216 Merritt, 119 Gammill, 9077 Whitford & Hutchinson, 19961, 20492 Villamil, B. S. 873 Foxworthy, 28160 Fénix, 11243 McGregor. In forests at low and medium altitudes; sometimes planted in towns for ornamental purposes. Endemic.

Local names: Anívung (Ibn.); aníbong (Ibn., Ilk., Tag.); bahi (Man.); batikan (C. Bis.); datípan (Tag.); hagol (Bik.); lagípan (Pamp.); lulúg (Tag.); pugáhan (Tag.); pugúhan (Mbo.); patikan (P. Bis., Sul.); pola (Bag.); takípan (Sul.); tagípan (Pamp.).

CARYOTA MAJESTICA Linden in Ill. Hort. 28 (1881) 16, *nomen nudum*; Becc. in Perk. Frag. Fl. Philip. (1904) 48, Webbia 1 (1905) 335, Philip. Journ. Sci. 14 (1919) 338.

Luzon (Rizal), Merrill 1892, 8490. In forests at low and medium altitudes. Endemic.

Local name: Aníbong (Tag.).

This is perhaps *C. rumphiana* Mart. var. *philippinensis* Becc.; the material is imperfect.

CARYOTA MITIS Lour. Fl. Cochinch. (1790) 569, ed. Willd. (1793) 697; Becc. & Hook. f. Fl. Brit. Ind. 6 (1892) 423; Becc. in Philip. Journ. Sci. 14 (1919) 338, Leaf. Philip. Bot. 8 (1919) 3020.

Caryota sobolifera Wall. Cat. (1848) No. 8594; Naves Novis. App. (1882) 279.

Palawan, *Elmer* 12606, *F. B.* 3838, 4148 *Curran*. In forests near streams at low altitudes; cultivated in Manila for ornamental purposes. Burma and Indo-China to the Malay Peninsula, Sumatra, Java, and Borneo.

Local name: Bató (Tagb.).

CARYOTA RUMPHIANA Mart. var. **PHILIPPINENSIS** Becc. in Philip. Journ. Sci. 14 (1919) 337, Leaf. Philip. Bot. 8 (1919) 3019.

Caryota rumphiana Naves Novis. App. (1882) 274; Becc. in Webbia 1 (1905) 331, Perk. Frag. Fl. Philip. (1904) 48, vix Mart.

Caryota maxima Blume in Mart. Hist. Nat. Palm. 3 (1838) 195; Merr. Fl. Manila (1912) 123.

Luzon (Cagayan, Rizal, Laguna, Tayabas, Sorsogon), Mindoro, Negros, Mindanao (Agusan, Davao, Cotabato), *Elmer* 10940, 13623b, 9301, 14929, 15572, 15931, *Merrill* 2006, 8037, *Whitford* 1373, *F. B.* 10064, 19403, 17157, 17764 *Curran*, 19961 *Villamil*, 7850 *Curran* & *Merritt*, *B. S.* 13471 *Foxworthy* & *Ramos*, 11572 *Robinson*. In primary forests at low and medium altitudes.

Var. **OXYODONTA** Becc. in Philip. Journ. Sci. 14 (1919) 337.

Luzon (Laguna), *F. B.* 10045 *Curran*. In forested valleys at medium altitudes. The two varieties endemic, the species widely distributed in Malaya.

Local names: Aníhung (It.); anivung (Ibn.); bagsang (Bis.); kalalios (Bis.); lumang (Ibn.); pogáhan (Mbo.); pola (Bag.); pugáhan (Tag.); pulá (Mag.).

DOUBTFUL AND EXCLUDED SPECIES

CARYOTA GRIFFITHII Becc.; Naves Novis. App. (1882) 279.

CARYOTA PROPINQUA Blume; Naves l. c.

CARYOTA NANA Linden Ill. Hort. 28 (1881) 16, *nomen nudum* "Philippines."

CARYOTA SPECIOSA Linden l. c. *nomen nudum* "Philippines."

13. ARENGA Labillardière

ARENGA AMBONG Becc. in Philip. Journ. Sci. 2 (1907) Bot. 229, 14 (1919) 335.

Arenga tremula Becc. op. cit. 4 (1909) Bot. 612, 14 (1919) 335, quoad descr., excl. syn., non *Caryota tremula* Blanco.

Wallichia oblongifolia Becc. in Webbia 1 (1905) 328, non Griff.

Luzon (Tayabas, Sorsogon), Palawan, Basilan, Balabac, Cebu, Mindanao (Zamboanga), Sulu Archipelago, *Merrill* 5372, 8367, *Elmer* 12596, 16237, *Ahern* 653, *Copeland* 1681, *F. B.* 10213, 10280, 3542 *Curran*, 4032 *Hutchinson*, *B. S.* 735, 10816 *Foxworthy*, 1737 *McGregor*. In forests at low altitudes. Endemic.

Local names: Ambúng (Sul.); banísan (Sul.); bat-bát (Tagb.).

ARENGA PINNATA (Wurmb) Merr. Interpret. Herb. Amb. (1917) 119, Sp. Blancoanae (1918) 88; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 26, t. 78.

Saguerus pinnatus Wurm. in Verh. Bat. Genoots. 1 (1779) 351.

Borassus gomutus Lour. Fl. Cochinch. (1790) 618.

Arenga saccharifera Labill. in Mém. Inst. Paris 4 (1801) 209; F.-Vill. Novis. App. (1882) 280; Vidal Sinopsis Atlas (1883) 42, t. 93, f. C; Becc. in Perk. Frag. Fl. Philip. (1904) 48, Webbia 1 (1905) 328; Merr. Fl. Manila (1912) 126.

Sagrus gomutus Perr. in Mém. Soc. Linn. Paris 3 (1824) 142; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 306.

Caryota onusta Blanco Fl. Filip. (1837) 741, ed. 2 (1845) 511, ed. 3, 3 (1879) 143, t. 419.

Arenga gamuta Merr. in Philip. Journ. Sci. 9 (1914) Bot. 63. (There is no "*Saguerus gamuto* Houtt." as indicated in literature.)

Luzon (Rizal, Cavite, Bataan, Laguna, Tayabas), Polillo, Biliran, Mindanao. Planted here and there about towns, and abundant in some forested areas but never at any great distance from settled areas; generally planted in most islands and provinces. It has all the appearance of an introduced palm that has become naturalized. Sugar palm.

Local names: Bagátbat (C. Bis.); bagóbat (C. Bis.); bat-bát (C. Bis.); hibiók (P. Bis.); hidiók (Bik., P. Bis.); hiliók (Mbo.); ibiók (C. Bis.); idióg (C. Bis.); idiók (C. Bis.); igók (P. Bis.); irók (Sbl.); kábo-négro (Tag.); káong (Tag.); káuing (Tag.); onau (C. Bis.); rapítan (Ilk.); unau (C. Bis.).

ARENGA TREMULA (Blanco) Becc. in Philip. Journ. Sci. 4 (1909) Bot. 612, quoad syn. excl. descr.; Merr. Sp. Blancoanae (1918) 87; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 34, t. 9.

Caryota tremula Blanco Fl. Filip. (1837) 744, ed. 2 (1845) 512, ed. 3, 3 (1879) 144; Kunth Enum. 3 (1841) 549; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88.

Wallichia tremula Mart. Hist. Nat. Palm. 3 (1849) 315; Miq. Fl. Ind. Bat. 3 (1855) 34; Naves Novis. App. (1882) 280; Vidal Rev. Pl. Vasc. Filip. (1886) 279.

Arenga mindorensis Becc. in Perk. Frag. Fl. Philip. (1904) 48, Webbia 1 (1905) 329, Philip. Journ. Sci. 14 (1919) 355.

Saguerus mindorensis O. F. Cook in Invent. Seeds Import. U. S. Dept. Agr. 33 (1915) 10.

Didymosperma tremulum Wendl. & Drude in Kerch. Palm. (1878) 243.

Luzon (Bataan, Batangas, Laguna, Tayabas), Mindoro, Mindanao (Davao), Merrill 1790, Sp. Blancoanae 828, Elmer 18472, 11192, Baker 1807. F. B. 10203, 5469 Curran, 8768 Merritt, B. S. 12395 McGregor, 13102 Foxworthy & Ramos, 19468 Ramos. In thickets and secondary forests at low altitudes. Endemic.

Local names: Abígi (Bik.); abíki (Bik.); báris (Bag.); bat-bát (Tagb.); bilis (Bik.); dayáka (Tag.); dumáka (Tag.); dumayáka (Tag.); gumáka (Bik.); gumayáka (Tag.); rumáka (Bik.); tipon-tipon (Bik.).

DOUBTFUL AND EXCLUDED SPECIES

ARENGA MANILLENSIS Wendl. Index Palm. (1864) 3, *nomen nudum*; Naves Novis. App. (1882) 280.

Saguerus manillensis Lodd. l. c. in syn.

A *nomen nudum* and hence indeterminable; probably the same as one of the three species listed above.

ARENCA OBTUSIFOLIA Mart.; Naves Novis. App. (1882) 280.

A species of the Malay Peninsula allied to *A. pinnata* (Wurmb) Merr.; not known from the Philippines.

14. *ORANIA* Zippel

ORANIA DECIPIENS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 614, 14 (1919) 333.

Mindoro, *F. B. 4120 Merritt*. In primary forests at low altitudes. Endemic.

Var. *MINDANAOENSIS* Becc. in Philip. Journ. Sci. 14 (1919) 333.

Mindanao (Zamboanga), *F. B. 9179 Whitford & Hutchinson*. In primary forests at low altitudes. Endemic.

Var. *MONTANA* Becc. in Leaf. Philip. Bot. 8 (1919) 3018, Philip. Journ. Sci. 14 (1919) 333.

Mindanao (Agusan, Davao), *Elmer 11881, 13970*. In forests, altitude 800 to 1,000 m. Endemic.

Local name: Báñga (Tag., P. Bis., Bag., Mbo.).

ORANIA PALINDAN (Blanco) Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88, Philip. Journ. Sci. 1 (1906) Suppl. 32, Sp. Blancoanae (1918) 88; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 110, t. 42.

Caryota palindan Blanco Fl. Filip. ed. 2 (1845) 513, ed. 3, 3 (1879) 145.

Orania regalis Naves Novis. App. (1882) 279, non Blume.

Orania philippinensis Scheff. ex Becc. in Ann. Jard. Bot. Buitenz. 2 (1885) 156; Becc. in Webbia 1 (1905) 335, Philip. Journ. Sci. 14 (1919) 332.

Luzon (Cagayan, Union, Benguet, Zambales, Bataan, Rizal, Batangas, Laguna, Tayabas, Camarines), Samar, *F. B. 1610 Borden, 5818, 10410, 17068, 10708 Curran, 8413 Curran & Merritt, 2183 Meyer, 20988 Villamil, B. S. 12391 McGregor, 24107, 24131 Ramos, Elmer 61, 74, 18248, Merrill Sp. Blancoanae 144, Phil. Pl. 1900*. In forests at low and medium altitudes. Endemic.

Var. *SIBUYANENSIS* (Becc.) comb. nov.

Orania philippinensis Scheff. var. *sibuyanensis* Becc. in Leaf. Philip. Bot. 8 (1919) 3017, Philip. Journ. Sci. 14 (1919) 332.

Sibuyan, *Elmer 12066*. In primary forests, altitude about 225 m. Endemic.

Local names: Ambobáñga (Ibn.); báñga (Buk., P. Bis.); barangói (Tag., Pamp., Sbl.); niogniógan (Tag.); palindán (Tag.).

ORANIA PARAGUANENSIS Becc. in Webbia 1 (1905) 335; Philip. Journ. Sci. 14 (1919) 332.

Palawan, *Merrill 869, 9269, Phil. Pl. 1316*. In primary forests at low altitudes; probably not distinct from *O. palindan* Merr., which it exactly resembles in habit.

Local name: Báñga (Tagb.).

ORANIA RUBIGINOSA Becc. in Philip. Journ. Sci. 14 (1919) 333.

Luzon (Cagayan, Isabela, Tayabas), *B. S.* 10658, 10575 *McGregor*, 13390 *Ramos*, *F. B.* 17259, 17192 *Curran*. In forests at low altitudes. Endemic.

Local name: Kalahán (Ibn.).

EXCLUDED SPECIES

ORANIA MACROCLADUS Mart.; *Naves Novis. App.* (1882) 280.

15. HETEROSPATHE Scheffer

HETEROSPATHE ELATA Scheff. in *Ann. Jard. Bot. Buitenz.* 1 (1876) 141, 162; *Becc. Webbia* 1 (1905) 328, *Philip. Journ. Sci.* 14 (1919) 325, *Leaf. Philip. Bot.* 8 (1919) 3013; *Brown & Merr. in Philip. Bur. Forestry Bull.* 18 (1919) 86, *t.* 32.

Luzon (Cagayan, Laguna, Tayabas, Camarines), Mindoro, Masbate, Cebu, Samar, Panay, Siquijor, Dinagat, Bucas Grande, Mindanao (Davao, Zamboanga), *F. B.* 10352, 10481, 13238, 17064 *Curran*, 9684 *Merritt*, 19749 *Villamil*, 9030 *Whitford & Hutchinson*, *B. S.* 35243, 34624 *Ramos & Pascasio*, *Whitford* 1395, *Elmer* 11968, 18222, *Merrill* 2770. In forests at low altitudes, sometimes planted. Moluccas; introduced in Guam.

Local names: Balaniúg (Sub.); dagumáka (Ibn.); dayumáka (Ibn.); dumayáka (Ibn.); niogniógan (Tag.); sagasí (Tag.); sagisí (Bik., Tag., S. L. Bis.); salanióg (Bag.); tagisí (Bik.).

HETEROSPATHE NEGROSENSIS Becc. in *Philip. Journ. Sci.* 4 (1909) *Bot.* 611, 14 (1919) 326, *Leaf. Philip. Bot.* 8 (1919) 3014.

Luzon (Tayabas, Sorsogon), Negros, Panay, *Elmer* 9439, 10147, 17297, *Whitford* 1539, *F. B.* 13639 *Curran & Foxworthy*, *B. S.* 9443, 9358 *Robinson*, 30752, 28684 *Ramos & Edaño*. In forests at medium altitudes. Endemic.

HETEROSPATHE PHILIPPINENSIS Becc. in *Philip. Journ. Sci.* 4 (1909) *Bot.* 610, 14 (1919) 326, *Leaf. Philip. Bot.* 8 (1919) 3013.

Ptychoraphis philippinensis Becc. in *Ann. Jard. Bot. Buitenz.* 2 (1885) 90, *Malesia* 3 (1886) 109, *Webbia* 1 (1905) 47.

Rhopaloblaste sp. *Vidal Phan. Cuming. Philip.* (1885) 153.

Luzon (Apayao, Ifugao, Benguet, Nueva Vizcaya, Rizal, Laguna, Camarines, Sorsogon), Leyte, *F. B.* 10856, 10892 *Curran*, 18019 *Merritt*, 21752 *Fischer*, *B. S.* 19784, 19785 *McGregor*, 8198, 15260 *Ramos*, 28391 *Fénix*, 33606 *Ramos & Edaño*, *Elmer* 16519. In primary forests at medium altitudes, ascending to 1,600 m or higher. Endemic.

Local names: Sanakti (Ig.); tagisí (Bik.).

HETEROSPATHE SIBUYANENSIS Elm. *Leaf. Philip. Bot.* 8 (1919) 3014, *Philip. Journ. Sci.* 14 (1919) 325.

Sibuyan, *Elmer* 12350. In forests, altitude about 225 m. Endemic.

Local name: Bilis (C. Bis.).

16. ONCOSPERMA Blume

ONCOSPERMA GRACILIPES Becc. in *Philip. Journ. Sci.* 2 (1907) *Bot.* 228, 4 (1909) *Bot.* 610, 14 (1919) 330, *Leaf. Philip. Bot.* 8 (1919) 3017.

Luzon (Laguna, Tayabas, Sorsogon), Biliran, *Merrill 4010, Elmer 15223, Baker 3704, B. S. 18490, 22876 McGregor, 13213 Ramos, F. B. 10148, 10396 Curran, 26801 Mabesa*. In forests at low altitudes. Endemic.

Local name: Anibong (Tag.).

ONCOSPERMA HORRIDUM (Griff.) Scheff. in *Nat. Tijdschr. Nederl. Ind.* 32 (1871) 189; Becc. in *Philip. Journ. Sci.* 4 (1909) Bot. 610, 6 (1911) Bot. 230, 14 (1919) 330, *Leaf. Philip. Bot.* 8 (1919) 3017.

Areca horrida Griff. in *Calcutta Journ. Nat. Hist.* 5 (1845) 465.

Polillo, Mindanao (Agusan, Zamboanga, Davao), *Elmer 11876, 13886, Copeland 1626, B. S. 9277 Robinson*. In forests at low and medium altitudes, ascending to 800 m. Malay Peninsula and Archipelago.

Local names: Anibong (Tag., Mbo.); tanáian (Bag.).

ONCOSPERMA PLATYPHYLLUM Becc. in *Philip. Journ. Sci.* 4 (1909) Bot. 609, 14 (1919) 330.

Negros, *Whitford 1670, F. B. 17347 Curran*. In forests at low and medium altitudes. Endemic.

Local name: Anibong (P. Bis.).

ONCOSPERMA TIGILLARIA (Jack) Ridl. in *Journ. Str. Branch Roy. As. Soc.* 33 (1900) 173.

Areca tigillaria Jack *Malay Miscel.* 2⁷ (1820) 88.

Oncosperma filamentosa Blume *Rumphia* 2 (1836) 97, *t. 82, 103*; Becc. in *Philip. Journ. Sci.* 4 (1909) Bot. 610, 14 (1919) 330, *Leaf. Philip. Bot.* 8 (1919) 3016.

Areca nibung Mart. *Hist. Nat. Palm.* 3 (1838) 173, *t. 153*; Naves *Novis. App.* (1882) 279.

Palawan, *Elmer 12662, F. B. 3790 Curran*. In somewhat swampy places immediately back of the mangrove more or less subject to the influence of brackish water. Malay Peninsula and Archipelago.

Local name: Anibong (Bis., Tagb.).

17. ACTINORHYTIS Wendland and Drude

ACTINORHYTIS CALAPPARIA (Blume) Wendl. & Drude in *Linnaea* 39 (1875) 184; Becc. in *Leaf. Philip. Bot.* 8 (1919) 3010, *Philip. Journ. Sci.* 14 (1919) 324.

Areca calapparia Blume *Rumphia* 2 (1836) 70, in *nota, t. 100, f. 2*.

Seaforthia calapparia Mart. *Hist. Nat. Palm.* 3 (1849) 313.

Ptychosperma calapparia Miq. *Fl. Ind. Bat.* 3 (1855) 20; Naves *Novis. App.* (1882) 377.

Mindanao (Davao), *Elmer 11238*. Isolated trees in settled areas, altitude about 500 m, undoubtedly introduced. Widely distributed in the Malay Archipelago.

Local name: Tañgálo (Bag.).

18. PTYCHORAPHIS Beccari

PTYCHORAPHIS CAGAYANENSIS Becc. in *Philip. Journ. Sci.* 14 (1919) 326.

Heterospathe cagayanensis Becc. *op. cit.* 4 (1909) Bot. 611.

Luzon (Cagayan), *F. B. 12286 Klemme, 16757 Curran*. In primary forests at low altitudes. Endemic.

Local name: Dagumáka (Ibn.).

PTYCHORAPHIS ELMERI Becc. in Philip. Journ. Sci. 14 (1919) 328, Leaf. Philip. Bot. 8 (1919) 3010.

Heterospathe elmeri Becc. in Leaf. Philip. Bot. 2 (1909) 646.

Negros, *Elmer 9669*. In forests, altitude about 1,000 m. Endemic.

Local name: Bilisan (C. Bis.).

PTYCHORAPHIS INTERMEDIA Becc. in Leaf. Philip. Bot. 8 (1919) 3011, Philip. Journ. Sci. 14 (1919) 328.

Samar, Biliran, Mindanao (Agusan), *Elmer 13663*, *B. S. 18107 McGregor*, *24753 Ramos*. In forests 600 to 1,000 m and above. Endemic.

Local names: Lubilúbi (S. L. Bis.); marighoi (Mbo.).

PTYCHORAPHIS MICROCARPA Becc. in Philip. Journ. Sci. 14 (1919) 327.

Camiguin de Misamis, *B. S. 14678 Ramos*. In forests at medium altitudes. Endemic.

19. ADONIDIA Beccari

ADONIDIA MERRILLII (Becc.) Becc. in Philip. Journ. Sci. 14 (1919) 329; Brown & Merr. in Philip Bur. Forestry Bull. 18 (1919) 15, t. 2, 3.

Normanbya merrillii Becc. in Philip. Journ. Sci. 14 (1909) Bot. 606, t. 30, 31, Leaf. Philip. Bot. 8 (1919) 3009; Merr. Fl. Manila (1912) 127.

Actinorhytis calapparia Vidal Sinopsis Atlas (1883) 42, t. 94, f. B, non Wendl. & Drude.

Coron, Palawan, *Elmer 12708*, *Merrill 9415*. On limestone formations in open places, often subgregarious, and in sandy soil near the seashore at all altitudes. Cultivated in Manila for ornamental purposes since about 1875. A monotypic, endemic genus.

Local names: Búnga de china (Tag.); búnga de jolo (Tag.); oring-oring (Tagb.).

20. PINANGA Blume

PINANGA BARNESII Becc. in Webbia 1 (1905) 320, Philip. Journ. Sci. 3 (1908) Bot. 340, 6 (1911) Bot. 229, 14 (1919) 320, Leaf. Philip. Bot. 8 (1919) 3002.

Pinanga barnesii Becc. var. *macrocarpa* Becc. in Philip. Journ. Sci. 2 (1907) Bot. 227.

Babuyan Islands, Luzon (Cagayan, Union, Pampanga, Rizal, Bataan, Laguna, Camarines, Tayabas), Polillo, Catanduanes, Mindoro, Panay, Siargao, Mindanao, *F. B. 122 Barnes*, *2762 Meyer*, *6309*, *10757*, *17756 Curran*, *6853*, *11385 Merritt*, *B. S. 13248*, *13419 Ramos*, *6937 Robinson*, *10466*, *23202 McGregor*, *4144 Fénix*, *28525*, *31204*, *33700 Ramos & Edaño*, *13248*, *13419*, *13858*, *30528 Ramos*, *34862 Ramos & Pascasio*, *2003 Foxworthy*, *McGregor 275*, *Elmer 7924*, *9297*, *Whitford 799*. In primary forests chiefly at medium altitudes, ascending to 1,600 m. Endemic.

Local names: Abísi (Tag.); bubulin (Sub.); búngang-matsíng (Tag.); gasígan (Neg.); katidi (Ig.); sauag (Sub.).

PINANGA BASILANENSIS Becc. in Philip. Journ. Sci. 14 (1919) 322.

Mindanao (Zamboanga), Basilan, *B. S. 16119 Reillo*, *3662*, *36623 Ramos & Edaño*. In primary forests at low or medium altitudes. Endemic.

Local name: Buburis (Sub.).

PINANGA BATANENSIS Becc. in Philip. Journ. Sci. 3 (1908) Bot. 340, 14 (1919) 322.

Batan Islands (Batan), *B. S.* 3841 *Fénix*. Along mountain streams in forests. Endemic.

Local name: Dapiau (Iv.).

PINANGA COPELANDII Becc. in Webbia 1 (1905) 317, 14 (1919) 320, Leaflet. Philip. Bot. 8 (1919) 3002.

Luzon (Camarines, Sorsogon), Negros, Mindanao (Agusan, Davao), *Weber* 1134, *Copeland* 1283, *Elmer* 10467, 13875, *Whitford* 1669, *Merrill* 7281, *Phil. Pl.* 1594, *B. S.* 15918 *Fénix*, 23619 *Ramos*, *F. B.* 2415a *Cortez & Valderrama*. In damp primary forests at low and medium altitudes, ascending to 1,200 m. Endemic.

Local names: Abikí (Bik.); bagtoan (Mbo.); habikí (Bik.); hiliuásiu (P. Bis.); tambingálan (Mbo.); timbañgalan (Bag.).

PINANGA CURRANII Becc. in Philip. Journ. Sci. 2 (1907) Bot. 226, 14 (1919) 320.

Palawan, *F. B.* 3515 *Curran*. In primary forests at low altitudes. Endemic.

PINANGA GEONOMAEFORMIS Becc. in Philip. Journ. Sci. 4 (1909) Bot. 602, 14 (1919) 318.

Luzon (Rizal, Laguna, Tayabas), *F. B.* 10155 *Curran*, 26797 *Mabesa*, *B. S.* 20474 *Ramos*, 28565 *Ramos & Edaño*. In damp primary forests at low and medium altitudes. Endemic.

PINANGA HETEROPHYLLA Becc. in Philip. Journ. Sci. 14 (1919) 319.

Luzon (Sorsogon), Negros, *F. B.* 19346 *Curran*, 23635 *Ramos*. In primary forests, altitude 100 to 700 m. Endemic.

PINANGA INSIGNIS Becc. in Philip. Journ. Sci. 2 (1907) Bot. 223, 14 (1919) 322, Leaflet. Philip. Bot. 8 (1919) 3004.

Luzon (Apayao, Pangasinan, Nueva Vizcaya, Rizal, Laguna, Camarines, Tayabas, Sorsogon), Mindoro, Leyte, Mindanao (Agusan, Zamboanga, Surigao, Bukidnon), *Elmer* 8028, 13950, 14909, 15445, 17656, *Whitford* 1388, *Bolster* 222, *Wenzel* 534, *Merrill Phil. Pl.* 1135, *B. S.* 11233 *McGregor*, 33512 *Ramos & Edaño*, 15111, 13236 *Ramos*, 26067, 29969, 28237 *Fénix*, 25566 *Yates*, *F. B.* 9502 *Merritt*, 10491 *Curran*, 7791 *Curran & Merritt*. In primary forests at low and medium altitudes. Endemic.

Var. **GASTEROCARPA** Becc. in Philip. Journ. Sci. 14 (1919) 322.

Masbate, Negros, *F. B.* 17348 *Curran*, 11555 *Rosenbluth*, 13560 *Meyer & Foxworthy*. In primary forests at low and medium altitudes. The Luzon material placed here by Beccari I refer to the typical form of the species. Endemic.

Var. **LEPTOCARPA** Becc. l. c.

Negros, *Whitford* 1621. In primary forests at low and medium altitudes. Endemic.

Subsp. **LOHERIANA** Becc. l. c.

Luzon (Laguna), (*Loher 7092*), *B. S. 9789 Robinson*. In forests on Mt. Banahao, altitude about 1,000 m. Endemic.

Local names: Bañga (Tag.); saláuak (Sub.); sarámau (Tag.); saráuag (P. Bis., Mbo.); takón (Ibn.); tibanglán (Tag.); tibanglóan (Bik., S. L. Bis.).

PINANGA ISABELEMIS Becc. in *Philip. Journ. Sci.* 14 (1919) 318.

Luzon (Isabela), *B. S. 10660 McGregor*. In forests near the sea at low altitudes. Endemic.

PINANGA MACULATA *Porte* in *Ill. Hort.* 10 (1863) 92, *t. 361*; *Drude* in *Bot. Zeit.* 35 (1877) 636, *t. 5, f. 12-13*; *Becc. Malesia* 3 (1886) 145, *Webbia* 1 (1905) 325, *Perk. Frag. Fl. Philip.* (1904) 48, *Philip. Journ. Sci.* 14 (1919) 317; *C. H. Wright* in *Curtis's Bot. Mag.* IV 1 (1905) *t. 8011*.

Palawan, *Merrill 712*. In primary forests at an altitude of about 500 m, a juvenile sterile specimen, its identity with *Porte's* species problematical.

I have observed juvenile forms of *Pinanga* with mottled leaves in Laguna and Pampanga Provinces, Luzon; and in Mindoro. Endemic.

PINANGA MODESTA Becc. in *Philip. Journ. Sci.* 2 (1907) *Bot.* 225, 14 (1919) 318.

Bucas Grande, Mindanao (*Davao, Bukidnon, Lanao, Agusan, Zamboanga*), *Basilan, Clemens 487, Copeland, Merrill 8302, B. S. 15751 Fénix, 35098 Ramos & Pascasio, 39002 Ramos & Pascasio, F. B. 3987 Hutchinson, 4583 Mearns & Hutchison*. In primary forests at low and medium altitudes, ascending to 1,200 m. Endemic.

Local names: Lampigi (Lan.); mambuding (Yak.); tigahui (Buk.).

PINANGA NEGROSENSIS Becc. in *Leaf. Philip. Bot.* 2 (1909) 642, 8 (1919) 3006, *Philip. Journ. Sci.* 14 (1919) 322.

Negros, Dinagat, *Elmer 10030, B. S. 35182 Ramos & Pascasio*. In damp forested ravines, altitude about 900 m. Endemic.

PINANGA PHILIPPINENSIS Becc. in *Malesia* 3 (1887) 180, *Perk. Frag. Fl. Philip.* (1904) 48, *Webbia* 1 (1905) 324; *Merr. in Philip. Journ. Sci.* 1 (1906) *Suppl.* 32; *Becc. in Philip. Journ. Sci.* 14 (1919) 320, *Leaf. Philip. Bot.* 8 (1919) 3003; *Brown & Merr. in Philip. Bur. Forestry Bull.* 18 (1919) 112, *t. 44*.

Pinanga elmeri Becc. in *Webbia* 1 (1905) 322, *Philip. Journ. Sci.* 3 (1908) *Bot.* 341, 14 (1919) 320, *Leaf. Philip. Bot.* 8 (1919) 3004; *Merr. Philip. Journ. Sci.* 1 (1906) *Suppl.* 32, 2 (1907) *Bot.* 264.

Babuyan Islands, Luzon (*Cagayan, Ilocos Norte, Ifugao, Bontoc, Benguet, Pangasinan, Nueva Vizcaya, Bulacan, Rizal, Bataan, Zambales, Laguna, Tayabas, Sorsogon*), Mindoro, Leyte, Negros, Panay. Common on exposed ridges in the mossy forest, sometimes in ravines, altitude 800 to 1,800 m; represented by numerous collections. Endemic.

Local names: Búa-ti-báked (Ilk.); búa-ti-bákis (Ilk.); buñgañg-dakigan (Ilk.); gatilei (Bon.); katidai (Ig.); katildi (Ig.); lubiá (Tag.); mamá (Ilk.); matsing (Tag.).

PINANGA RIGIDA Becc. in Leaf. Philip. Bot. 2 (1909) 644, 8 (1919) 3005, Philip. Journ. Sci. 14 (1919) 321.

Luzon (Tayabas), Negros, *Elmer 10187*, *B. S. 28732 Ramos & Edaño*. In the mossy forest, altitude 1,200 to 1,800 m. Endemic.

Elmer 10147 is not cited by Beccari in the original description; this is pretty clearly referable to *Heterospathe negrosensis* Becc. The type, *Elmer 10187*, is clearly a *Pinanga*. This note is published in reference to Elmer's statement, Leaf. Philip. Bot. 8 (1919) 3006, that *Pinanga rigida* Becc. was probably referable to *Heterospathe* or *Ptychoraphis*.

PINANGA SAMARANA Becc. in Philip. Journ. Sci. 14 (1919) 321.

Samar, *B. S. 17535 Ramos*. In forests at low or medium altitudes. Endemic.

PINANGA SCLEROPHYLLA Becc. in Philip. Journ. Sci. 4 (1909) Bot. 603, 14 (1919) 322.

Mindoro, *F. B. 4468 Merritt*. In primary forests, altitude about 1,500 m. Endemic.

PINANGA SIBUYANENSIS Becc. in Philip. Journ. Sci. 14 (1919) 324, Leaf. Philip. Bot. 8 (1919) 3006.

Sibuyan, *Elmer 12425*. In forests along streams, altitude about 600 m. Endemic.

Local name: Tibaṅgán (C. Bis.).

PINANGA SPECIOSA Becc. in Webbia 1 (1905) 316, Philip. Journ. Sci. 14 (1919) 323, Leaf. Philip. Bot. 8 (1919) 3005.

Mindanao (Agusan, Lanao, Zamboanga, Davao), *Copeland 1265*, *Elmer 10484*, *13739*, *13941*, *14023*, *Clemens 374*, *F. B. 9271 Whitford & Hutchinson*, *28206 Elumir*. In primary forests, altitude 400 to 1,200 m. Endemic.

Local names: Banísan (Sub.); sakolon (Mbo.); saráuag (Bag.); sa-dáuag (Bag.).

PINANGA URDANETANA Becc. in Leaf. Philip. Bot. 8 (1919) 3008, Philip. Journ. Sci. 14 (1919) 321.

Mindanao (Agusan), Bukidnon, *Elmer 14137*, *B. S. 38612*, *38611*, *38610*, *38570 Ramos & Edaño*. On forested slopes, altitude about 1,400 m. Endemic.

Local names: Sagasí (Buk.); sagisí (Buk.); salang-ísog (Mbo.).

PINANGA UROSPERMA Becc. in Philip. Journ. Sci. 3 (1908) Bot. 341.

Babuyan Islands (Camiguin), Luzon (Apayao), *B. S. 4044*, *28038 Fénix*. On forested slopes at medium altitudes. Endemic.

Local name: Lagás (Ibn.).

PINANGA WOODIANA Becc. in Philip. Journ. Sci. 4 (1909) Bot. 604, 14 (1919) 322, Leaf. Philip. Bot. 8 (1919) 3003.

Luzon (Bontoc), Mindoro, Mindanao (Davao), *Merrill 5680*, *Elmer 10485*, *11334*, *B. S. 38078*, *40651 Ramos & Edaño*. In primary forests, altitude 1,050 to 1,350 m. Endemic.

Local names: Irár (Bag.); mumuing (Mang.).

21. ARECA Linnaeus

ARECA CATECHU Linn. Sp. Pl. (1753) 1189 (err. *cathecu*); Blanco Fl. Filip. (1837) 714, ed. 2 (1845) 494, ed. 3, 3 (1879) 129, t. 350; Miq. Fl. Ind. Bat. 3 (1855) 8; Vidal Sinopsis Atlas (1883) 42, t. 94, f. A; Naves Novis. App. (1882) 278; Becc. in Philip. Journ. Sci. 3 (1908) Bot. 340; 6 (1911) Bot. 229; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 32, Fl. Manila (1912) 128, Sp. Blancoanae (1918) 89; Becc. in Leaf. Philip. Bot. 8 (1919) 2997, Philip. Journ. Sci. 14 (1919) 304; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 20, t. 4, 5.

Areca alba "Rumph;" Naves Novis. App. (1882) 278.

Cultivated throughout the settled areas of the Philippines, in some places spontaneous; possibly a native of the Philippines, having been found once spontaneous in primary forests in Palawan. Old World Tropics generally, introduced in the New World.

Var. **BATANENSIS** Becc. in Philip. Journ. Sci. 14 (1919) 304.

Batan Islands, *B. S. 3824 Fénix*. Along a mountain stream. Endemic.

Var. **LONGICARPA** Becc. in Philip. Journ. Sci. 6 (1911) Bot. 229, 14 (1919) 304, t. 2, f. 2.

Polillo, *B. S. 10470 McGregor*. Probably from planted trees. Endemic.

Local names: Boá (Ilk., Ig., Ibn.); buá (Ibn.); buñga (Tag., Bis., Sul.); dapiau (Iv.); huá (It.); lúgos (Sul.); lúyos (Pamp.); pasa (Yak.) takob-tob (Bik.); vuá (Ibn.); va (It.).

Var. **SILVATICA** Becc. l. c. pl. 1, f. 2.

Palawan, *Merrill 9447*. In primary forests at a place where an old trail crosses a small stream near Lake Manguao. Endemic.

ARECA CALISO Becc. in Leaf. Philip. Bot. 8 (1919) 2998, Philip. Journ. Sci. 14 (1919) 310.

Luzon (Sorsogon), Biliran, Leyte, Samar, Mindanao (Agusan, Davao), *Elmer 11898, 13814, 14839, B. S. 18875 McGregor, 24130, 15388 Ramos, 24881 Edaño*. In forests, altitude 350 to 850 m. Endemic.

Local names: Kaliso (Bag.); sakolon (Mbo.).

ARECA CAMARINENSIS Becc. in Philip. Journ. Sci. 14 (1919) 309, t. 3, f. 3.

Luzon (Camarines), *F. B. 14268 Aguilar, B. S. 33829 Ramos & Edaño*. In forests at low altitudes. Endemic.

Local name: Monó (Bik.).

ARECA COSTULATA Becc. in Philip. Journ. Sci. 14 (1919) 312.

Leyte, Dinagat, *B. S. 15236 Ramos, 35242 Ramos & Pascasio*. In primary forests at low and medium altitudes. Endemic.

ARECA HUTCHINSONIANA Becc. in Philip. Journ. Sci. 14 (1919) 312.

Areca mammillata Becc. var. *mindanaoensis* Becc. op. cit. 4 (1909) Bot. 602.

Mindanao (Lanao, Zamboanga), *Basilan, F. B. 9141 Whitford & Hutchinson, 4019 Hutchinson, 23159 Agama, 25242 Alvarez, B. S. 36953, 37288 Ramos & Edaño, 38364 Brown.* In primary forests at low altitudes. Endemic.

Local names: Pisa (Yak.); pitha (Sub.); sambuláyan (Sub.).

ARECA IPOT Becc. in Leaf. Philip. Bot. 2 (1909) 639, 8 (1919) 2998, Philip. Journ. Sci. 6 (1911) Bot. 229 (var. *polillensis* Becc.) 14 (1919) 308, t. 3, f. 4; Merr. Sp. Blancoanae (1918) 89.

Areca catechu Linn. var. *humilis* Blanco Fl. Filip. (1837) 716, ed. 2 (1845) 495, ed. 3, 3 (1879) 120; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 88.

Areca glandiformis Naves Novis. App. (1882) 278 (prob.!), non Lam.

Luzon (Laguna, Tayabas), Polillo, *Merrill Sp. Blancoanae 844, F. B. 10178 Curran, B. S. 6938, 9712 Robinson, 10469, 10463 McGregor.* Planted in and about towns, and along streams in forests at low and medium altitudes. Endemic.

Local names: Buṅgaṅgípot (Tag.); ípod (Tag.); ípot (Tag.); maṅgípod (Tag.); saksíg (Tag.) saksík (Tag.).

ARECA MACROCARPA Becc. in Philip. Journ. Sci. 4 (1909) Bot. 601, 14 (1919) 306, t. 3, f. 2.

Mindanao (Zamboanga), *F. B. 9103 Whitford & Hutchinson.* Probably but a form or variety of *Areca catechu* Linn. Endemic.

ARECA MAMMILLATA Becc. in Philip. Journ. Sci. 2 (1907) Bot. 220.

Areca vidaliana Becc. op. cit. 222, 14 (1919) 312.

Palawan, *Merrill Phil. Pl. 1464, F. B. 3816 Curran, B. S. 21575 Escritor, 619 Foxworthy.* In swampy forests at low altitudes, and also on forested slopes. Endemic.

Local names: Boga (Tagb.); pita (Tagb.).

ARECA PARENS Becc. in Philip. Journ. Sci. 14 (1919) 307, t. 3, f. 1.

Luzon (Camarines), *F. B. 21712 Peñas, Soriano, & Abellanosa.* Along streams at low altitudes. Endemic.

Local name: Takobtób (Bik.).

ARECA WHITFORDII Becc. in Philip. Journ. Sci. 2 (1907) Bot. 219, 14 (1919) 306, Leaf. Philip. Bot. 8 (1919) 2997.

Luzon (Sorsogon), Mindoro, *Whitford 1872, Elmer 16240.* In semi-swampy forests at low altitudes. Endemic.

Var. **LUZONENSIS** Becc. in Philip. Journ. Sci. 14 (1919) 306.

Luzon (Cagayan, Nueva Vizcaya), *B. S. 11384 McGregor, F. B. 17770 Curran.* Wild in forests and in river swamps at low and medium altitudes. Endemic.

Local name: Buṅgaṅg-gúbat (Tag.).

EXCLUDED SPECIES

ARECA TRIANDRA Roxb.; Naves Novis. App. (1882) 278.

22. ELAEIS Jacquin

ELAEIS GUINEENSIS Jacq. Select. Am. (1763) 280, *t.* 172; Merr. Fl. Manila (1912) 126; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 84, *t.* 30, 31.

Cultivated for ornamental purposes, and producing fruits abundantly. A native of tropical Africa. African oil palm.

23. COCOS Linnaeus

COCOS NUCIFERA Linn. Sp. Pl. (1753) 1188; Blanco Fl. Filip. (1837) 716, ed. 2 (1845) 495 (*nucigera*), ed. 3, 3 (1879) 123, *t.* 364; Naves Novis. App. (1882) 282; Vidal Sinopsis Atlas (1883) 42, *t.* 94, *f.* D; Miq. Fl. Ind. Bat. 3 (1855) 64 (incl. var. *macrocarpa* Miq. var. *saccharina* Miq.); Merr. Fl. Manila (1912) 125, Sp. Blancoanae (1918) 89, 90.

Cocos mamillaris Blanco Fl. Filip. (1837) 722 (*mammilaris*), ed. 2 (1845) 499, ed. 3, 3 (1879) 123.

Very extensively cultivated in the Philippines, especially in those regions where the dry season is not too prolonged; one of the most important agricultural crops of the Archipelago. Pantropic in cultivation, probably a native of Polynesia; certainly not a native of the Philippines. Coconut.

Local names: Coco (Sp.); cocotéro (Sp.); gira-gira (Sbl.); í-ing (It.); iniúg (Ibn.); lobí (S. L. Bis.); lubí (C. Bis.); nióg (Ilk., Bik., Bis., Ibn., Tag.); ñgot-ñgot (Pamp., Sbl.); oñgót (Ibn.); ponlaing (Sub.); punláing (Yak.).

24. NYPA Wurmbe

(*Nipa* Thunb.)

NYPA FRUTICANS Wurmbe in Verh. Bat. Genoots. 1 (1779) 350; Miq. Fl. Ind. Bat. 3 (1855) 150; Naves Novis. App. (1882) 284; Vidal Sinopsis Atlas (1883) 42, *t.* 94, *f.* C; Blanco Fl. Filip. ed. 2 (1845) 461, ed. 3, 3 (1879) 68, *t.* 386; Becc. in Webbia 1 (1905) 357; Merr. Fl. Manila (1912) 125, Sp. Blancoanae (1918) 90; Brown & Merr. in Philip. Bur. Forestry Bull. 18 (1919) 98, *t.* 38-41 (mostly as *Nipa*).

Nipa fruticans Thunb. in Vet. Akad. Nya Handl. Stockh. 3 (1782) 231.

Nipa fruticosa Lam. ex Perr. in Mem. Soc. Linn. Paris 3. (1824) 132; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 306.

Nipa litoralis Blanco Fl. Filip. (1837) 662.

Along tidal streams in brackish swamps throughout the Philippines, in favorable habitats gregarious over large areas. India, through Malaya to tropical Australia.

Local names: Anípa (Ibn.); lasá (Tag.); nipá (Sp.-Fil.); paúid (Tag.); pinóg (It.); pinók (Ibn.); sagá (Sbl.); sasá (Pamp., Tag.), tatá (Ibn.).

EXCLUDED GENERA AND SPECIES

BENTINCKIA RENDA Mart.; Naves Novis. App. (1882) 280.

BORASSUS DICHOTOMA White; Naves op. cit. 281.

BORASSUS FLABELLIFORMIS Linn.; Naves op. cit. 280.

- CALYPTROCALYX SPICATUS Blume; Naves op. cit. 280.
- CHAMAEROPS EXCELSA Thunb.; Naves op. cit. 282.
- CERATOLOBUS GLAUDESCENS Blume; Naves op. cit. 273.
- DRYMOPHLOEUS APPENDICULATUS Becc.; Naves op. cit. 277.
- IGUANURA sp.; Vidal Rev. Pl. Vasc. Filip. (1886) 279; Ceron Cat. Pl. Herb. (1892) 174.
- KENTIA MOLUCCANA Becc.; Naves l. c.
- LODOICEA SEYCHELLARUM Labill.; Naves op. cit. 281.
- NENGA LATISECTA Blume; Naves op. cit. 277.
- PHOLIDOCARPUS IHUR Blume; Naves op. cit. 281.
- PLECTOCOMIA ELONGATA Mart.; Naves op. cit. 274.
- PTYCHOSPERMA COCHINCHINENSIS Miq.; Naves op. cit. 277.
- PTYCHOSPERMA KUHLLI Miq.; Naves l. c.
- PTYCHOSPERMA PUNICEA Miq.; Naves l. c.
- PTYCHOSPERMA RUMPHII Blume; Naves l. c.
- PTYCHOSPERMA SAXATILIS Blume; Naves l. c.
- PTYCHOSPERMA VESTIARIA Miq.; Naves l. c.
- RHAPIS ARUNDINACEA Llanos in Mem. Acad. Cienc. Madrid III 2 (1857) 499; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100; Naves Novis. App. (1882) 282, *nomen nudum*.
- RHAPIS FLABELLIFORMIS L'Hérit.; Naves Novis. App. (1882) 282.
- This is occasionally cultivated in Manila gardens as a pot plant, but it never produces flowers here.
- RHAPIS JAVANICA Blume; Naves l. c.

The above twenty-two species were credited to the Philippines by Naves, but none of the genera has known indigenous representatives in the Archipelago. A few of the species in the list now occur as recently introduced and cultivated forms.

Since about 1905 a large number of palms have been introduced into Manila for cultivation as ornamental species. Some of these are now mature and flower and fruit freely; others have not as yet produced flowers. None of them occurs outside of cultivation, and for the most part they are as yet rare in cultivation. The list of these recently introduced and cultivated palms includes such species as: *Acoelorrhaphe wightii* Wendl., *Archontophoenix alexandrae* H. Wendl. & Drude, *Attalea cohune* HBK., *Chrysalidocarpus lutescens* Wendl., *Cocos plumosa* Hook., *Coccothrinax garberi* Sarg., *Cyrtostachys lakka* Becc., *Dictyosperma alba* Wendl. & Drude, *Dypsis madagascariensis* Nichols, *Howea belmoreana* Becc., *Hypophorbe amaricaulis* Mart., *Latania commersonii* Gmel., *L. loddigesii* Mart., *Livistona australis* Mart., *L. chinensis* R. Br., *Martinezia caryotaefolia* HBK., *Oreodoxia regia* HBK., *Phoenix pusilla* Gaertn., *P. roebelenii* O'Brien, *P. rupicola* T. And., *Pinanga kuhlii* Blume, *Pritchardia gaudichaudii* Wendl., *P. pacifica* Seem. & H. Wendl., *Ptychosperma macarthurii*

H. Wendl., *Raphia ruffia* Mart., *Sabal adansonii* Guerns., *S. blackburneanum* Glazeb., *S. mauritiiforme* Griseb. & Wendl., *S. palmetto* Lodd., *Thrinax argentea* Lodd., *T. robusta* Wendl., and *Washingtonia filifera* Wendl.

ARACEAE

1. POTHOS Linnaeus

POTHOS ACUMINATISSIMUS Merr. in Philip. Journ. Sci. 11 (1916) Bot. 175.

Samar, B. S. 24125 Ramos. In primary forests at low altitudes. Endemic.

POTHOS CYLINDRICUS Presl Epim. (1851) 243; Walp. Ann. 3 (1853) 501; Schott Aroid. 1 (1853) 23, t. 48, Prodr. (1860) 569; Miq. Fl. Ind. Bat. 3 (1855) 180; Engl. in DC. Monog. Phan. 2 (1879) 86, Pflanzenreich 21 (1905) 36, f. 17, D, E; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 282; Naves Novis. App. (1882) 287.

Luzon (Camarines, Albay), Leyte, Panay, Mindanao (Davao), Cuming 914, Williams 2937, Piper 481, Wenzel 387, B. S. 22143 Ramos, 32451 McGregor. In primary forests at low and medium altitudes. Endemic. Local name: Palipí (Bik.).

POTHOS DOLICHOPHYLLUS Merr. in Philip. Journ. Sci. 11 (1916) Bot. 4.

Luzon (Sorsogon), Samar, B. S. 17628, 23708 Ramos. In primary forests at low altitudes. Endemic.

POTHOS HERMAPHRODITUS (Blanco) Merr. Sp. Blancoanae (1918) 90.

Batis hermaphrodita Blanco Fl. Filip. (1837) 791, ed. 2 (1845) 541, ed. 3, 3 (1879) 197.

Pothos longifolius Presl Epim. (1851) 242; Schott Aroid. 1 (1853) 22, t. 34, Prodr. (1860) 563; Walp. Ann. 3 (1853) 501; Miq. Fl. Ind. Bat. 3 (1856) 180; Naves Novis. App. (1882) 262; Engl. in DC. Monog. Phan. 2 (1879) 82, Pflanzenreich 21 (1905) 25, f. 9; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 282; Merr. Interpret. Herb. Amb. (1917) 124.

Pothos chapelieri Naves Novis. App. (1882) 286, non Schott.

Luzon (Pangasinan, Rizal, Laguna, Tayabas), Samar, Leyte, Panay, Siargao, Mindanao (Davao), Cuming 682, Merrill 2792, Sp. Blancoanae 57, Phil. Pl. 4, Elmer 18005, Wenzel 265, Loher 7045, B. S. 34871 Ramos & Pascasio, 13622, 24475 Ramos, 19211 Reillo, 16859 Serviñas, F. B. 24911 Mabesa. In primary forests at low and medium altitudes. Sumatra, Java, Celebes, Moluccas.

Local names: Apis (Tag.); hipan (Bik.); oro-óla (Bik.).

POTHOS INAEQUILATERUS (Presl) Engl. in DC. Monog. Phan. 2 (1879) 88, Pflanzenreich 21 (1905) 37; Naves Novis. App. (1882) 287.

Scindapsus inaequilaterus Presl Epim. (1851) 241; Walp. Ann. 3 (1853) 500.

Pothos cummingianus Schott in Oester. Bot. Wochenbl. (1855) 19, Prodr. (1860) 570; Walp. Ann. 5 (1858) 910 (*cummingianus*); Miq. Fl. Ind. Bat. 3 (1856) 181; Vidal Rev. Pl. Vasc. Filip. (1886) 282, Phan. Cuming. Philip. (1885) 154.

Luzon (Albay, Sorsogon), Samar, Leyte, *Cuming* 908, *Wenzel* 357, 1110, *Elmer* 17050 as *P. peninsularis* Elm., *B. S.* 24474 *Ramos*. In primary forests at low and medium altitudes. Endemic.

POTHOS LUZONENSIS (Presl) Schott in Oester. Bot. Wochenbl. (1855) 19, Aroid. 1 (1853) 25, t. 56, Prodr. (1860) 574; Miq. Fl. Ind. Bat. 3 (1856) 182; Walp. Ann. 5 (1858) 910; Naves Novis. App. (1882) 287; Engl. in DC. Monog. Phan. 2 (1879) 93, Pflanzenreich 21 (1905) 42; Vidal Rev. Pl. Vasc. Filip. (1886) 282, Phan. Cuming. Philip. (1885) 154.

Goniurus luzonensis Presl Epim. (1851) 244; Walp. Ann. 3 (1853) 500.

Luzon (Zambales, Rizal, Bataan, Camarines), *Cuming* 1045, *Merrill Phil. Pl.* 1529, *Loher* 7036, 7037, *F. B.* 5948 *Curran*. In primary forests at low and medium altitudes. Endemic.

Local name: Uarat-uárat (Bik.).

POTHOS OVATIFOLIUS Engl. in Pflanzenreich 21 (1905) 40.

Luzon (Tayabas, Camarines, Sorsogon), Samar, Mindanao (Agusan), *Ahern* 155, *Elmer* 14014, 15204, *B. S.* 24543 *Ramos*, 28549 *Ramos & Edaño*, *F. B.* 12247 *Curran*. In primary forests at low or medium altitudes. Endemic.

Local names: Kalíkip (Tag.); kalot-kágot (Bik.); maláang-lakop-lákop (S. L. Bis.).

POTHOS PHILIPPINENSIS Engl. in Pflanzenreich 21 (1905) 315; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 32.

Luzon (Apayao, Nueva Ecija, Bulacan, Bataan, Tayabas), Samar, *F. B.* 369 *Barnes*, *B. S.* 34031, 36414 *Ramos & Edaño*, 20742 *Escritor*, 28190 *Fénix*, 13054 *Ramos*, 1640 *Foxworthy*, *Whitford* 341, 1046bis, *Williams* 486. In primary forests at low and medium altitudes. Endemic.

Local names: Bagubalának (S. L. Bis.); baladaúan (Bik., Tag.); bayámi (Tag.); malagayámon (Tag.); oroalaypan (Bik.); tibátib (Tag.); tagoápi (Bik.).

POTHOS RUMPHII Schott Melet. 1 (1832) 21; Engl. Pflanzenreich 21 (1905) 38, f. 16; Naves Novis. App. (1882) 287; Merr. Interpret. Herb. Amb. (1917) 124.

Scindapsus rumphii Presl Epim. (1851) 241.

Luzon (Rizal, Laguna, Tayabas, Camarines, Albay, Sorsogon), Negros, Samar, Panay, Leyte, Mindanao, Basilan, *B. S.* 31078, 28585 *Ramos & Edaño*, 24144, 22132, 19367 *Ramos*, 15435 *Reillo*, *F. B.* 9656 *Curran*, *Loher* 7033, *Wenzel* 22, *Clemens* 685, *Merrill* 7309, *Weber* 1044, *Elmer* 15854 as *P. clemensii* Elm. Celebes, Moluccas, New Guinea.

Local names: Amlóng (Bik., Bis., Tag.); bislag (Mbo.); dílang-báka (Tag.); takúling (C. Bis.).

EXCLUDED SPECIES

POTHOS BARBERIANUS SCHOTT; Naves Novis. App. (1882) 287.

POTHOS BECCARIANUS Engl.; Naves l. c.

POTHOS GRACILIS Roxb.; Naves l. c.

POTHOS LEPTOSTACHYUS Schott; Naves l. c.

POTHOS LONGIPES Schott; Naves l. c.

POTHOS LOUREIRI Hook.; Naves l. c.

POTHOS MACROPHYLLUS De Vriese; Naves op. cit. 286.

POTHOS OXYPHYLLUS Miq.; Naves op. cit. 287.

POTHOS ROXBURGHII De Vriese; Naves op. cit. 286.

POTHOS SUMATRANUS Miq.; Naves op. cit. 287.

POTHOS VRIESEANUS Schott; Naves op. cit. 286.

POTHOS ZOLLINGERI Schott; Naves op. cit. 287.

POTHOS CELATOCAULIS N. E. Br. in Gard. Chron. 13 (1880) 200; Engl. in Bot. Jahrb. 25 (1898) 5, Pflanzenreich 37 (1908) 44.

This was described from Bornean material, and Engler, l. c., referred to it *Warburg 14603*, from Davao, Mindanao. Later, Pflanzenreich 21 (1905) 44, Engler admits the species as one of doubtful status from Borneo only. It is probably a *Rhaphidophora*, *R. celatocaulis* v. A. v. R. in Bull. Jard. Bot. Buitenz. 1 (1920) 382.

POTHOS SCANDENS Linn. Engl. Pflanzenreich 21 (1905) 26, quoad Philip.; Naves Novis. App. (1882) 286; Vidal Rev. Pl. Vasc. Filip. (1886) 177.

Vidal 1990 is referred here by Engler, who gives the range of the species as from India to Indo-China, Java, Sumatra, and Borneo. I suspect the Philippine specimen to be referable to *P. hermaphroditus* Merr. in which species the relative length and width of the petiole and lamina are exceedingly variable.

2. ANADENDRON Schott

ANADENDRON MONTANUM (Blume) Schott in Bonplandia 5 (1857) 45, Prodr. (1860) 391; Engl. in DC. Monog. Phan. 2 (1879) 97, Pflanzenreich 21 (1905) 49; Naves Novis. App. (1882) 287.

Calla montana Blume Cat. Gew. Buitenz. (1823) 62.

Mindanao (Zamboanga), *B. S. 37023 Ramos & Edaño*. In primary forests at low altitudes. Siam and Indo-China to the Malay Peninsula, Sumatra, Java, and Borneo.

EXCLUDED SPECIES

ANADENDRON AFFINE Schott; Naves Novis. App. (1882) 287.

3. POTHOIDIUM Schott

POTHOIDIUM LOBBIANUM Schott in Oester. Bot. Wochenbl. 7 (1857) 70, Aroid. 1 (1857) 26, t. 57, Gen. Aroid. (1858) 96, t. 96, Prodr. (1860) 576; Naves Novis. App. (1882) 287; Engl. in DC. Monog. Phan. 2 (1879) 95, Pflanzenreich 21 (1905) 46; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 32.

Northern Luzon (Cagayan) to Palawan and Mindanao, in most or all islands and provinces. In primary forests at low and medium altitudes. A monotypic genus. Celebes, Ternate.

Local names: Ariuan (Ibn.); balong-kahinai (P. Bis.); barálta (Tag.); magutapilak (P. Bis.); malagayáman (Sbl.); pakokalanġan (Mbo.); vara alta (Sp.).

4. ACORUS Linnaeus

ACORUS CALAMUS Linn. Sp. Pl. (1753) 324; Miq. Fl. Ind. Bat. 3 (1856) 175; Naves Novis. App. (1882) 287; Engl. in DC. Monog. Phan. 2 (1879) 216, Pflanzenreich 21 (1905) 309; Merr. Fl. Manila (1912) 130.

Luzon (Bontoc, Benguet), *Merrill Phil. Pl. 1789, Vanoverbergh 2585*. In swamps, altitude about 1,300 m, here native or thoroughly naturalized; cultivated here and there at low altitudes. North temperate regions generally.

Local names: Acóro (Sp.); bueng (Pamp.); dálau (Ilk.); dárau (Ilk.); déngau (Bon.); lubígan (Tag.).

ACORUS GRAMINEUS Soland. in Ait. Hort. Kew 1 (1789) 474; Engl. Pflanzenreich 21 (1905) 312; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 288.

Luzon (Laguna), *B. S. 14949 Ramos*. On rocks or ledges in beds of small streams at low or medium altitudes. Japan to southern China.

5. RHAPHIDOPHORA Hasskarl

RHAPHIDOPHORA ACUMINATA Merr. in Philip. Journ. Sci. 10 (1915) Bot. 265.

Luzon (Batangas, Sorsogon), Leyte, *Wenzel 1139, B. S. 23300 Ramos, 22669 Ramos & Deroy*. In primary forests at low altitudes. Endemic.

RHAPHIDOPHORA COPELANDII Engl. in Bot. Jahrb. 37 (1905) 115; Engl. ex Krause in Pflanzenreich 37 (1908) 49.

Rhaphidophora trinervia Elm. Leaf. Philip. Bot. 8 (1919) 3073.

Luzon (Benguet, Ifugao, Laguna), Mindanao (Davao), *Copeland 1193, Merrill 7700, Williams 2605, Elmer 18057, B. S. 19706 McGregor*. In forests and on ledges in thickets, altitudes 400 to 1,600 m. Endemic.

RHAPHIDOPHORA ELMERI Engl. & Krause in Engl. Bot. Jahrb. 44 (1910) Beibl. 101: 11.

Luzon (Tayabas, Sorsogon), Negros, *Elmer (9286, 9268), 10135a, 15116*. In forests at medium altitudes. Endemic.

RHAPHIDOPHORA GRANDIFOLIA Krause in Engl. Bot. Jahrb. 44 (1910) Beibl. 101: 11.

Negros, *Elmer 9464*. In forests. Endemic.

Local name: Dakoling (C. Bis.).

RHAPHIDOPHORA LAGUNENSIS Elm. Leaf. Philip. Bot. 8 (1919) 2072.

Luzon (Laguna), *Elmer 17812, Pascasio*. In the mossy forest, summit of Mt. Maquiling, altitude about 1,100 m. Endemic.

RHAPHIDOPHORA MERRILLII Engl. Bot. Jahrb. 37 (1905) 115; Engl. & Krause in Engl. Pflanzenreich 37 (1908) 51; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 32, 5 (1910) Bot. 335, Fl. Manila (1912) 129, Sp. Blancoanae (1918) 90.

Pothos pinnata Blanco Fl. Filip. (1837) 646, ed. 2 (1845) 450, ed. 3, 3 (1879) 48, t. 339, non Linn.

Epipremnum merrillii Engl. & Krause in Engl. Pflanzenreich 37 (1908) 137.

Luzon (Bontoc, Benguet, Union, Nueva Vizcaya, Zambales, Bataan, Rizal, Laguna, Tayabás, Sorsogon), Polillo, Palawan, Leyte, Negros, Mindanao, Merrill 9434, *Sp. Blancoanae* 715, *Elmer* 5539, 7296, 15135 as *Epipremnum glaucocephalum* Elm., *B. S.* 11292, 10370 *McGregor*, 6004, 5883 *Curran*. In thickets and forests at low and medium altitudes. Endemic.

Local names: Amiling (Is.); amlóng (Bik., Bis.); amólong (Ilk.); amúling (Is.); bagák (S. L. Bis.); bakág (Bik.); balikupkúp (Bis.); balision (Is.); bisako (Bis.); daila (Bis.); dibátib (Bis.); dukúp (Bon.); garban (Bis.); garusíba (Ibn.); gátgati (Bon.); gayáman (Sbl.); horoa (Bis.); kígau (Bon.); kilat (Sbl.); malapakpák-baláuai (Tag.); mamisi (Ig.); maragayáman (Sbl.); tabátib (Tag.); takóling (C. Bis.); takótin (Bis.); tampinbanal (Tag.); tibátib (Tag.).

RHAPHIDOPHORA MONTICOLA Krause in Engl. Bot. Jahrb. 44 (1910) Beibl. 101: 12.

Luzon (Laguna, Sorsogon), Negros, Leyte, *Elmer* 10098, 16790 as *R. bulusanensis* Elm., Merrill 7154, *Wenzel* 1160, *B. S.* 23455 *Ramos*, *F. B.* 21387 *Villamil*. In primary forests ascending into the mossy forest, altitude 700 to 1,200 m. Endemic.

RHAPHIDOPHORA PERKINSIAE Engl. in Engl. Bot. Jahrb. 37 (1905) 115; Engl. & Krause in Engl. Pflanzenreich 37 (1908) 37.

Luzon (Benguet, Bataan, Rizal, Laguna, Tayabas), *Elmer* 6305, *Loher* 7035, *Whitford* 1169, Merrill 9667, *F. B.* 26747 *Mabesa*, 7826 *Curran & Merritt*. In primary forests on trees and in thickets about limestone cliffs and boulders, altitude 1,000 to 1,600 m. Endemic.

Local names: Ámlong (Tag., Ig.); madulúnang (Ig.).

RHAPHIDOPHORA PHILIPPINENSIS Engl. & Krause in Engl. Pflanzenreich 37 (1908) 137.

Samar, Mindanao (Agusan, Lanao), *Clemens* 1229, *Elmer* 13776, *B. S.* 24405 *Ramos*. In forests at low and medium altitudes. Endemic.

Local name: Mananáu (S. L. Bis.).

RHAPHIDOPHORA RIGIDA Krause in Leaf. Philip. Bot. 6 (1914) 2284.

Luzon (Sorsogon), Negros, *Elmer* 10220, 15056. In primary forests, altitude about 1,250 m; apparently very near *R. philippinensis* Engl. & Krause.

RHAPHIDOPHORA STENOPHYLLA Elm. Leaf. Philip. Bot. 8 (1919) 3073.

Luzon (Laguna), *Elmer* 18449. Margins of forests at low altitudes. Endemic.

RHAPHIDOPHORA TODAYENSIS Krause in Leaf. Philip. Bot. 6 (1914) 2284.

Mindoro, Mindanao (Davao), *Elmer* 10849, *B. S.* 39735 *Ramos*. In forests at low altitudes. Endemic.

RHAPHIDOPHORA WARBURGII Engl. Bot. Jahrb. 37 (1905) 116; Engl. & Krause in Engl. Pflanzenreich 37 (1908) 51.

Luzon (Cagayan or Isabela), *Warburg* in herb. Berol. Probably from primary forests at low altitudes. Endemic.

EXCLUDED SPECIES

- RHAPHIDOPHORA DECURSIWA Schott; Naves Novis. App. (1882) 288.
 RHAPHIDOPHORA FALLAX Schott; Usteri Beitr. Ken. Philip. Veg. (1905) 130.
 RHAPHIDOPHORA LANCIFOLIA Schott; Naves Novis. App. (1882) 288.
 RHAPHIDOPHORA MONTANA Schott; Naves l. c.
 RHAPHIDOPHORA PEEPLA Schott; Naves l. c.
 RHAPHIDOPHORA PERTUSA Schott; Naves l. c.
 RHAPHIDOPHORA SYLVESTRIS Engl.; Naves l. c.

6. EPIPREMNUM Schott

EPIPREMNUM ELMERIANUM Engl. ex Engl. & Krause in Pflanzenreich 37 (1908) 66.

Leyte, *Elmer* 7295. In forests at low and medium altitudes. Endemic.

EPIPREMNUM PHILIPPINENSE Engl. & Krause in Engl. Pflanzenreich 37 (1908) 137.

Luzon (Laguna, Tayabas, Sorsogon), *Elmer* 7623, 9253, 14522, 15113 as *E. miniatum* Elm., 16422 as *E. sorsogonense* Elm., B. S. 10052, 20548, 23299 Ramos, 35419 Martelino & Edaño. In forests, altitude about 650 m. Endemic.

EPIPREMNUM PINNATUM (Linn.) Engl. Pflanzenreich 37 (1908) 60, f. 25.

Pothos pinnata Linn. Sp. Pl. ed. 2 (1763) 1374.

Epipremnum robinsonii Krause in Notizbl. Kgl. Bot. Gart. Berlin 5 (1912) 266.

Luzon (Tayabas), Polillo, Mindanao (Davao), *Whitford* 746, B. S. 15866 *Fénix*, *Robinson* 9181, *Elmer* 16061 as *E. multicephalum* Elm. In forests near the seashore. Tenasserim to New Guinea and Polynesia.

Local names: Lukmói (Tag.); tibátib (Tag.).

EPIPREMNUM TRUNCATUM Engl. & Krause in Engl. Pflanzenreich 37 (1908) 63.

Luzon (Camarines, Sorsogon), Biliran, Leyte, Mindanao (Agusan), *Elmer* 7291, 13729, 14721, *Merrill* 7343, F. B. 27244 *Palma*, B. S. 18925 *McGregor*. In primary forests at low altitudes. Endemic.

Local names: Lukmói (Tag.); pasagianit (Mbo.).

EXCLUDED SPECIES

EPIPREMNUM GIGANTEUM Schott; Naves Novis. App. (1882) 288.

7. EPIPREMNOPSIS Engler

EPIPREMNOPSIS HUEGELIANA (Schott) Engl. in Engl. & Krause Pflanzenreich 37 (1908) 138.

Rhaphidophora huegeliana Schott in Bonplandia 5 (1857) 45, Prodr. (1860) 384.

Epipremnum mirabile Naves Novis. App. (1882) 288; Vidal Rev. Pl. Vasc. Filip. (1886) 281; Engl. in Bot. Jahrb. 37 (1905) 119; Usteri Beitr. Ken. Philip. Veg. (1905) 130, non Schott.

Epipremnum medium Engl. in DC. Monog. Phan. 2 (1879) 250, quoad Philip.; Naves Novis. App. (1882) 289.

Epipremnopsis media Engl. ex Engl. & Krause in Pflanzenreich 37 (1908) 1, quoad Philip.

Batan and Babuyan Islands, Luzon (Bataan, Laguna, Camarines, Sorsogon), Catanduanes, Panay, Mindanao (Agusan, Zamboanga, Cotabato), *B. S. 35662 Ramos & Edaña, 22852 McGregor, 11716 Robinson, 3902 Féniz, Elmer 16052, 18223, Merrill 8071.* In forests at low and medium altitudes. Endemic.

Local names: Lusígut (Ilk.); takhib (Iv.).

8. SCINDAPSUS Schott

SCINDAPSUS AUREUS (Lind. & André) Engl. in Pflanzenreich 37 (1908) 80; Merr. Fl. Manila (1912) 139.

Pothos aurea Lindl. & André in Ill. Hort. (1880) 69, t. 381.

Cultivated for ornamental purposes, but never observed in flower. A native of the Solomon Islands.

SCINDAPSUS CURRANII Engl. & Krause in Pflanzenreich 37 (1908) 138.

Luzon (Cagayan, Laguna, Tayabas, Sorsogon), Mindoro, Negros, Camiguin de Misamis, *F. B. 8014, 8027 Curran & Merritt, 13711 Curran, 11376 Merritt, 26916 Mabesa, 25707 Foxworthy, B. S. 24064 Ramos, Elmer 18475, 14590, 16556.* In primary forests at low altitudes. Endemic.

Local name: Lukmói (Tag.).

SCINDAPSUS FALCIFOLIUS Engl. Bot. Jahrb. 37 (1905) 119; Engl. & Krause in Pflanzenreich 37 (1908) 79.

Jolo, Warburg in herb. Berol. Celebes.

SCINDAPSUS HEDERACEUS Schott in Bonplandia 5 (1857) 45; Naves Novis. App. (1882) 289; Engl. & Krause in Pflanzenreich 37 (1908) 76, 138, f. 32, E-N.

Scindapsus pictus Hassk. var. *argyraeus* Engl. Bot. Jahrb. 25 (1898) 13, Pflanzenreich 37 (1908) 80.

Scindapsus argyraea Engl. in DC. Monog. Phan. 2 (1879) 255; Naves Novis. App. (1882) 289 (*argyracea*).

Luzon (Laguna, Sorsogon), Catanduanes, Negros, Leyte, *F. B. 4274 Everett, 26436 Catalan, B. S. 15019, 12028, 30383 Ramos, 23151 McGregor, 29233 Ramos & Edaña, Elmer 15230.* In primary forests at low altitudes. Malay Peninsula and CochinChina to Borneo, Sumatra, and Java.

Scindapsus argyraea Engl. as represented by *Engl. Araceae exsic. 265* is a juvenile form with mottled leaves; this is common in Laguna Province, Luzon. In mature plants the leaves are usually not mottled.

Local names: Maragayáman (Ilk.); putoputóhan (Tag.).

EXCLUDED SPECIES

SCINDAPSUS PERTUSUS Schott; Vidal Cat. Pl. Prov. Manila (1880) 47.

9. SPATHIPHYLLUM Schott

SPATHIPHYLLUM COMMUTATUM Schott in Oester. Bot. Wochenbl. 7 (1857) 158, Prodr. (1860) 424, Ann. Mus. Bot. Lugd.-Bat. 3 (1863) 79; Engl. in DC. Monog. Phan. 2 (1879) 230; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 281; Naves

Novis. App. (1882) 288; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 264; Engl. & Krause in Pflanzenreich 37 (1908) 133, f. 55, A-G.

Luzon (Tayabas, Camarines, Albay, Sorsogon), Polillo, Biliran, Mindoro, Leyte, Mindanao, *B. S.* 33596, 38523 *Ramos & Edaño*, 10401, 18707 *McGregor*, 23571, 15385 *Ramos*, 6476 *Robinson*, *F. B.* 11415 *Merritt*, *Wenzel* 235, *Merrill* 5486, 4080, *Copeland* 1263, *Williams* 2364. In damp primary forests at low and medium altitudes. Celebes, Amboina.

All other species of this genus are confined to tropical America.

10. CYRTOSPERMA Griffith

CYRTOSPERMA MERKUSII (Hassk.) Schott in Oester. Bot. Wochenbl. (1859) 61, Prodr. (1860) 403; Engl. in DC. Monog. Phan. 2 (1879) 271, Pflanzenreich 48 (1911) 20, f. 7, G-K.

Lasia merkusii Hassk. Cat. Hort. Bogor. (1844) 59, Pl. Jav. Rar. (1848) 161.

Cyrtosperma griffithii Merr. in Philip. Journ. Sci. 2 (1907) Bot. 422 sphalm.

Luzon (Sorsogon), Mindoro, Samar, Leyte, Mindanao, *Elmer* 15234, *Merrill* 5218, *B. S.* 1515 *Bermejos*, 24267 *Ramos*, 6678 *Robinson*. In ravines along streams, probably often planted. Java and Borneo to New Guinea.

Local names: Galiáng (Bik); palau (C. Bis.); palauán (S. L. Bis., P. Bis.).

11. AMORPHOPHALLUS Blume

AMORPHOPHALLUS CAMPANULATUS (Roxb.) Blume ex Decne. in Ann. Mus. Hist. Nat. Paris 3 (1834) 366; Miq. Fl. Ind. Bat. 3 (1856) 201; Engl. in DC. Monog. Phan. 2 (1879) 309, Pflanzenreich 48 (1911) 76; Naves Novis. App. (1882) 289; Merr. Fl. Manila (1912) 130, Philip. Journ. Sci. 10 (1915) Bot. 290, Interpret. Herb. Amb. (1917) 127, Sp. Blancoanae (1918) 91.

Arum campanulatum Roxb. Pl. Coromandel 3 (1819) 69, t. 272.

Arum decurrens Blanco Fl. Filip. (1837) 656, ed. 2 (1845) 457, ed. 3, 3 (1879) 62.

Amorphophallus decurrens Kunth Enum. 3 (1841) 581; Miq. Fl. Ind. Bat. 3 (1856) 202; Engl. in DC. Monog. Phan. 2 (1879) 319, Pflanzenreich 48 (1911) 108; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 290.

Luzon, most or all provinces, Mindoro. In thickets and secondary forests, along roads, trails, etc., at low and medium altitudes in the settled areas, often common. India through Malaya to Polynesia.

Local names: Ánto (Bis.); ápon (Tag.); apong-ápong (Tag.); bagáng (Ibn.); bagóng (Bik., Sul.); oroi (Bis.); pamangkílon (Bis.); puñgápuñg (Tag.); tigi-ñga magmánto (Ilk.); tokod-banuá (Pamp.).

AMORPHOPHALLUS LONGISPATHEUS Engl. & Gehrm. in Engl. Pflanzenreich 48 (1911) 91.

Mindanao (Davao), *Williams* 2684. In thickets or forests at medium altitudes. Endemic.

AMORPHOPHALLUS LUZONIENSIS Merr. in Philip. Journ. Sci. 10 (1915) Bot. 289.

Luzon (Cagayan), *F. B.* 19560 *Curran*, *Weber*. In crevices of limestone ledges at low altitudes. Endemic.

AMORPHOPHALLUS MERRILLII Krause in Notizbl. Kgl. Bot. Gart. Berlin 5 (1912) 266.

Panay (Batbatan Island), Cavilli, *Merrill 7177, McGregor*. Abundant in primary forests on Cavilli, a small island in Sulu Sea. Endemic.

AMORPHOPHALLUS RIVIERI Durieu Cat. Jard. Bordeaux (1869) 12, var. **KONJAC** (C. Koch) Engl. in DC. Monog. Phan. 2 (1879) 313, Pflanzenreich 48 (1911) 85.

Amorphophallus konjac C. Koch in Berl. Allg. Gartenz. (1858) 166.

Luzon (Rizal, Bataan, Laguna), Mindoro, Palawan, Balabac, Panay, Mindanao, *McGregor 274, Elmer 18185, Merrill 9419, Whiford 1338, F. B. 17649 Curran, B. S. 26211 Ramos, 15641 Fénix*. In thickets and forests at low altitudes. Indo-China, Japan.

Local names: Bulañg-an (Mang.); puñgápung (Tag.).

AMORPHOPHALLUS VARIABILIS Blume in Rumphia 1 (1835) 146, t. 35, 37; Kunth Enum. 3 (1841) 33; Miq. Fl. Ind. Bat. 3 (1856) 200; Naves Novis. App. (1882) 290; Engl. in DC. Monog. Phan. 2 (1879) 313, Pflanzenreich 48 (1911) 80; Usteri Beitr. Ken. Philip. Veg. (1895) 130.

Luzon (Union), Mindanao (Davao), Basilan, *Copeland 1264, B. S. 15444 Reillo, 9999 Robinson*. In light forests at low altitudes. Java.

Local name: Bagóng (Bag.).

EXCLUDED SPECIES

AMORPHOPHALLUS BULBIFER Blume; Naves Novis. App. (1882) 290.

AMORPHOPHALLUS DUBIUS Blume; Naves op. cit. 289.

AMORPHOPHALLUS GIGANTEUS Blume; Naves op. cit. 290.

AMORPHOPHALLUS TITANUM Becc.; Naves op. cit. 289.

AMORPHOPHALLUS TUBERCULIGERA Schott; Naves l. c.

12. HOMALOMENA Schott

HOMALOMENA ELMERI Engl. ex Engl. & Krause in Pflanzenreich 55 (1912) 42.

Palawan, *Elmer 13229, F. B. 21586 Danao*. In forests along streams at low altitudes; scarcely distinct from *H. palawanensis* Engl. & Krause. Endemic.

HOMALOMENA PALAWANENSIS Engl. ex Engl. & Krause in Pflanzenreich 55 (1912) 39, f. 19.

Palawan, *B. S. 912 Foxworthy*. Margins of pools at low altitudes. Endemic.

HOMALOMENA PHILIPPINENSIS Engl. ex Engl. & Krause in Pflanzenreich 55 (1912) 55, f. 33.

Homalomena rubescens Kunth var. *latifolia* Engl. Bot. Jahrb. 55 (1898) 18.

Luzon (Isabela, Union, Pangasinan, Pampanga, Rizal, Bataan, Laguna, Sorsogon), Mindoro, Palawan, Panay, Leyte, Biliran, Negros, Mindanao. Along small streams in primary forests at low altitudes; common. Endemic.

Local names: Alipaiu (Tag.); alupayi (Tag.); hayau (Sub.); payau (Tag.); salet-nga-nalabaga (Ilk.); tahig (Bik.).

HOMALOMENA PYGMAEA (Hassk.) Engl. in Bot. Jahrb. 25 (1898) 18; Engl. & Krause in Engl. Pflanzenreich 55 (1912) 34, f. 15.

Aglaonema pygmaeum Hassk. in Hoev. & De Vriese Tijdschr. 9 (1842) 161, Cat. Hort. Bogor. (1844) 57.

Chamaecladon lanceolatum Miq. Fl. Ind. Bat. 3 (1856) 212, t. 40; Schott Gen. Aroid. (1858) t. 60.

Negros, Whitford 1591. Along stream beds, in cañons, altitude about 175 m. Malay Peninsula, Sumatra, Java, Borneo, and Celebes.

EXCLUDED SPECIES

HOMALOMENA ALBA Hassk.; Naves Novis. App. (1882) 290.

HOMALOMENA AROMATICA Schott; Naves l. c.

HOMALOMENA COERULESCENS Jungh.; Naves l. c.

HOMALOMENA MINOR Griff.; Naves l. c.

HOMALOMENA MIQUELIANA Schott; Naves l. c.

HOMALOMENA PROPINQUA Schott; Naves l. c.

HOMALOMENA RUBESCENS Kunth; Naves l. c.

HOMALOMENA RUBRA Hassk.; Naves l. c.

HOMALOMENA SAGITTIFOLIA Jungh.; Naves l. c.

HOMALOMENA CORDATUM Schott; Usteri Beitr. Ken. Philip. Veg. (1905) 130.

HOMALOMENA PALUDOSUM Hook. f.; Usteri l. c.

HOMALOMENA ROSTRATUM Griff.; Usteri l. c.

The last three species were credited to the Philippines on the basis of specimens collected in Negros by Usteri and determined by Engler. The latter author, however, does not credit them to the Philippines in his recent monographic treatment of the group.

13. SCHISMATOGLOTTIS Zollinger and Moritz

SCHISMATOGLOTTIS CALYPTRATA (Roxb.) Zoll. & Mor. Syst. Verz. (1854) 83; Naves Novis. App. (1882) 291; Usteri Beitr. Ken. Philip. Veg. (1905) 130; Merr. Interpret. Herb. Amb. (1917) 129; Engl. Pflanzenreich 55 (1912) 115 (var. *concolor* Hallier f.).

Luzon (Bataan, Cavite, Laguna, Tayabas, Albay), Samar, Leyte, Panay, Mindanao, Merrill 7603, 7273 (forma *grandifolia* Engl.), 7275, 6703, B. S. 24952 *Fénix*. In damp shaded ravines usually along streams on banks and cliffs at low and medium altitudes. Burma, through Malaya to New Guinea.

Local name: Alapáyi (Tag.).

- SCHISMATOGLOTTIS LATIFOLIA** Miq. Fl. Ind. Bat. 3 (1856) 214, var. **VIRIDIS** Engl. ex Engl. & Krause in Pflanzenreich 55 (1912) 117.
Schismatoglottis luzonensis Engl. Pflanzenreich 55 (1912) 121, non Engl. op. cit. 88.
Alocasia merrillii Engl. & Krause in Engl. Bot. Jahrb. 54 (1916) Beibl. 118: 124, Pflanzenreich 71 (1920) 79.

Luzon (Pampanga, Nueva Vizcaya, Laguna, Camarines), Mindanao, *B. S.* 2015 Foxworthy, 33830, 33783 Ramos & Edaña, 11248 McGregor, 6736 Robinson, *F. B.* 1243 Borden Copeland 1310, Baker 1034, Elmer 17829, Merrill 8072. In shaded ravines along streams on banks and cliffs at low and medium altitudes. Java, Celebes.

- SCHISMATOGLOTTIS LUZONENSIS** Engl. in Pflanzenreich 55 (1912) 88.

Luzon (Rizal), Panay, Merrill 5047, *B. S.* 30875 Ramos & Edaña. On wet ledges in small stream beds at low altitudes. Endemic.

- SCHISMATOGLOTTIS MERRILLII** Engl. in Pflanzenreich 55 (1912) 89.

Luzon (Bontoc, Benguet), *B. S.* 12679 Fénix, 8772 McGregor, *F. B.* 5099 Curran. On ledges and cliffs in ravines along streams, altitude 1,200 to 1,600 m. Endemic.

Local names: Atúped (Ilk.); píhau (Ig.); píkau (Bon.); picháu (Ig.).

- SCHISMATOGLOTTIS MINDANAOANA** Engl. in Pflanzenreich 55 (1912) 103.

Mindanao (Agusan, Lanao, Davao), Merrill 7320, Elmer 13489, Clemens 1225. In damp primary forests along small streams at low and medium altitudes. Endemic.

- SCHISMATOGLOTTIS PUSILLA** Engl. in Bot. Jahrb. 1 (1881) 184, Pflanzenreich 55 (1912) 103, f. 64.

Leyte, Jagor 1018a in herb. Berol. Endemic.

- SCHISMATOGLOTTIS RAMOSII** Engl. in Pflanzenreich 55 (1912) 98.

Schismatoglottis rizalensis Engl. op. cit. 100.

Luzon (Bulacan, Rizal), *B. S.* 3351, 13023 Ramos. On cliffs and boulders along streams in shaded ravines at low altitudes. Endemic.

Local name: Talísig (Tag.).

- SCHISMATOGLOTTIS WARBURGIANA** Engl. in Bot. Jahrb. 55 (1898) 20, Pflanzenreich 55 (1912) 90, f. 57.

Mindanao (Davao), Warburg 14605 in herb. Berol. In forests up to 600 m altitude. Endemic.

EXCLUDED SPECIES

SCHISMATOGLOTTIS PICTA Schott; Naves Novis. App. (1882) 291.

SCHISMATOGLOTTIS RUPESTRIS Zoll. & Mor.; Naves l. c.

14. AGLAONEMA Schott

AGLAONEMA COMMUTATUM Schott Syn. (1856) 123, Prodr. (1860) 304, Gen. Aroid. (1859) t. 59; Regel Gartenflora (1865) t. 470; Engl. Pflanzenreich 64 (1919) 27, f. 10, G.

Aglaonema marantaefolium Blume var. *maculatum* Hook. in Curtis's Bot. Mag. III 21 (1865) t. 5500.

Aglaonema marantifolium Blume var. *commutatum* Engl. in DC. Monog. Phan. 2 (1879) 441.

Not uncommon in cultivation in Manila as an ornamental on account of its mottled leaves. Widely cultivated in tropical botanical gardens, probably a native of the Moluccas.

For some years I thought that this might be *Aglaonema maculatum* Blume. Blume's species is based entirely on *Dracunculo Luzonis primo seu indorum Gavay Sinbuga* Camel in Ray Hist. Nat. 3 App. (1704) 36, and Camel's specimens were from Leyte and apparently from wild plants. I now suspect *Aglaonema maculatum* Blume to be a form of *A. oblongifolium* Kunth. There is no evidence that *A. commutatum* Schott is of ancient cultivation in the Philippines.

AGLAONEMA CORDIFOLIUM Engl. Pflanzenreich 64 (1915) 29, f. 14, A-C.

Mindanao (Davao), (*Warburg 14604*). In primary forests at low altitudes. Endemic.

AGLAONEMA DENSINERVIUM Engl. Bot. Jahrb. 37 (1905) 134, Pflanzenreich 64 (1915) 13.

Negros, Leyte, Siquijor, Mindanao, Basilan, *Copeland 418*, *Piper 466*, *Clemens 476*, *Wenzel 1128*, *Williams 2429*, *Merrill 7347*, *8204*, *F. B. 4774*, *Mearns & Hutchinson, B. S. 26139 Fénix*, *15452 Reillo*. In primary forests at low and medium altitudes. Endemic.

AGLAONEMA HAENKEI (Presl) Schott in Bonplandia 7 (1859) 30, Prodr. (1860) 301; Naves Novis. App. (1882) 291; Engl. in DC. Monog. Phan. 2 (1879) 439, Pflanzenreich 64 (1915) 17; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 281; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 400.

Scindapsus haenkei Presl Epim. (1851) 240; Walp. Ann. 3 (1853) 500; Miq. Fl. Ind. Bat. 3 (1856) 183.

Caladium seguinum Presl Rel. Haenk. 1 (1827) 148, non Vent.

Babuyan Islands, Luzon (Apayao, Laguna, Cavite), Panay, *Merrill 4186*, *Elmer 17668*, *B. S. 4089 Fénix*, *35350 Martelino & Edaño*, *12404*, *32590 McGregor*, *10008 Ramos*, *F. B. 26047 Mabesa*. In primary forests chiefly along streams at low and medium altitudes. Celebes.

AGLAONEMA OBLONGIFOLIUM (Roxb.) Kunth Enum. 3 (1841) 55; Merr. Interpret. Herb. Amb. (1917) 129.

Calla oblongifolia Roxb. Hort. Beng. (1814) 65, Fl. Ind. ed. 2, 3 (1832) 516.

Aglaonema marantifolium Blume Rumphia 1 (1835) 153, t. 66; Engl. in DC. Monog. Phan. 2 (1879) 441, Pflanzenreich 64 (1915) 27, t. 10, f. A-F; Naves Novis. App. (1882) 291; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 33.

? *Aglaonema maculatum* Blume Rumphia 1 (1835) 154; Kunth Enum. 3 (1841) 56; Miq. Fl. Ind. Bat. 3 (1856) 217; Schott Syn. (1856) 123, Prodr. (1860) 306, Engl. in DC. Monog. Phan. 2 (1879) 442, Pflanzenreich 64 (1915) 34.

Northern Luzon (Cagayan) to Mindanao, in most or all islands and provinces, incl. *Elmer 16520* as *A. alpinum* Elm. In primary forests chiefly along streams; the leaves are sometimes mottled. Amboina, Banda.

Local names: Damping-banál (Tag.); pamitágen (Mang.); pauangkílon (C. Bis.).

AGLAONEMA PHILIPPINENSE Engl. in Bot. Jahrb. 25 (1898) 21, Pflanzenreich 64 (1915) 29.

Aglaonema latifolium Engl. Bot. Jahrb. 37 (1905) 134.

Luzon (Laguna, Tayabas, Camarines), Panay, Mindanao (Davao), Gregory 86, B. S. 18139 Robinson, 23084, 23088, 22847, 23077 McGregor. In forests at low altitudes. Endemic.

EXCLUDED SPECIES

AGLAONEMA MODESTUM Schott ex Engl. in DC. Monog. Phan. 2 (1879) 442, Pflanzenreich 64 (1915) 29, f. 13.

The type of this is a specimen collected by Gaudichaud supposedly in Luzon, but which in all probability came from Macao; the species occurs in southern China, but has not been detected in the Philippines. See Merrill in Philip. Journ. Sci. 13 (1918) Bot. 133.

AGLAONEMA PICTUM Kunth; Naves Novis. App. (1882) 291.

AGLAONEMA SIMPLEX Schott; Naves Novis. App. (1882) 291; ? Vidal Rev. Pl. Vasc. Filip. (1886) 281.

Vidal's doubtful record for this Bornean species was based on his No. 1996 from San Mateo, Rizal Province, Luzon. I have not seen the specimen but suspect it to be the same as *A. haenkei* Schott or *A. oblongifolium* Kunth.

15. ALOCASIA Necker

ALOCASIA ATROPURPUREA Engl. Pflanzenreich 71 (1920) 132.

Luzon (Ifugao, Mt. Polis), B. S. 19663 McGregor. In the mossy forest at medium altitudes. Endemic.

ALOCASIA CULIONENSIS Engl. in Engl. Bot. Jahrb. 37 (1905) 135; Engl. & Krause Pflanzenreich 71 (1920) 102.

Luzon (Pangasinan), Mindoro, Culion, Palawan, Panay, Merrill 555, 9819, 9824, 9507, B. S. 29883 Fénix, 31264 Ramos & Edaño, 39602 Ramos. In primary forests at low altitudes. Endemic.

To be compared with *A. longiloba* Miq.

ALOCASIA HETEROPHYLLA (Presl) Merr. in Philip. Journ. Sci. 3 (1908) Bot. 220, Fl. Manila (1912) 133; Engl. & Krause Pflanzenreich 71 (1920) 101, f. 22.

Caladium heterophyllum Presl Rel. Haenk. 1 (1827) 148.

Colocasia ? *heterophylla* Kunth Enum. 3 (1841) 40; Miq. Fl. Ind. Bat. 3 (1856) 204; Schott Prodr. (1860) 139.

Alocasia warburgii Engl. in Engl. Bot. Jahrb. 25 (1898) 25; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 33.

Alocasia manilensis Engl. Bot. Jahrb. 25 (1898) 23; Engl. & Krause Pflanzenreich 71 (1920) 95, f. 19, E, F.

Luzon (Ilocos Norte, Pampanga, Rizal, Bataan, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Alabat, Panay, Negros, Samar, Jolo. In primary forests at low and medium altitudes. *Alocasia manilensis* Engl. is merely a juvenile form. Celebes.

Local name: Naroknarok (Mbo.).

ALOCASIA MACRORRHIZA (Linn.) Schott Melet. (1832) 18; Engl. in DC. Monog. Phan. 2 (1879) 502; Naves Novis. App. (1882) 293; Merr. Interpret. Herb. Amb. (1917) 130, Sp. Blancoanae (1918) 91; Engl. & Krause Pflanzenreich 71 (1920) 84, f. 15.

Arum macrorrhizon Linn. Sp. Pl. (1753) 965.

Calla maxima Blanco Fl. Filip. (1837) 658.

Arum grandifolium Blanco op. cit. ed. 2 (1845) 458, ed. 3, 3 (1879) 63, t. 177.

Calla badian Blanco Fl. Filip. (1837) 658.

Alocasia indica Naves Novis. App. (1882) 293, Usteri Beitr. Ken. Philip. Veg. (1905) 130; Merr. Fl. Manila (1912) 133, Engl. & Krause Pflanzenreich 71 (1920) 87, quoad Philip., non Schott.

Throughout the Philippines in old clearings and secondary forests at low and medium altitudes (incl. *Elmer 17381* as *A. talipan* Elm.), the older plants often with a tall stout caudex. India to Malaya; planted for ornamental purposes in other tropical countries.

Local names: Abá (Ibn.); aba-ába (Ig.); badiáng (Tag., Bis.); bagiáng (Bis.); bíra (Ilk.); bíga (Tag., Ilk., Bis., Pamp.); bilbíla (Bon.); gabi (Bik.); galiáng (Bis.); gandus (Pamp.); malabíga (Tag.); ragiáng (Bis.); sininába (Ilk.); talípan (Bik.); taliáng (Bis.).

ALOCASIA MAQUILINGENSIS Merr. in Philip. Journ. Sci. 13 (1918) Bot 3.

Luzon (Laguna, Sorsogon), *Baker 868*, *Merrill 7153*, *Elmer 17955*, *16181* as *A. vulcanica* Elm., *B. S. 6737 Robinson*, *16926 Servinas*. On slopes in primary forests, on Mt. Maquiling, altitude 100 to 600 m. Endemic.

ALOCASIA MICHOLITZIANA Sander in Gard. Chron. 51 (1912) Suppl. XV, f. 9; N. E. Br. in Curtis's Bot. Mag. IV 9 (1913) t. 8522; Engl. & Krause Pflanzenreich 71 (1920) 89.

Luzon (Apayao, Benguet), *Merrill 9744*, *B. S. 28024*, *28421 Fénix*. In thickets about limestone boulders, altitude about 1,300 m. Endemic.

ALOCASIA PORTEI (Schott) Engl. & Becc. in DC. Monog. Phan. 2 (1879) 645; Naves Novis. App. (1882) 294; Merr. Fl. Manila (1912) 132. *Schizocasia portei* Schott in Bonplandia 10 (1862) 148; Engl. in DC. Monog. Phan. 2 (1879) 495; Engl. & Krause Pflanzenreich 71 (1920) 117.

Alocasia indica Schott var. *heterophylla* Engl. in DC. Monog. Phan. 2 (1879) 502.

Alocasia indica Schott var. *diversifolia* Engl. Pflanzenreich 71 (1920) 88.

Luzon (Bataan, Laguna, Camarines), *Lyon*, *Holman*. In old clearings at medium altitudes. Endemic.

Local name: Badiáng (Bik.).

ALOCASIA REVERSA N. E. Br. in Gard. Chron. III 8 (1890) 38; Hook. f. in Curtis's Bot. Mag. III 52 (1896) t. 7498; Usteri Beitr. Ken. Philip. Veg. (1905) 130; Engl. & Krause Pflanzenreich 71 (1920) 95.

This was based on cultivated specimens supposedly of Philippine origin. I have seen no specimens. Endemic.

ALOCASIA SANDERIANA Bull. Cat. (1894) 8, *fig.*; Merr. Fl. Manila (1912) 132; Engl. & Krause Pflanzenreich 71 (1920) 104.
Schizocasia sanderiana Engl. in Bot. Jahrb. 25 (1898) 26.

Mindanao (Misamis, Agusan), *Merrill 7276, F. B. 4762 Mearns & Hutchinson*. In damp primary forests at low altitudes. A very beautiful species common in greenhouse cultivation in Europe and America. Endemic.

ALOCASIA SINUATA N. E. Br. in Gard. Chron. (1885) 678; Engl. & Krause Pflanzenreich 71 (1920) 102.

This was described from cultivated specimens supposedly of Philippine origin; *Merrill 9568* from Palawan possibly represents it. Endemic.

ALOCASIA WENZELII Merr. in Philip. Journ. Sci. 13 (1918) Bot. 2.

Biliran, Samar, Leyte, Mindanao (Agusan), *Wenzel 97, B. S. 24266 Ramos, 18929 McGregor, F. B. 7615 Hutchinson*. In forests at low altitudes; perhaps a gigantic form of *A. zebrina* C. Koch & Veitch. Endemic.
 Local name: Handúroi (S. L. Bis.).

ALOCASIA ZEBRINA C. Koch & Veitch in Wochenschr. Gärt. Pflanz. 6 (1863) 331, Fl. des Serres 15: t. 1541; Naves Novis. App. (1882) 294; Engl. in DC. Monog. Phan. 2 (1879) 507; Merr. Fl. Manila (1912) 133; Engl. & Krause Pflanzenreich 71 (1920) 105.

Luzon (Laguna, Tayabas, Sorsogon), *Whitford 645, Elmer 17790, 14911, B. S. 6731 Robinson*. In primary forests at low and medium altitudes. Endemic.

Local name: Badiáng (Bik.).

EXCLUDED SPECIES

ALOCASIA ALBA Schott; Naves Novis. App. (1882) 293.

ALOCASIA CUCULLATA Schott; Naves l. c.

ALOCASIA LONGILOBA Miq.; Naves op. cit. 294.

ALOCASIA LOWII Hook. f.; Naves l. c.

ALOCASIA MONTANA Schott; Naves op. cit. 293.

ALOCASIA ODORA C. Koch; Naves op. cit. 294.

ALOCASIA PUBERA Schott; Naves op. cit. 293.

16. COLOCASIA Schott

COLOCASIA ESCULENTUM (Linn.) Schott Melet. 1 (1832) 18; Merr. Fl. Manila (1912) 134, Interpret Herb. Amb. (1917) 131, Sp. Blancoanae (1918) 92.

Arum esculentum Linn. Sp. Pl. (1753) 965.

Arum colocasia Linn. l. c.

Colocasia antiquorum Schott Melet. 1 (1832) 18, Naves Novis. App. (1882) 292; Engl. in DC. Monog. Phan. 2 (1879) 491; Hook. f. in Curtis's Bot. Mag. III 50 (1894) t. 7364; Engl. & Krause in Engl. Pflanzenreich 71 (1920) 65.

Calla gaby Blanco Fl. Filip. (1837) 659.

Caladium esculentum Vent. Jard. Cels. (1800) sub. t. 30; Blanco Fl. Filip. ed. 2 (1845) 459, ed. 3, 3 (1879) 64.

Caladium colocasia W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 208.

Caladium violaceum Desf. Cat. Hort. Paris ed. 3 (1829) 7; Naves Novis. App. (1882) 292.

Generally cultivated throughout the Philippines, but not a native of the Archipelago. It rarely produces flowers in the Philippines. A number of varieties are recognized by Filipinos. Pantropic in cultivation. Taro.

Local names: Abá (Ilk.); áua (Ilk.); abálong (Bis.); amóang (Bon.); gábi (Tag.); pising (Bon.); dagmái (Bis.); kimpói (Bis.); lagbái (Tag., Bis.); linsá (Bik.); lubíngan (If.); natóng (Bik.).

EXCLUDED SPECIES

COLOCASIA AFFINIS Schott; Naves Novis. App. (1882) 293.

COLOCASIA FALLAX Schott; Naves l. c.

COLOCASIA INDICA Engl.; Naves l. c.

17. CALADIUM Ventenat

CALADIUM BICOLOR (Ait.) Vent. Jard. Cels. (1800) t. 30; Naves Novis. App. (1882) 291; Merr. Fl. Manila (1912) 131; Engl. & Krause in Engl. Pflanzenreich 71 (1920) 31.

Arum bicolor Ait. Hort. Kew 3 (1789) 316.

Cultivated for its variegated leaves; occasional in the settled areas as an escape. A native of tropical America.

Local names: Corazón de maría (Sp.); linsáng-pulá (Bik.).

18. TYPHONIUM Schott

TYPHONIUM DIVARCATUM (Linn.) Decne. in Nouv. Ann. Mus. Paris 3 (1834) 367; Engl. in DC. Monog. Phan. 2 (1879) 611; Naves Novis. App. (1882) 295 et in Blanco Fl. Filip. ed. 3 (1877-83) t. 330, var. MOTLEYANUM (Schott) Engl. in DC. Monog. Phan. 2 (1879) 612.

Typhonium motleyanum Schott Prodr. (1860) 106; Usteri Beitr. Ken. Philip. Veg. (1905) 130; Engl. Pflanzenreich 73 (1920) 115.

Introduced and cultivated only; occasional in old gardens, but scarcely established. India to the Moluccas.

TYPHONIUM FLAGELLIFORME (Lodd.) Blume in Wall. Cat. (1832) No. 8928; Engl. Pflanzenreich 73 (1920) 112, f. 16.

Arum flagelliforme Lodd. Bot. Cab. (1819) t. 396.

Arum cuspidatum Blume Cat. Gew. Buitenz. (1823) 101.

Typhonium cuspidatum Decne. in Nouv. Ann. Mus. Paris 3 (1834) 367; Engl. in DC. Monog. Phan. 2 (1879) 616; Naves Novis. App. (1882) 295; Merr. Fl. Manila (1912) 131, Sp. Blancoanae (1918) 92.

Arum divaricatum Blanco Fl. Filip. (1837) 657, ed. 2 (1845) 458, ed. 3, 3 (1879) 62, non Linn.

Luzon (Ilocos Norte, Cavite, Rizal), *Merrill 4171, 7128, 7111, Sp. Blancoanae 676, 970, B. S. 1369 Robinson*. In waste places about settlements, along trails, roads, etc., at low altitudes. India to Indo-China through Malaya to Timor, and the Marianne Islands.

EXCLUDED SPECIES

TYPHONIUM TRILOBATUM Schott; Naves Novis. App. (1882) 295.

This, like *T. divaricatum* Decne., has apparently been cultivated in Manila, as it is represented in the Kew herbarium by *Loher 2428*, from Manila. It is not a native of the Philippines.

19. ARISAEMA Martius

ARISAEMA CUSPIDATUM (Roxb.) Engl. in DC. Monog. Phan. 2 (1879) 536, Pflanzenreich 73 (1920) 158.

Arum cuspidatum Roxb. Fl. Ind. ed. 2, 3 (1832) 506.

Arisaema roxburghii Kunth Enum. 3 (1841) 18; Hook. f. Fl. Brit. Ind. 6 (1893) 497.

Palawan, *Merrill 9339*. In primary forests at low altitudes. Assam, Malay Peninsula, Penang.

ARISAEMA POLYPHYLLUM (Blanco) Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 90, Philip. Journ. Sci. 1 (1906) Bot. 33, 2 (1907) Bot. 264, 5 (1910) Bot. 336, Sp. Blancoanae (1918) 92; Engl. Pflanzenreich 73 (1920) 182.

Calla polyphylla Blanco Fl. Filip. (1837) 659.

Caladium digitatum Blanco op. cit. ed. 2 (1845) 459, ed 3, 3 (1879) 64.

Arisaema cumingii Schott Syn. Aroid. (1856) 28 (*cunningii*), Prodr. (1860) 52; Miq. Fl. Ind. Bat. 3 (1856) 198; Engl. in DC. Monog. Phan. 2 (1879) 558; Naves Novis. App. (1882) 294; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 281.

Arisaema sarasinorum Engl. Bot. Jahrb. 37 (1905) 143.

Luzon (Ilocos Norte, Ifugao, Bontoc, Benguet, Rizal, Bataan, Laguna, Tayabas), Mindoro, Panay, Negros, Mindanao. In forests and in open places, altitude 1,000 to 2,400 m, in the mossy forest often growing as an epiphyte on mossy trunks. Celebes.

Local names: Bía (Ibn.); bíla (Bon.); igigin-di-gáyang (Bon.); takin-di-gáyang (Bon.); tigtigki (Ig.).

Var. ANGUSTIFOLIUM Merr. in Philip. Journ. Sci. 5 (1910) Bot. 336; Engler l. c.

Luzon (Bontoc, Benguet, Zambales), *Merrill 6472, Elmer 6330, F. B. 18387 Alvarez*. In the pine region and in forests, altitude 800 to 2,000 m. Endemic.

EXCLUDED SPECIES

ARISAEMA FIMBRIATUM Mast. in Gard. Chron. II 2 (1884) 680, 689, f. 119; Hook. f. in Curtis's Bot. Mag. III 46 (1890) t. 7150.

This was originally described from cultivated specimens supposed to have come from the Philippines; it is a native of the Langkawi Islands, north of Penang, and nothing remotely resembling it is known from the Philippines.

ARISAEMA LAMINATUM Blume; Naves Novis. App. (1882) 294.

20. CRYPTOCORYNE Fischer

CRYPTOCORYNE APONOGETIFOLIA Merr. in Philip. Journ. Sci. 14 (1919) 370.

Negros, Panay, *B. S. 31119 Ramos & Edaña, F. B. 19411 Curran*. Submerged in shallow water of streams at low altitudes. Endemic.

CRYPTOCORYNE PYGMAEA Merr. in Philip. Journ. Sci. 14 (1919) 371.

Mindanao (Zamboanga), *Merrill 8147, Copeland*. In crevices of ledges in stream beds in damp shaded ravines, altitude about 300 m. Endemic.

CRYPTOCORYNE USTERIANA Engl. ex Usteri Beitr. Ken. Philip. Veg. (1905) 130, Engl. Bot. Jahrb. 54 (1916) Beibl. 118: 125; Pflanzenreich 73 (1920) 240, f. 60.

Guimaras, (*Usteri*). I have seen only the type, which is preserved in the Berlin herbarium. Endemic.

EXCLUDED SPECIES

CRYPTOCORYNE CILIATA Fisch.; Naves Novis. App. (1882) 295.

21. PISTIA Linnaeus

PISTIA STRATIOTES Linn. Sp. Pl. (1753) 963; Blanco Fl. Filip. (1837) 651, ed. 2 (1845) 454, ed. 3, 3 (1879) 55, t. 468; Presl Epim. (1851) 240; Miq. Fl. Ind. Bat. 3 (1856) 218; Engl. in DC. Monog. Phan. 2 (1879) 634; Naves Novis. App. (1882) 296; Vidal Phan. Cuming. Philip. (1885) 154, Rev. Pl. Vasc. Filip. (1886) 281; Merr. Fl. Manila (1912) 130, Sp. Blancoanae (1918) 93; Engl. Pflanzenreich 73 (1920) 259, f. 63.

Pistia cumingii Klotz. in Abh. Akad. Berlin (1854) 354; Walp. Ann. 5 (1858) 913.

Pistia stratiotes Linn. var. *cuneata* Engl. in DC. Monog. Phan. 2 (1879) 634.

Northern Luzon to Mindanao. Floating in shallow water of lakes and slow streams at low altitudes; often abundant. Pantropic.

Local names: Alaluán (Ibn.); dagáylo (Mbo.); darahíro (Bik.); daraído (Bik.); daraíro (Bik.); kaiápo (Bis.); kiápo (Tag.); kiúpu (Mag.); loloán (Ilk.).

EXCLUDED GENERA AND SPECIES

CHAMAECLADON OVALIFOLIUM Schott; Naves Novis. App. (1882) 290.

CHAMAECLADON PYGMAEUM Engl.; Naves op. cit. 291.

CUSUARIA MARANTIFOLIA Schott; Naves op. cit. 289.

DIEFFENBACHIA SEGUINE Schott; Naves op. cit. 291.

LAGENANDRA TOXICARIA Dalz.; Naves op. cit. 295.

LASIA SPINOSA Thwaites; Naves op. cit. 289.

THEIOPHONIUM CRENATUM Blume; Naves op. cit. 295.

THEIOPHONIUM WIGHTII Engl.; Naves l. c.

PIPTOSPATHA sp.; Usteri Beitr. Ken. Philip. Veg. (1905) 130.

LEMNACEAE

1. SPIRODELA Schleiden

SPIRODELA POLYRRHIZA (Linn.) Schleid. in *Linnaea* 13 (1839) 392; Hegelm. *Monogr. Lemn.* (1868) 151, 156; *Engl. Bot. Jahrb.* 21 (1895) 284; Merr. *Fl. Manila* (1912) 134.

Lemna polyrrhiza Linn. *Sp. Pl.* (1753) 970; Naves Novis. *App.* (1882) 296; Usteri *Beitr. Ken. Philip. Veg.* (1905) 131.

Northern Luzon to Mindanao, *F. B. 11622 Curran, Merrill 5121, 4445*. Floating on stagnant fresh water at low and medium altitudes. Temperate and tropical regions of both hemispheres.

2. LEMNA Linnaeus

LEMNA PAUCICOSTATA Hegelm. *Lemn.* (1868) 294, *Engl. Bot. Jahrb.* 21 (1895) 294; Usteri *Beitr. Ken. Philip. Veg.* (1905) 131; Merr. *Fl. Manila* (1912) 134, *Sp. Blancoanae* (1918) 93.

Lemna gibba Blanco *Fl. Filip.* (1837) 672, ed. 2 (1845) 468, ed. 3, 3 (1879) 78; Naves Novis. *App.* (1882) 296, non Linn.

Lemna oligorrhiza Naves Novis. *App.* (1882) 296, non Kurz.

Luzon to Mindanao, floating on stagnant fresh water at low and medium altitudes, often abundant. Widely distributed in the warmer parts of both hemispheres.

Local names: Inaliá (Tag.); liá (Tag.); liyá (Tag.).

LEMNA TRISULCA Linn. *Sp. Pl.* (1753) 970; Naves Novis. *App.* (1882) 296; Hegelm. *Lemn.* (1868) 134, *Engl. Bot. Jahrb.* 21 (1895) 293.

Luzon (Benguet), Mindanao (Lanao), *Clemens, B. S. 8782 McGregor*. Floating on stagnant fresh water at medium altitudes. Widely scattered in both hemispheres.

EXCLUDED SPECIES

LEMNA MINOR Linn.; Naves Novis. *App.* (1882) 296.

3. WOLFFIA Horkel

WOLFFIA ARRHIZA (Linn.) Wimm. *Fl. Schles.* (1857) 140; Hegelm. *Monogr. Lemn.* (1868) 124, *Engl. Bot. Jahrb.* 21 (1895) 301; Naves Novis. *App.* (1882) 296.

Lemna arrhiza Linn. *Mant.* 2 (1771) 294.

Luzon (Pampanga, Bulacan, Rizal), *Merrill 4244*. Floating on stagnant fresh-water pools at low altitudes. Warmer parts of both hemispheres.

Local names: Dugmán (Bis.); liá (Tag.).

EXCLUDED SPECIES

WOLFFIA DELILII Schleid.; Naves Novis. *App.* (1882) 296.

WOLFFIA SHLEIDENI Miq.; Naves l. c.

FLAGELLARIACEAE

1. JOINVILLEA Gaudichaud

JOINVILLEA BORNEENSIS Becc. *Nelle Foreste di Borneo* (1902) 198.

Joinvillea malayana Ridley in *Journ. Str. Branch Roy. As. Soc.* 44 (1905) 199; Merr. in *Philip. Journ. Sci.* 1 (1906) Suppl. 181.

Palawan, Jolo, B. S. 575 *Foxworthy*, Merrill 9515, Clemens 9374. In forests and on ridges, altitude 650 to 1,000 m. Malay Peninsula, Borneo. Local name: Odyúng (Tagb.). *Sumatra!*

2. FLAGELLARIA Linnaeus

FLAGELLARIA INDICA Linn. Sp. Pl. (1753) 333; Blanco Fl. Filip. ed. 2 (1845) 196, ed. 3, 1 (1877) 347; Miq. Fl. Ind. Bat. 3 (1856) 249; Naves Novis. App. (1880) 263; Vidal Rev. Pl. Vasc. Filip. (1886) 279; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 87, Sp. Blancoanae (1918) 93.

Flagellaria philippinensis Elm. Leaf. Philip. Bot. 1 (1908) 274.

Batan Islands to Mindanao and Palawan, in all or most islands and provinces. In secondary forests at low and medium altitudes; common. Tropical Africa, tropical Asia through Malaya to tropical Australia and the Marianne Islands.

Flagellaria philippinensis Elm. is a form with very large leaves; all intergrades between it and the typical form occur.

Local names: Anuád (Ilk.); aráyan (Tag.); auái (Iv.); auái-si-gáyang (Is.); baling-uái (Tag., Pamp.); boboáya (Mbo.); hoág-uái (Bik.); huág (S. L. Bis., Mbo.); húak (Bis.); iñgúal (Ilk.); iñgúla (Tag.); inúal (Pang.); kala-uái (Ibn.); kala-uaiuai (Ibn.); ouág-oái (Bik.); ouag-ouág (Mbo.); páua (P. Bis.); sagákap (P. Bis.); tauá (P. Bis.); tinúung (Ibn.); uág (Sul., Bis., Bag., Bik.); uái-ti-uák (Ilk.); uák (Bis.); venag-aíang (Is.).

3. HANGUANA Blume

HANGUANA MALAYANA (Jack) Merr. in Philip. Journ. Sci. 10 (1915) Bot. 3.

Veratrum ? malayanum Jack Malay Miscel. 1 (1820) 25; Hook. Bot. Miscel. 2 (1831) 74.

Hanguana kassintu Blume Enum. Pl. Jav. (1827) 15, ed. 2 (1830) 15.

Veratonia malayana Miq. Fl. Ind. Bat. 3 (1859) 553.

Susum malayanum Planch. ex Hook. f. Fl. Brit. Ind. 6 (1892) 391.

Mindoro, Palawan, Mindanao (Lanao, Surigao), Merrill 7221, 9498, Piper 529, Clemens 1134a. In thickets back of the mangrove swamps, ascending to 900 m in forests. Malay Peninsula and Archipelago.

CENTROLEPIDACEAE

1. CENTROLEPIS Labillardière

CENTROLEPIS PHILIPPINENSIS Merr. in Philip. Journ. Sci. 2 (1907) Bot. 264.

Mindoro, Merrill 6160, Phil. Pl. 87. In open heaths, altitude about 2,400 m. Endemic.

This was also credited to Borneo, by Turrill, but Miss Gibbs later described the Bornean form as *C. kinabaluensis* Gibbs, Phytogeog. Dutch New Guinea (1917) 99. Only one other species of the family is known north of Australia, this being a species of *Centrolepis* in Indo-China.

XYRIDACEAE

1. XYRIS Linnaeus

XYRIS COMPLANATA R. Br. Prodr. (1810) 256; Naves Novis. App. (1880) 268; Hallier f. in Nova Guinea 8 (1913) 903, Beihefte Bot. Centralbl. 34² (1916) 45.

Xyris anceps Merr. in Philip. Journ. Sci. 5 (1910) Bot. 172, non Lam.

Luzon (Isabela), Semirara, *B. S. 8065 Ramos, Merrill 4151*. In open grasslands at low altitudes. Malay Archipelago to Australia.

Local name: Gumi-gúmi (Tag.).

XYRIS PAUCIFLORA Willd. Phytogr. (1794) 2, t. 1; Miq. Fl. Ind. Bat. 3 (1859) 529; Naves Novis. App. (1880) 268; Merr. Fl. Manila (1912) 135.

Luzon (Pangasinan, Tarlac, Bulacan, Rizal), *Loher 6985, Merrill 3628, Phil. Pl. 463, B. S. 4911 Ramos*. In open grasslands at low altitudes. India to China and Formosa through Malaya to Australia.

EXCLUDED SPECIES

XYRIS INDICA Linn.; Naves Novis. App. (1880) 268.

XYRIS OPERCULATA Labill.; Naves l. c.

ERIOCAULONACEAE

1. ERIOCAULON Linnaeus

ERIOCAULON ALATUM H. Lecomte in Journ. de Bot. 21 (1908) 104, f. 2; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 232, Fl. Manila (1912) 136.

Luzon (Rizal), *Merrill 7362, B. S. 1831 Ramos, 19270 Reillo*. In open wet grasslands at low altitudes. Indo-China.

ERIOCAULON ALPESTRE Hook. f. & Th. ex Koern. in Miq. Ann. Mus. Bot. Lugd.-Bat. 3 (1867) 163; Ruhl. in Engl. Pflanzenreich 13 (1903) 95; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 291.

Mindanao (Davao), *Copeland 1431*. In open wet places, altitude about 1,800 m. India to Japan south to Indo-China.

ERIOCAULON BREVIPEDUNCULATUM Merr. in Philip. Journ. Sci. 2 (1907) Bot. 265.

Mindoro, *Merrill 6214*. On seepage slopes in open heaths on Mt. Halcon, altitude about 2,400 m. Borneo.

ERIOCAULON CINEREUM R. Br. Prodr. (1810) 254; Benth. Fl. Austral. 7 (1878) 193; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 291, Fl. Manila (1912) 136.

Eriocaulon sieboldianum Sieb. & Zucc. ex Steud. Syn 2 (1855) 272; Ruhl. in Engl. Pflanzenreich 13 (1903) 111, f. 15 A-G.

Luzon (Rizal), *Merrill Phil. Pl. 293, 1276, B. S. 24072 Ramos*. In fallow wet rice lands at low altitudes. Africa to Japan, Malaya, and Australia.

Local name: Dundunsúg (P. Bis.).

ERIOCAULON DEPAUPERATUM Merr. in Philip. Journ. Sci. 5 (1910) Bot. 336.

Luzon (Lepanto, Benguet), *Merrill 4520, 6590, B. S. 40528 Ramos & Edaño*. In pools and small shallow ponds, altitude 2,200 to 2,800 m. Endemic.

ERIOCAULON ECHINULATUM Mart. in Wall. Pl. As. Rar. 3 (1832) 28; Ruhl. in Engl. Pflanzenreich 13 (1903) 106.

Culion, *B. S. 41325 Ramos*. In open grassy places at low altitudes. Tanasserim to Indo-China and southern China.

ERIOCAULON MERRILLII Ruhl. in Perk. Frag. Fl. Philip. (1904) 136; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 291, Fl. Manila (1912) 136.

Luzon (Nueva Vizcaya, Nueva Ecija, Tarlac, Zambales, Rizal, Sorsogon), Polillo, Culion, Sibuyan, Mindanao (Lanao), *Merrill 572, 7122, 3624, Elmer 14350, 12248, B. S. 14149 McGregor, 9031 Robinson*. In open wet places, old rice paddies, along small streams, etc., at low altitudes. Endemic. Distinct from *E. truncatum* Ham.?

Local name: Boibotánis (P. Bis.).

ERIOCAULON NIGRICEPS Merr. in Philip. Journ. Sci. 10 (1915) Bot. 290.

Luzon (Lepanto, Benguet), *B. S. 5544 Ramos, 31747 Santos, Merrill 4646, 7748, Elmer 6617, Williams 946, 1004*. In open wet places, altitude 1,400 to 2,300 m. Endemic.

ERIOCAULON TRUNCATUM Mart. in Wall. Pl. As. Rar. 3 (1832) 29; Steud. Syn. 2 (1859) 270; Ruhl. in Engl. Pflanzenreich 13 (1903) 107; Naves Novis. App. (1882) 299.

Luzon (Abra, Ifugao, Bontoc, Nueva Vizcaya, Rizal, Camarines), *B. S. 33721 Ramos & Edaño, 24089 Ramos, 19980, 20172, 20149 McGregor, 19267 Reillo*. In open wet places, rice paddies, etc., at low and medium altitudes. India to southern China and Malaya.

Cuming 2326, credited to the Philippines by Ruhland, was from Malacca.

Local names: Bauaṅ-bauaṅgan (Bik.); dasdasnuk (Bon.).

EXCLUDED SPECIES

ERIOCAULON LONGISSIMUM Usteri Beitr. Ken. Philip. Veg. (1905) 131.

This is a misprint for *E. longifolium* Nees, a species unknown from the Philippines.

ERIOCAULON SEXANGULARE Linn.; Naves Novis. App. (1882) 299; Miq. Fl. Ind. Bat. 3 (1857) 523; Ruhl. in Engl. Pflanzenreich 13 (1903) 110.

ERIOCAULON SETACEUM Linn.; Naves l. c.

BROMELIACEAE

1. ANANAS Tournefort

ANANAS COMOSUS (Linn.) Merr. Interpret. Herb. Amb. (1917) 133, Sp. Blancoanae (1918) 94.

Bromelia comosa Linn. in Stickman Herb. Amb. (1754) 21, Amoen. Acad. 4 (1759) 130.

Bromelia ananas Linn. Sp. Pl. (1753) 285; Blanco Fl. Filip. (1837) 230, ed. 2 (1845) 162, ed. 3, 1 (1877) 291, t. 453.

Bromelia pigna Perr. in Mém. Soc. Linn. Paris 3 (1824) 103; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 304.

Ananas sativus Schultes f. Syst. 7² (1830) 1283; Merr. Fl. Manila (1912) 137.

Ananassa sativa Lindl. Bot. Reg. (1827) sub. t. 1063; Miq. Fl. Ind. Bat. 3 (1859) 584; Naves Novis. App. (1880) 256; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 24.

Ananassa ananas Karst. Deutsch. Fl. (1880-83) 466.

Widely cultivated in the Philippines; in some regions, Palawan, thoroughly naturalized. A native of tropical America, now cultivated in all tropical countries. Pineapple.

Local names: Apangdán (Bon.); pangdán (Bon.); piña (Sp.).

EXCLUDED GENERA

BILLBERGIA ZEBRINA Lindl.; Naves Novis. App. (1880) 257.

PITCAIRNIA BRACHIATA Cham.; Naves l. c.

COMMELINACEAE

1. POLLIA Thunberg

POLLIA MACROPHYLLA Benth. Fl. Austral. 7 (1878) 90; C. B. Clarke in DC. Monog. Phan. 3 (1881) 123; Naves Novis. App. (1880) 269; Ceron Cat. Pl. Herb. (1892) 173.

Philippines, fide C. B. Clarke, who cites specimens from *Cuming* and *Llanos*. These, however, need critical comparison with *P. sorzogonensis* (E. Meyer) Steud. New Guinea, Solomon Islands, tropical Australia.

POLLIA SORZOGONENSIS (E. Meyer) Steud. Nomencl. ed. 2, 2 (1841) 368; Miq. Fl. Ind. Bat. 3 (1859) 541; C. B. Clarke in DC. Monog. Phan. 3 (1881) 126; Naves Novis. App. (1880) 269; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 278; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 34, 3 (1908) Bot. 400, Sp. Blancoanae (1918) 94.

Aclisia sorzogonensis E. Meyer in Presl Rel. Haenk. 1 (1827) 138, t. 25.

Lechea minor Blanco Fl. Filip. (1837) 52, ed. 2 (1845) 35, ed. 3, 1 (1877) 65, non Linn.

Aclisia cumingiana Klotz. in Hassk. Commel. Ind. (1870) 47.

Northern Luzon (Cagayan) to Palawan and Mindanao. Common in primary forests at low and medium altitudes. Southern China to Java and the Moluccas.

Local names: Alikbángon (Tag.); lohod-lóhod (Bis.); luho-lúho (S. L. Bis.); salibángung (Bag.).

POLLIA SUMATRANA Hassk. Commel. Ind. (1870) 56; C. B. Clarke in DC. Monog. Phan. 3 (1881) 125; Naves Novis. App. (1880) 269; Vidal Rev. Pl. Vasc. Filip. (1886) 278.

Luzon (Rizal, Bulacan, Laguna), Polillo, Biliran, Samar, Leyte, Mindanao, Piper 433, Elmer 7153, B. S. 6724 Robinson, 15702 Fénix, 18667 McGregor, 1088, 24425 Ramos. In primary forests at low and medium altitudes. Sumatra.

Clarke's description is based on Cuming's Philippine plant. I do not know whether or not it is identical with Hasskarl's type.

Local names: Aregbabáŋgon (S. L. Bis.); idog (Mbo.); mantauási (Buk.).

POLLIA THYRSIFLORA (Blume) Steud. Nomencl. ed. 2, 2 (1841) 368; C. B. Clarke in DC. Monog. Phan. 3 (1881) 124; Naves Novis. App. (1880) 269; Vidal Rev. Pl. Vasc. Filip. (1886) 278; Hallier f. in Nova Guinea 8 (1913) 907, Beihefte Bot. Centrabl. 34² (1916) 51.

Tradescantia thyrsoflora Blume Enum. Pl. Jav. 1 (1827) 6.

Pollia philippinensis Elm. Leaf. Philip. Bot. 1 (1908) 275.

Luzon (Cagayan to Sorsogon) to Palawan and Mindanao, *Elmer* 9162, 13545, 15329, *Wenzel* 1166, *Merrill* 9268, 4007, *Loher* 6981, *F. B.* 19623 *Curran*, *B. S.* 25857 *Yates*, 12651 *Fénix*. In primary forests at low and medium altitudes. Andaman Islands to Java and Celebes.

Local names: Talibáŋgung (Bag.); sapilko (Ig.).

EXCLUDED SPECIES

POLLIA ACLISIA Hassk.; Naves Novis. App. (1880) 268.

POLLIA CRISPATA Benth.; Naves op. cit. 269.

POLLIA JAPONICA Thunb.; Naves op. cit. 268.

2. COMMELINA Plumier

COMMELINA BENGHALENSIS Linn. Sp. Pl. (1753) 41; Miq. Fl. Ind. Bat. 3 (1859) 533; C. B. Clarke in DC. Monog. Phan. 3 (1881) 159; Naves Novis. App. (1880) 269; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 400, Fl. Manila (1912) 138, Sp. Blancoanae (1918) 94. *Commelina polygama* Blanco Fl. Filip. (1837) 25 (*poligama*), ed. 2 (1845) 18, ed. 3, 1 (1877) 34, t. 13, non Roth.

Throughout the Philippines in open grasslands and waste places in the settled areas at low and medium altitudes, often common. Tropical Africa and Asia to Japan and Malaya.

Local names: Alibáŋgon (Tag.); bias-bías (Pamp.); kabiláu (Bis.); kuhási (Iv.); kulkul-lási (Ilk.); sabiláu (Bis.); sambiláu (Bis.); uligbóŋgon (Tag.).

COMMELINA CHAMISSONIS Klotz. ex C. B. Clarke in DC. Monog. Phan. 3 (1881) 186; Naves Novis. App. (1880) 270.

The type is a specimen from Luzon, collected by *Chamisso*, probably from either Cavite or Batangas Province or possibly from the vicinity of Manila. I have seen no specimens I can with certainty refer to it, but some may have been distributed sub *C. nudiflora* Linn. Australia.

COMMELINA NUDIFLORA Linn. Sp. Pl. (1753) 4; C. B. Clarke in DC. Monog. Phan. 3 (1881) 144; Naves Novis. App. (1880) 269; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 400, Fl. Manila (1912) 138.

Tradescantia cristata Naves in Blanco Fl. Filip. ed. 3 (1877-83) t. 83, non Linn.

Throughout the Philippines in open grasslands, waste places, etc., at low and medium altitudes, often common. Pan-tropic.

Local names: Alikbáŋgon (Tag.); baŋgar-an-laláki (If.); gatiláng (Bon.) katkataúang (Bon.); kitkit-aúang (Bon.); kohási (Iv.); kolási (Ilk.); kulkul-lási (Ilk.).

COMMELINA UNDULATA R. Br. Prodr. (1810) 270; C. B. Clarke in DC. Monog. Phan. 3 (1881) 179; Naves Novis. App. (1880) 270; Ceron Cat. Pl. Herb. Manila (1892) 173.

Philippines fide C. B. Clarke l. c. India to southern China, Riu Kiu Islands, Moluccas, tropical Australia.

EXCLUDED SPECIES

COMMELINA BARBATA Lam.; Naves Novis. App. (1880) 270.

COMMELINA COMMUNIS Linn.; Naves l. c.

COMMELINA OBLIQUA Ham.; Naves l. c.

COMMELINA SALICIFOLIA Roxb.; Naves op. cit. 269.

COMMELINA SUBULATA Roth; Naves l. c.

COMMELINA SUFFRUTICOSA Blume; Naves op. cit. 270.

3. ANEILEMA R. Brown

ANEILEMA AZUREUM Merr. in Philip. Journ. Sci. 10 (1915) Bot. 292.

Luzon (Bulacan), Semirara, Merrill 4134, B. S. 21669 Ramos. In open wet grasslands at low altitudes. Endemic.

ANEILEMA GIGANTEUM (Vahl) R. Br. Prodr. (1810) 271; Naves Novis. App. (1880) 271; C. B. Clarke in DC. Monog. Phan. 3 (1881) 212; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 294.

Commelina gigantea Vahl Enum. 2 (1806) 177.

Semirara, Mindanao (Bukidnon), Merrill 4132, B. S. 31335 Escritor, 39210 Ramos & Edaña, 26138 Fénix, F. B. 26553 Rola. In open grasslands at low and medium altitudes. Tropical Africa and Asia through Malaya to tropical Australia.

ANEILEMA HUMILE Merr. in Philip. Journ. Sci. 13 (1918) Bot. 4.

Alabat, Merrill 10459. On wet slopes in primary forests at low altitudes. Endemic.

ANEILEMA MALABARICUM (Linn.) Merr. in Philip. Journ. Sci. 7 (1912) Bot. 232, Fl. Manila (1912) 138.

Tradescantia malabarica Linn. Sp. Pl. ed. 2 (1762) 412.

Commelina nudicaulis Burm. f. Fl. Ind. (1768) 17, t. 8, f. 1.

Commelina nudiflora Linn. Mant. 1 (1767) 177, non Sp. Pl. (1753).

Aneilema nudiflorum R. Br. Prodr. (1810) 271; Miq. Fl. Ind. Bat. 3 (1859) 537; Naves Novis. App. (1880) 271, t. 467; C. B. Clarke in DC. Monog. Phan. 3 (1881) 210; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 278.

Commelina trichocolea Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 448; Walp. Ann. 1 (1849) 885.

Throughout the Philippines. Common in open grasslands, waste places, etc., at low and medium altitudes. India to China, the Riu Kiu Islands, and Malaya.

Local names: Aligbāṅgon (Pang.); aligbāṅgon-laláki (Tag.); alitbāṅgun (Tag.); faṅgau (Bon.); olikbāṅgon (Tag.); takdāyan (Ig.).

ANEILEMA PLATYPHYLLUM Merr. in Philip. Journ. Sci. 10 (1915) Bot. 293.

Luzon (Pangasinan, Rizal), B. S. 985, 3342, 20985 Ramos, 17900 Otañes. Along streams in shaded places at low altitudes. Endemic.

ANEILEMA SPIRATUM (Linn.) R. Br. ex C. B. Clarke in DC. Monog. Phan. 3 (1881) 207; Naves Novis. App. (1880) 270.
Commelina spirata Linn. Mant. 2 (1771) 176.

This is credited to the Philippines by C. B. Clarke on the basis of a specimen from Lake Bay, Luzon, collected by *Baume* and preserved in the Paris herbarium. This specimen may be *A. malabaricum* (Linn.) Merr., or I may have included *A. spiratum* R. Br. in my conception of *A. malabaricum* (Linn.) Merr. India to Indo-China, China, and Java.

ANEILEMA VERSICOLOR Dalz. in Hook. Journ. Bot. Kew Miscel. 3 (1851) 136; C. B. Clarke in DC. Monog. Phan. 3 (1881) 208; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 232, Fl. Manila (1912) 139.

Luzon (Rizal), Culion, *Merrill 3644, 518, Phil. Pl. 1980, 763, Loher 6986, B. S. 10877 Ramos, 19171 Reillo*. In open wet grasslands, fallow rice paddies, etc., at low altitudes. India.

ANEILEMA VITIENSE Seem. Fl. Vit. (1865-73) 314, t. 96; C. B. Clarke in DC. Monog. Phan. 3 (1881) 220; Naves Novis. App. (1880) 271.
 Philippines, *Moseley*, fide C. B. Clarke. Fiji, Samoa.

Var. **PETIOLATA** C. B. Clarke l. c.

Northern Luzon (Cagayan) to Palawan and Mindanao, in most islands and provinces, *Merrill 4862, 5332, 5119, 7134, Phil. Pl. 33, Weber 1022, Elmer 13557, 15366, 18053, Wenzel 4, Whitford 973, B. S. 12409 McGregor, 18081 Robinson, 15701 Fénix, 29057, 30928 Ramos & Edaño*. In damp shaded ravines usually along small streams at low and medium altitudes, ascending to 1,400 m. The variety endemic, the species in Samoa and Fiji.

Aneilema scaberrima (Blume) Kunth has been credited to the Philippines by Hallier f., Beihefte Bot. Centralbl. 34² (1916) 50, as *A. protensum* Wall., but I suspect that the specimen he examined is *A. vitiense* Seem.

Local names: Balúkid (Buk.); dalidígan (Mbo.).

EXCLUDED SPECIES

ANEILEMA DIMORPHUM Dalz.; Naves Novis. App. (1880) 270.

ANEILEMA GLAUCUM Thw.; Naves l. c.

ANEILEMA HAMILTONIANUM Wall.; Naves l. c.

ANEILEMA HERBACEUM Wall.; Naves op. cit. 270.

ANEILEMA LOUREIRII Hance; Naves l. c.

ANEILEMA MEDIUM R. Br.; Naves l. c.

ANEILEMA MONADELPHUM Kunth; Naves op. cit. 271.

ANEILEMA SCAPIFLORUM Wight; Naves op. cit. 270.

ANEILEMA SINICUM Lindl.; Naves op. cit. 271.

4. FORRESTIA A. Richard

FORRESTIA HISPIDA Less. & A. Rich. Sert. Astrolab. (1834) 2, t. 1, var. **PHILIPPINENSIS** (Merr.) comb. nov.

Forrestia philippinensis Merr. in Govt. Lab. Publ. (Philip.) 35 (1906) 5, Philip. Journ. Sci. 2 (1907) Bot. 266.

Forrestia hispida Less. & A. Rich. var. *glabrior* Hallier f. in Nova Guinea 8 (1913) 906, Beihefte Bot. Centralbl. 34² (1916) 46.

Luzon (Laguna, Tayabas, Camarines, Albay, Sorsogon), Polillo, Catanduanes, Mindoro, Leyte, Negros, Panay, Samar, Mindanao (Agusan, Bukidnon, Lanao), Basilan. In thickets and old clearings in the vicinity of primary forests and in primary forests at low and medium altitudes; represented by numerous collections. The variety in Borneo, the species in New Guinea.

Local names: Arigbabáñgon (S. L. Bis.); lohod-lohód (C. Bis.); malitágo (Mbo.); mani-utús (Lan.); salimbágun (Bag.); tahig-tahíg (Bik.).

EXCLUDED SPECIES

FORRESTIA GLABRATA Hassk.; Naves Novis. App. (1880) 271.

FORRESTIA MARGINATA Hassk.; Naves l. c.

FORRESTIA MOLLIS Hassk.; Naves l. c.

5. CYANOTIS D. Don

CYANOTIS AXILLARIS (Linn.) D. Don Prodr. Fl. Nepal. (1825) 46; R. & S. Syst. 7² (1830) 1154; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 447; C. B. Clarke in DC. Monog. Phan. 3 (1881) 244; Naves Novis. App. (1880) 272; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 278; Merr. Fl. Manila (1912) 140.

Commelina axillaris Linn. Sp. Pl. (1753) 42.

Northern Luzon to Palawan and Mindanao, in most islands and provinces. In clearings, open places along streams, rice paddies, etc., at low and medium altitudes; often common. India to China through Malaya to Australia.

Local names: Alikbáñgon (Tag.); alithbáñgon (Tag.); kulásin-marinték (Pang.); sabiláu (P. Bis.).

CYANOTIS CAPITATA (Blume) C. B. Clarke in DC. Monog. Phan. 3 (1881) 243; Naves Novis. App. (1880) 272.

Tradescantia capitata Blume Enum. Pl. Jav. (1827-28) 6.

Northern Luzon to Palawan and Mindanao, *Clemens 773, Copeland 740, Merrill 803, 4426, 8154, 5462, 1804, F. B. 3453 Hutchinson, B. S. 26076 Fénix, 23850 Ramos, 32555 McGregor*. Along streams in shaded ravines, in damp thickets, etc., at low and medium altitudes; really distinct from *C. moluccana* (Roxb.) Merr.? Japan to Indo-China and Java.

CYANOTIS CRISTATA (Linn.) R. & S. Syst. 7² (1830) 1150; Miq. Fl. Ind. Bat. 3 (1859) 544; C. B. Clarke in DC. Monog. Phan. 3 (1881) 247; Naves Novis. App. (1880) 272; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 278; Merr. Fl. Manila (1912) 140, Sp. Blancoanae (1918) 95.

Commelina cristata Linn. Sp. Pl. (1753) 42.

Tradescantia cristata Linn. Syst. ed. '12 (1767) 233; Blanco Fl. Filip. (1837) 231, ed. 2 (1845) 163, ed. 3, 1 (1877) 293.

Zygomenes cristata W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 404.

Luzon (most or all provinces), Basilan. In old clearings, open places along streams, about cliffs, etc., at low and medium altitudes. India to Mauritius and Malaya.

Local names: Alikbáñgon (Tag.); kulásing-akabaleg (Pang.).

CYANOTIS MOLUCCANA (Roxb.) Merr. in Philip. Journ. Sci. 2 (1907) Bot. 266, Fl. Manila (1912) 139.

Commelina moluccana Roxb. Fl. Ind. 1 (1820) 176.

Cyanotis uniflora Hassk. Commel. Ind. (1870) 104; C. B. Clarke in DC. Monog. Phan. 3 (1881) 242; Naves Novis. App. (1880) 272; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 34.

Luzon (Bontoc, Pampanga, Rizal, Bataan, Laguna, Sorsogon), Mindoro, Leyte, Mindanao, *Baker 500, Wenzel 542, Merrill 4206, 6227, 7132, Williams 249*. On bowlders in ravines, along small streams, etc., at low and medium altitudes. Java, Borneo, Moluccas.

Not always readily distinguishable from *C. capitata* Clarke.

CYANOTIS PEDUNCULATA Merr. in Philip. Journ. Sci. 10 (1915) Bot. 266.

Leyte, *Wenzel 1024*. In forests, altitude about 500 m. Endemic.

EXCLUDED SPECIES

CYANOTIS PAPILIONACEA R. & S.; Naves Novis. App. (1880) 272.

CYANOTIS TUBEROSA R. & S.; Naves l. c.

CYANOTIS VAGA R. & S.; Naves l. c.

6. FLOSCOPA Loureiro

FLOSCOPA SCANDENS Lour. Fl. Cochinch. (1790) 193; C. B. Clarke in DC. Monog. Phan. 3 (1881) 265; Naves Novis. App. (1880) 272; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 278; Merr. Sp. Blancoanae (1918) 95.

Tradescantia rufa Presl Rel. Haenk. 1 (1827) 138.

Floscopa rufa Hassk. in Miq. Pl. Jungh. (1852) 151; Miq. Fl. Ind. Bat. 3 (1859) 542.

Tradescantia geniculata Blanco Fl. Filip. (1837) 232, ed. 3, 1 (1877) 294, non Jacq.

Northern Luzon (Cagayan) to the Calamian Islands and Mindanao, in most or all islands and provinces. In wet places along small streams at low and medium altitudes. India to China through Malaya to tropical Australia.

Local names: Aligbángon (Tag.); babiláu (S. L. Bis.); kúmpai (P. Bis.); púgad-labúyo (Tag.); sambiláu (S. L. Bis.).

7. RHOEO Hance

RHOEO DISCOLOR (L'Hérit.) Hance in Walp. Ann. 3 (1853) 660; Naves Novis. App. (1882) 273, Blanco Fl. Filip. ed. 3 (1877-83) t. 84; Merr. Fl. Manila (1912) 139, Sp. Blancoanae (1918) 94.

Tradescantia discolor L'Hérit. Sert. Angl. (1788) 8, t. 12; Blanco Fl. Filip. (1837) 232, ed. 2 (1845) 163, ed. 3, 1 (1877) 294, t. 84.

Planted for ornamental purposes, sometimes long-persistent after cultivation but scarcely established. A native of tropical America.

8. ZEBRINA Schnizlein

ZEBRINA PENDULA Schnizl. in Bot. Zeit. 7 (1849) 870; C. B. Clarke in DC. Monog. Phan. 3 (1881) 318; Merr. in Philip. Journ. Sci. 7 (1912) Bot. 232, Fl. Manila (1912) 140.

Introduced from tropical America and cultivated as an ornamental, now more or less established in coconut groves, etc., at medium altitudes. Wandering Jew.

PONTEDERIAACEAE

1. EICHORNIA Kunth

EICHORNIA CRASSIPES (Mart.) Solms in DC. Monog. Phan. 4 (1883) 527.

Pontederia crassipes Mart. Nov. Gen. Sp. Pl. 1 (1824) 9, t. 4.

Eichornia speciosa Kunth Enum. Pl. 4 (1843) 131.

Piaropus crassipes Britton in Ann. N. Y. Acad. Sci. 7 (1893) 241.

Introduced about 1912 as an ornamental, now very abundant in shallow water of lakes and in slow streams in central Luzon, occurring also in northern Luzon and in Negros. Within a few years to be expected throughout the Philippines. A native of tropical America, now pantropic. Water hyacinth.

2. MONOCHORIA Presl

MONOCHORIA HASTATA (Linn.) Solms in DC. Monog. Phan. 4 (1883) 523; Merr. Fl. Manila (1912) 141, Sp. Blancoanae (1918) 95.

Pontederia hastata Linn. Sp. Pl. (1753) 288.

Monochoria hastaeifolia Presl Rel. Haenk. 1 (1827) 128; Miq. Fl. Ind. Bat. 3 (1859) 548; Naves Novis. App. (1880) 267.

Pontederia vaginalis Blanco Fl. Filip. (1837) 255, ed. 2 (1845) 178, ed. 3, 1 (1877) 320, t. 466, non Burm. f.

Monochoria dilatata Kunth Enum. 4 (1843) 134; Naves Novis. App. (1880) 268.

Luzon to Mindanao, *Elmer 14319, Merrill 4088, Phil. Pl. 1121, Sp. Blancoanae 419, McGregor 93, Clemens 179, B. S. 1456 Mangubat*. In open wet lands, swamps, etc., at low and medium altitudes. India to southern China and Malaya.

Local names: Gabi-gábi (Bis); gabi-gabihan (Tag.); kosol-kósol (Bis.); payau-páyau (Bis).

MONOCHORIA VAGINALIS (Burm. f.) Presl Rel. Haenk. 1 (1827) 128; Miq. Fl. Ind. Bat. 3 (1859) 548; Naves Novis. App. (1880) 267; Solms in DC. Monog. Phan. 4 (1883) 524; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 278; Merr. Fl. Manila (1912) 141, Interpret. Herb. Amb. (1917) 135.

Pontederia vaginalis Burm. f. Fl. Ind. (1768) 80; Linn. Mant. 2 (1771) 222; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 446.

Monochoria ovata Kunth Enum. 4 (1843) 665; Naves Novis. App. (1880) 268.

Throughout the Philippines at low and medium altitudes; common in open wet places, rice paddies, swamps, etc. Tropical and subtropical Asia to Malaya.

Local names: Biga-bigáan (Tag.); bil-lagut (Ilk.); gabi-gábi (P. Bis., C. Bis.); gabing-uák (Tag.) hahalung (If.); hakhaklung (If.); kalabúa (Tag.); lagtáng (Bik.); lapalápa (Ilk.); saksaklung (Ig.); saksakong (Bon.); upi-úpi (Bik.).

Var. PAUCIFLORA (Blume) comb. nov.*Pontederia pauciflora* Blume Enum. Pl. Jav. 1 (1827) 32.*Pontederia plantaginea* Roxb. Fl. Ind. ed. 2, 2 (1832) 123.*Monochoria pauciflora* Kunth Enum. 4 (1843) 135; Naves Novis. App. (1880) 268.*Monochoria plantaginea* Kunth l. c.; Naves l. c.*Monochoria vaginalis* Presl, var. *plantaginea* Solms in DC. Monog. Phan. 4 (1883) 524.

Luzon (Ilocos Norte, Bontoc, Benguet, Nueva Vizcaya), *Merrill 4616*, *Phil. Pl. 1723*, *Elmer 5953*, *B. S. 31813 Santos, 11371 McGregor, 14109 Robinson, F. B. 16531 Curran & Merritt, 15516 Merritt & Darling*. In swamps, rice paddies, etc., altitude 1,000 to 2,000 m. India to China and Java.

The family Philydraceae has been credited to the Philippines on the basis of a *Cuming* specimen so labeled; Caruel in DC. Monog. Phan. 3 (1881) 3; Naves Novis. App. (1880) 268. *Cuming's* specimen was from the Malay Peninsula, not from the Philippines. No representative of the family is known from the Archipelago.

JUNCACEAE

1. JUNCUS Linnaeus

JUNCUS BUFONIUS Linn. Sp. Pl. (1753) 328; Buchenau in Engl. Pflanzenreich 25 (1906) 105; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 294.

Luzon (Benguet), *Merrill 7794*. Along the bridle trails in the mossy forest on Mt. Santo Tomas, altitude 1,900 m; probably introduced. Warmer parts of both hemispheres.

JUNCUS EFFUSUS Linn. Sp. Pl. (1753) 326; Buchenau in Engl. Pflanzenreich 25 (1906) 135.

Juncus communis E. Meyer Junc. Gen. Monogr. Spec. (1819) 20; Naves Novis. App. (1882) 273.

Luzon (Benguet, Bontoc), Mindanao (Davao). In open swampy places, altitude 1,400 to 2,300 m. Warmer parts of both hemispheres, especially common in the North Temperate Zone.

Local names: Badili (Ig.); balili (Bon.); piñgót (Bon.); sudsúd (Ig.).

JUNCUS PRISMATOCARPUS R. Br. Prodr. (1810) 259, var. **LESCHE-NAULTII** (J. Gay) Buchenau in Engl. Bot. Jahrb. 6 (1885) 137, Engl. Pflanzenreich 25 (1906) 180.

Juncus leschenaultii J. Gay ex Laharpe in Mém. Soc. Linn. Paris 3 (1827) 205; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 176.

Luzon (Lepanto, Benguet), *Merrill 4517, 4739, Williams 1974bis, B. S. 2791, 4261 Mearns*. In cold open swamps, altitude 1,500 to 2,300 m. India and Ceylon to Kamchatka and Japan, to southern China, Australia, and New Zealand.

2. LUZULA de Candolle

LUZULA CAMPESTRIS (Linn.) DC. Fl. Fr. 3 (1805) 161; Naves Novis. App. (1882) 273; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 337, in nota; Buchenau in Engl. Pflanzenreich 25 (1906) 83, f. 52-55.

Juncus campestris Linn. Sp. Pl. (1753) 329.

Luzon (Benguet), *B. S.* 8426 McGregor, 31787 Santos. On slopes in pine forests, at Pauai, extending into the mossy forest along trails, altitude 2,000 to 2,400 m. Temperate regions in both hemispheres.

LUZULA EFFUSA Buchenau Krit. Verz. Junc. (1880) 53, 88, Engl. Pflanzenreich 25 (1906) 61; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 337.

Luzon (Benguet), *Merrill* 6490. In the upper border of the mossy forest and in the summit grasslands on Mt. Pulong, altitude about 2,700 m. Himalayan region, China.

STEMONACEAE

1. STEMONA Loureiro

STEMONA PHILIPPINENSIS Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 16.

Masbate, *Merrill* 3061. On ridges in thickets of forests at low altitudes. Endemic.

Local name: Sigid (P. Bis.).

STEMONA TUBEROSA Lour. Fl. Cochinch. (1790) 404; C. H. Wright in Journ. Linn. Soc. Bot. 32 (1896) 495; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 16, Interpret. Herb. Amb. (1917) 135.

Roxburghia gloriosoides Roxb. Pl. Coromandel 1 (1795) 29, t. 32.

Luzon, without definite locality, Palawan, Leyte, Negros, *Merrill* 792, *Loher* 6969, *Wenzel* 1746, *Eskridge*. In thickets and forests at low altitudes; rare or local. India to Formosa and southward to the Moluccas.

Local name: Lubilubí (C. Bis.).

LILIACEAE

1. IPHIGENIA Kunth

IPHIGENIA INDICA (Linn.) Kunth Enum. 4 (1843) 213; Hook. f. Fl. Brit. Ind. 6 (1892) 357; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 176.

Melanthium indicum Linn. Mant. 2 (1771) 226.

Luzon (Cavite, Rizal, Laguna), Mindanao (Bukidnon, Davao), *Weber* 1462, *Williams* 2962, *Loher* 1913. In open grasslands at low and medium altitudes; rare or local. India to Australia.

2. THYSANOTUS R. Brown

THYSANOTUS CHINENSIS Benth. Fl. Hongk. (1861) 372; Hallier f. in Nova Guinea 8 (1914) 991.

Thysanotus chrysanthorus F. Muell. Fragm. 5 (1866) 202; Naves Novis. App. (1880) 266; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 277.

Luzon (Ilocos Norte, Ilocos Sur, Bontoc, Lepanto, Benguet, Pangasinan), *Cuming* 981, *Merrill* 4451, 7750, *Williams* 971, 1455, *B. S.* 4905, 27527 *Ramos*, *F. B.* 4836, 15618 *Curran*. On open grassy slopes at medium altitudes, ascending to 1,600 m. Malay Peninsula, Indo-China, southern China, New Guinea, tropical Australia.

3. DIANELLA Lamarck

DIANELLA CAERULEA Sims in Curtis's Bot. Mag. 15 (1801) t. 505; Red. Lil. 2 (1805) t. 79; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 266, 5 (1910) Bot. 337; Hallier f. in Nova Guinea 8 (1914) 993.

Luzon (Cagayan, Ilocos Norte, Bontoc, Benguet, Pangasinan, Bulacan, Rizal, Bataan, Nueva Vizcaya, Laguna, Tayabas, Camarines, Sorsogon), Catanduanes, Polillo, Mindoro, Leyte, Negros, Mindanao. In forests at medium and higher altitudes, ascending to 2,200 m; represented by numerous collections. New Guinea, New Caledonia, Queensland, Fiji.

Local names: Abláas (Bag.); bariu-báriu (P. Bis.); duñgau (Ig.); oyon-eyon (P. Bis.).

DIANELLA ENSIFOLIA (Linn.) DC. in Red. Lil. 1 (1802) *t. 1*; Naves Novis. App. (1880) 264; Vidal Rev. Pl. Vasc. Filip. (1886) 277, Phan. Cuming. Philip. (1885) 153; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 266, 3 (1908) Bot. 401, 5 (1910) Bot. 337; Hallier f. in Nova Guinea 8 (1914) 992.

Dracaena ensifolia Linn. Mant. 1 (1767) 63.

Dianella robusta Elm. Leaf. Philip. Bot. 5 (1913) 1806.

Palawan, Panay, Jolo, Bucas Grande, Mindanao, *B. S. 35067, 34801 Ramos & Pascasio, 30995 Ramos & Edaño, 21562 Escritor, Elmer 12900, Merrill 5317, 9338, Clemens 9377*. In light forests at low altitudes. India to the Mascarene Islands, China, Formosa, Malaya, tropical Australia, and Polynesia.

DIANELLA JAVANICA (Blume) Kunth Enum. 5 (1850) 52; Hallier f. in Nova Guinea 8 (1914) 992.

Rhuacophila javanica Blume Enum. 1 (1827) 14; Ridl. in Journ. Linn. Soc. Bot. 42 (1914) 166.

Dianella austro-caledonica Seem. Fl. Vit. (1865-73) 312; Lauterb. in Engl. Bot. Jahrb. 50 (1913) 294.

Batan Islands, Luzon (Ifugao, Bontoc, Lepanto, Benguet, Zambales, Tayabas, Camarines, Albay), Mindoro, Mindanao (Davao), *B. S. 31837 Santos, 8478, 17671, 8873 McGregor, 5408 Ramos, 37636 Ramos & Edaño, Cuming 929, Merrill 5504, 6585, Williams 2556, 1274, 1533, Elmer 5797, Copeland 1035*. On open slopes, in ravines, and in forests, altitude 1,000 to 2,400 m. Malay Peninsula, Sumatra, Java, Borneo, Celebes, Moluccas, New Guinea, New Caledonia.

Local names: Apilug (Ig.); sapiláu (Ig.); talobatub (Bon.); uráya (Ig.).

DOUBTFUL AND EXCLUDED SPECIES

DIANELLA BANCANA Miq.; Naves Novis. App. (1880) 264.

DIANELLA MONTANA Blume; Naves l. c.

DIANELLA PHILIPPENSIS Perr. in Mém. Soc. Linn. Paris 3 (1824) 112; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 305.

"Especie nouvelle de Mindanao" Perrottet. *A nomen nudum*.

DIANELLA REVOLUTA R. Br.; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Supp. 1: 445.

Meyen's Jala Jala specimen, so identified by Schauer, is probably referable to the species I have interpreted as *D. caerulea* Sims.

4. ALOE Tournefort

ALOE VERA Linn. Sp. Pl. (1753) 320; Berger in Engl. Pflanzenreich 33 (1908) 228, f. 84, A-D; Merr. Sp. Blancoanae (1918) 95.

Aloe humilis Blanco Fl. Filip. (1837) 256, ed 2 (1845) 179, ed. 3, 1 (1877) 321, t. 95, non Linn.

Aloe barbadensis Mill. Gard. Dict. ed. 8 (1768) No. 2; Naves Novis. App. (1880) 266.

Cultivated for ornamental and medicinal purposes, *Merrill Sp. Blancoanae* 1005. Subtemperate and tropical regions generally, often cultivated.

Local names: Dílang-boáia (Bis.); dílang-hálo (Bis.); sábila (Tag.); sábila-píña (Tag.).

EXCLUDED SPECIES

ALOE UMBELLATA DC.; Naves Novis. App. (1880) 266.

5. ALLIUM Tournefort

No species of this genus occurs naturally in the Philippines, although several are cultivated. These include the onion, *Allium cepa* Linn., locally known as *cebollas* or *sibuyas*, *lasoná* (Ilk.); the garlic, *Allium sativum* Linn., locally known as *ájos* and *báuang*; and *A. tuberosum* Roxb., a form locally known as *cuchái* or *ganda*. The last is the form described by Blanco as *Allium tricoccum* (non Ait.) and by Loureiro as *Allium angulosum* (non Linn.). Other species credited to the Philippines by Naves Novis. App. (1880) 267 are *Allium fistulosum* Linn., *A. jacquemontii* Kunth, *A. ascalonicum* Linn., and *A. uliginosum* Don = *A. tuberosum* Roxb.; I have seen no Philippine specimens of the first three, and it is probable that they do not occur in the Philippines even in cultivation.

6. LILIUM Tournefort

LILIUM LONGIFLORUM Thunb. in Trans. Linn. Soc. 2 (1794) 333; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 400.

Y'Ami and Batan Islands, *Worcester, B. S. 3774 Fénix*. On open grassy slopes at low altitudes. Japan, China, Formosa.

Local name: Vonitan (Iv.).

LILIUM PHILIPPINENSE Baker in Gard. Chron. (1873) 1141, f. 243, Journ. Linn. Soc. Bot. 14 (1874) 228 (*philippense*); Curtis's Bot. Mag. III 32 (1876) t. 6250; Naves Novis. App. (1880) 266; Vidal Rev. Pl. Vasc. Filip. (1886) 277; Merr. Philip. Journ. Sci. 5 (1910) Bot. 337.

Luzon (Bontoc, Benguet, Pangasinan), *B. S. 31851 Santos, 8791 McGregor, 4349, 3493 Mearns, 8291 Ramos, Merrill 4802, Williams 1132, Elmer 6389*. On open slopes in the pine region, altitude 1,100 to 2,300 m. Formosa.

Local names: Kanyon (Ilk.); luplúpak (Ig.); soyásoi (Ig.); tuktukpáu (Bon.); tubtubkau (Ig.).

7. YUCCA Dillenius

YUCCA ALOIFOLIA Linn. Sp. Pl. (1753) 319; Naves Novis. App. (1880) 266; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 32, Fl. Manila (1912) 145.

Planted here and there for ornamental purposes, but not naturalized. A native of tropical America.

EXCLUDED SPECIES

YUCCA GLORIOSA Linn.; Naves Novis. App. (1880) 266.

YUCCA SUPERBA Haw.; Naves l. c.

8. TAETSIA Medicus

(*Cordyline* Commerson)

TAETSIA FRUTICOSA (Linn.) Merr. Interpret. Herb. Amb. (1917) 137, Sp. Blancoanae (1918) 96.

Convallaria fruticosa Linn. in Stickman Herb. Amb. (1754) 16, Amoen. Acad. 4 (1759) 126, Syst. ed. 10 (1759) 894.

Asparagus terminalis Linn. Sp. Pl. ed. 2 (1762) 450.

Dracaena terminalis Rich. in Lam. Encycl. 2 (1786) 324; Blanco Fl. Filip. (1837) 263, ed. 2 (1845) 183, ed. 3, 1 (1877) 328, t. 98.

Cordyline terminalis Kunth in Abh. Acad. Berlin (1820) 30, Enum. 5 (1850) 25; Miq. Fl. Ind. Bat. 3 (1855) 558; Naves Novis. App. (1880) 265; Merr. Fl. Manila (1912) 145.

Widely cultivated for ornamental purposes, but not a native of the Archipelago. Pantropic in cultivation, probably a native of Malaya.

Local names: Bastón de San José (Sp.); daṅgá (Ilk.); dang-ṅgá (Bon.); donglá (If.); kiláa (Bis.); kilála (Bik.); sagilála (Tag.); tókor-pári (Pamp.); tónkod-obísipo (Tag.); túngkod-pári (Tag.).

9. PLEOMELE Salisbury

PLEOMELE ANGUSTIFOLIA (Roxb.) N. E. Br. in Kew Bull. (1914) 277; Merr. Interpret. Herb. Amb. (1917) 137, Sp. Blancoanae (1918) 96.

Dracaena angustifolia Roxb. Hort. Beng. (1814) 24, Fl. Ind. ed. 2, 2 (1832) 155; Naves Novis. App. (1880) 264; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 35, 3 (1908) Bot. 400.

Pandanus inermis Blanco Fl. Filip. ed. 2 (1845) 537, ed. 3, 3 (1879) 184, non Roxb.

Batan Islands and northern Luzon to Palawan and Mindanao, in most or all islands and provinces. Chiefly in primary forests at low and medium altitudes; represented by more than 50 individual collections. India through Malaya to Australia.

Local names: Gigiman (Pang.); malasambál (Tag.); pasiú (Ibn.); saranakatmano (Bag.); taligbúhuk (Tag.); uhaṅgo-nu-kurang (Iv.).

PLEOMELE MULTIFLORA (Warb.) comb. nov.

Dracaena multiflora Warb. ex Sarasin Reisen in Celebes 1 (1905) 136, t. 4.

Batan Islands, Luzon (Nueva Ecija, Tayabas), Polillo, Mindoro, Palawan, Ticao, Masbate, Sulu Archipelago, Merrill 9424, 5402 B. S. 26434 Ramos & Edaño, 10407 McGregor, 39508 Ramos, F. B. 18815 Foxworthy, De Mesa, & Villamil. Chiefly on limestone formations along the seashore. Celebes.

EXCLUDED SPECIES

DRACAENA ACUMINATA Thunb.; Naves Novis. App. (1880) 264.

DRACAENA DRACO Linn.; Naves op. cit. 265.

10. CORDYLINE Royen ex Adanson

(Sansevieria Thunberg)

CORDYLINE ROXBURGHIANA (Schultes) comb. nov.

Sansevieria roxburghiana Schultes Syst. 7 (1829) 357; Naves Novis.

App. (1880) 265; N. E. Br. in Kew Bull. (1915) 227, f. 12, D. E.

Sansevieria zeylanica Roxb. Pl. Coromandel. 2 (1798) 43, t. 184, quoad fig.; Naves Novis. App. (1880) 265, Fl. Filip. ed. 3 (1877-83) t. 422; Miq. Fl. Ind. Bat. 3 (1859) 559; Merr. Fl. Manila (1912) 144, non Willd.*Cordylina hyacinthoides* W. F. Wight in Contr. U. S. Nat. Herb. 9 (1905) 249, non *Aloe hyacinthoides* Linn.

Often cultivated, but in many regions naturalized, occurring in thickets and hedges at low and medium altitudes. A native of tropical Asia, now pantropic in cultivation. Bowstring hemp.

Local names: Aspi-aspi (Pamp.); baniat (Is.); bontót-pálos (Tag.); dildíla (Ilk.); kakaróhai (Ibn.); pakaróhai (Ibn.); rabo de león (Sp.); rabo de tigre (Sp.); sígre (Is.); sináua (Tag.); tígi (Ibn.); tigre (Sp.).

EXCLUDED SPECIES

SANSEVIERIA GRANDICUSPIS Haw.; Naves Novis. App. (1880) 265.

SANSEVIERIA GUINEENSIS Willd.; Naves l. c.

11. ASPARAGUS Tournefort

ASPARAGUS COCHINCHINENSIS (Lour.) Merr. in Philip. Journ. Sci. 15 (1919) 230.

Melanthium cochinchinense Lour. Fl. Cochinch. (1790) 216.*Asparagus lucidus* Lindl. Bot. Reg. (1844) Misc. 29, var. *dolichocladus* Merr. & Rolfe in Philip. Journ. Sci. 3 (1908) Bot. 96.

Luzon (Benguet), *Loher* 1928, without data, but probably from above 1,200 m altitude. Japan, China, Formosa.

The above is the only indigenous species of the genus. Introduced and occasionally cultivated species include the common asparagus (Sp. espárrago) *Asparagus officinalis* Linn., *A. plumosus* Baker, and *A. sprengeri* Regel; see Merrill, Fl. Manila (1912) 143. Naves has credited *Asparagus declinatus* Linn. and *A. racemosus* Willd. to the Philippines, Novis. App. (1880) 264, but both were apparently admitted on erroneous identifications.

12. DISPORUM Salisbury

DISPORUM LUZONIENSE Merr. in Philip. Journ. Sci. 5 (1910) Bot. 338.

Disporum pullum Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 182, non Salisb.

Luzon (Lepanto, Bontoc, Benguet), *Merrill* 4857, 6619, *Phil. Pl.* 899, *B. S.* 31675 *Santos*, 5942 *Ramos*, 8483 *McGregor*, 4316 *Mearns*. In the mossy forest, altitude 1,700 to 2,400 m. Endemic.

13. LIRIOPE Loureiro

LIRIOPE BRACHYPHYLLA Merr. in Philip. Journ. Sci. 2 (1907) Bot. 266.

Mindoro, *Merrill* 5710. On seepage slopes in open heath on Mt. Halcon, altitude about 2,400 m. Endemic.

LIRIOPE GRAMINIFOLIA (Linn.) Baker in Journ. Linn. Soc. Bot. 17 (1879) 499.

Dracaena graminifolia Linn. Syst. ed. 12 (1767) 246.

Liriope spicata Lour. Fl. Cochinch. (1790) 201; C. H. Wright in Journ. Linn. Soc. Bot. 36 (1903) 79.

Ophiopogon spicatus Ker in Bot. Reg. t. 593; Naves Novis. App. (1880) 264.

Luzon (Ilocos Norte, Benguet), *F. B. 968 Barnes, B. S. 33196 Ramos*. On open slopes, altitude about 1,400 m. Japan to Manchuria, southward to Hongkong and Formosa.

14. OPHIOPOGON Ker

OPHIOPOGON JAPONICUS (Linn. f.) Ker in Curtis's Bot. Mag. (1807) t. 1063; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 35, 5 (1910) Bot. 338.

Convallaria japonica Linn. f. Suppl. (1781) 204.

Luzon (Bontoc, Lepanto, Benguet, Nueva Ecija, Zambales, Pampanga, Bataan, Laguna, Tayabas), Mindanao (Bukidnon, Davao). Chiefly in the mossy forest, altitude 850 to 2,400 m; represented by numerous collections. Japan to southern China and Formosa.

Local names: Langigit (Ig.); takaáu (Bon.); uli-úli (Bag.).

OPHIOPOGON MALAYANUS Ridl. in Journ. Str. Branch Roy. As. Soc. 41 (1904) 34.

Jolo, *Clemens 9374*. On forested slopes, altitude about 700 m. Malay Peninsula, Borneo.

15. ALETRIS Linnaeus

ALETRIS SPICATA (Thunb.) Franch. in Journ. de Bot. 10 (1896) 199; Merr. in Philip. Journ. Sci. 1 (1908) Suppl. 182, 5 (1910) Bot. 338.

Hypoxis spicata Thunb. Fl. Jap. (1784) 136.

Aletris japonica Lamb. in Trans. Linn. Soc. 10 (1811) 407, non Thunb.

Luzon (Abra, Bontoc, Benguet), *Merrill 4685, 4801, Clemens 9178, Vanoverbergh 568, B. S. 7242 Ramos, 31668 Santos*. On open grassy slopes in the pine region, altitude 1,600 to 2,300 m. Japan, China, Formosa.

Local name: Salenganga (Ig.).

16. SMILAX Tournefort

SMILAX BRACTEATA Presl Rel. Haenk. 1 (1827) 131; Kunth Enum. 5 (1850) 251; Miq. Fl. Ind. Bat. 3 (1859) 566; A. DC. Monog. Phan. 1 (1878) 197; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 87, Fl. Manila (1912) 142, Sp. Blancoanae (1918) 97.

Smilax pseudochina Blanco Fl. Filip. (1837) 795, ed. 2 (1845) 548, ed. 3, 3 (1879) 204, non Linn.

Smilax fistulosa Blanco op. cit. 796, 549, 205; A. DC. Monog. Phan. 1 (1878) 211.

Smilax divaricata Blanco op. cit. 795, 548, 206, p. p.; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 87.

Smilax blancoi Kunth Enum. 5 (1850) 262; A. DC. Monog. Phan. 1 (1878) 211.

Smilax indica Naves Novis. App. (1880) 262, non Vitm.

Luzon (Benguet, Pangasinan, Bulacan, Rizal, Bataan, Laguna, Tayabas), *Merrill 2548, Sp. Blancoanae 572, Phil. Pl. 313, B. S. 17777 Otanes*. Common in thickets and secondary forests at low and medium altitudes. Endemic.

Local names: Banág (Ilk., Tag., Ig.); banál (Ig.); barág (Pamp.); kamágsa-óbat (Tag.); hampás-tigbálang (Tag.); kolót-bábui (Tag.); rónas (Pamp.).

SMILAX CHINA Linn. Sp. Pl. (1753) 1029; Naves Novis. App. (1880) 261; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 267, 5 (1910) Bot. 339.

Luzon (Bontoc, Lepanto, Benguet, Zambales), Mindoro, Negros, *B. S. 31754 Santos, 5413, 5733 Ramos, 8411 McGregor, F. B. 17382 Curran, Merrill 4749, 4497, 6126, 6211, 6140, 6552, 6494, 7043, Williams 1340, 2008*. In the mossy forest, altitude 1,600 to 2,400 m. Japan, China.

Local names: Buánal (Ig.); palitpit (Bon.).

SMILAX ELMERI Merr. in Philip. Journ. Sci. 13 (1918) Bot. 6.

Smilax reticulata Elm. Leaf. Philip. Bot. 8 (1915) 2740, non Desv., nec Heer.

Luzon (Sorsogon), Bucas Grande, Mindanao (Bukidnon, Agusan), *Elmer 13684, 16132, B. S. 35065 Ramos & Pascasio, 38761 Ramos & Edaño*. On forested ridges up to 1,000 m altitude. Endemic.

Local name: Simbanag (Mbo.).

SMILAX ERECTA Merr. in Philip. Journ. Sci. 13 (1918) Bot. 5.

Luzon (Tayabas, east coast), *B. S. 28560, 28472 Ramos & Edaño*. On forested slopes at low or medium altitudes. Endemic.

SMILAX LEUCOPHYLLA Blume Enum. Pl. Jav. 1 (1827) 18; Naves Novis. App. (1880) 263; Stapf in Trans. Linn. Soc. Bot. 4 (1894) 242; Ceron Cat. Pl. Herb. Manila (1892) 172; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 96, *Sp. Blancoanae* (1918) 97.

Smilax latifolia Blanco Fl. Filip. ed. 2 (1845) 548, ed. 3, 3 (1879) 204, non R. Br.

Smilax vicaria Kunth Enum. 5 (1850) 262; A. DC. Monog. Phan. 1 (1878) 212; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 35, 2 (1907) Bot. 267, 436.

Smilax macrophylla Naves Novis. App. (1880) 262, non Roxb.

Luzon (Benguet, Pangasinan, Pampanga, Rizal, Bataan, Laguna), Mindoro, Culion, Palawan, Mindanao, *Merrill 3771, 1713, 659, 5579, Sp. Blancoanae 951, 690, Elmer 12613, Williams 1555, B. S. 20554, 10888 Ramos, 22776 McGregor, 12664, 15865 Fénix, 1992 Foxworthy, F. B. 1877 Ahern's collector*. Chiefly in primary forests at medium altitudes. Malay Peninsula, Java, Borneo, Moluccas.

Local names: Banág (Tagb.); banál (Ig.); kaguno (Neg.); kámot-kabág (Tag.); sarsaparílang-putí (Tag.); sípit-uláng (Tag.).

SMILAX LOHERI Merr. in Philip. Journ. Sci. 10 (1915) Bot. 294.

Luzon (Nueva Ecija, Rizal), *Loher 6968, B. S. 26506 Ramos & Edaño, 40708 Ramos*. In the mossy forest at or above 900 m altitude.

Local name: Kiting-kiting (Neg.).

SMILAX LUCIDA Merr. in Philip. Journ. Sci. 13 (1918) Bot. 5.

Luzon (Rizal, Tayabas, east coast), *B. S. 26611 Ramos & Edaño, 40787 Ramos*. In forests at low or medium altitudes. Endemic.

Local name: Báru (Ibn.).

SMILAX LUZONENSIS Presl Rel. Haenk. 1 (1827) 131; Kunth Enum. 5 (1850) 261; Miq. Fl. Ind. Bat. 3 (1859) 566; Merr. in Philip. Journ. Sci. 10 (1915) Bot. 295.

Smilax helferi A. DC. Monog. Phan. 1 (1878) 176, quoad. syn. Presl; Naves Novis. App. (1880) 262.

Luzon (Ilocos Norte), Panay, Catanduanes, Mindanao (Bukidnon, Lanao), *B. S. 33327, 30228 Ramos, 35611 Martelino & Edaño, 38849 Ramos & Edaño, Clemens 835, 905, Presl!* in herb. Prague. In forests at medium altitudes. Endemic.

SMILAX PYGMAEA Merr. in Philip. Journ. Sci. 5 (1910) Bot. 339.

Luzon (Benguet), *Merrill 6598, B. S. 8902*. In the summit grasslands on Mt. Pulog, altitude about 2,800 m. Endemic.

SMILAX VERRUCULOSA Merr. in Philip. Journ. Sci. 5 (1910) Bot. 173.

Luzon (Ifugao, Bontoc, Benguet), *Topping 13, Merrill 7742, Elmer 5820, 8572, Williams 1046, B. S. 19702 McGregor, 37672 Ramos & Edaño, 2810 Mearns, F. B. 15617 Curran*. In thickets about limestone cliffs, in forests and ravines, altitude 1,300 to 1,600 m. Endemic.

Local names: Baná (Bon.); pauá (Bon.).

SMILAX WILLIAMSII Merr. in Philip. Journ. Sci. 5 (1910) Bot. 173.

Luzon (Tayabas, east coast), Mindanao (Lanao, Bukidnon, Davao), *Williams 2579, Clemens 751, B. S. 29139, 38760 Ramos & Edaño*. In thickets or forests at medium altitudes. Endemic.

Local name: Banúg (Bag.).

EXCLUDED SPECIES

SMILAX ASPERA Linn.; Naves Novis. App. (1880) 262.

SMILAX BARBATA Wall.; Naves op. cit. 263.

SMILAX BAUHINIODES Kunth; Naves op. cit. 262.

SMILAX BLUMEI A. DC.; Naves op. cit. 263.

SMILAX FEROX Wall.; Naves op. cit. 262.

SMILAX JAVENSIS A. DC.; Naves l. c.

SMILAX GRANDIFOLIA Regel; Naves l. c.

SMILAX KLOTZSCHII Kunth; Naves l. c.

SMILAX LAEVIS Wall.; Naves op. cit. 261.

SMILAX MACRABUCHA Duch.; Naves op. cit. 263.

SMILAX MACROCARPA Blume; Naves op. cit. 262.

SMILAX ODORATISSIMA Blume; Naves l. c.; Stapf in Trans. Linn. Soc. Bot. 4 (1894) 242.

SMILAX OVALIFOLIA Roxb.; Naves op. cit. 263.

SMILAX POLYACANTHA Wall.; Naves op. cit. 262.

SMILAX WIGHTII A. DC.; Naves l. c.

SMILAX ZEYLANICA Linn.; Naves l. c.

EXCLUDED GENERA

HETEROSMILAX BORNEENSIS A. DC.; Naves Novis. App. (1880) 261.

SCILLA sp.; Naves op. cit. 266.

TULIPA GESNERIANA Linn.; Naves op. cit. 266.

XEROTES LONGIFOLIA R. Br.; Naves op. cit. 273.

AMARYLLIDACEAE

1. ATAMOSCO Greene

ATAMOSCO ROSEA (Lindl.) Greene Pittonia 3 (1897) 188.

Zephyranthes rosea Lindl. in Bot. Reg. 10 (1824) t. 821; Merr. Fl. Manila (1912) 147.

Habranthus versicolor Herb. in Curtis's Bot. Mag. t. 2485; Naves Novis. App. (1880) 255, Fl. Filip. ed. 3 (1877-83) t. 271, 375.

Widely distributed in cultivation in and about towns, sometimes persisting from previous cultivation but scarcely naturalized. A native of tropical America.

2. CRINUM Linnaeus

CRINUM ASIATICUM Linn. Sp. Pl. (1753) 292; Blanco Fl. Filip. (1837) 251, ed. 2 (1845) 175, ed. 3, 1 (1877) 314, t. 168; Miq. Fl. Ind. Bat. 3 (1859) 580; Naves Novis. App. (1880) 254; Merr. in Philip. Journ. Sci. 1 (1906) Bot. 35, Fl. Manila (1912) 147, Sp. Blancoanae (1918) 98.

Crinum giganteum Blanco Fl. Filip. ed. 2 (1845) 175, ed. 3, 1 (1877) 315, non Andr.

Haemanthus pubescens Blanco Fl. Filip. (1837) 253, non Linn. f.

Throughout the Philippines along sandy seashores, sometimes planted inland. India to Malaya and western Polynesia.

Local names: Agabáhan (Bis.); agákong (Bon.); bákong (Tag., Bag., Ilk., Sbl.); biliba (Sub.); kábong (Bik.); palagúkon (Bis.); saknib (Bon.); salibángbang (Bis.); tebá (Bon.).

CRINUM GRACILE E. Meyer in Presl Rel. Haenk. 1 (1827) 120; Kunth Enum. 5 (1850) 566; Naves Novis. App. (1880) 255; Baker Handb. Amaryl. (1888) 81.

Crinum cumingii Baker in Gard. Chron. II 16 (1881) 72, Handb. Amaryl. (1888) 81; Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 275.

Luzon (Nueva Ecija, Laguna), Mindoro, Mindanao, *Cuming 1882, Presl! in herb. Prague, Merrill 4063, McGregor, 152, 228, Elmer 17667, Baker 3725, Bolster 303.* In primary forests at low altitudes. Endemic.

CRINUM LATIFOLIUM Linn. Sp. Pl. (1753) 291; Merr. Fl. Manila (1912) 148.

Cultivated in towns for ornamental purposes, but scarcely naturalized. A native of India.

CRINUM ZEYLANICUM Linn. Syst. ed. 12 (1767) 236; Baker Handb. Amaryll. (1888) 87; Merr. Fl. Manila (1912) 148.
Crinum ornatum Herb. Amaryll. (1837) 262; Naves Novis. App. (1880) 255.

Cultivated for ornamental purposes but nowhere naturalized. A native of tropical Asia.

Local names: Lilio (Sp.); lirio (Tag.).

EXCLUDED SPECIES

CRINUM AMABILE Donn; Naves Novis. App. (1880) 254.

CRINUM AMOENUM Roxb.; Naves l. c.

CRINUM DEFIXUM Ker; Naves l. c.

CRINUM PRATENSE Hub.; Naves l. c.

CRINUM SUMATRANUM Roxb.; Naves l. c.

3. HYMENOCALLIS Salisbury

HYMENOCALLIS LITTORALE (Jacq.) Salisb. in Trans. Hort. Soc. 1 (1812) 338; Merr. Sp. Blancoanae (1918) 98.

Pancratium littorale Jacq. Select. Stirp. Amer. (1763) 99.

Pancratium illyricum Blanco Fl. Filip. (1837) 251, ed. 2 (1845) 176, ed. 3, 1 (1877) 316, t. 411 (as *Hymenocallis adnata* Herb.), non Linn.

Pancratium maritimum Blanco op. cit. 252, 177, 316; Naves Novis. App. (1880) 256; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 86, non Linn.

Hymenocallis adnata Herb. Amaryll. (1837) 215; Naves Novis. App. (1880) 256.

Hymenocallis tenuiflora Herb. Amaryll. Suppl. (1837) 44; Merr. Fl. Manila (1912) 149.

Cultivated in Manila and other towns, occasional in waste places, etc., but apparently propagating only by its bulbs. A native of tropical America.

4. EUCHARIS Planchon and Linden

EUCHARIS GRANDIFLORA Planch. & Linden Fl. des Serres I 9 (1853-54) 255; Merr. Fl. Manila (1912) 149.

This occurs in the Philippines only as a cultivated ornamental in Manila and in a few of the larger towns. A native of South America. The Amazon lily.

5. EURYCLES Salisbury

EURYCLES AMBOINENSIS (Linn.) Lindl. in Loud. Encycl. Pl. (1829) 242; Miq. Fl. Ind. Bat. 3 (1859) 582; Naves Novis. App. (1880) 255; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 276; Merr. Fl. Manila (1912) 149, Sp. Blancoanae (1918) 99.

Pancratium amboinense Linn. Sp. Pl. (1753) 291; Blanco Fl. Filip. (1837) 252, ed. 2 (1845) 177, ed. 3, 1 (1877) 317, t. 406.

Eurycles sylvestris Salisb. in Trans. Hort. Soc. 1 (1812) 337.

Luzon (Cavite, Rizal, Laguna), Mindoro, Palawan. In secondary forests at low altitudes, never far from settlements and hence probably introduced; often cultivated for ornamental purposes. Malay Peninsula and Archipelago to tropical Australia.

Local names: Abod (Bis.); abur (Bis.); dausum (Bis.); katañgal (Bis.); kosol (Bis.); panabor (Bis.); taliunúd (Bik.); talaonor (Bis.); tambal (Tag.); tanual (Bis.).

6. PANCRATIUM Linnaeus

PANCRATIUM ZEYLANICUM Linn. Sp. Pl. (1753) 290; Blanco Fl. Filip. (1837) 253, ed. 2 (1845) 177, ed. 3, 1 (1877) 317, *t.* 321; Naves Novis. App. (1880) 256; Merr. Fl. Manila (1912) 148, Sp. Blancoanae (1918) 99.

Luzon (Union, Camarines), Palawan, Biliran, *Merrill Sp. Blancoanae* 378, *B. S. 18534 McGregor, 881 Foxworthy*. In coconut and abaca plantations, thickets, etc., sometimes planted. Certainly not a native of the Philippines. India to Malaya.

Local names: Baoang-báoang (Bis.); hagobaóa (Bis.); katoñgal (Tag.).

EXCLUDED SPECIES

PANCRATIUM MALABATHRICUM Herb.; Naves Novis. App. (1880) 256.

7. HIPPEASTRUM Herbert

HIPPEASTRUM MINIATUM Herb. Amaryll. Suppl. (1837) 31; Naves Novis. App. (1880) 255; Merr. Sp. Blancoanae (1918) 99.

Amaryllis atamasco Blanco Fl. Filip. (1837) 254, ed. 2 (1845) 178, ed. 3, 1 (1877) 319, *t.* 359; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 85, non Linn.

Luzon (Union, Rizal, Laguna, Tayabas), Panay, Mindanao, Siassi, *Merrill Sp. Blancoanae* 1048, *Elmer 17438b*. Introduced and planted for ornamental purposes, but more or less naturalized in coconut groves, etc. A native of tropical America.

Local name: Lirio (Tag.).

8. POLIANTHES Linnaeus

POLIANTHES TUBEROSA Linn. Sp. Pl. (1753) 316; Blanco Fl. Filip. (1837) 259, ed. 2 (1845) 181, ed. 3, 1 (1877) 323; Naves Novis. App. (1880) 254; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 85, Fl. Manila (1912) 146, Sp. Blancoanae (1918) 99.

Cultivated for its fragrant flowers but nowhere naturalized. A native of tropical America. Tuberose.

Local names: Azucéna (Sp.); nárdó (C. Bis.).

9. AGAVE Linnaeus

AGAVE CANTALA Roxb. Hort. Beng. (1814) 25; Berger Agav. (1915) 236; Merr. Interpret. Herb. Amb. (1917) 144, Sp. Blancoanae (1918) 99.

Furcraea cantala Haw. Syn. Pl. Succul. Suppl. (1819) 42.

Agave cantula Roxb. Fl. Ind. ed. 2, 2 (1832) 167; Naves Novis. App. (1880) 253; Merr. Fl. Manila (1912) 146.

Agave americana Blanco Fl. Filip. (1837) 258, ed. 2 (1845) 180, ed. 3, 1 (1877) 322, *t.* 96; Naves Novis. App. (1880) 253; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 23; Usteri Beitr. Ken. Philip. Veg. (1905) 129, non Linn.

Introduced from Mexico at an early date in colonial history, now widely distributed in the Philippines, in some provinces extensively grown for its fiber. Pantropic.

Local names: magéi (all dialects); píta (C. Bis.).

AGAVE SISALANA Perrine ex Engelm. in Trans. Acad. Sci. St. Louis 3 (1875) 314; Berger Agav. (1915) 230. f. 66.

Introduced in the early part of the present century, and now widely planted for its fiber. A native of tropical America. Sisal.

EXCLUDED SPECIES

AGAVE MEXICANA Lam.; Naves Novis. App. (1880) 253.

10. CURCULIGO Gaertner

CURCULIGO AGUSANENSIS Elm. Leaf. Philip. Bot. 5 (1913) 1645.

Luzon (Sorsogon), Mindanao (Zamboanga, Agusan), *Elmer 13584, 17005, B. S. 36910 Ramos & Edaño*. In forests at low altitudes. Endemic.

Local names: Kuyabáng (Sub.); pagátpat (Mbo.); tayábong (Sub.).

CURCULIGO BREVIPEDUNCULATA Elm. Leaf. Philip. Bot. 5 (1913) 1647.

Palawan, Balabac, Panay, *Elmer 13164, Merrill 5373, Phil. Pl. 1318, B. S. 32576 McGregor*. In forests at low altitudes. Endemic.

CURCULIGO CAPITULATA (Lour.) O. Kuntze Rev. Gen. Pl. (1891) 703; Merr. Interpret. Herb. Amb. (1917) 142.

Leucojum capitulatum Lour. Fl. Cochinch. (1790) 199.

Curculigo recurvata Dry. in Ait. Hort. Kew ed. 2, 2 (1811) 253; Naves Novis. App. (1880) 253; Merr. in Philip. Journ. Sci. 3 (1908) Bot. 401; Elm. Leaf. Philip. Bot. 5 (1913) 1645.

Batan Islands, Luzon (Bontoc, Benguet, Laguna, Tayabas, Sorsogon, Bili-ran, Leyte, Panay, Siargao, Mindanao), *Merrill 9670, Phil. Pl. 847, Elmer 6202, 7771, 18196, Williams 1150, B. S. 35389 Ramos & Edaño, 31962 Santos, 15299, 23571 Ramos, 18867 McGregor, 14006 Robinson, F. B. 4893 Curran*. In thickets and forests, chiefly at medium altitudes, ascending to 1,800 m. India to southern China through Malaya to tropical Australia.

Local names: Abang-ábang (Bik.); atukgan (Ig.); katuágan (Ig.); talauángit (Bag.); tayúbang (Bon.); tokán (Ig.); tolábang (If.).

CURCULIGO GLABRA Merr. in Philip. Journ. Sci. 2 (1907) Bot. 267; C. B. Rob. op. cit. 6 (1911) Bot. 196; Elm. Leaf. Philip. Bot. 5 (1913) 1645.

Luzon (Sorsogon), Polillo, Alabat, Mindoro, *Merrill 5750, 10449, Elmer 14990, B. S. 10366 McGregor*. In primary forests at low and medium altitudes. Endemic.

CURCULIGO ORCHIOIDES Gaertn. Fruct. 1 (1788) 63, t. 16; Elm. Leaf. Philip. Bot. 5 (1913) 1645; Merr. Sp. Blancoanae (1918) 100.

Gethyllis acaulis Blanco Fl. Filip. (1837) 260, ed. 2 (1845) 181, ed. 3, 1 (1877) 324.

Luzon (Bontoc, Pangasinan, Rizal, Sorsogon), Mindoro, Semirara, Palawan, Biliran, Panay, Mindanao, *Merrill 4144*, *Sp. Blancoanae 260*, *Elmer 16484*, *Williams 2927*, *Copeland 472*, *B. S. 32413*, *18637 McGregor*, *26122 Fénix*, *1520 Bermejos*, *6739 Robinson*, *21440 Escritor*, *17858 Otanes*. In open grasslands, chiefly in cogonales (*Imperata* areas), at low and medium altitudes. India to Java.

Local names: Sulsulitík (Bon.); taloáñgi (Bag.); tataluañgi (Buk.).

CURCULIGO WEBERI Elm. Leaf. Philip. Bot. 5 (1913) 1646.

Samar, Mindanao (Agusan, Lanao), *Elmer 14304*, *Weber 1081*, *Clemens 1225*, *Taylor 285*, *B. S. 17657 Ramos*. In forests at low and medium altitudes. Endemic.

Local name: Bongbón (C Bis.).

EXCLUDED SPECIES

CURCULIGO ENSIFOLIA R. Br.; Naves Novis. App. (1880) 253.

CURCULIGO LATIFOLIA Dry.; Naves l. c.

11. HYPOXIS Linnaeus

HYPOXIS AUREA Lour. Fl. Cochinch. (1790) 200.

Hypoxis franquevillei Miq. Fl. Ind. Bat. 3 (1859) 586; Naves Novis. App. (1880) 253.

Hypoxis flava Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 85 sphalm.

Luzon (Bontoc, Benguet, Nueva Ecija), Mindanao (Lanao, Bukidnon), *Elmer 6585*, *Merrill 247*, *4453*, *Phil. Pl. 848*, *Williams 1063*, *B. S. 15775 Fénix*, *31818 Santos*. In open grasslands at medium and higher altitudes, ascending to 2,300 m. India to Japan southward to Java.

Local name: Kitkitlí (Ig.).

EXCLUDED GENERA AND SPECIES

AMARYLLIS CARNEA Schultes; Naves Novis. App. (1880) 255.

AMARYLLIS FORMOSISSIMA Linn.; Naves l. c.

AMARYLLIS VITTATA Ait.; Naves l. c.

FURCRAEA GIGANTEA Vent.; Naves op. cit. 253.

FURCRAEA RIGIDA Haw.; Naves op. cit. 254.

HERMIONE TAZETTA Haw.; Naves op. cit. 256.

TACCACEAE

1. TACCA Forster

TACCA ELMERI Krause in Leaf. Philip. Bot. 6 (1914) 2283.

Palawan, Leyte, *Elmer 12679*, *Weber 1522*. In light forests at low altitudes. Endemic.

Local name: Odang-ódang (Tagb.).

TACCA PALMATA Blume Enum. Pl. Jav. 1 (1827) 83; Miq. Fl. Ind. Bat. 3 (1859) 577; Naves Novis. App. (1880) 257; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 35, Fl. Manila (1912) 150, *Sp. Blancoanae* (1918) 100.

Tacca montana Schultes Syst. 7¹ (1829) 168; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 276.

Tacca vesicaria Blanco Fl. Filip. (1837) 261.

Tacca palmata Blanco Fl. Filip. ed. 2 (1845) 182, ed. 3, 1 (1877) 325.

Tacca rumphii Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 442; Miq. Fl. Ind. Bat. 3 (1859) 577; Naves Novis. App. (1880) 257.

Luzon (Pangasinan, Nueva Ecija, Bulacan, Rizal, Bataan, Laguna), Semirara, Panay. In thickets and secondary forests at low altitudes. Malay Archipelago.

Local names: Kanálong (Bis.); magsalóro (Bis.); payung-payúṅgan (Tag.).

TACCA PINNATIFIDA Forst. Char. Gen. (1776) 70, *t.* 35; Presl Rel. Haenk. 1 (1827) 148; Blanco Fl. Filip. ed. 2 (1845) 182, ed. 3, 1 (1877) 327; Miq. Fl. Ind. Bat. 3 (1859) 577; Naves Novis. App. (1880) 257; Hook. f. in Curtis's Bot. Mag. III 49 (1893) *t.* 7299, 7300; Merr. Fl. Manila (1912) 150, Sp. Blancoanae (1918) 101.

Tacca gaogao Blanco Fl. Filip. (1837) 262, 856.

Luzon (Ilocos Norte) to Palawan and Mindanao, in most islands and provinces. Chiefly in sandy soil in thickets near the sea. Tropical Africa and Asia through Malaya to Australia and Polynesia.

Local names: Gau-gáu (Tag.); kanóbong (Bis.); panarién (Ilk.); tam-bóbon (Sbl.); tayóbong (Bis.); yabyában (Tag.).

EXCLUDED SPECIES

TACCA ASPERA Roxb.; Naves Novis. App. (1880) 258.

TACCA LANCAEFOLIA Zoll.; Naves l. c.

DIOSCOREACEAE

1. DIOSCOREA Plumier

DIOSCOREA ALATA Linn. Sp. Pl. (1753) 1033; Blanco Fl. Filip. (1837) 799, ed. 2 (1845) 550, ed. 3, 3 (1879) 207; Miq. Fl. Ind. Bat. 3 (1859) 572; Naves Novis. App. (1880) 259; Merr. Fl. Manila (1912) 151, Interpret. Herb. Amb. (1917) 146, Sp. Blancoanae (1918) 102; Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 39.

Cultivated throughout the Philippines for its edible tubers, but certainly not a native of the Archipelago. Pantropic in cultivation, a native of the Old World Tropics.

Local names: fpoi (Ig.); kinampái (Bis.); lúktu (If.); úbi (Tag.), Bik., Ibn., Ilk., It., Bis.).

DIOSCOREA BULBIFERA Linn. Sp. Pl. (1753) 1033; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 445; Miq. Fl. Ind. Bat. 3 (1859) 574; Naves Novis. App. (1880) 258; Ceron Cat. Pl. Herb. Manila (1892) 171; Merr. Fl. Manila (1912) 151; Prain & Burkill in Leaf. Philip. Bot. 5 (1913) 1596, Journ. As. Soc. Beng. II 10 (1914) 26.

Luzon (Bontoc, Union, Nueva Ecija, Batangas, Bulacan, Rizal, Bataan, Zambales, Sorsogon), Masbate, Leyte, Mindanao. In thickets at low

and medium altitudes; represented by numerous collections. India to China and Malaya.

Local names: Aribúkbuk (Ilk.); bayag-kabáyo (Tag.); dadákan (Bag.); pulúgan (Bik.); ubí-ubíhan (Tag.); utóñg-utóñgan (Tag.).

DIOSCOREA CUMINGII Prain & Burkill in Journ. As. Soc. Beng. II 4 (1908) 449, 10 (1914) 25, Leaflet. Philip. Bot. 5 (1913) 1595.

Dioscorea inaequifolia Elmer ex Prain & Burkill in Leaflet. Philip. Bot. 5 (1913) 1595, Journ. As. Soc. Beng. II 10 (1914) 24.

Luzon (Apayao, Bontoc, Benguet, Lepanto, Rizal, Bataan, Laguna, Nueva Vizcaya, Rizal, Camarines, Sorsogon), Mindoro, Mindanao, *Cuming* 1469, *Elmer* 15333, 8692, *Merrill* 4455, 7683, *Phil. Pl.* 2034, *F. B.* 22956bis *Calycosa*, 3376 *Ahern's collector*, 69 *Barnes*, *B. S.* 8207 *Ramos*. In thickets and forests at low and medium altitudes, ascending to 1,400 m. Endemic.

Local names: Lima-limá (Tag.); pari (Bag.).

DIOSCOREA DIVARICATA Blanco Fl. Filip. (1837) 797, ed. 2 (1845) 550, ed. 3, 3 (1879) 207; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 35, 2 (1907) Bot. 436, Sp. Blancoanae (1918) 101.

Dioscorea batatas Naves Novis. App. (1880) 258, non Decne.

Dioscorea foxworthyi Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 34.

Luzon (Ilocos Norte, Pangasinan, Pampanga, Rizal, Bataan, Laguna), *B. S.* 22425, 33185 *Ramos*, 29960, 30092 *Fénix*, 1558 *Foxworthy*, *F. B.* 2572 *Borden*, *Merrill Sp. Blancoanae* 391. In thickets and secondary and primary forests at low and medium altitudes. Endemic.

Local names: Baliákag (Bis.); bulói (Tag.); dulián (Ilk.); durián (Ilk.); áúyan (Pang.); kirói (Tag.); kobág (Tag.); pakít (Tag.); ubág (Tag.).

DIOSCOREA ELMERI Prain & Burkill in Leaflet. Philip. Bot. 5 (1913) 1594, Journ. As. Soc. Beng. II 10 (1914) 24 (incl. var. *vera* and var. *dubia*).

Batan Islands, Luzon (Laguna, Tayabas), Samar, *Elmer* 9156, 8265, *B. S.* 28717, 24196 *Ramos & Edaña*, 3659 *Fénix*. In forests at low altitudes; really distinct from *D. cumingii* Prain & Burkill? Endemic.

DIOSCOREA ESCULENTA (Lour.) Burkill in Gard. Bull. Straits Sett. 1 (1917) 396; Merr. Interpret. Herb. Amb. (1917) 147, Sp. Blancoanae (1918) 102.

Oncus esculentus Lour. Fl. Cochinch. (1790) 194.

Dioscorea papillaris Blanco Fl. Filip. (1837) 801, ed. 2 (1845) 552, ed. 3, 3 (1879) 200.

Dioscorea tugui Blanco Fl. Filip. (1837) 800.

Dioscorea sativa Blanco Fl. Filip. ed. 2 (1845) 551, ed. 3, 3 (1879) 209, Naves Novis. App. (1880) 259, non Linn.

Dioscorea fasciculata Roxb. Fl. Ind. ed. 2, 2 (1832) 801; Naves Novis. App. (1880) 260; Merr. Fl. Manila (1912) 151.

Dioscorea tiliaefolia Kunth Enum. 5 (1840) 401; Miq. Fl. Ind. Bat. 3 (1859) 576; Naves Novis. App. (1880) 260; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 276.

Dioscorea aculeata Naves Novis. App. (1880) 260; Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 19, Leaflet. Philip. Bot. 5 (1913) 1593 (var. *tiliaefolia* Prain & Burkill), non Linn.

Batan Islands, Luzon (Cagayan, Benguet, Union, Pangasinan, Nueva Vizcaya, Bulacan, Rizal, Bataan, Laguna), Semirara, *Merrill 3805, 2727, 2709, 4147, Sp. Blancoanae 677, 202, Phil. Pl. 325, Elmer 6421, B. S. 32265, 14381, 23076 McGregor, 21120 Escritor, 21720 Ramos, F. B. 3280 Ahern's collector, 24879 Mabesa*. In thickets and secondary forests at low altitudes, also often cultivated. India to Malaya and Polynesia.

Local names: Anég (Ibn.); bóga (Ilk.); dúkai (Iv.); kamigíng (Bik.); lúttu (Ibn.); tóngó (Tag.) túgi (Tag., Ilk.); túngo (Tag.).

DIOSCOREA FILIFORMIS Blume Enum. Pl. Jav. 1 (1827) 22; Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 38.

Dioscorea myriantha Kunth Enum. 5 (1850) 382; Miq. Fl. Ind. Bat. 3 (1859) 571; Naves Novis. App. (1880) 258; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 276; Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 38.

Luzon (Nueva Ecija, Rizal, Bataan, Batangas, Laguna), Culion, Palawan, Mindanao (Jolo), *Merrill 466, 857, 8058, 3167, 3120, 8244, Baker 460, B. S. 22266 Ramos, 209, 330 Bermejos, F. B. 6036 Zschokke*. In thickets and in ravines at low altitudes. Java, Celebes, Moluccas.

DIOSCOREA FLABELLIFOLIA Prain & Burkill in Leaf. Philip. Bot. 5 (1913) 1593, Journ. As. Soc. Beng. II 10 (1914) 12.

Luzon (Laguna, Sorsogon), Catanduanes, Mindoro, *Loher 7027, Elmer 9095, 16677, B. S. 30278, 23564, 39412 Ramos, 22830 McGregor*. In forests at low and medium altitudes. Borneo.

DIOSCOREA GRATA Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 35.

Luzon (Rizal), *Loher 7012, 7017*, without data as to habitat. Endemic.

DIOSCOREA HISPIDA Dennst. Schlüssel Hort. Malabar. (1818) 33; Merr. Interpret. Herb. Amb. (1917) 148, *Sp. Blancoanae* (1918) 103.

Dioscorea hirsuta Blume Enum. Pl. Jav. 1 (1827) 21; Miq. Fl. Ind. Bat. 3 (1859) 575; Naves Novis. App. (1880) 260.

Dioscorea daemona Roxb. Hort. Beng. (1814) 72, *nomen nudum*, Fl. Ind. ed. 2, 3 (1832) 805; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 86, Fl. Manila (1912) 151.

Dioscorea triphylla Blanco Fl. Filip. (1837) 799, ed. 2 (1845) 551, ed. 3, 3 (1879) 208; Miq. Fl. Ind. Bat. 3 (1859) 573; Naves Novis. App. (1880) 259; Prain & Burkill in Journ. As. Soc. Bengal II 10 (1914) 25, Leaf. Philip. Bot. 5 (1913) 1595, non Linn.

Throughout the Philippines in thickets and forests at low and medium altitudes. India to southwestern China and Formosa through Malaya to New Guinea.

Local names: Bagai (Mbo.); gáyos (Bis.); kalút (Tag., Pamp., Sbl.); karót (Ilk.); karóti (Sul.); káyos (Tag.); kolót (Bis.); korót (S. L. Bis.); kulót (Sbl.); mamó (Bik.); namí (Tag.); orkót (Bis.).

DIOSCOREA LOHERI Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 33; Merr. *Sp. Blancoanae* (1918) 101.

Dioscorea sp. (*cobag na quiroi*) Blanco Fl. Filip. (1837) 798, ed. 2 (1845) 550, ed. 3, 3 (1879) 206.

Luzon (Rizal), *Merrill Sp. Blancoanae* 634, *Phil. Pl.* 6, 1498, *B. S.* 12529 Ramos. In thickets at low altitudes. This is perhaps referable to *D. divaricata* Blanco, the plants from not fully grown tubers. Endemic.

Local names: Kirói (Tag.); kobág (Tag.).

DIOSCOREA LUZONENSIS Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 444; Miq. Fl. Ind. Bat. 3 (1859) 571; Naves Novis. App. (1880) 258; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 276; Merr. in Philip. Journ. Sci. 5 (1910) Bot. 339; Prain & Burkill in Leaf. Philip. Bot. 5 (1913) 1597, Journ. As. Soc. Beng. II 10 (1914) 28.

Luzon (Bontoc, Lepanto, Ilocos Norte, Union, Nueva Vizcaya, Bulacan, Rizal, Batangas, Laguna, Tayabas), *Cuming* 779, *Elmer* 18290, 18466, *Merrill Sp. Blancoanae* 117, *Phil. Pl.* 1070, *F. B.* 24909 Mabesa, 13218 Curran, 13336 Tamesis, 3293 Ahern's collector, *B. S.* 13734, 37240, 21752, 1380 Ramos. In thickets at low and medium altitudes. Endemic.

Local names: Aribúbu (Ilk.); kamángeg (Ilk.); kirini (Tag.); mayatbáng (Tag.); pakít (Tag.); ubág (Tag.).

DIOSCOREA MERRILLII Prain & Burkill in Leaf. Philip. Bot. 5 (1913) 1598, Journ. As. Soc. Beng. II 10 (1914) 35.

Mindoro, Mindanao (Davao), *Merrill* 5657, *Elmer* 11924. In old clearings and in thickets, altitude about 700 m. Endemic.

DIOSCOREA NUMMULARIA Lam. Encycl. 3 (1789) 231; Naves Novis. App. (1880) 258; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 276; Prain & Burkill in Leaf. Philip. Bot. 5 (1913) 1599, Journ. As. Soc. Beng. II 10 (1914) 35.

Luzon (Bontoc to Sorsogon, in most or all provinces), Polillo, Leyte, Camiguin de Misamis, Mindanao, Basilan. In thickets and forests at low and medium altitudes.

Var. **GLAUCA** Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 35.

Luzon (Benguet), *Elmer* 5638. Scarcely distinguishable from the type. The species in Celebes, Moluccas, New Guinea to Prince of Wales Island.

Local names: Banán (Bag.); malabalukbúk-dágis (Pamp.); pakít (Tag.); pinót (Tag.); úbing-básol (Tag.).

DIOSCOREA PENTAPHYLLA Linn. Sp. Pl. (1753) 1032; Blanco Fl. Filip. (1837) 802, ed. 2 (1845) 552, ed. 3, 3 (1879) 210; Miq. Fl. Ind. Bat. 3 (1859) 574; Naves Novis. App. (1880) 259; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 35, 3 (1908) Bot. 401, Fl. Manila (1912) 152, Sp. Blancoanae (1918) 103; Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 23.

Luzon (Rizal, Bataan, Laguna), Panay, *Merrill Sp. Blancoanae* 458, *F. B.* 1971 Ahern's collector, *B. S.* 12176 Ramos. In thickets at low and medium altitudes. India to Yunnan, southward through Malaya to New Guinea.

Local names: Kasi (Ig.); lima-limá (Tag.) sapáng (Bis.).

The Philippine form is apparently referable to var. *malaica* Prain & Burkill l. c.

DIOSCOREA PEPEROIDES Prain & Burkill in Elm. Leaf. Philip. Bot. 5 (1913) 1597, Journ. As. Soc. Beng. II 10 (1914) 28 (var. *vera* Prain & Burkill).

Luzon (Benguet, Rizal), Golo, Palawan, *Elmer* 6399, 6400, *Merrill* 9441, 6512, 11552, 11555, *Loher* 7007, 7010. In thickets up to 1,300 m altitude, in Palawan about limestone formations at sea level.

Var. **SAGITTIFOLIA** Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 28.

Luzon (Nueva Vizcaya), *B. S.* 8178 *Ramos*. In thickets at low or medium altitudes. The species and this variety endemic, another variety in Indo-China.

DIOSCOREA SOROR Prain & Burkill in Leaf. Philip. Bot. 5 (1913) 1598, Journ. As. Soc. Beng. II 10 (1914) 34 (var. *vera* Prain & Burkill).

Luzon (Lepanto, Bataan), *Elmer* 6737, *Bona*, *B. S.* 5941 *Ramos*. In thickets at low and medium altitudes. Endemic. Probably the same as *D. divaricata* Blanco.

Var. **GLAUCA** Prain & Burkill in Journ. As. Soc. Beng. II 10 (1914) 34.

Luzon (Bulacan), *Yoder*. In thickets at low altitudes. Endemic.

EXCLUDED SPECIES

DIOSCOREA EBURNEA Lour.; Naves Novis. App. (1880) 260.

DIOSCOREA GLABRA Roxb.; Naves op. cit. 258.

DIOSCOREA JAPONICA Thunb.; Naves op. cit. 259.

DIOSCOREA OPPOSITIFOLIA Linn.; Naves op. cit. 258.

DIOSCOREA PYRIFOLIA Kunth Enum. 5 (1850) 384; Miq. Fl. Ind. Bat. 3 (1859) 571; Naves Novis. App. (1880) 258.

This record was based on *Cuming* 2314 as Philippine, but this number was from Malacca; the species does not occur in the Philippines.

DIOSCOREA SATIVA Linn.; Naves op. cit. 259.

DIOSCOREA SPICULATA Blume; Naves l. c.

DIOSCOREA VULGARIS Miq.; Naves l. c.

2. STENOMERIS Planchon

STENOMERIS CUMINGIANA Becc. in Nuov. Giorn. Bot. Ital. 2 (1870) 8, t. 2; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 70; Rolfe in Journ. Bot. 23 (1885) 215, Rev. Pl. Vasc. Filip. (1886) 278; Taubert in Engl. Bot. Jahrb. 15 (1893) Beibl. 38: 2.

Luzon (Laguna, Sorsogon), Mindoro, Sibuyan, Leyte, Mindanao (Zamboanga), *Cuming* 1739 in herb. Webb., *Wenzel* 1210, 962, 718, *Elmer* 12064, *F. B.* 26909 *Mabesa*, *B. S.* 39429 *Ramos*. In forests at low and medium altitudes. Endemic.

STENOMERIS DIOSCOREAEFOLIA Planch. in Ann. Sci. Nat. Bot. III 18 (1852) 320; Walp. Ann. 6 (1860) 42; Becc. in Nuov. Giorn. Bot. Ital. 2 (1870) 8; Scheff. in Nat. Tijdschr. Nederl. Ind. 34 (1874) 70; Vidal Phan. Cuming. Philip. (1885) 153, Rev. Pl. Vasc. Filip. (1886) 276; Rolfe in Journ. Bot. 23 (1885) 216; Taubert in Engl. Bot. Jahrb. 15 (1893) Beibl. 38: 2; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 268.

Luzon (Rizal, Laguna, Tayabas, Albay, Sorsogon), Catanduanes, Mindoro, Leyte, Biliran, Dinagat, Mindanao, *Cuming 375, Merrill 5775, 8301, Phil. Pl. 427, Elmer 9098, Baker 3383, B. S. 1005, 30339, 23514 Ramos, 35192 Ramos & Pascasio*. In thickets, old clearings, etc., at low and medium altitudes. Endemic.

STENOMERIS WALLISII Taubert in Engl. Bot. Jahrb. 15 (1893) Beibl. 38: 2.

Luzon, *Wallis!* in herb. Berol. Endemic.

The genus has one other known species in Borneo and the Malay Peninsula.

EXCLUDED GENUS

RAJANA sp.; Naves Novis. App. (1880) 260.

IRIDACEAE

1. BELAMCANDA Adanson

BELAMCANDA CHINENSIS (Linn.) DC. in Red. Lil. 3 (1802) t. 121; Usteri Beitr. Ken. Philip. Veg. (1905) 129; C. B. Rob. in Philip. Journ. Sci. 6 (1911) Bot. 196; Merr. Fl. Manila (1912) 152.

Ixia chinensis Linn. Sp. Pl. (1753) 36.

Pardanthus chinensis Ker in König & Sims Ann. Bot. 1 (1805) 247;

Naves Novis. App. (1880) 252, Fl. Filip. ed. 3 (1877-83) t. 376.

Planted for ornamental purposes but nowhere naturalized. A native of southeastern Asia, now cultivated in most warm countries.

Local name: Abanico (Sp.).

2. PATERSONIA R. Brown

PATERSONIA LOWII Stapf in Trans. Linn. Soc. Bot. 4 (1894) 241, t. 20, f. 7-9; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 268.

Mindoro, *Merrill 5507*. In an open heath on Mt. Halcon, altitude about 2,400 m. Borneo. The genus a typical Australian one.

3. ELEUTHERINE Herbert

ELEUTHERINE PALMIFOLIA (Linn.) Merr. in Philip. Journ. Sci. 7 (1912) Bot. 233, Fl. Manila (1912) 153, Sp. Blancoanae (1918) 104.

Sisyrinchium palmifolium Linn. Mant. 1 (1767) 122; Naves Novis. App. (1880) 252.

Eleutherine plicata Herb. in Bot. Reg. 29 (1843) sub t. 57.

Antholyza meriana Blanco Fl. Filip. (1837) 24, ed. 2 (1845) 18, ed. 3, 1 (1877) 33, t. 100; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 85, non Linn.

A native of tropical America, introduced and now naturalized in some parts of the Philippines, occurring in hemp plantations and, occasionally, in waste places about towns, etc.; also planted for ornamental purposes.

Local names: Ahos-áhos (C. Bis.); hagusáhis (S. L. Bis.).

EXCLUDED GENERA AND SPECIES

ANTHOLYZA AETHIOPICA Linn.; Naves Novis. App. (1880) 252.

IRIS SUSIANA Linn.; Naves op. cit. 253.

GLADIOLUS SEGETUM Ker; Naves l. c.

WATSONIA MERIANA Ker; Naves l. c.

MUSACEAE

1. MUSA Linnaeus

MUSA ERRANS (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 390, t. 17, f. 6-8; Merr. Sp. Blancoanae (1918) 105.

Musa troglodytarum Linn. var. *errans* Blanco Fl. Filip. (1837) 247, ed. 2 (1845) 172, ed. 3, 1 (1877) 310.

Musa amboinensis Naves Novis. App. (1880) 218, non Rumph.

Luzon (Apayao, Bulacan, Bataan, Rizal, Laguna), *Merrill Sp. Blancoanae* 873, Teodoro, B. S. 28268 *Fénix*. In forests at low and medium altitudes. Endemic.

Local names: Pau-uá (Ibn.); ságing-matsíng (Tag.); ságing-na-ligau (Tag.).

Var. **BOTOAN** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 391, t. 7, f. 5-10; Merr. Sp. Blancoanae (1918) 106.

Musa troglodytarum Blanco Fl. Filip. (1837) 246, ed. 2 (1845) 172, ed. 3, 1 (1877) 310, t. 89, non Linn.

Luzon (Rizal), *Merrill Sp. Blancoanae* 217, Teodoro. Planted in most parts of the Philippines; really referable to *M. errans* as a variety?

Local names: Baláiang (Ilk.); butúan (Tag.); butúhan (Tag.); vul-lūngán (Ibn., It.).

MUSA GLAUCA Roxb. Pl. Coromandel 3 (1819) t. 300; Naves Novis. App. (1880) 218; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 387; Merr. Sp. Blancoanae (1918) 104.

Musa troglodytarum Linn. var. *dolioformis* Blanco Fl. Filip. (1837) 855, ed. 2 (1845) 174, ed. 1 (1877) 312.

Luzon (Cavite, Laguna), *Merrill Sp. Blancoanae* 946, 537. In ravines, etc., at low and medium altitudes. Burma, Pegu.

Local name: Virgen (Sp.).

MUSA PARADISIACA Linn. Sp. Pl. (1753) 1043; Miq. Fl. Ind. Bat. 3 (1859) 588; Naves Novis. App. (1880) 215; Merr. Fl. Manila (1912) 154; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 412; K. Schum. in Engl. Pflanzenreich 1 (1900) 19, f. 3.

The plantains are those bananas which are edible after being cooked and very inferior or inedible raw. A number of forms or varieties occur in the Philippines, which from a species standpoint are scarcely to be distinguished from *Musa sapientum* Linn., the bananas. Following custom the forms are distributed between the two Linnean binomials. It is by no means certain that the numerous varieties which have been proposed by Blanco, Teodoro, and Quisumbing are correctly distributed as between *Musa paradisiaca* Linn. and *M. sapientum*.

I am of the opinion that no more weight should be placed on the numerous forms of the banana than on similar forms of other exceedingly variable cultivated plants such as the orange, the apple, rice, the common bean, sugar cane, etc., because relatively speaking the differences in cultivated bananas are certainly no greater than those in the cultivated forms of the other plants mentioned above. No attempt has been made to correlate these Philippine forms or varieties with extra-Philippine ones although some, perhaps many, of our cultivated bananas are certainly identical with

those of tropical Asia and Malaya. Quisumbing, Phil. Agr. Review 12³ (1919) 1-73 t. 1-30, described and figured a number of varieties from extra-Philippine plants, cultivated at Los Baños, which have not been included in this enumeration. They are *Musa sapientum* Linn. var. *gladiata* Quis., origin unknown; var. *oanga* Quis., from New Guinea; var. *klum* Quis., from India; var. *misi* Quis., from Samoa; var. *saloor* Quis., from Borneo; var. *chuntara* Quis., from Siam; var. *pop* Quis., from Borneo; var. *maori* Quis., from Raratonga; var. *sicsec* Quis., from New Guinea; var. *languma* Quis., from New Guinea; var. *radja* Quis., from Java (= *Musa regia* Rumph.!); var. *nam* Quis., apparently from Indo-China; var. *yanakonban* Quis., from India; var. *rachidis* Quis., from the Malay Peninsula; var. *cochinchinensis* Quis., from Indo-China (= *Musa odorata* Lour.!); var. *katali* Quis., from India; var. *khai* Quis., from Siam; var. *krie* Quis., origin unknown; var. *yan* Quis., origin unknown; var. *hazara* Quis., from India; var. *maduranga* Quis., from India; *Musa paradisiaca* Linn. var. *enosa* Quis., from Java (= *Musa corniculata* Rumph.!); and var. *tiparot* Quis., origin unknown.

Var. **MAGNA** Blanco Fl. Filip. (1837) 244, ed. 2 (1845) 171, ed. 3, 1 (1877) 307; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 413; Merr. Sp. Blancoanae (1918) 108; Quis. in Philip. Agr. Review 12³ (1919) 67, t. 12, f. 212, 213.

Planted here and there for its fruit. Endemic?

Local name: Tundók (Tag.).

Var. **MAXIMA** Blanco Fl. Filip. (1837) 245, ed. 2 (1845) 171, ed. 3, 1 (1877) 308; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 414; Merr. Sp. Blancoanae (1918) 109.

Planted here and there for its fruit, *Merrill Sp. Blancoanae* 920. Endemic?

Local names: Batábia (Tag.); matábia (Tag.).

Var. **SUBRUBEA** Blanco Fl. Filip. (1837) 245, ed. 2 (1845) 171, ed. 3, 1 (1877) 307; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 414, t. 16, f. 1-5; Merr. Sp. Blancoanae (1918) 109.

Planted here and there. Endemic?

Teodoro's figure and description were apparently based on material of Cuban origin and hence may well not represent the form Blanco described.

MUSA SAPIENTUM Linn. Syst. ed. 10 (1759) 1303; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 393.

Musa paradisiaca Linn. subsp. *sapientum* O. Kuntze Rev. Gen. Pl. 1 (1892) 692; K. Schum. in Engl. Pflanzenreich 1 (1900) 20.

The bananas, distinguishable from the plantains (*Musa paradisiaca*) chiefly in the consistency and characters of the mature fruits. A number of forms are found in the Philippines in cultivation, many in all probability introduced from other parts of Malaya and from tropical Asia.

Var. **ANGAO** Quis. in Philip. Agr. Review 12³ (1919) 58, t. 5, f. 30; t. 7, f. 69; t. 9, f. 126; t. 11, f. 192, 193; t. 26, f. 2.

Mindanao (Misamis). Cultivated at Los Baños.

Local name: Añgau (C. Bis.).

Var. **BACA** Quis. in Philip. Agr. Review 12³ (1919) 51, t. 5, f. 23; t. 7, f. 60; t. 9, f. 118; t. 11, f. 176, 177.

Luzon (Cavite). Cultivated at Los Baños.

Local name: Súngai-báka (Tag.).

Var. **BINUTIG** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 401, t. 10, f. 1-5.

Teodoro's specimens were apparently from Leyte, according to material prepared by him from plants cultivated at Los Baños.

Local name: Binutig (S. L. Bis.).

Var. **CANARA** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 406, t. 10, f. 6-10.

Cultivated at Los Baños, origin unknown but probably Philippine, and very likely from Camarines Province, where *kanára* is a well-known banana.

Var. **CANAYA** Quis. in Philip. Agr. Review 12³ (1919) 44, t. 4, f. 18; t. 6, f. 54; t. 9, f. 111; t. 10, f. 162, 163; t. 20, f. 2.

Mindanao (Surigao). Cultivated at Los Baños.

Local name: Kanáya (C. Bis.).

Var. **CINEREA** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 397, t. 13, f. 1-5; Merr. Sp. Blancoanae (1918) 106.

Musa paradisiaca Linn. var. *cinerea* Blanco Fl. Filip. (1837) 250, ed. 2 (1845) 175, ed. 3, 1 (1877) 313.

One of the commonest bananas in the Philippines, *Merrill Sp. Blancoanae* 926.

Local names: Latúndan (Tag.); letóndal (Tag.); tórdan (Tag.).

Var. **COMPRESSA** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 408, t. 7, f. 1-5; Merr. Sp. Blancoanae (1918) 108.

Musa paradisiaca Linn. var. *compressa* Blanco Fl. Filip. (1837) 240, ed. 2 (1845) 168, ed. 3, 1 (1877) 304.

A very common banana in most parts of the Philippines.

Local names: Bisco (Tag.); obispo (Tag.); sabá (Tag.).

This is probably better placed under *M. paradisiaca* Linn.

Var. **DINALAGA** Quis. in Philip. Agr. Review 12³ (1919) 55, t. 5, f. 31; t. 7, 66; t. 9, f. 122; t. 11, f. 184, 185; t. 26, f. 1.

Luzon (Tayabas). Cultivated at Los Baños.

Local name: Dinalága (Tag.).

Var. **DOOL** Quis. in Philip. Agr. Review 12³ (1919) 35, t. 4, f. 6; t. 6, f. 44; t. 9, f. 101; t. 10, f. 142, 143; t. f. 15, f. 1.

Mindanao (Bukidnon). Cultivated at Los Baños.

Local name: Duol (Buk.).

Var. **EDA** Quis. in Philip. Agr. Review 12³ (1919) 37, t. 4, f. 9; t. 6, f. 47; t. 9, f. 104; t. 10, f. 148, 149; t. 15, f. 1.

Mindanao (Surigao). Cultivated at Los Baños.

Local name: Eda-án (C. Bis.).

- Var. **FLABELLATA** Quis. in Philip. Agr. Review 12³ (1919) 66, t. 5, f. 35; t. 8, f. 78, 87; t. 12, f. 208, 209; t. 29, f. 2.
Luzon (Laguna, Bulacan). Cultivated at Los Baños.
Local names: Binendito (Tag.); inabaníko (Tag.).
Perhaps better placed under *M. paradisiaca* Linn.
- Var. **GALATAYAN** Quis. in Philip. Agr. Review. 12³ (1919) 40, t. 4, f. 13; t. 6, f. 50; t. 9, f. 108; t. 10, f. 154, 155; t. 18, f. 1.
Luzon (Cavite). Cultivated at Los Baños.
Local name: Galatáyan (Tag.).
- Var. **DARYAO** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 403, t. 8, f. 1-5.
Leyte, *Teodoro*, cultivated at Los Baños.
Local name: Daryau (S. L. Bis.).
- Var. **GARANGAO** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 401, t. 10, f. 6-10.
Leyte, *Teodoro*, cultivated at Los Baños.
Local name: Garañgau (S. L. Bis.).
- Var. **GLABERRIMA** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 399, t. 15, f. 1-5; Merr. Sp. Blancoanae (1918) 106.
Musa paradisiaca Linn. var. *glaberrima* Blanco Fl. Filip. (1837) 245, ed. 2 (1845) 171, ed. 3, 1 (1877) 308.
Luzon (Rizal, Laguna), *Merrill Sp. Blancoanae* 865, *Teodoro*.
Local names: Dinuguan (Tag.); durugó (Tag.); galámai-señora (Tag.).
- Var. **GLAUCA** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 402, t. 11, f. 6-10; Merr. Sp. Blancoanae (1918) 107.
Musa paradisiaca Linn. var. *glauca* Blanco Fl. Filip. (1837) 250, ed. 2 (1845) 175, ed. 3, 1 (1877) 312.
Luzon (Rizal), *Merrill Sp. Blancoanae* 912.
Local names: Bingti-kohol (Tag.); tinalong (Tag.); veinte-kohol (Tag.).
- Var. **GRANDIS** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 410, t. 14, f. 1-3.
Luzon (Laguna). Occasionally planted.
Local name: Sabáng-ilóko (Tag.).
- Var. **HUMILIS** (Perr.).
Musa humilis Perr. in Mém. Soc. Linn. Paris 3 (1824) 131; C. B. Rob. in Philip. Journ. Sci. 3 (1908) 306; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 393, t. 18, f. 1-5.
Mindanao (Zamboanga, Cotabato). A peculiar form with small obovoid fruits.
Local name: Pula-pulá (Mag.).
- Var. **KINAMAY** Quis. in Philip. Agr. Review 12³ (1919) 57, t. 5, f. 32; t. 7, f. 67; t. 9, f. 125; t. 11, f. 190, 191; t. 23, f. 1.
Luzon (Bulacan). Cultivated at Los Baños.
Local name: Kinamáí (Tag.).

Var. **INARNIBAL** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 406, t. 9, f. 1-5.

Luzon (Laguna), *Teodoro*.

Local name: Inarnibal (Tag.).

Var. **LACATAN** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 405, t. 11, f. 1-5; Merr. Sp. Blancoanae (1918) 107.

Musa paradisiaca Linn. var. *lacatan* Blanco Fl. Filip. (1837) 243, ed. 2 (1845) 170, ed. 3, 1 (1877) 305, t. 88.

A widely cultivated banana and one of the most popular varieties in the Philippines.

Local names: Bañgáran (Bik.); lakatán (Tag., S. L. Bis.).

Var. **LONGA** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 407; Merr. Sp. Blancoanae (1918) 108.

Musa paradisiaca Linn. var. *longa* Blanco Fl. Filip. (1837) 245, ed. 2 (1845) 171, ed. 3, 1 (1877) 308.

Known only from Blanco's description.

Local name: Kinauáyan (Tag.).

Var. **PUMILA** (Blanco).

Musa paradisiaca Linn. var. *pumila* Blanco Fl. Filip. (1837) 244, ed. 2 (1845) 171, ed. 3, 1 (1877) 306.

Musa cavendishii Lamb. var. *pumila* Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 412.

Musa sapientum Linn. var.; Merr. Sp. Blancoanae (1918) 108.

Occasionally planted; perhaps a dwarfed form of the *bunġulan*.

Local name: Tampóhin (Tag.).

Var. **PADILAT** Quis. in Philip. Agr. Review 12³ (1919) 63, t. 5, f. 38; t. 7, f. 71; t. 8, f. 88; t. 12, f. 202, 203; t. 28, f. 1.

Babuyan Islands (Dalupiri). Cultivated at Los Baños.

Local name: Padilat (Ilk.).

Var. **PAMOTION** Quis. in Philip. Agr. Review 12³ (1919) 49, t. 5, f. 21; t. 7, f. 59; t. 9, f. 116; t. 11, f. 172, 173; t. 22, f. 1.

Mindanao (Surigao). Cultivated at Los Baños.

Local name: Pamotíon (C. Bis.).

Var. **PELIPIA** Quis. in Philip. Agr. Review 12³ (1919) 42, t. 4, f. 14; t. 6, f. 51; t. 9, f. 109; t. 10, f. 158, 159; t. 18, f. 2.

Mindanao (Surigao). Cultivated at Los Baños.

Local name: Pelípiá (C. Bis.).

Var. **PRINCIPE** Quis. in Philip. Agr. Review 12³ (1919) 50, t. 5, f. 22; t. 7, f. 61; t. 9, f. 117; t. 11, f. 174, 175; t. 22, f. 2.

Philippines, without locality. Cultivated at Los Baños.

Var. **PUTIAN** Quis. in Philip. Agr. Review 12³ (1919) 36, t. 4, f. 8; t. 6, f. 46; t. 9, f. 103; t. 10, f. 146, 147; t. 15, f. 2.

Cultivated at Los Baños, from the Visayan Islands.

Local name: Putían (Bis.).

- Var. **RAINES** Quis. in Philip. Agr. Review 12³ (1919) 55, t. 5, f. 27; t. 7, f. 65; t. 9, f. 124; t. 11, f. 186, 187; t. 25, f. 1.
 Polillo. Cultivated at Los Baños.
 Local name: Raines-na-putí, (Tag.).
- Var. **SAROCSOC** Quis. in Philip. Agr. Review 12³ (1919) 39, t. 4, f. 11; t. 6, f. 49; t. 9, f. 104; t. 10, f. 152, 153; t. 17, f. 1.
 Luzon (Camarines). Cultivated at Los Baños.
 Local names: Saroksók (Bik.); sinaroksók (Bik.).
- Var. **SISION** Quis. in Philip. Agr. Review 12³ (1919) 35, t. 4, f. 7; t. 6, f. 45; t. 9, f. 102; t. 10, f. 144, 145.
 Panay (Capiz). Cultivated at Los Baños.
 Local name: Sisíon (P. Bis.).
- Var. **SUAVEOLENS** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 400, t. 14, f. 6-10; Merr. Sp. Blancoanae (1918) 106.
Musa paradisiaca Linn. var. *suaveolens* Blanco Fl. Filip. (1837) 244, ed. 2 (1845) 171, ed. 3, 1 (1877) 306.
 One of the most popular bananas in the Philippines, *Merrill Sp. Blancoanae* 928.
 Local name: Buñgúlan (Tag.).
- Var. **TERNATENSIS** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 404, t. 7, f. 1-5; Merr. Sp. Blancoanae (1918) 107.
Musa paradisiaca Linn. var. *suaveolens* Blanco Fl. Filip. (1837) 243, ed. 2 (1845) 170, ed. 3, 1 (1877) 305.
 Occasionally cultivated, *Merrill Sp. Blancoanae* 216.
 Local names: Glória (Tag., Ilk., Ibn.); ternáte (Tag.).
- Var. **TOMBAK** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 407; Merr. Sp. Blancoanae (1918) 108; Quis. in Philip. Agr. Review 12³ (1919) 33, t. 4, f. 4; t. 6, f. 42; t. 9, f. 99; t. 10, f. 138, 139; t. 14, f. 2.
Musa paradisiaca Linn. var. *tombak* Blanco Fl. Filip. (1837) 246, ed. 2 (1845) 171, 3, 1 (1877) 307.
 Cultivated here and there, *Merrill Sp. Blancoanae* 953.
 Local names: Goyóran (Tag.); tinumbága (Tag.); tómbak (Tag.).
- Var. **TULDOC** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 407, t. 10, f. 1-5.
 Luzon (Rizal). Cultivated by Teodoro at Los Baños.
 Local name: Tuldók (Tag.).
- Var. **TUDLONG** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 402, t. 11, f. 6-10.
 Leyte, cultivated by Teodoro at Los Baños.
 Local name: Tudlong (S. L. Bis.).
- Var. **VIOLACEA** (Blanco) Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 398, t. 5, f. 6-10; Merr. Sp. Blancoanae (1918) 106.
Musa paradisiaca Linn. var. *violacea* Blanco Fl. Filip. (1837) 245, ed. 2 (1845) 171, ed. 3, 1 (1877) 307.
 Cultivated here and there.
 Local name: Morádo (Sp.).

MUSA TEXTILIS Née in Anal. Cienc. Nat. 4 (1801) 123; Hook. Lond. Journ. Bot. 7 (1848) 268; Baker in Ann. Bot. 7 (1893) 211; K. Schum. in Engl. Pflanzenreich 1 (1900) 19; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 199; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 388, t. 18, f. 6-10; Merr. Sp. Blancoanae (1918) 105.

Musa abaca Perr. in Mém. Soc. Linn. Paris 3 (1824) 130; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 305.

Musa troglodytarum Linn. var. *textoria* Blanco Fl. Filip. (1837) 247, ed. 2 (1845) 173, ed. 3, 1 (1877) 311.

Musa mindanaensis Rumph.; Miq. Fl. Ind. Bat. 3 (1859) 588; Naves Novis. App. (1880) 218.

Abacá or Manila hemp of commerce. Very extensively cultivated in those provinces from central Luzon to Mindanao where the dry season is not pronounced. In Luzon the chief abacá provinces are Camarines, Albay, and Sorsogon. The islands of Samar and Leyte are very important producing centers, as are also parts of Mindanao, especially Davao. Endemic.

Local name: Abaká (most dialects). The species is variable, and sundry forms have distinctive local names.

MUSA URANOSCOPOS Lour. Fl. Cochinch. (1790) 645, excl. syn. Rumph.

Musa coccinea Andr. Bot. Rep. t. 47; Naves Novis. App. (1880) 218; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 199; Teodoro op. cit. 10 (1915) Bot. 388.

Planted here and there for ornamental purposes on account of its very showy red bracts. A native of southeastern Asia.

DOUBTFUL AND EXCLUDED SPECIES

MUSA ALPHURICA Rumph.; Naves Novis. App. (1880) 217.

MUSA CORNICULATA Rumph.; Naves l. c.

MUSA ORNATA Roxb.; Naves op. cit. 218.

MUSA SIMIARUM Rumph.; Naves op. cit. 217.

MUSA URANOSCOPOS Rumph.; Naves op. cit. 218.

MUSA CHAPARA Perr. in Mém. Soc. Linn. Paris 3 (1825) 131; C. B. Rob. in Philip. Journ. Sci. 3 (1908) Bot. 305; Teodoro op. cit. 10 (1915) Bot. 415.

Unrecognizable except that it is a form or variety of *Musa sapientum* Linn.

MUSA ENSETE Gmel.; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 387.

This occurs on the grounds of the College of Agriculture at Los Baños as a recently introduced and cultivated species and has no claims to be considered as Philippine.

MUSA NIGRA Perr. in Mém. Soc. Linn. Paris 3 (1824) 131; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 415.

Unrecognizable except that it is a form or variety of *Musa sapientum* Linn.

MUSA PARADISIACA Linn. var. *ULNARIS* Blanco Fl. Filip. (1837) 246, ed. 2 (1845) 172, ed. 3, 1 (1877) 309; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 414; Merr. Sp. Blancoanae (1918) 107.

A purely imaginary variety, the description based on hearsay data, probably inspired by one of the *tundók* varieties.

MUSA CAVENDISHII Lamb.; Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 410, incl. var. *hawaiiensis* Teodoro l. c. t. 15, f. 6-10.

This is generally considered as a distinct species, but I believe it to be merely a variety of *Musa sapientum* Linn. Teodoro's descriptions were based on recently introduced and cultivated specimens.

MUSA SAPIENTUM Linn. var. **CUBENSIS** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 397, t. 11, f. 6-10.

The description was based on material originating in Cuba.

MUSA SAPIENTUM Linn. var. **AMERICANA** Teodoro in Philip. Journ. Sci. 10 (1915) Bot. 398, t. 7, f. 6-10.

This variety was based on material of Cuban origin.

2. RAVENALA Adanson

RAVENALA MADAGASCARIENSIS Gmel. Syst. (1791) 567; Naves Novis. App. (1880) 219; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 199; Merr. Fl. Manila (1912) 155; K. Schum. in Engl. Pflanzenreich 1 (1900) 29, f. 6.

Planted in Manila and in other large towns for ornamental purposes, a native of Madagascar. The so-called traveler's palm.

3. HELICONIA Linnaeus

HELICONIA BIHAI Linn. Mant. 2 (1771) 211; Naves Novis. App. (1880) 215; K. Schum. in Engl. Pflanzenreich. 1 (1900) 36.

Heliconiopsis amboinensis Miq. Fl. Ind. Bat. 3 (1859) 590; Naves l. c.

Planted occasionally for ornamental purposes. A native of tropical America, naturalized in some parts of the Old World.

EXCLUDED GENUS

STRELITZIA REGINAE Banks Naves Novis. App. (1880) 219.

ZINGIBERACEAE

1. ZINGIBER Adanson

ZINGIBER APOENSE Elm. Leaf. Philip. Bot. 8 (1915) 2914.

Mindanao (Davao), *Elmer* 10545, 11750. In forests, 1,000 to 1,300 m. Endemic.

Local name: Bogóbog (Bag.).

ZINGIBER BREVIFOLIUM N. E. Br. in Gard. Chron. 26 (1886) 390, non K. Schum. (1904).

Described from cultivated specimens said to have originated in the Philippines. Otherwise unknown. Endemic.

ZINGIBER BULUSANENSE Elm. Leaf. Philip. Bot. 8 (1919) 2993.

Luzon (Sorsogon), *Elmer* 16914, 15178. In primary forests, altitude about 450 m. Endemic.

ZINGIBER CASSUMUNAR Roxb. in *Asiat. Research* 11 (1810) 347, *t. 5*, *Fl. Ind.* 1 (1820) 49; Vidal Phan. Cuming. Philip. (1885) 152, *Rev. Pl. Vasc. Filip.* (1886) 274; Naves Novis. App. (1880) 220; Ridl. in *Govt. Lab. Publ. (Philip.)* 35 (1906) 84.

Mindanao (Misamis), (*Cuming 1605*). I have seen no Philippine material representing this species. Pantropic in cultivation, a native of tropical Asia.

ZINGIBER MOLLE Ridl. in *Leaf. Philip. Bot.* 2 (1909) 607, *nomen nudum*, *Philip. Journ. Sci.* 4 (1909) Bot. 168.

Luzon (Bontoc, Lepanto, Benguet, Pampanga, Tayabas), *F. B.* 4886, 4914 *Curran*, 14490 *Darling*, *B. S.* 5364 *Ramos*, 1991, 1985 *Foxworthy*, *Merrill* 4843. In ravines along streams and in forests, altitude 1,100 to 1,600 m. Endemic.

Local name: Kaláuing (Bon.).

ZINGIBER NEGROSENSE Elm. *Leaf. Philip. Bot.* 8 (1915) 2917.

Negros, Mindanao (Davao), *Elmer* 10421, 11768. In forested ravines, altitude about 1,050 m. Endemic.

ZINGIBER OFFICINALE Rosc. in *Trans. Linn. Soc.* 7 (1807) 348, *Monandr.*

Pl. (1828) *t. 88*; *Miq. Fl. Ind. Bat.* 3 (1859) 593; Naves Novis. App. (1880) 220; K. Schum. in *Engl. Pflanzenreich* 20 (1904) 170, *f. 23*; *Merr. Fl. Manila* (1912) 159, *Sp. Blancoanae* (1918) 110.

Amomum zingiber Linn. *Sp. Pl.* (1753) 1; *Blanco Fl. Filip.* (1837) 2, ed. 2 (1845) 2, ed. 3, 1 (1877) 3, *t. 131*.

Zingiber blancoi Hassk. in *Flora* 47 (1864) 20.

Widely cultivated although not on an extensive scale, and nowhere naturalized. Pantropic in cultivation, a native of tropical Asia. Ginger.

Local names: Agát (Pang.); baseng (Ilk.); gengibre (Sp.); laíal (Sbl.); laíya (If.); layá (Ilk., Bon., Ibn., It.); lúya (Tag.).

ZINGIBER PUBISQUAMA Ridl. in *Philip. Journ. Sci.* 4 (1909) Bot. 169.

Mindanao (Lanao), *Clemens* 1163a. In open grasslands, cogon (*Imperata*) type; altitude about 650 m. Endemic.

ZINGIBER SYLVATICUM Elm. *Leaf. Philip. Bot.* 8 (1915) 2919.

Luzon (Cagayan, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Catanduanes, Negros, *Elmer* 9843, 18004, 14992, 14936, *B. S.* 19456, 30521, 22048 *Ramos*, 13076 *Foxworthy*, 10237, 10238 *McGregor*, 9198 *Robinson*. In forests chiefly at medium altitudes, ascending to 1,300 m. Endemic.

ZINGIBER ZERUMBET (Linn.) Sm. *Exot. Bot.* 2 (1804) 103, *t. 112*;

Miq. Fl. Ind. Bat. 3 (1859) 593; Naves Novis. App. (1880) 220; K. Schum. in *Engl. Pflanzenreich* 20 (1904) 172, *f. 24*; Ridl. in *Philip. Journ. Sci.* 4 (1909) Bot. 199; *Merr. Fl. Manila* (1912) 159, *Sp. Blancoanae* (1918) 111.

Amomum zerumbet Linn. *Sp. Pl.* (1753) 1; *Blanco Fl. Filip.* (1837) 2, ed. 2 (1845) 2, ed. 3, 1 (1877) 3, *t. 370*.

Zingiber cassumunar Naves in *Blanco Fl. Filip.* ed. 3 (1877-83), *t. 370*, non Roxb.

Zingiber zerumbet Sm. var. *magnum* Elm. *Leaf. Philip. Bot.* 8 (1919) 2922.

Northern Luzon to Mindanao, in most or all islands and provinces. In thickets in and near settlements at low and medium altitudes. A native of tropical Asia, now pantropic.

Local names: Balauág (Bik.); barik (Tag.); lampúyang (Buk.); langkauás (Tag.); layág sa-súlug (Bik.); luiang-ósiu (Tag.); tamohílang (Buk.); tumbóng-áso (Tag.).

EXCLUDED SPECIES

ZINGIBER AMERICANS Blume; Naves Novis. App. (1880) 221.

ZINGIBER AQUOSUM Blume; Naves l. c.

ZINGIBER ELATUM Roxb.; Naves l. c.

ZINGIBER GRACILE Jack; Naves l. c.

ZINGIBER GRAMINEUM Blume; Naves l. c.

ZINGIBER INFLEXUM Blume; Naves l. c.

ZINGIBER MARGINATUM Roxb.; Naves l. c.

ZINGIBER ODORIFERUM Blume; Naves l. c.

ZINGIBER OVOIDEUM Blume; Naves op. cit. 220.

ZINGIBER PURPUREUM Rosc.; Naves l. c.

ZINGIBER ROSEUM Roxb.; Naves op. cit. 221.

ZINGIBER SQUARROSUM Roxb.; Naves l. c.

2. LANGUAS Koenig

(*Alpinia* auctt., non Linnaeus*)

LANGUAS APOENSIS (Elm.) comb. nov.

Alpinia apoensis Elm. Leaff. Philip. Bot. 8 (1915) 2888.

Catanduanes, Mindanao (Davao), *Elmer 11889*, *B. S. 30490 Ramos*. In primary forests up to an altitude of 1,200 m. Endemic.

Local name: Otin-ótin (Bag.).

LANGUAS BRACHYANTHA (Merr.) comb. nov.

Alpinia brachyantha Merr. in Philip. Journ. Sci. 10 (1915) Bot. 296.

Luzon (Cagayan), *Weber 1571*, *B. S. 13920 Ramos*, *F. B. 19617 Curran*. In forests at low or medium altitudes. Endemic.

LANGUAS BREVILABRIS (Presl) comb. nov.

Alpinia brevibras Presl Rel. Haenk. 1 (1827) 110, *t. 17*; Naves Novis. App. (1880) 225; Ridl. in Philip. Journ. Sci. 4 (1919) Bot. 190; K. Schum. in Engl. Bot. Jahrb. 27 (1899) 276, *Pflanzenreich* 20 (1904) 314.

Alpinia fraseriana Usteri Beitr. Ken. Philip. Veg. (1905) 128, non Oliv.

* The genus *Alpinia* of Linnaeus was based wholly on a tropical American plant, *Alpinia racemosa* Linn. = *Renealmia racemosa* A. Rich., and is thus identical with *Renealmia* Linn. f. (1781), which it should replace. As between the two generic names, no matter which one be applied to the American group, *Alpinia* is clearly not applicable to the Indo-Malayan genus long known under this name.

Batan Islands and northern Luzon to Mindanao. In most islands and provinces. Common in primary forests at low and medium altitudes.

Local names: Ápat (Ig.); bagongbóng (Bis.); bánai (Pamp.); gagad-no-kóran (Iv.); katkátan (Pamp.); katólan (Bis.); katótang (Bis.); luyalúya (Tag.); malatalbák (Tag.); panáuil (Bon.); parapád (Ig.); tagbák (Bik., Tag., C. Bis.); tagbák-gúbat (Tag.); tagbák-matsíng (Tag.); tagbás (C. Bis.); talbák (Tag.); tugís (S. L. Bis.).

LANGUAS COPELANDII (Ridl.) comb. nov.

Alpinia copelandii Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 164.

Mindanao (Zamboanga), *Copeland*. In forests at low altitudes. Endemic.

LANGUAS CUMINGII (K. Schum.) comb. nov.

Alpinia cumingii K. Schum. in Engl. Pflanzenreich 20 (1904) 315;

Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 187.

Alpinia chinensis Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl.

Vasc. Filip. (1886) 274, non Rosc.

Luzon (Albay), *Cuming* 859. Of this I have seen only a few detached flowers. Endemic.

LANGUAS DIVERSIFOLIA (Elm.) comb. nov.

Alpinia diversifolia Elm. Leaf. Philip. Bot. 8 (1915) 2919.

Vanoverberghia diversifolia Elm. op. cit. 2913.

Luzon (Benguet), *Elmer* 8853. In damp ravines, altitude about 1,000 m. Endemic.

LANGUAS FLABELLATA (Ridl.) comb. nov.

Alpinia flabellata Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 188.

Batan Islands and northern Luzon to Mindanao, *Merrill Phil. Pl. 831, B. S. 31885 Santos, 2718, 2871 Mearns*. Not always easily distinguishable from *A. brevilabris* Presl. In forests at medium altitudes ascending to 2,000 m. Endemic.

Var. **MAJOR** (Ridl.) comb. nov.

Alpinia flabellata Ridl. var. *major* Ridl. l. c.

Range and habitat of the species and doubtfully distinct even as a variety.

Local names: Biláu (Sub.); tagbák-gúbat (Bik.); tagbák-laláki (Tag.); tuká-máya (Tag.).

LANGUAS FOXWORTHYI (Ridl.) comb. nov.

Alpinia foxworthyi Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 189;

Elm. Leaf. Philip. Bot. 8 (1915) 2887.

Palawan, *B. S. 772 Foxworthy, Elmer 12930*. In forests at low altitudes. Endemic.

LANGUAS GLABRESCENS (Ridl.) comb. nov.

Alpinia glabrescens Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 185.

Batan Islands (Batan, Sabtan), *B. S. 3679 Fénix, 3141 Mearns, 10156 McGregor*. On slopes at low altitudes. Endemic unless identical with some Formosan species represented by *Faurie* 975 of his 1913 collection.

Local name: Gagád (Iv.).

LANGUAS GRAMINEA (Ridl.) comb. nov.

Alpinia graminea Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 189; Elm. Leaf. Philip. Bot. 8 (1919) 2966.

Luzon (Cagayan, Nueva Ecija, Laguna, Tayabas, Sorsogon), Catanduanes, Negros, Panay, *F. B. 7267, 11208 Everett, B. S. 26359, 30678 Ramos & Edaño, 25411 Yates, 28375 Fénix, 30239 Ramos, Elmer 16228*. In forests up to 800 m altitude. Endemic.

Perhaps but a narrow-leaved form of *L. brevibrabis* (Presl) Merr.

Local names: Barañgin (Ibn.); lampíngit (Neg.); tagbák (P. Bis.).

LANGUAS HAENKEI (Presl) comb. nov.

Alpinia haenkei Presl Symb. 1 (1832) 66, t. 43; K. Schum. in Engl. Pflanzenreich 20 (1904) 341; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 86, Philip. Journ. Sci. 4 (1909) Bot. 184.

Alpinia malaccensis Presl Rel. Haenk. 1 (1827) 110; Naves Novis. App. (1880) 225; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 275, non Rosc.

Alpinia philippinensis Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 86; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 37.

Batan Islands and northern Luzon to Palawan and Mindanao, in most or all islands and provinces. In primary forests at low and medium altitudes, often common, represented by over 60 individual collections. Endemic.

Local names: Bagumbúng (Tag.); barápat (Ig.); bírao-bírao (Sul.); púnan (C. Bis.); sakiápág (Neg.); simiónan (Buk.); tagbák-babáe (Tag.); tagbák-laláki (Tag.); tagusáhis (P. Bis.) talbák (Tag.); tamotámo (Sbl.); tukáng-máya (Tag.); kalaueg (Ig.).

LANGUAS ILLUSTRIS (Ridl.) comb. nov.

Alpinia illustris Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 185.

Mindoro, Palawan, *B. S. 514 Foxworthy, 39485 Ramos*. In forests along stream at low altitudes; very similar to *A. malaccensis* Rosc. Endemic.

LANGUAS LONGIPETIOLATA (Elm.) comb. nov.

Alpinia longipetiolata Elm. Leaf. Philip. Bot. 8 (1919) 2969.

Luzon (Sorsogon), *Elmer 16167, 16954, B. S. 23441 Ramos*. In forests up to 600 m altitude. Endemic.

LANGUAS MACROSCAPHIS (K. Schum.) comb. nov.

Alpinia macroscaphis K. Schum. in Engl. Bot. Jahrb. 27 (1899) 284, Pflanzenreich 20 (1904) 337, f. 40, B. C.

Mindanao (prob. Davao), (*Warburg*). The type is preserved in the Berlin herbarium. I have seen no specimens. Endemic.

LANGUAS MOLLIS (Presl) comb. nov.

Alpinia mollis Presl Rel. Haenk. 1 (1827) 110; Naves Novis. App. (1880) 225; K. Schum. in Engl. Pflanzenreich 20 (1904) 368; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 191.

"Hab. in insula Luzon." The type was collected by Haenke. It is known only from the original description and is a species of doubtful status. Endemic.

LANGUAS MUSAEFOLIA (Ridl.) comb. nov.

Alpinia musaefolia Ridl. in Leaf. Philip. Bot. 2 (1909) 604, Philip. Journ. Sci. 4 (1909) Bot. 186.

Negros, *Elmer 9539*. In forests, altitude about 1,200 m. Endemic.
Local name: Lunhás (C. Bis.).

LANGUAS PENDULIFLORA (Ridl.) comb. nov.

Alpinia penduliflora Ridl. in Leaf. Philip. Bot. 2 (1909) 570, Philip. Journ. Sci. 4 (1909) Bot. 187.

Leyte, *Elmer 7288*. Probably from forests at low altitudes. Endemic.

LANGUAS PUBIFLORA (Benth.) comb. nov.

Hellenia pubiflora Benth. in Hook. Lond. Journ. Bot. 2 (1843) 235.

Alpinia pubiflora K. Schum. in Engl. Pflanzenreich 20 (1904) 313, f. 39, F, G; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 188; Elm. Leaf. Philip. Bot. 8 (1919) 2965.

Luzon (Tayabas, Sorsogon), Guimaras, Camiguin de Misamis, Mindanao, *B. S. 34699 Ramos & Pascasio, 14728 Ramos, F. B. 9100 Whitford & Hutchison, 34 Ritchie, Elmer 10456, 16546, Copeland 319, Piper 606*. In forests at low and medium altitudes. New Guinea, Caroline Islands.

Local names: Tagbág (Sul.); tagbák (P. Bis.).

LANGUAS PULCHELLA (K. Schum.) comb. nov.

Alpinia pulchella K. Schum. in Engl. Bot. Jahrb. 27 (1899) 276, t. 2, f. E, Pflanzenreich 20 (1904) 315; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 189.

Mindanao (Davao), (*Warburg*). I have seen no material representing this species. New Guinea.

LANGUAS PYRAMIDATA (Blume) comb. nov.

Alpinia pyramidata Blume Enum. Pl. Jav. 1 (1827) 58; Naves Novis. App. (1880) 225; Ridl. in Elm. Leaf. Philip. Bot. 2 (1909) 604, Philip. Journ. Sci. 4 (1909) Bot. 186; Elm. Leaf. Philip. Bot. 8 (1919) 2967.

Alpinia galanga Naves Novis. App. (1880) 225, vix Sw.

Alpinia galanga Sw. var. *pyramidata* K. Schum. in Engl. Pflanzenreich 20 (1904) 316.

Luzon (Rizal, Laguna, Camarines, Sorsogon), Panay, Leyte, Mindanao (Lanao, Agusan), Palawan, *Elmer 16352, Wenzel 1407, Clemens 612, B. S. 21556 Escritor, 23865 Fénix, 35279 Martelino & Edaño*. In old clearings, etc., sometimes planted. Himalayan region through Malaya to the Moluccas.

Local names: Langkauás (Tag., Bis., Bik.); lankauás (Tag., Pamp., Bis., Mbo.); pal-la (Mand.).

LANGUAS ROMBLONENSIS (Elm.) comb. nov.

Alpinia romblonensis Elm. Leaf. Philip. Bot. 8 (1915) 2889.

Romblon, Samar, *Elmer 12050, B. S. 21238 Escritor* (from Romblon, not from Panay as labeled), *17548 Ramos*. Along streams at low altitudes. Endemic.

Local name: Tagbák (P. Bis.).

LANGUAS ROSEA (Elm.) comb. nov.

Alpinia rosea Elm. Leaf. Philip. Bot. 8 (1915) 2890.

Sibuyan, *Elmer 12509*. In dense forests, altitude about 1,350 m. Endemic.

LANGUAS RUFA (Presl) comb. nov.

Hellenia rufa Presl Rel. Haenk. 1 (1827) 114, t. 21; Miq. Fl. Ind. Bat. 3 (1859) 604.

Alpinia rufa Naves Novis. App. (1880) 226; K. Schum. in Engl. Bot. Jahrb. 27 (1899) 293, Pflanzenreich 20 (1904) 361; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 190, Leaf. Philip. Bot. 2 (1909) 604, 8 (1915) 2887.

Luzon (Laguna, Camarines, Tayabas, Sorsogon), Polillo, Catanduanes, Samar, Leyte, Panay, Negros, Bucas, Mindanao, Basilan. In forests at low and medium altitudes. Endemic.

Local names: Bihai (C. Bis.); kaukos (Bag.); tagbág (Mbo.); tagbák (Tag.).

LANGUAS SIBUYANENSIS (Elm.) comb. nov.

Alpinia sibuyanensis Elm. Leaf. Philip. Bot. 8 (1915) 2891.

Sibuyan, *Elmer* 12316. In forests along streams, altitude about 225 m. Endemic.

Local name: Bobonan (P. Bis.).

LANGUAS SPECIOSA (Wendl.) comb. nov.

Zerumbet speciosum Wendl. Sert. Hann. (1798) t. 19.

Alpinia nutans Rosc. in Sm. Exot. Bot. 2 (1805) t. 106; Naves Novis. App. (1880) 226.

Alpinia cernua Naves Novis. App. (1880) 226, Fl. Filip. ed. 3 (1877-83) t. 464, non Sims.

Alpinia speciosa K. Schum. Fl. Kaiser-Wilhelmsl. (1887) 29, Engl. Bot. Jahrb. 27 (1899) 284, t. 3, f. A, B, Pflanzenreich 20 (1904) 339, f. 40 D-F.

Occasionally cultivated for ornamental purposes, but not a native of the Archipelago and nowhere naturalized. China, Japan; cultivated in other warm countries.

LANGUAS SUBFUSICARPA (Elm.) comb. nov.

Alpinia subfuscarpa Elm. Leaf. Philip. Bot. 8 (1915) 2892.

Mindanao (Davao), *Elmer* 10506. In damp primary forests, altitude about 1,200 m. Endemic.

Local name: Bosadak (Bag.).

LANGUAS TRACHYASCUS (K. Schum.) comb. nov.

Alpinia trachyascus K. Schum. in Engl. Pflanzenreich 20 (1904) 336; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 185.

Mindanao (Davao), *Warburg* 14514 in herb. Berol. I have seen only the type which has the facies of *Alpinia haenkei* Presl. Endemic.

LANGUAS VANOVERBERGHII (Merr.) comb. nov.

Alpinia vanoverberghii Merr. in Philip. Journ. Sci. 7 (1912) Bot. 75.

Luzon (Bontoc), *Vanoverbergh* 573. On slopes, altitude about 1,350 m. Endemic.

Local names: Kalaúin (Bon.); kalaúing (Bon.); takáng (Bon.).

LANGUAS VULCANICA (Elm.) comb. nov.

Alpinia vulcanica Elm. Leaf. Philip. Bot. 8 (1919) 2971.

Luzon (Camarines, Sorsogon), Bucas Grande, *Elmer* 16168, B. S. 33864 Ramos & Edaña, 35051 Ramos & Pascasio, 23428 Ramos. In primary forests at low and medium altitudes, ascending to 800 m. Endemic.

LANGUAS WENZELII (Merr.) comb. nov.

Alpinia wenzelii Merr. in Philip. Journ. Sci. 9 (1914) Bot. 353.

Leyte, Wenzel 623. In forests, altitude about 500 m. Endemic.

EXCLUDED SPECIES

ALPINIA ARCTIFLORA F.-Muell.; Naves Novis. App. (1880) 227.

ALPINIA CAPITELLATA Jack; Naves op. cit. 226.

ALPINIA CHINENSIS Rosc.; Naves op. cit. 225.

ALPINIA COERULEA Benth.; Naves op. cit. 227.

ALPINIA ELATIOR Jack; Naves op. cit. 226.

ALPINIA GIGANTEA Blume; Naves op. cit. 225.

ALPINIA JAVANICA Blume; Naves op. cit. 226.

ALPINIA MUTICA Roxb.; Naves l. c.

ALPINIA RUBRA (Miq.) Naves l. c.

ALPINIA SCABRA (Blume) Naves l. c.

ALPINIA SPICATA Roxb.; Naves l. c.

ALPINIA STACHYOIDES Hance; Naves l. c.

3. KOLOWRATIA Presl**KOLOWRATIA CONGESTA** (Elm.) comb. nov.

Alpinia congesta Elm. Leaf. Philip. Bot. 8 (1919) 2968.

Luzon (Sorsogon), Elmer 16144. On forested slopes, altitude about 450 m. Endemic.

KOLOWRATIA ELEGANS Presl Rel. Haenk. 1 (1827) 113, t. 20; Miq. Fl. Ind. Bat. 3 (1859) 610; Naves Novis. App. (1880) 228; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 86, Philip. Journ. Sci. 4 (1909) Bot. 181; Merr. Fl. Manila (1912) 159, Sp. Blancoanae (1918) 110.

Renealmia gracilis Blanco Fl. Filip. (1837) 1.

Renealmia exaltata Blanco op. cit. ed. 2 (1845) 1, ed. 3, 1 (1877) 2, t. 1, non Linn.

Hellenia gracilis Hassk. in Flora 47 (1864) 19.

Alpinia gracilis Rolfe in Journ. Linn. Soc. Bot. 21 (1884) 316; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 275.

Alpinia elegans K. Schum. in Engl. Bot. Jahrb. 27 (1899) 288, Pflanzenreich 20 (1904) 352, f. 41 C; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 84, Philip. Journ. Sci. 1 (1906) Suppl. 37.

Luzon (Apayao, Amburayan, Lepanto, Nueva Vizcaya, Pampanga, Bulacan, Nueva Ecija, Bataan, Rizal, Laguna, Tayabas, Sorsogon), Polillo, Mindoro, Leyte. In thickets along streams at low and medium altitudes; represented by numerous collections. Endemic.

Local names: Bagombón (Tag.); katkátan (Bis.); katótang (Bis.); salbák (Tag.); tagbák (Tag.); tugbák (Tag.).

KOLOWRATIA ERUCAEFORMIS Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 182.

Bucas, Merrill 5258. On dry slopes at the borders of forests at low altitudes. Endemic.

The genus is confined to the Philippines.

4. LEPTOSOLENA Presl

LEPTOSOLENA HAENKEI Presl Rel. Haenk. 1 (1827) 111, t. 28; Naves Novis. App. (1880) 227; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 181.

Alpinia leptosolenia K. Schum. in Engl. Pflanzenreich 20 (1904) 312.

Leptosolenia insignis Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 84, Philip. Journ. Sci. 4 (1909) 181; Elm. Leaf. Philip. Bot. 8 (1915) 2907.

Luzon (Ilocos Norte, Bontoc, Benguet, Union, Nueva Vizcaya, B. S. 8211, 8172, 32863 Ramos, 21093 Escritor, 37396 Ramos & Edaño, Elmer 6428, Lete 84, Vanoverbergh 384. In thickets and on open slopes at low and medium altitudes. A monotypic endemic genus.

Local names: Bánai (Ilk.); kaláuin (Bon.); panáuil (Bon.).

5. ADELMERIA Ridley

(*Elmeria* Ridley, non Rydberg)

ADELMERIA ALPINA Elm. Leaf. Philip. Bot. 8 (1915) 2885.

Mindanao (Davao), Elmer 10534, 10642. In forests, altitude 1,200 to 2,100 m. Endemic.

Local names: Lurasising (Bag.); pusadók (Bag.).

ADELMERIA GIGANTIFOLIA (Elm.) Elm. Leaf. Philip. Bot. 8 (1919) 2963.

Zingiber gigantifolium Elm. op. cit. 8 (1915) 2916.

Luzon (Tayabas, Sorsogon), Elmer 9282, 14603. In forests on talus slopes and along small streams, altitude 500 to 700. Endemic.

ADELMERIA OBLONGA Merr. in Philip. Journ. Sci. 9 (1914) Bot. 443.

Adelmeria albida Elm. Leaf. Philip. Bot. 8 (1919) 2964.

Luzon (Bontoc, Benguet, Sorsogon), Panay, Negros, Elmer 6255, 15912, Vanoverbergh 3108, Eskridge, B. S. 30884 Ramos & Edaño. In forests at low and medium altitudes, ascending to 1,500 m. Endemic.

ADELMERIA PARADOXA (Ridl.) Merr. in Philip. Journ. Sci. 9 (1914) Bot. 444.

Hornstedtia paradoxa Ridl. in Govt. Lab. Publ. (Philip.) 35 (1905) 85.

Elmeria bifida Ridl. in Leaf. Philip. Bot. 2 (1909) 568.

Adelmeria bifida Ridl. op. cit. 603; Philip. Journ. Sci. 4 (1909) Bot. 179.

Luzon (Apayao, Bontoc, Benguet, Nueva Ecija, Pangasinan, Rizal, Laguna, Tayabas), Panay, Mindanao (Lanao), Elmer 6629, 18145, 8568, 7706, Merrill 4786, 7517, Loher 7001, B. S. 29652, 26458, 30675, 38032 Ramos & Edaño, F. B. 15854 Merritt. In the mossy forest and in damp forested ravines, altitude 1,000 to 1,800 m. Endemic.

Local name: Parápat (Ig.)

ADELMERIA PINETORUM (Ridl.) Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 180.

Elmeria pinetorum Ridl. in Leaf. Philip. Bot. 2 (1909) 569.

Luzon (Bontoc, Benguet), *Elmer 8548, Vanoverbergh 453*. In thickets and on open slopes about limestone boulders, altitude 1,300 to 1,500 m. Endemic.

Local names: Alauín (Bon.); barápat (Ig.).

6. PLAGIOSTACHYS Ridley

PLAGIOSTACHYS CORRUGATA Elm. Leaf. Philip. Bot. 8 (1915) 2908.

Mindanao (Davao), *Elmer 11130*. In forests, altitude about 450 m. Endemic.

The flowers are unknown; probably no *Plagiostachys*.

PLAGIOSTACHYS ELEGANS Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 178.

Mindanao (Lanao), *Clemens*. In forests, altitude about 900 m. Endemic.

PLAGIOSTACHYS ESCRITORII Elm. Leaf. Philip. Bot. 8 (1915) 2910, (1919) 2989.

Luzon (Laguna, Sorsogon), Samar, Mindanao, *Elmer 10516, 10544, 17501, 18071, 16216, 18477a, 16062, B. S. 15919 Fénix, 24242 Ramos, 21504 Escritor, 37290, 37262 Ramos & Edaño*. In forests at medium altitudes, ascending to 1,300 m. Endemic.

Local names: Bosadak (Bag.); pupundia (Sub.).

PLAGIOSTACHYS PARVIFLORA (Presl) Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 178.

Amomum parviflorum Presl Rel. Haenk. 1 (1827) 112, t. 19; Naves Novis. App. (1880) 224.

Alpinia parviflora Rolfe in Journ. Linn. Soc. Bot. 21 (1884) 316 quoad syn. Presl, Vidal Phan. Cuming. Philip. (1885) 152, Rev Pl. Vasc. Filip. (1886) 275, p. p.; K. Schum. in Engl. Bot. Jahrb. 27 (1899) 299, Engl. Pflanzenreich 20 (1904) 368.

Luzon (Sorsogon, Leyte, Panay, Mindanao (Agusan), *Elmer 16215, Wenzel 305, Merrill 7318, B. S. 31256 Ramos*. In primary forests at low altitudes. Endemic.

PLAGIOSTACHYS PHILIPPINENSIS Ridl. in Leaf. Philip. Bot. 2 (1909)

571, Philip. Journ. Sci. 4 (1909) Bot. 177; C. B. Rob. op. cit. 6 (1911) Bot. 197; Elm. Leaf. Philip. Bot. 9 (1919) 2989.

Plagiostachys ridleyi Elm. Leaf. Philip. Bot. 8 (1915) 2911, (1919) 2989.

Luzon (Laguna, Tayabas, Camarines), Polillo, Catanduanes, Sibuyan, Panay, Samar, Mindanao, *B. S. 33614, 31245, 31239, 29272 Ramos & Edaño, 24157 Ramos, 11253 McGregor, Baker 3457, Clemens 855, Elmer 7915, Williams 2402, Elmer 12209*. In primary forests at low and medium altitudes. Endemic.

Local name: Tagbák (S. L. Bis., Mbo.).

PLAGIOSTACHYS ROLFEI (K. Schum.) Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 179.

Alpinia rolfei K. Schum. in Engl. Pflanzenreich 20 (1904) 368.

Alpinia parviflora Rolfe in Journ. Linn. Soc. Bot. 21 (1884) 316, p. p. quoad Cuming 1327.

Luzon (Cagayan, not Albay!), *Cuming 1327*. Not seen. Endemic.

7. *VANOVERBERGHIA* Merrill

VANOVERBERGHIA SEPULCHREI Merr. in Philip. Journ. Sci. 7 (1912) Bot. 76; Elm. Leaf. Philip. Bot. 8 (1915) 2913, (1919) 2990.

Luzon (Ifugao, Bontoc, Benguet, Sorsogon), Panay, *Vanoverbergh* 956, *Sandkuhl* 257, *Elmer* 8560, 17383, *F. B.* 947 *Barnes*, *B. S.* 12913 *Fénix*, 30734, 30134 *Ramos & Edaño*. In ravines along streams at medium altitudes, ascending to 1,500 m. A monotypic endemic genus.

Local names: Agbáb (Bon.); barápat (Ig.); chakchákil (Ig.).

8. *AMOMUM* Linnaeus

AMOMUM BULUSANENSE Elm. Leaf. Philip. Bot. 8 (1919) 2973.

Luzon (Sorsogon), *Elmer* 16395. In primary forests, altitude about 600 m. Endemic.

AMOMUM CONOIDEUM (Ridl.) Elm. Leaf. Philip. Bot. 8 (1919) 2972.

Hornstedtia conoidea Ridl. in Leaf. Philip. Bot. 2 (1909) 605, Philip. Journ. Sci. 4 (1909) Bot. 175.

Luzon (Laguna, Sorsogon), Negros, Mindanao (Davao, Lanao), *Elmer* 10246, 16924, 17896, 11131, *Clemens*, *Barrett*. In damp forested ravines at medium altitudes. Endemic.

Valeton, in Bull. Jard. Bot. Buitenz. III 3 (1921) 174, 176, cites *Elmer* 10246, the type of this species, as possibly representing *Hornstedtia alliacea* Valeton and as *H. deliana* Valeton, the former definitely known from Java and Sumatra, the latter from Sumatra. Ridley's specific name is the oldest.

AMOMUM DALICAN (Elm.) comb. nov.

Hornstedtia dalican Elm. Leaf. Philip. Bot. 8 (1915) 2906.

Mindanao (Davao), *Elmer* 11626, 11759. Along small streams in forests, altitude about 1,100 m. Endemic.

Local name: Dalikán (Bag.).

AMOMUM DEUTERAMOMUM K. Schum. in Engl. Bot. Jahrb. 27 (1899) 313, *Pflanzenreich* 20 (1904) 237; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 173.

Mindanao (Davao), (*Warburg* 15735). In forests. Not seen. Endemic.

AMOMUM ELEGANS Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 84, Philip. Journ. Sci. 4 (1909) Bot. 172.

Luzon (Bataan), Mindanao (Lanao), *Whitford* 300, 207, *Clemens* 1061, *F. B.* 3033 *Borden*. In primary forests at low and medium altitudes. Endemic.

Local names: Luya-luyáhan (Tag.); tagbák-gúbat (Tag.).

AMOMUM FUSIFORME Ridl. in Leaf. Philip. Bot. 2 (1909) 604, *nomen nudum*, Philip. Journ. Sci. 4 (1909) Bot. 171.

Negros, Samar, Mindanao (Zamboanga, Surigao), *Bolster* 331, 224, *Elmer* 9509, *B. S.* 24159 *Ramos*, 36690, 37001 *Ramos & Edaño*. In primary forests at low and medium altitudes. Endemic.

Local names: Loya-lóya (C. Bis.); mali-máli (Sub.); satagbák (C. Bis.).

AMOMUM IROSINENSIS (Elm.) comb. nov.*Hornstedtia irosinensis* Elm. Leaf. Philip. Bot. 8 (1919) 2981.*Hornstedtia peninsula* Elm. op. cit. 2983.

Luzon (Sorsogon), *Elmer* 16332, 16263, 16455. In forested ravines, altitude 350 to 500 m. Endemic.

AMOMUM LEPICARPUM Ridl. in Leaf. Philip. Bot. 2 (1909) 605, Philip.

Journ. Sci. 4 (1909) Bot. 174.

Hornstedtia lepicarpa Elm. Leaf. Philip. Bot. 8 (1919) 2979.*Hornstedtia subviridis* Elm. op. cit. 2987.

Luzon (Laguna, Sorsogon), Negros, Mindanao (Davao), *Elmer* 10044, 17176, 17869, 10835, 16437, 17788. In forests at low and medium altitudes. Endemic.

AMOMUM LINEARIFOLIUM Elm. Leaf. Philip. Bot. 8 (1919) 2975.

Luzon (Sorsogon), *Elmer* 16926. In thickets bordering forests, altitude about 300 m. Endemic.

AMOMUM LOHERI K. Schum. in Engl. Pflanzenreich 20 (1904) 247;

Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 171.

Luzon, without locality, *Loher* 675. I saw Loher's specimen at Kew in 1907 and at that time made a note that it was very similar to *A. trilobum* Ridl. Endemic.

AMOMUM LOPHOPHORUM (Ridl.) Elm. Leaf. Philip. Bot. 8 (1919) 2973.*Hornstedtia lophophora* Ridl. in Leaf. Philip. Bot. 2 (1909) 607,

Philip. Journ. Sci. 4 (1909) Bot. 176.

Negros, Mindanao (Davao), *Elmer* 10365, 10519. In forested ravines, altitude about 1,200 m. Endemic.

AMOMUM LUZONENSE Elm. Leaf. Philip. Bot. 8 (1919) 2976.

Luzon (Sorsogon), Biliran, (*Elmer* 15645), *B. S.* 18696 *McGregor*, det. *Elmer*. In secondary forests at low or medium altitudes. Endemic.

AMOMUM MICROCHEILA (Ridl.) comb. nov.*Hornstedtia microcheila* Ridl. in Leaf. Philip. Bot. 2 (1909) 606,

Philip. Journ. Sci. 4 (1909) Bot. 176.

Negros, *Elmer* 10279. On seepage banks along streams, altitude about 900 m. Endemic.

AMOMUM MINDANAENSE Elm. Leaf. Philip. Bot. 8 (1915) 2894.

Mindanao (Davao), *Elmer* 10822. In primary forests, altitude about 1,100 m. Endemic.

Local name: Tanaulak (Bag.).

AMOMUM MURICARPUM Elm. Leaf. Philip. Bot. 8 (1915) 2896.*Hornstedtia muricarpa* Elm. op. cit. 8 (1919) 2980.

Mindanao (Cotabato, Zamboanga, Davao), *Elmer* 10947, *B. S.* 15846 *Fénix*, 11555 *Robinson*, 36603 *Ramos & Edaña*. In primary forests at low and medium altitudes. Endemic.

Local names: Lalañgod (Sub.); panikabán (Bag.); túgis (Buk.).

AMOMUM PALAWANENSE Elm. Leaf. Philip. Bot. 8 (1915) 2897.

Palawan, *Elmer* 12795. In forests along streams at low altitudes. Very similar to *A. mindanaense* Elm. Endemic.

AMOMUM PANDANICARPUM Elm. Leaf. Philip. Bot. 8 (1915) 2899.

Hornstedtia pandanicarpa Elm. op. cit. 8 (1919) 2979.

Hornstedtia sorsogonensis Elm. op. cit. 2985.

Luzon (Sorsogon), Panay, Mindanao (Bukidnon, Davao), *Elmer* 10508, 16223, 16755, *B. S.* 31184, 38758 *Ramos & Edaño*. In primary forests at low and medium altitudes, ascending to 1,100 m. Endemic.

Local name: Dalikán (Bag.).

AMOMUM PHILIPPINENSE (Ridl.) comb. nov.

Hornstedtia philippinensis Ridl. in Govt. Lab. Publ. (Philip.) 35

(1906) 86 p. p., Leaf. Philip. Bot. 2 (1909) 605, Philip. Journ.

Sci. 4 (1909) Bot. 175; Elm. Leaf. Philip. Bot. 8 (1915) 2905,

(1919) 2981.

Luzon (Laguna, Tayabas, Sorsogon), Mindoro, Palawan, Biliran, Negros, Panay, Leyte, Mindanao, *Elmer* 9281, 9280, 11973, 16713, 17903, 7285, *Merrill* 9762, *Williams* 2888, *B. S.* 804 *Foxworthy*, 34554 *Ramos & Pascasio*, 31123 *Ramos & Edaño*, *F. B.* 3737 *Merritt*. In forests at low and medium altitudes. Endemic.

The fruit described by Ridley does not belong to this species; see *Elmer*, Leaf. Philip. Bot. 8 (1919) 2980, for the true fruit characters.

Local names: Dalikán (Mbo.); lulutúkoan (Sub.); parápad (Ig.); tagbák (P. Bis.); tumbák (Neg.).

AMOMUM PROPINQUUM Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906)

84, Philip. Journ. Sci. 4 (1909) Bot. 172; Elm. Leaf. Philip. Bot.

8 (1915) 2894; *Merr. Sp. Blancoanae* (1918) 111.

Amomum echinatum Blanco Fl. Filip. ed. 2 (1845) 3, ed. 3, 1 (1877) 4, non Willd.

Amomum aculeatum Naves Novis. App. (1880) 224, non Roxb.

Hornstedtia propinqua Elm. Leaf. Philip. Bot. 8 (1919) 2979.

Luzon (Benguet, Batangas, Laguna, Tayabas, Sorsogon), Masbate, Biliran, Mindanao (Davao), *Elmer* 6284, 15900, 15614, 18456, *Merrill Sp. Blancoanae* 925, *F. B.* 1704 *Clark*. In forests at low and medium altitudes. Endemic.

Local name: Túgis (S. L. Bis.).

AMOMUM PUBESCENS (Ridl.) comb. nov.

Amomum lepicarpum Ridl. var. *pubescens* Ridl. in Leaf. Philip. Bot. 2 (1909) 604.

Hornstedtia pubescens Elm. op. cit. 8 (1919) 2979.

Negros, *Elmer* 10384. In forests, altitude about 1,100 m. Endemic.

AMOMUM PUBIMARGINATUM Elm. Leaf. Philip. Bot. 8 (1915) 2900.

Mindanao (Davao), *Elmer* 10546. In primary forests, altitude about 1,300 m. Endemic.

Local name: Tabák-bukalod (Bag.).

AMOMUM PURPUREUM (Elm.) comb. nov.

Hornstedtia purpurea Elm. Leaf. Philip. Bot. 9 (1919) 2984.

Luzon (Sorsogon), *Elmer* 16819, 17023. In densely forested ravines, altitude about 450 m. Endemic.

AMOMUM TRILOBUM Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 85, Philip. Journ. Sci. 4 (1909) Bot. 173.

Luzon (Ilocos Norte, Bontoc, Bulacan, Pampanga, Laguna, Rizal), Palawan, Panay, *Bolster* 67, *Elmer* 17566, *B. S.* 4602, 4655 *Ramos*, 817 *Foxworthy*, *F. B.* 12698 *Rosenbluth & Tamesis*. In primary forests at low and medium altitudes. Endemic.

Local name: Tagbák (Tag.).

AMOMUM WARBURGII K. Schum. in Engl. Pflanzenreich 20 (1904) 257; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 171.

Costus warburgii K. Schum. in Engl. Bot. Jahrb. 27 (1899) 346.

Mindanao (probably Davao), *Warburg*. The type, a mere scrap, has distinctly the appearance of *Costus*; to be compared with *Costus clemensiae* Ridl. A species of very indefinite status. Endemic.

EXCLUDED SPECIES

AMOMUM CARDAMOMUM Linn.; Naves Novis. App. (1880) 223.

AMOMUM CILIATUM Blume; Naves l. c.

AMOMUM DEALBATUM Roxb.; Naves l. c.

AMOMUM GRACILE Blume; Naves l. c.

AMOMUM HATUANUM (Rumph.) Naves op. cit. 224.

AMOMUM HETERANTHUM Blume; Naves op. cit. 223.

AMOMUM MAXIMUM Roxb.; Naves op. cit. 224.

AMOMUM VILLOSUM Lour.; Naves op. cit. 223.

9. PHAEOMERIA Lindley

PHAEOMERIA SPECIOSA (Blume) comb. nov.

Elettaria speciosa Blume Enum. Pl. Jav. (1827) 51.

Alpinia magnifica Rose. Monandr. Pl. (1828) t. 75.

Phaeomeria magnifica K. Schum. in Engl. Pflanzenreich 20 (1904) 262.

Phaeomeria imperialis Lindl. Nat. Syst. ed. 2 (1836) 446; Ridl. in

Philip. Journ. Sci. 4 (1909) Bot. 177; Elm. Leaf. Philip. Bot. 8 (1915) 2907.

Mindanao (Davao, Bukidnon), *Elmer* 10652, *Williams* 2943, *Wester* 49. In ravines, probably introduced. Malay Peninsula, Java, Celebes.

Local name: Tikála (Buk.).

10. HEDYCHIUM Koenig

HEDYCHIUM CORONARIUM Koenig in Retz. Obs. 3 (1783) 73; Miq. Fl.

Ind. Bat. 3 (1859) 608; Naves Novis. App. (1880) 227; Vidal Phan.

Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 274;

K. Schum. in Engl. Pflanzenreich 20 (1904) 44, f. 8, A, C; Ridl.

in Philip. Journ. Sci. 4 (1909) Bot. 164; Merr. Fl. Manila (1912)

157; Elm. Leaf. Philip. Bot. 8 (1919) 2978.

Hedychium lingulatum Hassk. Pl. Jav. Rar. (1848) 135; Naves Novis.

App. (1880) 227.

Introduced and cultivated for ornamental purposes, in some regions in the southern Philippines naturalized. In Mindanao certainly of prehistoric introduction, in Luzon of recent introduction. A native of India, now pantropic, cultivated and naturalized.

Local names: Bánai (Bis.); dansúli (Buk.); kámia (Tag., Bik., Sp., C. Bis.); katkátan (Bis.); katotang (Bis.).

HEDYCHIUM PHILIPPINENSE K. Schum. in Engl. Pflanzenreich 20 (1904) 47; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 164; Elm. Leaf. Philip. Bot. 8 (1915) 2903.

Hedychium mindanaense Elm. Leaf. Philip. Bot. 8 (1915) 2904, (1919) 2978.

Luzon (Rizal, Laguna, Tayabas, Sorsogon), Polillo, Negros, Leyte, Mindanao, Jolo, *Elmer* 10673, 7909, 15968, *Loher* 6996, *Clemens* 747, *B. S.* 22897 *McGregor*, 35015 *Ramos & Pascasio*, 27947 *Ocampo*, 20777 *Escritor*, 15206, 10962, 10003 *Ramos*, *F. B.* 4252 *Everett*. In damp forests at low and medium altitudes, ascending to 1,100 m; often epiphytic or pseudoeiphytic. Endemic.

Local names: Dainsúli (Sub.); kaláuag (Bag.); tagbák (P. Bis.).

EXCLUDED SPECIES

HEDYCHIUM HASSELTII Blume; Naves Novis. App. (1880) 227.

11. KAEMPFERIA Linnaeus

KAEMPFERIA GALANGA Linn. Sp. Pl. (1753) 2; Naves Novis. App. (1880) 222; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 165; K. Schum. in Engl. Pflanzenreich 20 (1904) 77; Merr. Fl. Manila (1912) 157, Sp. Blancoanae (1918) 110.

Kaempferia rotunda Blanco Fl. Filip. (1837) 5, ed. 2 (1845) 4, ed. 3, 1 (1877) 7; Naves Novis. App. (1880) 222; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 84, non Linn.

Kaempferia latifolia Hornem. Hassk. in Flora 47 (1864) 22, quoad syn. Blanco.

Luzon (Bontoc, Rizal), Mindanao (Bukidnon), *Merrill Sp. Blancoanae* 942, *B. S.* 26075 *Fénix*, 4560 *Ramos*. In open grasslands at low and medium altitudes. India through Malaya to the Moluccas.

Local names: Disól (Ilk.); dosó (Bon.); dosól (Sbl.); dotó (Bon.); dusó (Tag.); dusóg (Tag.); dusól (Tag.); gisól (Tag., Pamp.); kisól (Buk., Bis.); kosól (Bis.); kusól (Pamp.).

KAEMPFERIA PHILIPPINENSIS Merr. in Philip. Journ. Sci. 10 (1915) Bot. 296.

Luzon (Laguna), *B. S.* 14952 *Ramos*. In forests along streams at low or medium altitudes. Endemic.

EXCLUDED SPECIES

KAEMPFERIA LATIFOLIA Don; Naves Novis. App. (1880) 223.

KAEMPFERIA PANDURATA Roxb.; Naves l. c.

12. **BOESENBERGIA** O. Kuntze*(Gastrochilus* Wallich, non Don)**BOESENBERGIA LONGIPETIOLATA** (Ridl.) comb. nov.*Gastrochilus longipetiolatus* Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 166.Mindanao (Zamboanga), *F. B. 9110 Whitford & Hutchinson, B. S. 36798 Ramos & Edaña*. In primary forests at low altitudes. Endemic.

Local name: Bubúnad (Sub.).

13. **CURCUMA** Linnaeus**CURCUMA LONGA** Linn. Sp. Pl. (1753) 2; Blanco Fl. Filip. (1837) 5, ed. 2 (1845) 4, ed. 3, 1 (1877) 6, t. 3; Naves Novis. App. (1880) 221; K. Schum. in Engl. Pflanzenreich 20 (1904) 108; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 167.*Curcuma xanthorrhiza* Naves in Blanco Fl. Filip. ed. 3 (1877-83) t. 3, Novis. App. (1880) 222, non Roxb.Widely distributed in the Philippines in and about towns, sometimes in open waste places, sometimes planted, *Merrill Sp. Blancoanae 917, Elmer 11933*. A native of India, now pantropic. Turmeric.

Local names: Añgai (Pamp.); barák (Kuy.); diláu (Tag.); duláu (S. L. Bis.); kalabága (Bis.); kaláuag (Mbo., Bis.); kinambói (Bis.); kulálo (Pamp.); kulyáu (Ilk.); kúnig (Ilk.); kúnik (Ibn.); lampúyang (P. Bis.); láuag (Sub.); lúyang-dilaú (Tag.); pañgas (Pamp.); salampauyan (Bag.).

CURCUMA ZEDOARIA (Berg.) Rosc. Monandr. Pl. (1828) t. 109; K. Schum. in Engl. Pflanzenreich 20 (1904) 110; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 84, Philip. Journ. Sci. 4 (1909) Bot. 166; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 433; Fl. Manila (1912) 158, Sp. Blancoanae (1918) 109; Elm. Leaf. Philip. Bot. 8 (1919) 2977 (err. *zedoaria*).*Amomum zedoaria* Berg. Mat. Med. (1788) 41.*Curcuma zerumbet* Roxb. Pl. Coromandel 2 (1798) t. 101; Naves Novis. App. (1880) 222.*Costus nigricans* Blanco Fl. Filip. (1837) 3, ed. 2 (1845) 3, ed. 3, 1 (1877) 5.*Costus luteus* Blanco Fl. Filip. (1837) 4, ed. 2 (1845) 3, ed. 3, 1 (1877) 6.*Roscoea nigro-ciliata* Hassk. in Flora 47 (1864) 21.*Roscoea lutea* Hassk. l. c.

In open waste places in and near towns throughout the Philippines, often abundant; certainly introduced. India to Malaya.

Local names: Alimpuyás (C. Bis.); alimpuying (C. Bis.); barák (Tag.); bolon (Tag.); gánda (Sbl.); kónik (Ilk.); konikó (Bon.); lampúyang (P. Bis.); langkúas (Ilk.); luya-luyáhan (Tag.); tamahíba (Tag.); tamahílan (Bik.); tamó (Pamp., Tag.); tamokánsi (Tag.); únig (If.).

EXCLUDED SPECIES

CURCUMA AERUGINOSA Roxb.; Naves Novis. App. (1880) 222.**CURCUMA AUSTRALASICA** Hook. f.; Naves op. cit. 221.**CURCUMA CAESIA** Roxb.; Naves op. cit. 222.**CURCUMA PURPURASCENS** Blume; Naves l. c.**CURCUMA VIRIDIFLORA** Roxb.; Naves op. cit. 221.

14. GLOBBA Linnaeus

GLOBBA AUREA Elm. Leaf. Philip. Bot. 8 (1915) 2902.

Busuanga, Palawan, *Elmer* 13243, *Merrill* 7199, 9342, *B. S.* 21569 *Escritor*, 19339 *Weber*. In thickets and in rock crevices along streams at low and medium altitudes. Endemic.

GLOBBA BREVIFOLIA K. Schum. in Engl. Pflanzenreich 20 (1904) 145; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 160.

Globba parviflora Presl var. *brevifolia* K. Schum. in Engl. Bot. Jahrb. 27 (1899) 330.

Luzon (Zambales), Corregidor, *Warburg* 14066, *F. B.* 13216 *Curran*. In thickets or forests at low or medium altitudes. Endemic.

Apparently very close to *G. campsophylla* K. Schum.

GLOBBA CAMPSOPHYLLA K. Schum. in Engl. Pflanzenreich 20 (1904) 145; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 83, Philip. Journ. Sci. 4 (1909) Bot. 159.

Globba parviflora Vidal Phan. Cuming. Philip. (1885) 152; K. Schum. in Engl. Bot. Jahrb. 27 (1899) 330, non Presl.

Luzon (Nueva Ecija, Rizal, Tayabas, Bulacan, Bataan), Catanduanes, Panay, Samar, Mindanao, *Cuming* 1090, *Elmer* 11284, *Merrill* 4177, 5207, *Phil. Pl.* 334, *F. B.* 1461 *Ahern's* collector, 7208 *Curran*, *B. S.* 1110, 24277, 19444, 30296 *Ramos*, 6132 *Robinson*, 24975 *Fénix*, 21392 *Escritor*. In damp thickets and forests, on and about cliffs, etc., at low and medium altitudes. Endemic.

Local names: Luya-luyáhan (Tag.); malasabután (S. L. Bis.); tangbák (Tag.).

GLOBBA GRACILIS K. Schum. in Engl. Pflanzenreich 20 (1904) 145; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 161, Leaf. Philip. Bot. 2 (1909) 605.

Mindanao (Misamis, Davao), *Warburg* 14580 in herb. Berol., *F. B.* 4752 *Mearns & Hutchinson*. In primary forests, altitude 300 to 1,000 m. Endemic.

GLOBBA LATIFOLIA Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 161.

Mindoro, Panay, Mindanao (Zamboanga), *Merrill* 4064, *Williams* 2334, *B. S.* 30976 *Ramos & Edaña*, 32197 *McGregor*. In damp primary forests at low and medium altitudes. Endemic.

GLOBBA LEUCOCARPA Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 161.

Luzon (Benguet), Polillo, Mindoro, Leyte, *Elmer* 6465, 7298, *Merrill* 1789, *McGregor* 201. In damp primary forests, on shaded cliffs, etc., at low and medium altitudes. Endemic.

GLOBBA MARANTINA Linn. Mant. 2 (1771) 170; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 427; Presl Rel. Haenk. 1 (1827) 115; Llanos Frag. Pl. Filip. (1851) 7; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 2, t. 351; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 274; Naves Novis. App. (1880) 219; K. Schum. in Engl. Pflanzenreich 20 (1904) 156; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 162; Merr. Fl. Manila (1912) 156, Sp. Blancoanae (1918) 111.

Globba barthei Gagnep. in Bull. Soc. Bot. France 48 (1901) 208, t. 7, f. 1-4; K. Schum. in Engl. Pflanzenreich 20 (1904) 159; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 83.

Globba ectobolus K. Schum. in Engl. Pflanzenreich 20 (1904) 156, f. 19, L.

Globba heterobracteata K. Schum. op. cit. 159, f. 19, H.

Luzon (Union, Nueva Ecija, Pangasinan, Pampanga, Bulacan, Cavite, Rizal, Bataan), Palawan, Panay, Negros. In thickets in and near towns at low and medium altitudes, often abundant. Certainly introduced. Malay Peninsula and Archipelago to New Guinea and the Bismarck Archipelago.

Local names: Bangliu (Ilk.); barák (Tag.); luyan-luyáan (Pang.).

GLOBBA MERRILLII Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 83, Philip. Journ. Sci. 4 (1909) Bot. 160.

Luzon (Bataan), *Merrill 3869, Whitford 481, F. B. 1598 Borden*. Along streams in primary forests and on exposed forested ridges up to 1,000 m altitude; not easily distinguishable from *G. campsophylla* K. Schum. Endemic.

GLOBBA PARVIFLORA Presl Rel. Haenk. 1 (1827) 115; Miq. Fl. Ind. Bat. 3 (1859) 592; Naves Novis. App. (1880) 220; K. Schum. in Engl. Pflanzenreich 20 (1904) 146; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 83, Philip. Journ. Sci. 4 (1909) Bot. 159; Elm. Leaf. Philip. Bot. 8 (1919) 2977.

Luzon (Rizal, Albay, Sorsogon), Leyte, Mindanao, *Merrill 2783, 2782, Elmer 15666, 15381, B. S. 16503 Reillo, F. B. 2910 Mearns, 12453 Danao*. In forests at low and medium altitudes. Endemic.

GLOBBA PYRAMIDATA Gagnep. in Bull. Soc. Bot. France 48 (1901) 204, t. 4, f. 7-10; K. Schum. in Engl. Pflanzenreich 20 (1904) 146; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 158.

Mindanao (Agusan), *Weber 1128*. In primary forests at low altitudes. Indo-China.

GLOBBA USTULATA Gagnep. in Bull. Soc. Bot. France 48 (1901) 208, t. 7, f. 5-7; K. Schum. in Engl. Pflanzenreich 20 (1904) 158; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 163.

Balabac (*Marche 227B*). The type is preserved in the Paris Museum; I have seen no specimens. A variety in Indo-China fide Gagnepain.

EXCLUDED SPECIES

GLOBBA AURANTIACA Miq.; Naves Novis. App. (1880) 219.

GLOBBA CAREYANA Roxb.; Naves op. cit. 220.

GLOBBA CILIATA Jack; Naves op. cit. 219.

GLOBBA LEUCANTHA Miq.; Naves l. c.

GLOBBA MACULATA Blume; Naves l. c.

GLOBBA PATENS Miq.; Naves l. c.

GLOBBA SALTATORIA Rosc.; Naves l. c.

GLOBBA STROBILIFERA Zoll.; Naves l. c.

15. *COSTUS* Linnaeus

COSTUS CLEMENSAE Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 193.

Mindanao (Zamboanga, Lanao), *Clemens 1173*, *B. S. 36578 Ramos & Edaño*. On forested slopes, altitude about 700 m. Endemic.

Local name: Tiulasi-tiulasi (Sub.).

COSTUS GLABRA (K. Schum.) comb. nov.

Costus speciosus Sm. var. *glabra* K. Schum. in Engl. Bot. Jahrb. 27 (1899) 343, *Pflanzenreich* 20 (1904) 398.

Herba spiralis II laevis Rumph. Herb. Amb. 6: 143, t. 64, f. 2.

Costus speciosus Elm. Leaflet. Philip. Bot. 8 (1919) 2977, non Sm.

Luzon (Tayabas, Albay, Sorsogon), Polillo, Samar, Leyte, Panay, *Wenzel 227*, *Elmer 14707*, *Whitford 830*, *B. S. 6299*, *6926 Robinson*, *35478 Martelino & Edaño*, *24820*, *24811 Edaño*, *15302 Ramos*. In forests, old clearings, abacá plantations, etc., at low and medium altitudes. Moluccas.

COSTUS HIRSUTUS Blume Enum. Pl. Jav. 1 (1827) 61.

Costus speciosus Sm. var. *hirsuta* K. Schum. in Engl. Bot. Jahrb. 27 (1899) 343, *Pflanzenreich* 20 (1904) 398.

Luzon (Bontoc, Benguet, Union, Batangas), *Vanoverbergh 1392*, *Cuzner 21*, *Lete 264*. In thickets and forests, altitudes 300 to 1,200 m. India to southern China and Formosa through Malaya to New Guinea.

COSTUS SERICEUS Blume Enum. Pl. Jav. 1 (1827) 62; Naves Novis. App. (1880) 227.

Costus speciosus Sm. var. *argyrophyllus* Wall. ex Baker in Hook. f. Fl. Brit. Ind. 6 (1892) 249; Ridl. in Govt. Lab. Publ. (Philip.) 35 (1906) 84, Philip. Journ. Sci. 4 (1909) Bot. 192.

Costus speciosus Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 274; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 192, non Sm.

Luzon (Benguet, Nueva Vizcaya, Bataan, Rizal, Cavite, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Masbate, Negros, Panay, Cebu, Leyte, Mindanao, Basilan, Jolo. Common in primary forests, old clearings, etc., at low and medium altitudes. Burma, through Malaya to New Guinea.

Local names: Tiúási (Sub.); tubong-usá (Bik.).

COSTUS SPECIOSUS (Koenig) Sm. in Trans. Linn. Soc. 1 (1791) 249; Miq. Fl. Ind. Bat. 3 (1859) 610; Naves Novis. App. (1880) 227, in Blanco Fl. Filip. ed. 3 (1877-83) t. 442; K. Schum. in Engl. *Pflanzenreich* 20 (1904) 398; Merr. Fl. Manila (1912) 160.

Banksia speciosa Koenig in Retz. Obs. 3 (1783) 75.

Occasionally cultivated for ornamental purposes. This very large-flowered form is a native of the Malay Peninsula and is, I believe, specifically distinct from the common wild Philippine form. It is cultivated in various parts of Malaya and in some other tropical countries.

EXCLUDED SPECIES

COSTUS POTIERAE F.-Muell.; Naves Novis. App. (1880) 227.

EXCLUDED GENERA

ELETTARIA ALBA Blume; Naves Novis. App. (1880) 224.

ELETTARIA ANTHOIDES Teysm.; Naves l. c.

- ELETTARIA CARDAMOMUM Maton; Naves l. c.
 ELETTARIA COCCINEA Blume; Naves l. c.
 ELETTARIA FOETENS Blume; Naves l. c.
 ELETTARIA HEMISPHAERICA Blume; Naves l. c.
 ELETTARIA MACROCEPHALA Miq.; Naves op. cit. 225.
 ELETTARIA MINUTA Blume; Naves op. cit. 224.
 ELETTARIA MOLLIS Blume; Naves l. c.
 ELETTARIA PINANGA Miq.; Naves op. cit. 225.
 ELETTARIA PUNICEA (Roxb.) Naves op. cit. 224.
 ELETTARIA RUBRA Miq.; Naves op. cit. 225.
 ELETTARIA SOLARIS Blume; Naves op. cit. 224.
 ELETTARIA SPECIOSA Blume; Naves l. c.
 ELETTARIA TOMENTOSA Miq.; Naves op. cit. 225.
 ELETTARIA VILLOSA Miq.; Naves l. c.
 ROSCOEA PURPUREA Spreng.; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 99; Naves Novis. App. (1880) 223.

CANNACEAE

1. CANNA Linnaeus

- CANNA COCCINEA Mill. Gard. Dict. ed. 8 (1768) No. 3; Miq. Fl. Ind. Bat. 3 (1859) 613; Naves Novis. App. (1880) 229; Merr. Fl. Manila (1912) 161; Kränzl. in Engl. Pflanzenreich 56 (1912) 60.

Cultivated for ornamental purposes, but nowhere naturalized. A native of Central and South America.

- CANNA FLACCIDA Salisb. Ic. Stirp. Rar. (1791) t. 2; Merr. Fl. Manila (1912) 162 (*C. flaccida* Salisb. x *iridiflora* Ruiz & Pav.); Kränzl. in Engl. Pflanzenreich 56 (1912) 49.

Widely cultivated for its showy flowers, but not naturalized and apparently never or very rarely producing seeds. A native of tropical America.

Local names: Bandera española (Sp.); kambút bandera española (Ilk.).

- CANNA GLAUCA Linn. Sp. Pl. (1753) 1; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 198; Merr. Fl. Manila (1912) 162; Kränzl. in Engl. Pflanzenreich 56 (1912) 54.

Cultivated for ornamental purposes, Merrill 648, 6690, 6691. A native of tropical America.

- CANNA INDICA Linn. Sp. Pl. (1753) 1; Blanco Fl. Filip. (1837) 6, ed. 2 (1845) 5, ed. 3, 1 (1877) 8, t. 4; Naves Novis. App. (1880) 229; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 198; Merr. op. cit. 5 (1910) Bot. 340, Fl. Manila (1912) 161, Sp. Blancoanae (1918) 111. *Canna orientalis* Rosc. Monandr. Pl. (1828) t. 12; Kränzl. in Engl. Pflanzenreich 56 (1912) 47. ?

Throughout the Philippines in the settled areas, occurring in waste places in and near settlements. A native of tropical America, now pan-tropic

Local names: Balunsáyng (Bis.); baṅgali (Bik.); kakuentásan (Tag.); kiuiṅgan (If.); kolintásan (Bis.); kuentas-kuentásan (Tag.); lása (Iv.); saging-ságing (Tag.); tapuráṅga (Bis.); t́kas (Tag.); t́kas-t́kas (Tag., P. Bis.); t́kis-t́kis (Tag.); tukas-t́kas (Tag.).

CANNA REEVESII Lindl. Bot. Reg. 23 (1837) *t.* 2004; Kränzl. in Engl. Pflanzenreich 56 (1912) 50, *f.* 10, A-C.

Cultivated for ornamental purposes, but nowhere spontaneous, *Merrill 6692*, det. Kränzl. A native of tropical America.

CANNA SPECIOSA Rosc. Monandr. Pl. (1828) *t.* 17; Naves Novis. App. (1880) 229; Merr. Fl. Manila (1912) 161; Kränzl. in Engl. Pflanzenreich 56 (1912) 44, *f.* 8 D.

Occasional in open waste places, naturalized, *Merrill 6689*. A native of tropical America.

Local name: Kambút-ṅganalabága (Ilk.).

CANNA WARCEWICZII Dietr. in Allg. Gart. Zeit. 19 (1851) 290, fig.; Merr. Fl. Manila (1912) 162; Kränzl. in Engl. Pflanzenreich 56 (1912) 64.

Cultivated for ornamental purposes, but nowhere established. A native of tropical America.

Local name: Kambut-ṅga-naṅgísit (Ilk.).

EXCLUDED SPECIES

CANNA MONTANA Blume; Naves Novis. App. (1880) 229.

CANNA EDULIS Ker.; Naves l. c.

CANNA PEDUNCULATA Sims; Naves l. c.

CANNA RUBICAULIS Link; Naves l. c. = *C. edulis* Ker.

CANNA FLAVESCENS Link; Naves l. c.

All our forms of *Canna* are introduced plants in the Philippines. Two or three characteristic species are thoroughly naturalized, especially *Canna indica* Linn., which was introduced at an early date in colonial history. A very large number of garden forms, mostly hybrids, have been introduced from Europe and America since the beginning of the present century, but it is impossible to determine their botanical origins, as the necessary data are lacking; few of these hybrids ever produce seeds here, but are reproduced by means of their rhizomes.

MARANTACEAE

1. **DONAX** Loureiro

DONAX CANNAEFORMIS (Forst. f.) K. Schum. in Engl. Bot. Jahrb. 15 (1893) 440; Rolfe in Journ. Bot. 45 (1907) 243; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 193; Valetton ex Merr. Interpret. Herb. Amb. (1917) 166; Merr. Sp. Blancoanae (1918) 112.

Thalia cannaeformis Forst. f. Prodr. (1786) 1.

Maranta arundinacea Blanco Fl. Filip. (1837) 7, ed. 2 (1845) 5, ed. 3, 1 (1877) 9, *t.* 5, non Linn.

Clinogyne grandis Benth. & Hook. f. ex Rolfe in Journ. Bot. 23 (1885) 465; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 275.

Donax arundastrum K. Schum. in Engl. Pflanzenreich 11 (1902) 33; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 37, Forestry Bureau Bull. 1 (1903) 16; Perk. Frag. Fl. Philip. (1904) 68, t. 1 f. G-J, non Lour.

Actoplanes cannaeformis K. Schum. in Engl. Pflanzenreich 11 (1902) 34 p. p. excl. syn. Miq.

Phrynium dichotomum Roxb. in Asiat. Research 11 (1810) 324; Schauer in Nov. Act. Acad. Nat. Cur. 19 (1843) Suppl. 1: 427.

Maranta dichotoma Wall. Cat. (1832) No. 6615; Naves Novis. App. (1880) 228; Miq. Fl. Ind. Bat. 3 (1859) 611; Usteri Beitr. Ken. Philip. Veg. (1905) 128.

Maranta grandis Miq. Fl. Ind. Bat. Suppl. (1860-1) 616; Naves Novis. App. (1880) 228.

Batan Islands and northern Luzon to Palawan and Mindanao, in all or most islands and provinces. Common in secondary forests, especially along streams at low and medium altitudes. Java and Borneo to New Guinea, the Aru and Admiralty Islands, New Hebrides, and the Marianne Islands.

Local names: Alaró (Bis.); aratán (Gad.); babán (Chab.); bambán (Ilk.; Bis., Mbo., Tag., Sul., Buk.); banbán (Tag., Bik., Bis., Bag., Ibn.); bankólid (Mbo.); baras-barásan (Tag.); bonbón (Tag.); buabán (Bag.); daromáka (Ilk.); darumáka (Ilk.); garomáka (Ilk.); lankuás (Ilk.); mambán (Bik., S. L. Bis.); manbán (Tag.); matalbák (Tag.); matapál (Isn.); miní (Ig.); niní (Ig.); niník (Iv.).

2. PHRYNIUM Willdenow

PHRYNIUM PHILIPPINENSE Ridl. in Leaflet. Philip. Bot. 2 (1909) 569, Philip. Journ. Sci. 4 (1909) Bot. 195.

Phrynium capitatum Presl Rel. Haenk. 1 (1827) 109; Naves Novis. App. (1880) 228, non Willd.

Phrynium densiflorum Naves Novis. App. (1880) 228; Vidal Phan. Cuming. Philip. (1885) 152, Rev. Pl. Vasc. Filip. (1886) 275, non Blume.

Babuyan Islands, Luzon (Isabela, Nueva Vizcaya, Rizal, Laguna, Taya-bas, Albay, Sorsogon), Alabat, Polillo, Mindoro, Palawan, Panay, Leyte, Negros, Mindanao. In forests at low and medium altitudes; scarcely distinguishable from *P. capitatum* Willd. Endemic.

Local names: Bogíghík (Mbo.); hagíghík (Bis.); hagithít (Tag.); lutatában (P. Bis.); okos (Bag.).

EXCLUDED SPECIES

PHRYNIUM LATIFOLIUM Blume; Naves Novis. App. (1880) 228.

PHRYNIUM MAXIMUM Blume; Naves l. c.

PHRYNIUM PUBINERVE Blume; Naves l. c.

Some species of *Calathea* have been introduced into the Philippines, and are occasionally found in cultivation for ornamental purposes.

3. MARANTA Linnaeus

MARANTA ARUNDINACEA Linn. Sp. Pl. (1753) 2; Naves Novis. App. (1880) 228; K. Schum. in Engl. Pflanzenreich 11 (1902) 125; Perk. Frag. Fl. Philip. (1904) 72, t. 2; Merr. in Govt. Lab. Publ. (Philip.) 6 (1904) 29, Fl. Manila (1912) 163; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 197.

Maranta ramosissima Wall. Pl. As. Rar. 3 (1832) 51, t. 286; Naves Novis. App. (1880) 228.

Widely distributed in the Philippines in cultivation for its starchy rhizomes. A native of tropical America, now pantropic. Arrowroot.

Local names: Ararú (Ibn., It.); arorú (Tag.); arurú (Tag.); báí (Iv.); galamáka (Bon.); sagú (Bik., Ilk.).

Var. **VARIEGATA** (N. E. Br.) Ridl. in Gard. Chron. III 9 (1891) 73; Merr. Fl. Manila (1912) 163.

Phrynium variegatum N. E. Br. in Ill. Hort. 33 (1886) 125, t. 606.

Cultivated for ornamental purposes.

4. MONOPHRYNium K. Schumann

MONOPHRYNium CONGESTUM Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 197.

Luzon (Apayao, Tayabas, Laguna), *F. B.* 9548 *Curran*, *B. S.* 28082, 28245 *Fénix*. In primary forests at low or medium altitudes. Endemic.

Local names: Aláuan (Ibn.); arúpi (Ibn.).

MONOPHRYNium FASCICULATUM (Presl) K. Schum. in Engl. Pflanzenreich 11 (1902) 68; Perk. Frag. Fl. Philip. (1904) 69, t. 1, f. A-F; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 196.

Calathea fasciculata Presl Rel. Haenk. 1 (1827) 108, t. 10, f. 1; Naves Novis. App. (1880) 229.

Phrynium fasciculatum Horan. Monogr. (1862) 11.

Luzon (Cagayan, Laguna, Tayabas), Samar, Leyte, Biliran, Mindanao (Surigao), *B. S.* 34797 *Ramos & Pascasio*, 28589 *Ramos & Edaño*, 24136, 13356 *Ramos*, 22837 *McGregor*, *Elmer* 9148, *Wenzel* 378. In primary forests at low and medium altitudes. Halmahera. The only representative of the genus known outside of the Philippines.

Local names: Hagithít (Tag.); hagithít-na-daytoi (S. L. Bis.).

MONOPHRYNium SIMPLEX Elm. Leaf. Philip. Bot. 1 (1908) 276; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 196.

Luzon (Laguna, Tayabas), *Elmer* 7611, 9147, 17502. On forested slopes near streams, altitude about 750 m. Endemic.

5. PHACELOPHRYNIUM K. Schumann

PHACELOPHRYNIUM BRACTEOSUM (Warb.) K. Schum. in Engl. Pflanzenreich 11 (1902) 123; Perk. Frag. Fl. Philip. (1904) 71, t. 3, f. F-O; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 268; Ridl. op. cit. 4 (1909) Bot. 195.

Phrynium bracteosum Warb. ex K. Schum. l. c. in syn.

Luzon (Sorsogon), Mindoro, Samar, Leyte, Biliran, Mindanao (Agusan, Lanao), *Merrill* 7288, 5202, 5232, *Elmer* 16145, 15240, *Clemens* 423, *Wenzel*

232, *B. S.* 34564 *Ramos & Pascasio*, 24732 *Edaño*, 15229, 17456, 39780 *Ramos*, 18818 *McGregor*. In primary forests at low and medium altitudes. Borneo.

PHACELOPHRYNIUM CYLINDRICUM Merr. in Philip. Journ. Sci. 13 (1918) Bot. 269.

Catanduanes, Panay, *B. S.* 30503 *Ramos*, 31271 *Ramos & Edaño*. In primary forests at low altitudes. Endemic.

PHACELOPHRYNIUM INTERRUPTUM (Warb.) K. Schum. in Engl. Pflanzenreich 11 (1902) 121; Perk. Frag. Fl. Philip. (1904) 70, t. 3, f. A-E; Ridl. in Philip. Journ. Sci. 4 (1909) Bot. 194.

Phrynium interruptum Warb. ex K. Schum. l. c. in syn.

Luzon (Ilocos Norte, Benguet, Bulacan, Rizal, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Samar, Panay, Leyte, Mindanao (Agusan, Davao). In forests at low and medium altitudes, represented by about fifty individual collections. Endemic.

Local names: Bambán (Tag.); hagikhiik (S. L. Bis.); hagitit (Tag., Bik.); kokos (Bag.); nítni (Ig.).

BURMANNIACEAE

1. BURMANNIA Linnaeus

BURMANNIA CLEMENTIS Schltr. in Philip. Journ. Sci. 1 (1906) Suppl. 305.

Mindanao (Lanao), *Clemens* 24. In grasslands, altitude about 650 m. Endemic.

BURMANNIA COELESTIS D. Don Prodr. Fl. Nepal. (1825) 44; Naves Novis. App. (1880) 252; Merr. in Philip. Journ. Sci. 4 (1909) Bot. 250.

Burmammia azurea Griff. Notul. 3 (1851) 236, Ic. Pl. As. (1851) t. 272, f. 1; Becc. Malesia 1 (1878) 242, t. 15, f. 1-3.

Luzon (Pangasinan), *B. S.* 4913 *Ramos*. Probably in open places at low altitudes. India to Borneo and the Caroline Islands.

BURMANNIA LONGIFOLIA Becc. Malesia 1 (1878) 244; Naves Novis. App. (1880) 252; Rendle in Journ. Bot. 34 (1896) 355; Merr. in Philip. Journ. Sci. 2 (1907) Bot. 268.

Mindoro, Negros, Camiguin de Misamis, Mindanao (Agusan, Davao), *Merrill* 5741, *Elmer* 13690, *B. S.* 14428 *Ramos*, *F. B.* 13611 *Curran & Foxworthy*. On ridges in wet mossy forests, altitude 1,200 to 1,600 m. Malay Peninsula, Borneo, Amboina, New Guinea.

Local names: Bagumbáan (Mbo.); pamaláuan (Bag.).

EXCLUDED GENERA AND SPECIES

BURMANNIA DISTICHA Linn.; Naves Novis. App. (1880) 252.

BURMANNIA SUMATRANA Miq.; Naves l. c.

BURMANNIA TUBEROSA Becc.; Naves l. c.

GYMNOSIPHON APHYLLUM Blume; Naves l. c.

SARCOSIPHON CLANDESTINUM Blume; Naves l. c.

APOSTASIACEAE *

1. NEUWIEDIA Blume

NEUWIEDIA VERATRIFOLIA Blume in Hoen. & De Vriese Tijdschr. Nat. Gesch. 1 (1834) 142, in Ann. Sci. Nat. II 2 (1834) 94; Reichb. f. in Bonpl. 5 (1857) 58, in nota; Miq. Fl. Ind. Bat. 3 (1859) 748; Naves Novis. App. (1882) 251; Rolfe in Journ. Linn. Soc. 25 (1889) 231, 241; Hook. f. in Hook. Icon. Pl. 20 (1891) *t.* 1987; Rolfe in Orch. Rev. 4 (1896) 329; Kränzl. Orch. Gen. et Sp. 1 (1897) 3, p. p.; Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 5, *fig.* 3; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 15; Ames in Philip. Journ. Sci. 2 (1907) Bot. 311; J. J. Sm. in Bull. Jard. Buitenz. II 14 (1914) 1.

Mindoro, *Merrill 5681*. Mt. Halcon, in forest at 750 m altitude. In Java and Borneo.

EXCLUDED OR SYNONYMOUS SPECIES

NEUWIEDIA ZOLLINGERI Reichb. f. apud Naves Novis. App. (1882) 251; Rolfe in Journ. Linn. Soc. 25 (1889) 231; Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 5.

This Javan and Bornean species is credited to the Philippines solely on the basis of Naves's record.

2. APOSTASIA Blume

APOSTASIA WALLICHII R. Br. in Wall. Pl. Asiat. Rar. 1 (1830) 75, *t.* 84; Miq. Fl. Ind. Bat. 3 (1859) 748; Rolfe in Journ. Linn. Soc. 25 (1889) 239; Hook. f. Fl. Brit. Ind. 6 (1890) 175; Ridl. in Journ. Linn. Soc. 32 (1896) 415; Kränzl. Orch. Gen. et Sp. 1 (1897) 7; Trim. Handb. Fl. Ceyl. 4 (1898) 238; Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 7, *fig.* 4; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 18, Fig.-Atlas (1908) *fig.* 3, in Bull. Dept. Agr. Ind. Néerl. 13 (1907) 1; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 232; Ames in Philip. Journ. Sci. 2 (1907) Bot. 311.

Mesodactylus deflexa Wall. Pl. Asiat. Rar. 1 (1830) 74.

Luzon (Camarines), Sibuyan, Mindoro, Palawan, *F. B. 18750 Darling, Merrill 5521, 5639, Elmer 13141*. Terrestrial, up to 1,000 m altitude. In India, Ceylon, Malay Peninsula, Sumatra, Java, Borneo, and New Guinea.

EXCLUDED OR SYNONYMOUS SPECIES

APOSTASIA ODORATA Blume apud Naves Novis. App. (1882) 251; Rolfe in Journ. Linn. Soc. 25 (1889) 239.

The Philippine record, repeated by Rolfe, is based wholly on the earlier one of Naves; the species is known only from Java.

* The enumeration of the Apostasiaceae and the Orchidaceae was prepared by Prof. Oakes Ames, who was materially assisted in the bibliographic research connected therewith by Mr. F. Tracy Hubbard.— E. D. M.

ORCHIDACEAE *

1. CORDULA Rafinesque

CORDULA ARGUS (Reichb. f.) Rolfe in Orch. Rev. 20 (1912) 2.

Cypripedium argus Reichb. f. in Gard. Chron. (1873) 608, (1874) 710, III 7 (1890) 259; Hook. f. in Bot. Mag. 101 (1875) t. 6175; Naves Novis. App. (1882) 251; Veitch Man. Orch. Pl. pt. 4 (1889) 11; Sander Reichenbachia I 2 (1890) 77, t. 83; Ceron Cat. Pl. Herb. Manila (1892) 169; Kränzl. Orch. Gen. et Sp. 1 (1897) 56.

Cypripedium barbatum Lindl. var. *argus* Vos. in Belg. Hort. 25 (1875) 57.

Cypripedium pitcherianum Manda in Am. Flor. 3 (1887) 178.

Cypripedium moensianum Hort. ex L. Linden in Lindenia 3 (1887) 69, t. 129.

Cypripedium argus Reichb. f. var. *moensii* Hort. ex Veitch Man. Orch. Pl. pt. 4 (1889) 11.

Cypripedium argus Reichb. f. subvar. *nigricans* Veitch Man. Orch. pl. pt. 4 (1889) 11.

Paphiopedilum argus Stein Orchideenb. (1892) 453; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 40, in Engl. Pflanzenreich IV. 50 (1903) 90; Ames Orch. 1 (1905) 63, 2 (1908) 23, 5 (1915) 10.

Paphiopedilum argus Stein var. *moensii* Stein Orchideenb. (1892) 450; Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 91; Ames Orch. 2 (1908) 24.

Paphiopedilum argus Stein var. *nigricans* Hort. ex Stein Orchideenb. (1892) 454; Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 91; Ames Orch. 2 (1908) 24.

Luzon (Bontoc, Benguet, Tayabas), Negros (Oriental Negros), *Van-overbergh* 262, 1186, 4042, *Elmer* 6296, 9433, *Lyon* 158, *B. S.* 32015 *Santos*, 20735 *Escritor*. Grassy slopes at 1,300 to 2,200 m altitude.

CORDULA BARBATA (Lindl.) Rolfe in Orch. Rev. 20 (1912) 2.

Cypripedium barbatum Lindl. in Bot. Reg. 27 (1841) Misc. 53, 28 (1842) t. 17; Hook. in Bot. Mag. 72 (1846) t. 4234; Miq. Fl. Ind. Bat. 3 (1859) 737; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 237; Veitch Man. Orch. Pl. pt. 4 (1889) 12, text cut; Hook. f. Fl. Brit. Ind. 6 (1890) 174; Ridl. in Journ. Linn. Soc. 32 (1896) 414; Kränzl. Orch. Gen. et Sp. 1 (1897) 62; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 229.

Paphiopedilum barbatum Pfitz. in Pringsh. Jahrb. 19 (1888) 159, in Engl. & Prantl Pflanzenfam. 2, Abt. 6 (1888) 84, in Engl. Bot. Jahrb. 19 (1894) 40, in Engl. Pflanzenreich IV. 50 (1903) 91; Ames Orch. 2 (1908) 24, 5 (1915) 10.

Palawan, *B. S.* 640, 645 *Foxworthy*. Mt. Victoria, in rocky forest at 930 to 1,000 m altitude. In Malay Peninsula and western Siam.

* See preceding footnote.

CORDULA CILIOLARIS (Reichb. f.) Rolfe in Orch. Rev. 20 (1912) 2.

Cypripedium ciliolare Reichb. f. in Gard. Chron. II 18 (1882) 488, 20 (1883) 46; Rodigas in Ill. Hort. 31 (1884) 127, t. 530; Veitch Man. Orch. Pl. pt. 4 (1889) 16; Kränzl. Orch. Gen. et Sp. 1 (1897) 64.

Paphiopedilum ciliolare Stein Orchideenb. (1892) 462, fig. 145; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 40, in Engl. Pflanzenreich IV. 50 (1903) 89; Ames Orch. 2 (1908) 23, 5 (1915) 11, in Philip. Journ. Sci. 6 (1911) Bot. 39.

Luzon (Bontoc), Dinagat, Mindanao (Surigao, Agusan), *B. S.* 37996 Ramos & Edaña, 34586 Ramos & Pascasio, *Lyon* 142, *Weber* 110. Forest slopes at 310 to 1,830 m altitude.

CORDULA HAYNALDIANA (Reichb. f.) Rolfe in Orch. Rev. 20 (1912) 2.

Cypripedium haynaldianum Reichb. f. Xen. Orch. 2 (1874) 222, 3 (1881) 33, t. 212, in Gard. Chron. II 7 (1877) 272, 752; Hook. f. in Bot. Mag. 103 (1877) t. 6296; Naves Novis. App. (1882) 251; Veitch Man. Orch. Pl. pt. 4 (1889) 28; Kränzl. Orch. Gen. et Sp. 1 (1897) 71.

Paphiopedilum haynaldianum Stein Orchideenb. (1892) 470; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 41, in Engl. Pflanzenreich IV. 50 (1903) 66, fig. 31 A-C; Ames Orch. 2 (1908) 22, 5 (1915) 10, in Philip. Journ. Sci. 6 (1911) Bot. 39; Memml. in Orchis 14 (1920) 18.

Luzon (Kalinga, Tarlac, Rizal), *Wallis s. n.*, *Lyon* 49, *Boettcher s. n.*, *Ramos s. n.* Sea level to 1,000 m altitude.

CORDULA PHILIPPINENSIS (Reichb. f.) Rolfe in Orch. Rev. 20 (1912) 2.

Cypripedium philippinense Reichb. f. in Bonpl. 10 (1862) 335; Veitch Man. Orch. Pl. pt. 4 (1889) 42, text cut; Kränzl. Orch. Gen. et Sp. 1 (1897) 46; Rolfe in Orch. Rev. 22 (1914) 253, fig. 29.

Cypripedium laevigatum Batem. in Bot. Mag. 91 (1865) t. 5508; Dombraïn in Paxt. Floral Mag. 5 (1866) t. 298; Morr. in Belg. Hort. 17 (1867) 102, t. 6; Van Houtte in Fl. des Serres 17 (1869) 83, t. 1760-61; Batem. Second Cent. Orch. Pl. (1867) t. 101; Naves Novis. App. (1882) 251; Ceron Cat. Pl. Herb. Manila (1892) 169.

Cypripedium roebbelenii Reichb. f. in Gard. Chron. II 20 (1883) 684; Kränzl. Xen. Orch. 3 (1893) 113, t. 265, Orch. Gen. et Sp. 1 (1897) 47.

Selenipedium laevigatum May in Rev. Hort. 57 (1885) 301.

Cypripedium cannartianum Hort ex L. Linden in Lindenia 3 (1887) 93, t. 141.

Cypripedium roebbelenii Reichb. f. var. *cannartianum* L. Linden in Lindenia 3 (1887) 93, in syn.

Paphiopedilum laevigatum Pfitz. in Engl. & Prantl Pflanzenfam. 2, Abt. 6 (1888) 84.

Cypripedium philippinense Reichb. f. var. *platytaenium* Desbois Bull. Orchidéene (1890), *Cypriped.* (1898) 285.

Phapiopedilum philippinense Stein Orchideenb. (1892) 480; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 41, in Engl. Pflanzenreich IV. 50 (1903) 61, fig. 28 A-E; Ames Orch. 2 (1908) 19, 5 (1915) 10.

Paphiopedilum roebelinii Stein Orchideenb. (1892) 480, 482, in syn.; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 41 (as *roëbbelinii*), in Engl. Pflanzenreich IV. 50 (1903) 62.

Paphiopedilum philippinense Stein var. *platytaenium* Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 61; Ames Orch. 2 (1908) 20.

Paphiopedilum philippinense Stein var. *cannartianum* Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 62 (as *cannaertianum*); Ames Orch. 2 (1908) 20; Memml. in Orchis 14 (1920) 18.

Paphiopedilum philippinense Stein var. *roebbelenii* Kerchove apud Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 62, in syn.; Ames Orch. 2 (1908) 20.

Cypripedium philippinense Reichb. f. var. *roebbelenii* Veitch apud Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 62, in syn.

Palawan (Apulit Island), *Merrill 9423*.

2. HERMINIUM R. Brown

HERMINIUM ANGUSTIFOLIUM (Lindl.) Benth. ex Hook. f. Fl. Brit. Ind. 6 (1890) 129; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 332, t. 434; Kränzl. Orch. Gen. et Sp. 1 (1901) 553; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 25, Fig.-Atlas (1908) fig. 7; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 197; Ames Orch. 2 (1908) 25, 5 (1915) 12; Merr. in Philip. Journ. Sci. 13 (1918) Bot. 135.

Aceras angustifolia Lindl. in Wall. Cat. (1832) No. 7061, *nomen*; Lindl. in Bot. Reg. 18 (1832) sub t. 1525; Lindl. Gen. & Sp. Orch. (1835) 282; Wight Icon. 5¹ (1852) 11, t. 1691.

Aceras longicuris C. Wright ex A. Gray in Mem. Am. Acad. II 6 (1858) 411.

Luzon (Bontoc, Lepanto, Benguet). Reasonably common in this region at about 2,000 to 2,300 m altitude. In British India, China, Japan, and Java.

3. HABENARIA Willdenow

HABENARIA ALAGENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 312, Orch. 5 (1915) 12.

Habenaria luzonensis Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1550.

Luzon (Tayabas, Sorsogon), Catanduanes, Mindoro, Leyte, Mindanao (Camiguin Island), *Elmer 7254, 7636a, 15474, 15748, B. S. 28510 Ramos & Edaño, 14440 Ramos, Merrill 5803, Wenzel 0971, 0972, 0984, Ramos s. n.* Terrestrial, apparently near streams, up to 3,000 m in Mindoro.

HABENARIA ANGUSTATA (Blume) Ktze. Rev. Gen. Pl. 2 (1891) 664; Ames in Philip. Journ. Sci. 2 (1907) Bot. 312; Orch. 2 (1908) 26, in Merr. in Journ. Roy. Asiatic Soc. Straits Branch, Special No. (1921) 137.

Mecosa angustata Blume Bijdr. (1825) 404, fig. 1.

Platanthera angustata Lindl. Gen. & Sp. Orch. (1835) 290; Reichb. f. in Bonpl. 5 (1857) 34; Miq. Fl. Ind. Bat. 3 (1859) 714; Naves Novis. App. (1882) 251; Kränzl. Orch. Gen. et Sp. 1 (1899) 647; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 29, Fig.-Atlas (1908) fig. 12.

Platanthera amesiana Schltr. in Fedde Repert. 9 (1911) 212.

Habenaria amesiana Ames Orch. 5 (1915) 12.

Luzon (Benguet, Bataan, Laguna, Albay), Negros (Oriental Negros), Mindanao (Camiguin Island), B. S. 31983 Santos, 14442 Ramos, 2948 Mearns, Ramos & Edaño s. n., Whitford 162, Elmer 10002, Merrill 7494. In Java and Borneo.

HABENARIA ARISTULIFERA Reichb. f. in Bonpl. 3 (1855) 213; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 273; Kränzl. in Engl. Bot. Jahrb. 16 (1892) 84, Orch. Gen. et Sp. 1 (1897) 231; Ames Orch. 2 (1908) 32, 5 (1915) 13.

Habenaria cumingii Kränzl. Orch. Gen. et Sp. 1 (1898) 262; Ames Orch. 5 (1915) 13.

Luzon (Rizal, Laguna), Alabat, Mindanao (Davao), Cuming 2091, Wilkes Exped., B. S. 12530, 14590, 16957 Ramos, Weber 283, Merrill 10435, Taylor 9, 10.

HABENARIA BISAETA Ames Orch. 2 (1908) 30, text cut, 5 (1915) 13.

Habenaria tipulifera Par. & Reichb. f. apud Vidal Phan. Cuming. Philip. (1885) 151; Vidal Rev. Pl. Vasc. Filip. (1886) 273; Ames Orch. 5 (1915) 15.

Luzon (Bontoc, Benguet), Cuming 2095, Vanoverbergh 788, 1359, 1480, R. S. Williams 1936, Merrill 4325, F. B. 4887 Curran, B. S. 4461 Mearns, Ramos & Edaño 57, 73. Terrestrial in mountain regions at about 1,300 to 2,300 m altitude.

HABENARIA BOADANENSIS Ames Orch. 7 (1922) 153.

Luzon (Benguet), Ramos & Edaño 78. Mt. Boadan, on open slopes at 2,000 m altitude.

HABENARIA CONGESTA Ames Orch. 2 (1908) 38, text cut, 5 (1915) 13; Merr. Fl. Manila (1912) 167.

Luzon (Lepanto), Panay (Antique), Guimaras, Busuanga, Culion, Mindanao (Zamboanga), Jolo, Merrill 4479, Yoder 26, s. n., B. S. 18008 Robinson, 41397 Lopez, 41333 Ramos, Hallier s. n., F. Link s. n. Apparently in dry open situations at low altitude.

HABENARIA COPELANDII Ames Orch. 2 (1908) 29, text cut, 5 (1915) 13.

Luzon (Bontoc, Nueva Ecija, Bataan, Rizal, Cavite, Laguna, Batangas), Panay, Culion, Coron, Vanoverbergh 1320, 3664, 4050, Copeland 1403, Merrill 4172, F. B. 3243 Ahern's collector, B. S. 4588, 22398, 41147, 41321 Ramos, 6139 Robinson, 1248 Mangubat, 26316, 26489 Ramos & Edaño, 32444 McGregor, Native collector s. n. At about 15 to 1,250 m altitude.

HABENARIA CURRANII Ames in Philip. Journ. Sci. 7 (1912) Bot. 2, Orch. 5 (1915) 13.

Luzon (Cagayan, Benguet), F. B. 17138 Curran, B. S. 12578 Fenix. Lalloc, edge of forest at 50 m altitude, and Sablan.

HABENARIA DELESSERTIANA Kränzl. in Ann. Conserv. Jard. Bot. Genève 1 (1897) 108, Orch. Gen. et Sp. 1 (1897) 233; Ames in Philip. Journ. Sci. 7 (1912) Bot. 2, Orch. 5 (1915) 13.

Luzon (Laguna), Mindanao (Lanao), *Cuming 2086, pro parte, B. S. 12407* McGregor, *M. S. Clemens 865*. Calauan and Camp Keithley.

HABENARIA DIPHYLLA (Nim.) Dalz. in Hook. Kew Journ. Bot. 2 (1850) 262; Reichb. f. in Walp. Ann. 3 (1852) 588; Hook. f. Fl. Brit. Ind. 6 (1890) 151; Kränzl. Orch. Gen. et Sp. 1 (1898) 346; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 185, *t. 137*; Ames Orch. 2 (1908) 37, 5 (1915) 13.

Liparis diphyllus Nim. in Grah. Cat. Bomb. Pl. (1839) 252.

Habenaria jerdoniana Wight Icon. 5¹ (1852) 14, *t. 1715*; Kränzl. in Engl. Bot. Jahrb. 16 (1892) 154.

Habenaria sutteri Reichb. f. in Linnaea 25 (1852) 229.

Habenaria cordata Naves Novis. App. (1882) 250; Merr. Sp. Blancoanae (1918) 21, non R. Br.

Habenaria ahernii Schltr. in Engl. Bot. Jahrb. 53 (1915) 502.

Luzon (Ilocos Norte, Rizal, Laguna), *F. B. 3241 Ahern's collector, Ramos 1071, 4668, B. S. 21331, 32754 Ramos, Native collector s. n.* In India.

HABENARIA ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1550, Orch. 5 (1915) 14.

Mindanao (Davao), *Elmer 11845a*. Todaya, Mt. Apo.

HABENARIA FALCIGERA Reichb. f. in Linnaea 41 (1876) 100; Kränzl. in Engl. Bot. Jahrb. 16 (1892) 153, Orch. Gen. et Sp. 1 (1898) 345; Ames Orch. 2 (1908) 36, 5 (1915) 14.

Luzon (Bontoc, Benguet), *Wallis s. n., Vanoverbergh 1239, B. S. 8792* McGregor, *4332 Mearns, F. B. 4884 Curran*.

HABENARIA GOODYEROIDES D. Don Prodr. Fl. Nepal. (1825) 25; Hook. f. Fl. Brit. Ind. 6 (1890) 161 (excl. syn. *P. grandis* Blume); King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 326, *t. 430*; Schltr. in K. Schum. & Lauterb. Nachtr. Fl. Deutsch. Schutzgeb. (1905) 79; Ames Orch. 2 (1908) 27, 5 (1915) 14, 6 (1920) 8; Schltr. in Engl. Bot. Jahrb. 45, Beibl. No. 104 (1911) 3, in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 5; Merr. Fl. Manila (1912) 167; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 138.

Herminium goodyeroides Lindl. in Wall. Cat. (1832) No. 7066, *nomen*.

Peristylus goodyeroides Lindl. Gen. & Sp. Orch. (1835) 299, in Bot. Reg. 26 (1840) Misc. 81; Ceron Cat. Pl. Herb. Manila (1892) 169; Grant Orch. Burma (1898) 329 (excl. syn. *P. grandis* Blume); Kränzl. Orch. Gen. et Sp. 1 (1898) 503 (excl. syn. *P. grandis* Blume); J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 32, Fig.-Atlas (1908) *fig. 14*, in Nova Guinea 12 (1915) Bot. 178.

Platanthera goodyeroides Vidal Phan. Cuming. Philip. (1885) 151.

Habenaria lauterbachii Kränzl. in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 239, *nomen*.

Luzon (Bontoc, Union, Benguet, Bataan, Rizal, Cavite, Laguna), Panay (Antique), Mindanao (Bukidnon, Lanao). A fairly common species in

the Philippines, especially in Luzon, up to 1,300 m altitude in Bontoc. In India, Yunnan, Assam, Burma, Andaman Islands, Sumatra, Java, Borneo, New Guinea, and Formosa.

HABENARIA HALCONENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 312, Orch. 5 (1915) 14.

Platanthera halconensis Schltr. in Fedde Repert. 9 (1911) 212.

Mindoro, Merrill 5835. Mt. Halcon, terrestrial on open heath at 2,650 m altitude.

HABENARIA HYSTRIX Ames Orch. 2 (1908) 35, 5 (1915) 14.

Dissorhynchium muricatum Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 434, t. 1, fig. A.

Habenaria muricata Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 273; Rolfe in Journ. Bot. 24 (1886) 59; Kränzl. in Engl. Bot. Jahrb. 16 (1892) 140, Orch. Gen. et Sp. 1 (1898) 326; Ames Orch. 1 (1905) 64, non Rodr. (1882).

Luzon (Bataan, Rizal, Cavite, Laguna, Batangas, Tayabas, Sorsogon), Siquijor, Dumarán, Mindanao (Zamboanga). A rather widely distributed species in the Philippines and fairly common in Luzon at about 120 to 300 m altitude. Also in British North Borneo.

HABENARIA INTRUDENS Ames Sched. Orch. 6 (1923) 1, fig. 1.

Habenaria lacertifera Ames Orch. 2 (1908) 39, 5 (1915) 14, non Benth.

Luzon (Benguet, Bulacan, Rizal), F. B. 5085 Curran, R. S. Williams 1933, B. S. 21706, 21999 Ramos. At 90 to 100 m altitude.

HABENARIA LEIBERGII Ames Orch. 2 (1908) 34, text cut, 5 (1915) 14, in Philip. Journ. Sci. 4 (1909) Bot. 600.

Luzon (Ilocos Norte, Bontoc, Bataan, Rizal), B. S. 32866 Ramos, 5266 Topping, Vanoverbergh 620, Leiberg 6112, Reillo s. n. Mountain regions of Luzon.

HABENARIA LINGULOSA Ames Orch. 2 (1908) 33, text cut, 5 (1915) 14.

Luzon (Bontoc, Benguet), Vanoverbergh 900, 1525, Merrill 4845. Near streams in mountain regions up to 1,400 m altitude.

HABENARIA MALINTANA (Blanco) Merr. Sp. Blancoanae (1918) 112; Ames Orch. 6 (1920) 276.

Thelymitra malintana Blanco Fl. Filip. (1837) 642, ed. 2 (1845) 447, ed. 3, 3 (1879) 40; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 85.

Habenaria pelorioides Par. & Reichb. f. in Trans. Linn. Soc. 30 (1874) 135, 139, t. 27, fig. A, 1-3; Reichb. f. Otia Bot. Hamb. 1 (1878) 39; Hook. f. Fl. Brit. Ind. 6 (1890) 166; Kränzl. in Engl. Bot. Jahrb. 16 (1892) 85, Orch. Gen. et Sp. 1 (1897) 234; Grant Orch. Burma (1898) 340; Ames Orch. 2 (1908) 31, 5 (1915) 15.

Habenaria trinervis Naves Novis. App. (1882) 250, non Wight.

Luzon (Bontoc, Benguet, Nueva Vizcaya, Pangasinan, Tarlac, Zambales, Bataan, Rizal). A common species in Luzon, in the forest, up to 1,300 m altitude in Bontoc. In Burma.

HABENARIA MEARNsii Ames in Philip. Journ. Sci. 7 (1912) Bot. 3, Orch. 5 (1915) 15.

Luzon (Benguet), *B. S. 4313 Mearns, 8815, 8835 McGregor*. Pauai and Mt. Pulog, at about 2,300 m altitude.

HABENARIA MILITARIS Reichb. f. var. **PHILIPPINENSIS** Ames Orch. 2 (1908) 40, 5 (1915) 15.

Mindanao (Zamboanga, Cotabato, Davao), *Copeland 1425, 1586, Merrill 8299, Weber 214, 287*. Terrestrial in damp soil at 60 to 1,000 m altitude. The variety endemic, the species a native of CochinChina and China.

HABENARIA PONEROSTACHYS Reichb. f. in Bonpl. 3 (1855) 213; Kränzl. in Engl. Bot. Jahrb. 16 (1892) 179, Orch. Gen. et Sp. 1 (1898) 385; Ames in Philip. Journ. Sci. 7 (1912) Bot. 4, Orch. 5 (1915) 15.

Leyte, Negros, Mindanao (Camiguin Island, Cotabato), *Cuming 2095, Wenzel 0636, Merrill 7022, B. S. 14415 Ramos, Weber 200*. Terrestrial in forest at 230 to 1,600 m altitude.

HABENARIA RETICULATA Ames Orch. 7 (1922) 154.

Luzon (Benguet), *Ramos & Edaño 108*. Mt. Pulogloko, on open pine slopes, at 2,000 m altitude.

HABENARIA ROBINSONII Ames in Philip. Journ. Sci. 7 (1912) Bot. 5, Orch. 5 (1915) 15.

Luzon (Laguna, Batangas, Tayabas), *Merrill 1095, 6306, B. S. 9666 Robinson, 25541 Yates, F. B. 29030 Sulit*.

HABENARIA ROSULATA Ames in Philip. Journ. Sci. 7 (1912) Bot. 5, Orch. 5 (1915) 15.

Luzon (Tayabas), *B. S. 13203 Foxworthy & Ramos*. Quinatacutan.

HABENARIA VANOVERBERGHII Ames Orch. 5 (1915) 15.

Luzon (Bontoc), *Vanoverbergh 3687*. In wet soil, at 1,300 m altitude.

HABENARIA WARBURGIANA Kränzl. Orch. Gen. et Sp. 1 (1898) 467; Ames Orch. 5 (1915) 16.

Luzon (Ilocos Norte, Bontoc, Zambales), *Warburg s. n., Vanoverbergh 567, 8654, 3923, B. S. 32841 Ramos*. Up to 1,300 m altitude in Bontoc.

HABENARIA ZEPHYRICA Ames Orch. 5 (1915) 16.

Mindanao (Lanao), *M. S. Clemens s. n., Lake Lanao, Camp Keithley*, terrestrial.

EXCLUDED OR SYNONYMOUS SPECIES

HABENARIA AHERNII Schltr. in Engl. Bot. Jahrb. 53 (1915) 502 = *Habenaria diphylla* (Nim.) Dalz.

HABENARIA AMESIANA Ames Orch. 5 (1915) 12 = *Habenaria angustata* (Blume) Ktze.

HABENARIA CORDATA Naves Novis. App. (1882) 250; Merr. Sp. Blancoanae (1918) 21 = *Habenaria diphylla* (Nim.) Dalz.

HABENARIA CUMINGII Kränzl. Orch. Gen. et Sp. 1 (1897) 262; Ames Orch. 5 (1915) 13 = *Habenaria aristulifera* Reichb. f.

- HABENARIA GRAMINEA Lindl. apud Naves Novis. App. (1882) 251.
- HABENARIA LACERTIFERA Ames Orch. 2 (1908) 39, 5 (1915) 14, non Benth.
= *Habenaria intrudens* Ames.
- HABENARIA LUZONENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1550 =
Habenaria alagensis Ames.
- HABENARIA MURICATA Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl.
Vasc. Filip. (1886) 273; Rolfe in Journ. Bot. 24 (1886) 59;
Kränzl. in Engl. Bot. Jahrb. 16 (1892) 140, Orch. Gen. et Sp. 1
(1898) 326; Ames Orch. 1 (1905) 64, non Rodr. (1882) = *Habe-*
naria hystrix Ames.
- HABENARIA PELORIOIDES Par. & Reichb. f. in Trans. Linn. Soc. 30 (1874)
135, 139, t. 27, fig. A, 1-3; Kränzl. in Engl. Bot. Jahrb. 16
(1892) 85, Orch. Gen. et Sp. 1 (1897) 234; Ames Orch. 2 (1908)
31, 5 (1915) 15 = *Habenaria malintana* (Blanco) Merr.
- HABENARIA RUMPHII Lindl. apud Naves Novis. App. (1882) 250.
- HABENARIA THISBE Vidal Phan. Cuming. Philip. (1885) 151 = *Oberonia*
thisbe Reichb. f.
- HABENARIA TIPULIFERA Par. & Reichb. f. apud Vidal Phan. Cuming.
Philip. (1885) 151; Vidal Rev. Pl. Vasc. Filip. (1886) 273; Ames
Orch. 5 (1915) 15 = *Habenaria bisaeta* Ames.
- HABENARIA TRINERVIS Naves Novis. App. (1882) 250, non Wight = *Ha-*
benaria malintana (Blanco) Merr.

4. DISPERIS Swartz

- DISPERIS PHILIPPINENSIS Schltr. in Fedde Repert. 9 (1911) 486;
Ames Orch. 5 (1915) 17.

Luzon (Pampanga, Batangas, Sorsogon), Merrill 4215, Curran & Mer-
ritt s. n., Elmer 17355. Terrestrial at 300 to 700 m altitude.

5. THELYMITRA Blume

- THELYMITRA JAVANICA Blume Bijdr. (1825) 419; Lindl. Gen. & Sp.
Orch. (1840) 519; Blume Fl. Jav. Orch. (1858) 116 (Orch. Arch.
Ind. 137), t. 44, fig. 3, t. 48A; Miq. Fl. Ind. Bat. 3 (1859) 737; J. J.
Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 45, Fig.-Atlas (1908) fig.
25; Ames Orch. 2 (1908) 41, 5 (1915) 18.

Luzon (Bontoc, Benguet), Vanoverbergh 1357, 1476, R. S. Williams
1944, Merrill 4848, F. B. 4998 Curran, B. S. 4325 Mearns, 31929 Santos,
Ramos & Edaño 61, 63. On open slopes at 1,350 to 2,300 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

- THELYMITRA MALINTANA Blanco Fl. Filip. (1837) 642, ed. 2 (1845) 447,
ed. 3, 3 (1879) 40; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905)
85 = *Habenaria malintana* (Blanco) Merr.

6. MICROTIS R. Brown

MICROTIS UNIFOLIA (Forst. f.) Reichb. f. Beitr. Syst. Pflanzenk. (1871) 62; Kränzl. in Engl. Bot. Jahrb. 6 (1885) 55; Ames Orch. 1 (1905) 65, 2 (1908) 41, 5 (1915) 18; J. J. Sm. Orch. Jav. Fig.-Atlas (1908) fig. 26; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340.

Ophrys unifolia Forst. f. Fl. Ins. Austral. Prodr. (1786) 59.

Epipactis porrifolia Sw. in Kongl. Svensk. Vet.-Akad. Nya Handl. 21 (1800) 233.

Microtis rara R. Br. Prodr. Nov. Holl. (1810) 321; Lindl. Gen. & Sp. Orch. (1840) 396; Hook. f. Fl. Tasm. 2 (1860) 24; Reichb. f. Beitr. Syst. Pflanzenk. (1871) 22.

Microtis porrifolia R. Br. Prodr. Nov. Holl. (1810) 320, by inference only; Spreng. Syst. Veg. 3 (1826) 713; Hook. f. Fl. N. Zeal. 1 (1853) 245; Benth. Fl. Austral. 6 (1873) 347; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 47.

Microtis banksii A. Cunn. ex Hook. in Bot. Mag. 62 (1835) sub t. 3377.

Microtis pulchella Lindl. Gen. & Sp. Orch. (1840) 395; Hook. f. Fl. Tasm. 2 (1860) 24, t. 118, non R. Br.

Microtis arenaria Lindl. Gen. & Sp. Orch. (1840) 396; Hook. f. Fl. Tasm. 2 (1860) 24.

Microtis frutetorum Schlechtendal in Linnaea 20 (1847) 568; Reichb. f. in Walp. Ann. 1 (1849) 807.

Microtis javanica Reichb. f. in Bonpl. 5 (1857) 36.

Luzon (Bontoc, Lepanto, Benguet), *Vanoverbergh* 743, 1356, 1479, *R. S. Williams* 1935, *Merrill* 4345, *Topping* 97, *B. S.* 31930a *Santos*, *Ramos* & *Edaño* 98, *F. B.* 16344 *Curran*, *Merritt*, & *Zschokke*. Grassy slopes up to 1,300 m altitude in Bontoc. In Java and Australia.

7. CORYBAS Salisbury

CORYBAS MERRILLII Ames Orch. 2 (1908) 42, 5 (1915) 19; Schltr. in Fedde Repert. 19 (1923) 20.

Corysanthes merrillii Ames Orch. 1 (1905) 65, text cut.

Luzon (Bataan, Laguna), *Merrill* 3871, *Whitford* 451, *Leiberg* 6074, *B. S.* 26664 *Brown*, *Elmer* 17939. Mt. Mariveles and Mt. Maquiling, terrestrial on mossy cliffs.

8. CRYPTOSTYLIS R. Brown

CRYPTOSTYLIS ARACHNITES (Blume) Hassk. Cat. Hort. Bogor. (1844) 48; Reichb. f. in Bonpl. 5 (1857) 36; Blume Fl. Jav. Orch. (1858) 112 (Orch. Arch. Ind. 133), t. 45, fig. 2; Hook. f. Fl. Brit. Ind. 6 (1890) 118; Ridl. in Journ. Linn. Soc. 32 (1896) 407; Trim. Handb. Fl. Ceyl. 4 (1898) 209; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 59, Fig.-Atlas (1908) fig. 37; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 225; Ames Philip. Journ. Sci. 2 (1907) Bot. 313, Orch. 2 (1908) 44, 5 (1915) 19, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 139.

Zosterostylis arachnites Blume Bijdr. (1825) 419, fig. 32; Lindl. Gen. & Sp. Orch. (1840) 446; Miq. Fl. Ind. Bat. 3 (1859) 721; Hook. in Bot. Mag. 89 (1863) t. 5381.

Zosterostylis zeylanica Lindl. Gen. & Sp. Orch. (1840) 446; Thw. Enum. Pl. Zeyl. (1864) 312.

Zosterostylis walkerae Wight Icon. 5¹ (1852) 20, t. 1748 IV.

Cryptostylis philippinensis Schltr. in Engl. Bot. Jahrb. 58 (1922) 54.

Luzon (Apayao, Rizal, Tayabas, Sorsogon), Polillo, Mindoro, Biliran, Leyte, Negros (Occidental Negros), Mindanao (Camiguin Island, Davao). A common, variable, and widely distributed species in the Philippines. Terrestrial, in forest at 30 to 1,000 m altitude. In India, Ceylon, Malay Peninsula, Java, and Borneo (Sarawak).

J. J. Smith, in Bull. Jard. Bot. Buitenz. III 3 (1921) 239, expresses doubt as to *B. S. 20462* being referable to *Cryptostylis arachnites*. A re-examination of all available specimens from the Philippines indicates that there is only one species present. I cannot separate this from *C. arachnites*. It is probable that Smith in studies of living material has been able to detect differences among the species of *Cryptostylis* allied with *C. arachnites* which are evanescent in herbarium specimens. *Cryptostylis arachnites* is a very variable species, flowers on the same raceme exhibiting a wide range of variation in the labellum.

EXCLUDED OR SYNONYMOUS SPECIES

CRYPTOSTYLIS PHILIPPINENSIS Schltr. in Engl. Bot. Jahrb. 58 (1922) 54 = *Cryptostylis arachnites* (Blume) Hassk.

9. GALEOLA Loureiro

GALEOLA ALTISSIMA (Blume) Reichb. f. Xen. Orch. 2 (1865) 77; Hook. f. Fl. Brit. Ind. 6 (1890) 98; Grant Orch. Burma (1895) 349; Ridl. in Journ. Linn. Soc. 32 (1896) 395; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 67, Fig.-Atlas (1908) fig. 44; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 202; Ames in Merr. in Journ. Roy. Asiatic Soc. Straits Branch, Special No. (1921) 139.

Cyrtosia altissima Blume Bijdr. (1825) 396, Rumphia 1 (1835) t. 70; Lindl. Gen. & Sp. Orch. (1830) 26.

Erythrorchis altissima Blume Rumphia 1 (1835) 200; Lindl. Gen. & Sp. Orch. (1840) 438, p. p.; Reichb. f. in Walp. Ann. 3 (1852) 602; Miq. Fl. Ind. Bat. 3 (1859) 720.

Haematorchis altissima Blume Rumphia 4 (1848) t. 200 B.

Luzon (Cagayan), *F. B. 17812* Curran, *B. S. 13985* Ramos. In Malay Peninsula, Burma, and Java.

GALEOLA KUHLLII Reichb. f. Xen. Orch. 2 (1865) 78; Naves Novis. App. (1882) 247.

Vanilla pterosperma Lindl. in Wall. Cat. (1832) No. 7402, *nomen*.

Erythrorchis altissima Lindl. Gen. & Sp. Orch. (1840) 438, p. p., non Blume.

Erythrorchis kuhlii Reichb. f. Xen. Orch. 2 (1862) t. 119.

Galeola hydra Reichb. f. Xen. Orch. 2 (1865) 77; Hook. f. Fl. Brit. Ind. 6 (1890) 89; Grant Orch. Burma (1895) 350; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 264, t. 351; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 68, Fig.-Atlas

(1908) *fig. 45*; Ames in Philip. Journ. Sci. 2 (1907) Bot. 313; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 202; Ames Orch. 2 (1908) 46.

Galeola pterosperma Schltr. in Engl. Bot. Jahrb. 45 (1910) 386; J. J. Sm. in Bull. Jard. Bot. Buitenz. II 9 (1913) 10; Ames Orch. 5 (1915) 19.

Babuyan Islands (Camiguin Island), Luzon (Rizal, Laguna, Tayabas, Sorsogon), Mindoro, Samar, Panay (Capiz), Leyte, Mindanao (Lanao, Davao). A common and widely distributed species in the Philippines at 75 to 180 m altitude. In Sikkim, Himalaya, Burma, Malay Peninsula, Java, and Sumatra.

The valid name of this species has been a source of difference of opinion. It has been most widely known as *Galeola hydra* Reichb. f., but in 1910 Schlechter proposed the combination *Galeola pterosperma* (Lindl.) Schltr. In doing so he overlooked the fact that the name-bringing synonym *Vanilla pterosperma* Lindl. is a *nomen nudum* and has no standing. The earliest valid name is *Galeola kuhlîi* Reichb. f., published in Xen. Orch. 2 (1865) 78, in which Reichenbach cites, in synonymy, the plate published prior to the description [Xen. Orch. 2 (1862) *t. 119*]. This plate was published as *Erythrorchis kuhlîi* and has full analytical drawings. According to the international rules adopted at Vienna in 1905 (Sect. 4, Art. 37) this plate constitutes valid publication.

GALEOLA PHILIPPINENSIS Ames Sched. Orch. 6 (1923) 5.

Luzon (Sorsogon), *Elmer 16122*.

EXCLUDED OR SYNONYMOUS SPECIES

GALEOLA HYDRA Reichb. f. Xen. Orch. 2 (1865) 77; Ames in Philip. Journ. Sci. 2 (1907) Bot. 313, Orch. 2 (1908) 46 = *Galeola kuhlîi* Reichb. f.

GALEOLA PTEROSPERMA Schltr. in Engl. Bot. Jahrb. 45 (1910) 386; Ames Orch. 5 (1915) 19 = *Galeola kuhlîi* Reichb. f.

10. VANILLA Swartz

VANILLA CALOPOGON Reichb. f. in Otia Bot. Hamb. 1 (1878) 40; Rolfe in Journ. Linn. Soc. 32 (1896) 459; Ames Orch. 5 (1915) 21, 6 (1920) 277.

Luzon (Ilocos Norte), *Cuming 2070, Loher 630, B. S. 32928 Ramos*.

VANILLA OVALIS Blanco Fl. Filip. ed. 2 (1845) 448, ed. 3, 3 (1879) 42; Ames Orch. 5 (1915) 21; Merr. Sp. Blancoanae (1918) 112.

Epidendrum vanilla Blanco Fl. Filip. (1837) 643, non Linn.

Vanilla majaijensis Blanco Fl. Filip. ed. 2 (1845) 593, ed. 3, 3 (1879) 43.

Vanilla planifolia Ait. apud Naves Novis. App. (1882) 248.

Vanilla aromatica Sw. apud Naves Novis. App. (1882) 248.

Vanilla philippinensis Rolfe in Journ. Linn. Soc. 32 (1896) 459; Ames Orch. 2 (1908) 47.

Luzon (Bulacan, Rizal, Laguna, Tayabas, Sorsogon), Sibuyan, Biliran, Mindanao (Dapitan, Lanao, Butuan), *Cuming 2132, Merrill Sp. Blancoanae 665, Holman s. n., Serrato s. n., Elmer 9259, 12534, 16035, F. B. 4585 Mearns & Hutchinson, 28614 Sulit, B. S. 19346 Wester, 18952 McGregor, M. S. Clemens s. n.*

EXCLUDED OR SYNONYMOUS SPECIES

- VANILLA ALBIDA Blume apud Naves Novis. App. (1882) 248.
- VANILLA APHYLLA Blume apud Naves Novis. App. (1882) 248.
- VANILLA AROMATICA Sw. apud Naves Novis. App. (1882) 248 = *Vanilla ovalis* Blanco.
- VANILLA PHILIPPINENSIS Rolfe in Journ. Linn. Soc. 32 (1896) 459; Ames Orch. 2 (1908) 47 = *Vanilla ovalis* Blanco.
- VANILLA PLANIFOLIA Ait. apud Naves Novis. App. (1882) 248 = *Vanilla ovalis* Blanco.
- VANILLA MAJAJIENSIS Blanco Fl. Filip. ed. 2 (1845) 593, ed. 3, 3 (1879) 43 = *Vanilla ovalis* Blanco.
- VANILLA SPECIOSA Boxall ex Vidal ex Naves Novis. App. (1882) 248, *nomen*.

11. LECANORCHIS Blume

- LECANORCHIS JAVANICA Blume Mus. Bot. Lugd.-Bat. 2 (1856) 188, Fl. Jav. Orch. (1858) 150 (Orch. Arch. Ind. 178), t. 63, fig. 1, t. 66A; Miq. Fl. Ind. Bat. 3 (1859) 718; J. J. Sm. in Nova Guinea 8 (1909) Bot. 136, 12 (1913) 7, 12³ (1915) 185, in Bull. Dept. Agr. Ind. Néerl. 43 (1910) 1; Ames Orch. 6 (1920) 277.
- Lecanorchis triloba* J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 19 (1908) 26, in Nova Guinea 8 (1909) Bot. 10, t. 3, fig. 9; Ames Orch. 5 (1915) 21.

Luzon (Laguna, Sorsogon), Leyte, Mindanao (Agusan), B. S. 22961 McGregor, Elmer 16369, Wenzel 0120, 0282, 0851, Weber 71. Terrestrial, in damp forest, at 150 to 320 m altitude. In Java and New Guinea.

J. J. Smith, in Nova Guinea 12³ (1915) 185, suggests that *Lecanorchis papuana* Schltr. is the same as this species.

EXCLUDED OR SYNONYMOUS SPECIES

- LECANORCHIS TRILOBA J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 19 (1908) 26; Ames Orch. 5 (1915) 21 = *Lecanorchis javanica* Blume.

12. APHYLLORCHIS Blume

- APHYLLORCHIS BENGUETENSIS Ames Orch. 2 (1908) 49, *text cut*, 5 (1915) 22.

Luzon (Benguet, Laguna), Mindanao (Agusan, Davao), F. B. 5086 Curran, B. S. 16565 Ramos, Serrato s. n., Elmer 11848, 13841.

- APHYLLORCHIS HALCONENSIS Ames Sched. Orch. 6 (1923) 7.

Mindoro, Ramos & Edaña s. n. Terrestrial at 1,000 m altitude.

- APHYLLORCHIS PALLIDA Blume Bijdr. (1825) fig. 77 (without description); Lindl. Gen. & Sp. Orch. (1840) 536 (mention only); Blume Mus. Bot. Lugd.-Bat. 1 (1849) 30, Fl. Jav. Orch. (1858) 43 (Orch. Arch. Ind. 52), t. 13, fig. 1 and t. 17D; Miq. Fl. Ind. Bat. 3 (1859) 721; Hook. f. Fl. Brit. Ind. 6 (1890) 117; Ridl. in Journ. Linn. Soc. 32 (1896) 405; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 71, Fig.-Atlas (1908) fig. 47; Ridl. Mat. Fl. Mal. Penin.

1 (1907) 205; Ames in Philip. Journ. Sci. 2 (1907) Bot. 313, Orch. 2 (1908) 50, 5 (1915) 22, 6 (1920) 11, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 139.

Mindoro, Merrill 5810, 5811. Terrestrial in humid forest, Mt. Halcon and vicinity, at 330 to 930 m altitude. In Malay Peninsula, Java, and Borneo.

13. EPIPOGUM Gmelin

EPIPOGUM ROSEUM (D. Don) Lindl. in Journ. Linn. Soc. 1 (1857) 177; Ames Orch. 2 (1908) 48, 5 (1915) 22.

Limodorum roseum D. Don Prodr. Fl. Nepal. (February, 1825) 30.

Galera nutans Blume Bijdr. (July, 1825) 416, fig. 3; Lindl. Gen. & Sp. Orch. (1840) 500; Blume Mus. Bot. Lugd.-Bat. 2 (1856) 187, Fl. Jav. Orch. (1858) 117 (Orch. Arch. Ind. 139) t. 52, fig. 3, t. 54E.

Ceratopsis rosea Lindl. Gen. & Sp. Orch. (1840) 384.

Podanthera pallida Wight Icon. 5¹ (1852) 22, t. 1759.

Galera rosea Blume Mus. Bot. Lugd.-Bat. 2 (1856) 188, Fl. Jav. Orch. (1858) 117 (Orch. Arch. Ind. 139).

Galera pallida Blume Fl. Jav. Orch. (1858) 117 (Orch. Arch. Ind. 139), in obs.

Epipogum nutans Reichb. f. in Bonpl. 5 (February, 1857) 36; Lindl. in Journ. Linn. Soc. 1 (1857) 177; Miq. Fl. Ind. Bat. 3 (1859) 718; Thw. Enum. Pl. Zeyl. (1864) 311; Benth. Fl. Austral. 6 (1873) 308; Hook. f. Fl. Brit. Ind. 6 (1890) 124; Fitzger. Austral. Orch. 2⁵ (1896) t.; Rolfe in Dyer. Fl. Trop. Afr. 7 (1897) 188; Trim. Handb. Fl. Ceyl. 4 (1898) 222; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 253, t. 335; Ames Orch. 1 (1905) 68; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 61, Fig.-Atlas (1908) fig. 39; Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 37; J. J. Sm. in Nova Guinea 12³ (1915) 185.

Epipogum guilfoylilii F. v. Muell. Fragm. 8 (1873) 30, comb. non faciet, ex Benth. Fl. Austral. 6 (1873) 308, in syn.

Luzon (Nueva Vizcaya, Laguna), Mindoro, Leyte, Negros (Oriental Negros), Mindanao (Lanao, Davao), Merrill 231, Copeland 1241, 5623, Gates 6317, B. S. 17915 Brown, Wenzel 0284, 0289, 0427, Elmer 9467, 10536, M. S. Clemens s. n., Weber 277, Ramos s. n. Terrestrial in forest, at 500 to 700 m altitude. In India, Ceylon, Java, New Guinea, Australia, and Africa.

EXCLUDED OR SYNONYMOUS SPECIES

EPIPOGUM NUTANS Reichb. f. in Bonpl. 5 (February, 1857) 36; Ames Orch. 1 (1905) 68 = *Epipogum roseum* (D. Don) Lindl.

14. NERVILIA Commerson ex Gaudichaud

NERVILIA ARAGOANA Gaudich. in Freyc. Voy. Uranie et Physic. Bot. (1829) 422, t. 35; Schltr. in Engl. Bot. Jahrb. 45 (1911) 405 (excl. syn. *Pogonia gracilis* Blume), in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 3, in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 43; J. J. Sm. in Nova Guinea 12³ (1915) 182; Ames Orch. 5 (1915) 23.

Pogonia flabelliformis Lindl. in Wall. Cat. (1832) No. 7400, *nomen*, Gen. & Sp. Orch. (1840) 415; Miq. Fl. Ind. Bat. 3 (1859) 715; Naves Novis. App. (1882) 248; Hook. f. Fl. Brit. Ind. 6 (1890) 121; Grant Orch. Burma (1895) 353; Ridl. in Journ. Linn. Soc. 32 (1896) 408; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 57, Fig.-Atlas (1908) fig. 36, Orch. Amb. (1905) 10; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 158, t. 125; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 203.

Pogonia carinata Wight Icon. 5¹ (1852) 15, t. 1720, non Lindl.

Luzon (Zambales, Rizal, Laguna), Mindanao (Lanao), *Sandkuhl s. n.*, B. S. 22683 Ramos, 17918 Brown, M. S. Clemens s. n., 546. Terrestrial, in forest. In India, Burma, Malay Peninsula, Java, Sumatra, and New Guinea, and said to extend to Samoa.

Pogonia nervilia Blume is commonly referred to this species as a synonym, but J. J. Smith [in Nova Guinea 8 (1809) 10] treats it as a valid species. Further study is needed to determine its status.

Pogonia gracilis Blume is treated by J. J. Smith [Fl. Buitenz. 6 (Orch. Jav.) (1905) 58] as a distinct species, but in his notes Smith states that he regards it as a weak form of *Pogonia flabelliformis* Lindl. which is conspecific with *Nervilia aragoana* Gaudich.

NERVILIA CRISPATA (Blume) Schltr. in K. Schum. & Lauterb. Nachtr. Fl. Deutsch. Schutzgeb. (1905) 82, in nota, in Engl. Bot. Jahrb. 45 (1911) 402, in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 38; Ames Orch. 5 (1915) 23.

Pogonia crispata Blume Mus. Bot. Lugd.-Bat. 1 (1849) 32, Fl. Jav. Orch. (1858) 127 (Orch. Arch. Ind. 151), t. 52, fig. 4A-C; Miq. Fl. Ind. Bat. 3 (1859) 716; Reichb. f. Xen. Orch. 2 (1863) t. 126, V, VI, 10, (1865) 87; Naves Novis. App. (1882) 248; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 56, Fig.-Atlas (1908) fig. 35.

Bolborchis javanica Zoll. & Mor. Syst. Verz. Java (1845-46) 89; Reichb. f. in Walp. Ann. 1 (1849) 775.

Coelogyne javanica Lindl. Fol. Orch. Coelog. (1854) 17.

Nervilia fimbriata Schltr. in K. Schum. & Lauterb. Nachtr. Fl. Deutsch. Schutzgeb. (1905) 82.

Palawan, Merrill 9288. Taytay, in forests on dry slopes. In Java and German New Guinea.

Pogonia prainiana King & Pantl. has been referred to this species, which would extend the range to India, if this reduction is correct.

NERVILIA DILATATA (Blume) Schltr. in Engl. Bot. Jahrb. 45 (1911) 402; Ames Orch. 5 (1915) 23, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 140.

Pogonia dilatata Blume Fl. Jav. Orch. (1858) 127 (Orch. Arch. Ind. 151), t. 10, fig. 4, 1 and 2; Ames Orch. 1 (1905) 66.

Luzon (Benguet), Bancalan, Elmer 6259, Merrill 7741, Weber 1574. Grassy hillsides at about 1,600 m altitude. In Borneo.

Although the determination has been carefully made, there is room for uncertainty as to the identity of the Philippine material with the Bornean species. In twenty years but three incomplete specimens have been obtained.

NERVILIA DISCOLOR (Blume) Schltr. in Engl. Bot. Jahrb. 45 (1911) 403; Ames Orch. 5 (1915) 23.

Cordyla discolor Blume Bijdr. (1825) 417.

Rophostemon discolor Blume Fl. Jav. Praef. (1828) VI; Lindl. Gen. & Sp. Orch. (1840) 453.

Pogonia discolor Blume Mus. Bot. Lugd.-Bat. 1 (1849) 32, Fl. Jav. Orch. (1858) 128 (Orch. Arch. Ind. 152.), t. 57, fig. 1; Miq. Fl. Ind. Bat. 3 (1859) 716; Hook. f. in Bot. Mag. 100 (1874) t. 6125; Naves Novis. App. (1882) 248.

Pogonia pudica Ames Orch. 2 (1908) 44, text cut.

Pogonia plicata Ames Orch. 2 (1908) 43, non Lindl.

Luzon (Bontoc, Laguna, Batangas), Panay (Antique), Mindanao (Davao), *Vanoverbergh* 459, 1201, 1460, *Elmer* s. n., 8180, 9346, *B. S.* 17917 *Brown*, *F. B.* 21501 *Tamesis*, 21331 *Foxworthy & Catalan*, *Copeland* 427. *R. S. Williams* 3088, *McLean* s. n., *McGregor* 5748. Grassy hillsides, at 350 to 1,240 m altitude. In Java.

15. STEREOSANDRA Blume

STEREOSANDRA JAVANICA Blume Mus. Bot. Lugd.-Bat. 2 (1856) 176, Fl. Jav. Orch. (1858) 27 (Orch. Arch. Ind. 31), t. 10, fig. 3, t. 11G; Miq. Fl. Ind. Bat. 3 (1859) 717; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 63, Fig.-Atlas (1908) fig. 41; Ames Orch. 1 (1905) 67, 2 (1908) 46, 5 (1915) 24; Schltr. in Engl. Bot. Jahrb. 45 (1911) 406.

Luzon (Benguet, Bataan, Rizal, Sorsogon), Mindanao (Zamboanga, Agusan), *Elmer* 6458, 17302, *Merrill* s. n., *B. S.* 12124 *Ramos*, 38349 *Brown*, *Weber* 300. In Java.

16. DIDYMOPLEXIS Griffith

DIDYMOPLEXIS PALLENS Griff. in Calc. Journ. Nat. Hist. 4 (1844) 383, t. 17; Kurz in Journ. Bot. 4 (1866) 40 (excl. syns. *Epiphanes* and *Gastrodia*); Hemsl. in Journ. Linn. Soc. 20 (1884) 308; Hook. f. Fl. Brit. Ind. 6 (1890) 122; Ridl. in Journ. Linn. Soc. 32 (1896) 408; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 260, t. 346; J. J. Sm. in Icon. Bogor. 2 (1903) 5, t. 101B, Fl. Buitenz. 6 (Orch. Jav.) (1905) 77, Fig.-Atlas (1908) t. 51; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 206; Ames Orch. 2 (1908) 51, 5 (1915) 24; Schltr. in Engl. Bot. Jahrb. 45 (1911) 408; Merr. Fl. Manila (1912) 165.

? *Leucorchis sylvatica* Blume Mus. Bot. Lugd.-Bat. 1 (1849) 31, Fl. Jav. Orch. (1858) 124 (Orch. Arch. Ind. 147) text cut; Miq. Fl. Ind. Bat. 3 (1859) 722.

? *Arethusa cristata* Griff. Notul. 3 (1851) 378, Icon. Pl. Asiat. 3 (1851) 343, 344.

? *Apetalon minutum* Wight Icon. 5¹ (1852) 22, t. 1758.

Epiphanes pallens Reichb. f. in Seem. Fl. Vit. (1868) 266, in nota.

? *Arethusa bengalensis* Hort. Bot. Calc. ex Hemsl. in Journ. Linn. Soc. 20 (1884) 311, in syn.

Luzon (Pampanga, Rizal, Laguna), *Bolster* 27, *Loher* 6002, *F. B.* 21333 *Foxworthy & Catalan*. Saprophyte, near brooks. In India, Malay Peninsula, and Java.

DIDYMOPLEXIS PHILIPPINENSIS Ames Orch. 5 (1915) 24.

Luzon (Camarines), Palawan, Leyte, *B. S. 22083* Ramos, Merrill 9409, Wenzel 0362. In forest at 6 to 300 m altitude.

17. **GASTRODIA** R. Brown

GASTRODIA JAVANICA (Blume) Lindl. Gen. & Sp. Orch. (1840) 384; Blume Mus. Bot. Lugd.-Bat. 2 (1856) 175, Fl. Jav. Orch. (1858) 121 (Orch. Arch. Ind. 145) *t. 52, fig. 1*; Miq. Fl. Ind. Bat. 3 (1859) 717; Thw. Enum. Pl. Zeyl. (1864) 311; Naves Novis. App. (1882) 247; Hook. f. Fl. Brit. Ind. 6 (1890) 123; Trim. Handb. Fl. Ceyl. 4 (1898) 221; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 75, Fig.-Atlas (1908) *fig. 50*; Schltr. in Engl. Bot. Jahrb. 45 (1911) 410; Ames in Philip. Journ. Sci. 7 (1912) Bot. 6, Orch. 5 (1915) 25.

Epiphanes javanica Blume Bijdr. (1825) 421, *fig. 4*.

Luzon (Pampanga, Tayabas), Mindoro, Palawan, Leyte, Mindanao (Davao), Merrill 4211, 7233, Elmer 7276, Weber 274, *B. S. 28718* Ramos & Edaño, Ramos s. n. Terrestrial, apparently more general in mountainous regions. In Ceylon (?) and Java.

18. **SPIRANTHES** L. C. Richard

SPIRANTHES SINENSIS (Pers.) Ames Orch. 2 (1908) 53, 5 (1915) 26, 6 (1920) 11; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 160, in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 357; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 140.

Ophrys spiralis Georgi Bemerk. Reise Russ. Reich. 1 (1775) 232, non Linn.

Aristotelea spiralis Lour. Fl. Cochinch. 2 (1790) 522, non *Spiranthes spiralis* Koch.

Epidendrum aristotelia Raeusch. Nomencl. Bot. ed. 3 (1797) 265, *nomen*.

Neottia spiralis δ Willd. Sp. Pl. 4 (1805) 74.

Neottia sinensis Pers. Syn. Pl. 2 (1807) 511; Poir. in Lam. Encyc. Suppl. 4 (1816) 85.

Neottia australis R. Br. Prodr. Nov. Holl. (1810) 319; Poir. in Lam. Encyc. Suppl. 4 (1816) 86; D. Don Prodr. Fl. Nepal (1825) 27.

Neottia flexuosa Sm. in Rees. Cycl. 24 (1819) No. 9.

Neottia parviflora Sm. in Rees Cycl. 24 (1819) No. 10, non Blume.

Neottia amoena Bieb. Fl. Taur. Cauc. 3 (1819) 606.

Spiranthes pudica Lindl. Collect. Bot. (1821) *t. 30*, in Bot. Reg. 10 (1824) sub *t. 823*; Spreng. Syst. Veg. 3 (1826) 708, p. p.

Neottia australis R. Br. β *chinensis* Lindl. in Bot. Reg. 7 (1822) *t. 602*.

Spiranthes australis Lindl. in Bot. Reg. 10 (1824) sub *t. 823*; Spreng. Syst. Veg. 3 (1826) 708; Lindl. Gen. & Sp. Orch. (1840) 464 (excl. syns. *Neottia strateumatica* R. Br., *Spiranthes strateumatica* Lindl., and *Orchis strateumatica* Linn.); A. Rich. in Ann. Sci. Nat.

- II 15 (1841) 78, Monogr. Orch. Nil-Gher. (1841) 10, 30; Reichb. f. Icon. Fl. Germ. 13-14 (Orch. Eur.) (1851) 152, *t.* 124 (476); Wight Icon. 5¹ (1851) 15, *t.* 1724; Hook. f. Fl. Tasm. 2 (1855) 15; Lindl. in Journ. Linn. Soc. 1 (1857) 178; Miq. Fl. Ind. Bat. 3 (1859) 722; Benth. Fl. Hongk. (1861) 360; Miq. Sumatra (1862) 275; Thw. Enum. Pl. Zeyl. (1864) 312; Hook. f. Handb. N. Zeal. Fl. (1864) 272; Miq. Prol. Fl. Jap. (1866-67) 141; Benth. Fl. Austral. 6 (1873) 314; Franch. & Sav. Enum. Pl. Jap. 2 (1879) 37; Fitzger. Austral. Orch. 1² (1880) *t.* 1; Naves Novis. App. (1882) 249; Kränzl. in Engl. Bot. Jahrb. 6 (1885) 55; Ridl. in Journ. Linn. Soc. 24 (1888) 392; Hook. f. Fl. Brit. Ind. 6 (1890) 102; Kränzl. in Notizbl. Bot. Gart. Berl. 2 (1898) 105; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik-Himal.) (1898) 278, *t.* 369; Trim. Handb. Ceyl. Fl. 4 (1898) 217; Palibin Consp. Fl. Korea pt. 3 (1901) 4 [in Act. Hort. Petrop. 19 (1901) 104]; Kränzl. in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 240; Ames Orch. 1 (1905) 69; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 81, Fig.-Atlas (1908) *fig.* 55; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 163; Nakai in Journ. Coll. Sci. Imp. Univ. Tokyo 31 (Fl. Korea 2) (1911) 225.
- Spiranthes parviflora* Lindl. in Bot. Reg. 10 (1824) sub *t.* 823.
- Spiranthes flexuosa* Lindl. in Bot. Reg. 10 (1824) sub *t.* 823; Spreng. Syst. Veg. 3 (1826) 708; Schltr. in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 352.
- Neottia crispata* Blume Bijdr. (1825) 406.
- Spiranthes amoena* Spreng. Syst. Veg. 3 (1826) 708 (excl. syn. *Spiranthes congesta* Lindl.); Lindl. in Wall. Cat. (1832) No. 7377; Bunge Enum. Pl. Chin. (1831) 63; Ledeb. Fl. Alt. 4 (1833) 173; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 160, in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 349.
- Neottia pudica* Sweet Hort. Brit. ed. 2 (1830) 485.
- Spiranthes wightiana* Lindl. in Wall. Cat. (1832) No. 7378, *nomen*; Schltr. in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 359.
- Calanthe australis* Ait. ex Loud. Hort. Brit. Suppl. 2 (1839) 615, *nomen*.
- Spiranthes longispicata* A. Rich. in Ann. Sci. Nat. II 15 (1841) 78, Monogr. Orch. Nil-Gher. (1841) 10, 30.
- Spiranthes densa* A. Rich. in Ann. Sci. Nat. II 15 (1841) 78, Monogr. Orch. Nil-Gher. (1841) 10, 31.
- Spiranthes indica* Steud. Nomencl. Bot. ed. 2, 2 (1841) 625.
- Spiranthes crispata* Zoll. & Mor. Syst. Verz. Java (1845-46) 89; Schltr. in Beihefte Bot. Centralb. 37, Abt. 2 (1920) 352.
- Spiranthes* Griff. Notul. 3 (1851) 384, Icon. Pl. Asiat. 3 (1851) *t.* 348.
- Spiranthes novaezealandiae* Hook. f. Fl. N. Zeal. 1 (1853) 243; Schltr. in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 355; Ames Orch. 7 (1922) 128, in obs.
- Spiranthes australis* Lindl. var. *pudica* Lindl. in Hook. Kew Journ. Bot. 7 (1855) 38.

- Spiranthes stylites* Lindl. in Journ. Linn. Soc. 1 (1857) 178; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 160, in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 358.
- Gyrostachys australis* Blume Fl. Jav. Orch. (1858) 108 (Orch. Arch. Ind. 129).
- Gyrostachys australis* Blume var. *amoena* Blume Fl. Jav. Orch. (1858) 108 (Orch. Arch. Ind. 129).
- Gyrostachys australis* Blume var. *wightiana* Blume Fl. Jav. Orch. (1858) 108 (Orch. Arch. Ind. 129).
- Gyrostachys australis* Blume var. *flexuosa* Blume Fl. Jav. Orch. (1858) 109 (Orch. Arch. Ind. 130) t. 38, fig. 2.
- Gyrostachys australis* Blume var. *parviflora* Blume Fl. Jav. Orch. (1858) 109 (Orch. Arch. Ind. 130).
- Gyrostachys australis* Blume var. *crispata* Blume Fl. Jav. Orch. (1858) 109 (Orch. Arch. Ind. 130) t. 37, fig. 3.
- Gyrostachys amoena* Blume Fl. Jav. Orch. (1858) 108 (Orch. Arch. Ind. 129), text cut only.
- Spiranthes australis* Lindl. var. *wightiana* Lindl. ex Hook. f. Fl. Brit. Ind. 6 (1890) 102.
- Gyrostachys stylites* Ktze. Rev. Gen. Pl. 2 (1891) 664.
- Spiranthes chinensis* Ames apud Schltr. in Notes Roy. Bot. Gard. Edinb. 24 (1912) 105.
- Spiranthes australis* Lindl. var. *suishanensis* Hayata Icon. Fl. Formosa 6 (1916) 86.
- Spiranthes suishanensis* Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 161.
- Gyrostachys crispata* Blume apud Schltr. in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 352, in syn.
- Gyrostachys flexuosa* Blume apud Schltr. in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 352, in syn.
- Gyrostachys parviflora* Blume apud Schltr. in Beihefte Bot. Centralbl. 37, Abt. 2 (1920) 352, in syn.

Luzon (Bontoc, Benguet), Mindanao (Agusan), Vanoverbergh 1269, 3668, 3922, Elmer 6623, R. S. Williams 1923, Merrill 4344, F. B. 4951 Curran, Weber 80. Terrestrial, at 360 to 1,500 m altitude. In India, Afghanistan, Ceylon, China, Formosa, Japan, Manchuria, Siberia, Java, Sumatra, Borneo, Australia, Tasmania, and New Zealand.

Spiranthes novaezelandiae Hook. f. does not appear to be a distinct species, but is probably a good geographic variety of *Spiranthes sinensis* differing in the structure of the column.

EXCLUDED OR SYNONYMOUS SPECIES

SPIRANTHES AUSTRALIS Lindl. in Bot. Reg. 10 (1824) sub t. 823; Naves Novis. App. (1882) 249; Ames Orch. 1 (1905) 69, *Spiranthes sinensis* (Pers.) Ames.

SPIRANTHES PUSILLA Miq. apud Naves Novis. App. (1882) 249.

Possibly a form of *Spiranthes sinensis* (Pers.) Ames.

19. *EPIPACTIS* (Haller) Boehmer, non Zinn *

EPIPACTES CLAUSA A. A. Eaton ex Ames Orch. 3 (1908) 41, *t.* 38;
Ames in Philip. Journ. Sci. 4 (1909) Bot. 663.

Goodyera clausa Schltr. in Engl. Bot. Jahrb. 45 (1911) 392; Ames
Orch. 5 (1915) 26.

Luzon (Lepanto, Benguet, Tayabas), Negros (Occidental Negros),
Merrill 4591, 7759, Elmer 8052, Banks *s. n.*

EPIPACTIS ELMERI Ames Sched. Orch. 6 (1923) 12, *fig.* 3.

Luzon (Sorsogon), Elmer 14884.

EPIPACTIS LUZONENSIS Ames comb. nov.

Goodyera luzonensis Ames Orch. 5 (1915) 26.

Luzon (Tayabas), B. S. 2465 Foxworthy, Merrill 7488. Mt. Banahao at
about 2,000 m altitude.

EPIPACTIS PHILIPPINENSIS Ames Orch. 2 (1908) 60, *text cut.*

Goodyera philippinensis Schltr. in Engl. Bot. Jahrb. 45 (1911) 392;
Ames Orch. 5 (1915) 27.

Luzon (Bontoc, Benguet, Zambales, Bataan, Cavite, Laguna), Mindoro.
Reasonably common in Luzon at 100 to 1,100 m altitude.

EPIPACTIS RAMOSII Ames Sched. Orch. 6 (1923) 14.

Goodyera ramosii Ames in Philip. Journ. Sci. 8 (1913) Bot. 409, Orch.
5 (1915) 27.

Mindanao, (Camiguin Island), B. S. 14421, 14446 Ramos. On a dead
tree, in mossy forest.

EPIPACTIS RUBICUNDA (Blume) A. A. Eaton in Proc. Biol. Soc. Wash.
21 (1908) 65; Ames in Philip. Journ. Sci. 4 (1909) Bot. 664.

Neottia rubicunda Blume Bijdr. (1825) 408.

Neottia grandis Blume Bijdr. (1825) 407; Lindl. Gen. & Sp. Orch.
(1840) 494.

Goodyera rubicunda Lindl. in Bot. Reg. 25 (1839) Misc. 61, Gen. &
Sp. Orch. (1840) 493, in Journ. Linn. Soc. 1 (1857) 184; Miq. Fl.
Ind. Bat. 3 (1859) 728; Naves Novis. App. (1882) 248; J. J. Sm.
Fl. Buitenz. 6 (Orch. Jav.) (1905) 121, Fig.-Atlas (1909) *fig.* 88;
Ames Orch. 5 (1915) 27.

Speiranthes grandis Hassk. Cat. Pl. Bogor. (1844) 47.

Goodyera rubens Blume Fl. Jav. Orch. (1858) 36 (Orch. Arch. Ind.
43) *t.* 9c, *figs.* 17-26, *t.* 9d, *fig.* 1; Hook. f. Fl. Brit. Ind. 6 (1890)
111; Ridl. in Journ. Linn. Soc. 32 (1896) 403, Mat. Fl. Mal. Penin.
1 (1907) 219.

Goodyera grandis Blume Fl. Jav. Orch. (1858) 36 (Orch. Arch. Ind.
43); Miq. Fl. Ind. Bat. 3 (1859) 728; Naves Novis. App. (1882)
248.

* Zinn's *Epipactis* as to apparent source of name is this, but as to majority
of species given is *Serapias* Linn.

Goodyera celebica Blume Fl. Jav. Orch. (1858) 36 (Orch. Arch. Ind. 43); Miq. Fl. Ind. Bat. 3 (1859) 728.

Orchiodes rubicundum Ktze. Rev. Gen. Pl. 2 (1891) 675.

Orchiodes celebicum Ktze. Rev. Gen. Pl. 2 (1891) 675.

Orchiodes grande Ktze. Rev. Gen. Pl. 2 (1891) 675.

Luzon (Laguna, Sorsogon), Panay (Capiz), Negros (Oriental Negros), Mindanao (Lanao, Cotabato, Davao); *Elmer* 10173, 11150, 14867, *B. S.* 31096, 31455 *Ramos & Edaña*, 6049 *Robinson*, *Weber* 201, *Cuming* s. n., *M. S. Clemens* s. n. Terrestrial in damp forest at 110 to 260 m altitude. In Malay Peninsula and Java.

EPIPACTIS VIRIDIFLORA (Blume) Ames Orch. 2 (1908) 61.

Neottia viridiflora Blume Bijdr. (1825) 408; Lindl. Gen. & Sp. Orch. (1840) 494.

Physurus viridiflorus Lindl. in Journ. Linn. Soc. 1 (1857) 180; Miq. Fl. Ind. Bat. 3 (1859) 734.

Goodyera viridiflora Blume Fl. Jav. Orch. (1858) 34 (Orch. Arch. Ind. 41), *t. 9c, fig. 2, 12-16*; Ames Orch. 5 (1915) 27.

Orchiodes viridiflorum Ktze. Rev. Gen. Pl. 2 (1891) 675.

Luzon (Benguet, Bataan), Negros (Oriental Negros), Mindanao (Davao), *R. S. Williams* 1085, *B. S.* 6191 *Robinson*, *F. B.* 13657 *Curran*, *Elmer* 9820, *Copeland* s. n. Epiphyte, up to 2,000 m altitude in Davao. In Java.

20. LEPIDOGYNE Blume

LEPIDOGYNE LONGIFOLIA Blume Fl. Jav. Orch. (1858) 78 (Orch. Arch. Ind. 94) *t. 25*; Miq. Fl. Ind. Bat. 3 (1859) 745; Naves Novis. App. (1882) 249; Ridl. in Journ. Linn. Soc. 32 (1896) 405; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 129, Fig.-Atlas (1909) *fig. 95*; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 224; J. J. Sm. in Nova Guinea 8 (1909) 18, *t. 6, fig. 17, 8³* (1912) 526, 12 (1913) 10; Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1553, Orch. 5 (1915) 28.

Neottia longifolia Blume Bijdr. (1825) 406; Lindl. Gen. & Sp. Orch. (1840) 476.

Speiranthus longifolia Hassk. Cat. Pl. Bogor. (1844) 47.

Lepidogyne sceptrum Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) 55.

Negros (Oriental Negros), Leyte, *Elmer* 9579, *Wenzel* 0399, 0789, 0850, 0904. Terrestrial in mountain regions, at 500 to 600 m altitude in Leyte. In Malay Peninsula, Java, and New Guinea.

Lepidogyne minor Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1922) 56 is probably conspecific.

21. DICEROSTYLIS Blume

DICEROSTYLIS LANCEOLATA Blume Fl. Jav. Orch. (1858) 98 (Orch. Arch. Ind. 116) *t. 38, fig. 1, 1-11*; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 117, Fig.-Atlas (1908) *fig. 85*; Ames Orch. 5 (1915) 28.

Hylophila lanceolata Miq. Fl. Ind. Bat. 3 (1859) 746; Hook. f. Fl. Brit. Ind. 6 (1890) 110; Ridl. in Journ. Linn. Soc. 32 (1896) 403, Mat. Fl. Mal. Penin. 1 (1907) 223.

Batan Islands, *B. S.* 3781 *Fenix*. Santo Domingo de Basco, near Mt. Iraya, in mossy forest. In Malay Peninsula and Java.

DICEROSTYLIS RUBRA (Ames) Schltr. in Engl. Bot. Jahrb. 45 (1911) 392; Ames Orch. 5 (1915) 28.

Hylophila rubra Ames in Philip. Journ. Sci. 2 (1907) Bot. 315, *nomen*, Orch. 2 (1908) 65, *text cut*.

Luzon (Laguna), Mindoro, Panay (Antique), Sibuyan, *F. B. 4378 Merritt, Elmer 12517, Baker 3573, B. S. 32201 McGregor*. Mt. Halcon and Mt. Giting-giting, terrestrial, up to 1,450 m altitude.

22. ERYTHRODES Blume

ERYTHRODES BOETTCHERI Ames Sched. Orch. 6 (1923) 8.

Luzon (Kalinga), *Boettcher s. n.* At 1,000 m altitude.

ERYTHRODES MERRILLII Ames Orch. 3 (1908) 79, *t. 54*, 5 (1915) 29.

Herpysma merrillii Ames in Philip. Journ. Sci. 2 (1907) Bot. 313.

Mindoro, *Merrill 5836*. Mt. Halcon, in damp ravine at 730 m altitude.

ERYTHRODES VRYDAGZYNOIDES Ames Orch. 5 (1915) 29.

Leyte, *Wenzel 0150, 0225*. Dagami, in forest at 60 m altitude.

ERYTHRODES WEBERI Ames Orch. 5 (1915) 30.

Mindanao (Lanao, Agusan), Jolo, *M. S. Clemens s. n., Weber 156, 156A, J. Clemens s. n.* Lake Lanao and Talacogon, terrestrial in shady forest at 10 m altitude. In Jolo at 830 m altitude.

ERYTHRODES WENZELII Ames Orch. 5 (1915) 31.

Palawan, Leyte, *Merrill 7218, Wenzel s. n.*

23. KUHLSSELTIA J. J. Smith

KUHLSSELTIA WHITEHEADII (Rendle) Ames Orch. 5 (1915) 32.

Zeuxine whiteheadii Rendle in Journ. Bot. 34 (1896) 358.

Haemaria merrillii Ames in Philip. Journ. Sci. 2 (1907) Bot. 315; Orch. 3 (1908) 21, *t. 30*.

Kuhlsseeltia merrillii Schltr. in Fedde Repert. 9 (1911) 437.

Luzon (Laguna), Mindoro, *B. S. 6567 Robinson, 19577 Ramos, Copeland s. n., Merrill 5819, 5840*. Mt. Banahao and Mt. Halcon, terrestrial in mossy forest at 1,400 to 2,000 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

KUHLSSELTIA MERRILLII Schltr. in Fedde Repert. 9 (1911) 437 = *Kuhlsseeltia whiteheadii* (Rendle) Ames.

24. CYSTORCHIS Blume

CYSTORCHIS APHYLLA Ridl. in Journ. Linn. Soc. 32 (1896) 400; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 93, Fig.-Atlas (1908) *fig. 64*; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 212; Ames in Philip. Journ. Sci. 2 (1907) Bot. 313, Orch. 2 (1908) 56, 5 (1915) 33.

Luzon (Benguet, Pampanga, Bataan), *Merrill s. n., 3135, R. S. Williams 1921, Elmer 7012, F. B. 7496 Curran*. Terrestrial, in forest. In Malay Peninsula and Java.

- CYSTORCHIS JAVANICA** Blume Fl. Jav. Orch. (1858) 73 (Orch. Arch. Ind. 87) *t.* 24, *fig.* 1, 1-16; Hook. f. Fl. Brit. Ind. 6 (1890) 97; Ames in Philip. Journ. Sci. 6 (1911) Bot. 39, Orch. 5 (1915) 33.
Etaeria javanica Blume Bijdr. (1825) 410.
Aetheria javanica Blume apud Lindl. Gen. & Sp. Orch. (1840) 491.
Hetaeria javanica Miq. Fl. Ind. Bat. 3 (1859) 725.
Cystorchis variegata Blume var. *purpurea* Ridl. in Journ. Linn. Soc. 32 (1896) 399; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 93, Fig.-Atlas (1908) *fig.* 63; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 212; J. J. Sm. in Nova Guinea 8 (1909) 14; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 3; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 140.

Mindanao (Davao), *Weber 286*. Magum-Compostela Trail, in dense forest at 45 m altitude. In Malay Peninsula, Sumatra, Java, Borneo, and Dutch New Guinea.

CYSTORCHIS LUZONENSIS Ames Orch. 5 (1915) 33.

Luzon (Laguna), Mindoro, *Merrill 5146*, *B. S. 17910 Brown*, *Quisumbing 1221*, *Ramos & Edaña s. n.* Mt. Maquiling and Mt. Halcon, terrestrial in forest at 800 to 1,600 m altitude.

25. **CYSTOPUS** Blume

CYSTOPUS MINDANAENSIS Ames Orch. 5 (1915) 34.

Mindanao (Camiguin Island), *B. S. 14422 Ramos*. Mt. Mahinog, on tree top.

CYSTOPUS PHILIPPINENSIS Ames Orch. 5 (1915) 35.

Luzon (Tayabas, Sorsogon), *B. S. 9446 Robinson*, *23636 Ramos*. Infanta and Lake Pulog, the latter in moss at 500 m altitude.

26. **ANOECTOCHILUS** Blume

ANOECTOCHILUS sp. Ames Orch. 5 (1915) 36.

Leyte, *Wenzel 0286*.

EXCLUDED OR SYNONYMOUS SPECIES

ANOECTOCHILUS ARGYRONEURUM Boxall ex Naves Novis. App. (1882) 250, *nomen*.

ANOECTOCHILUS BULLENII Boxall ex Naves Novis. App. (1882) 250, *nomen*.

ANOECTOCHILUS DAYI Boxall ex Naves Novis. App. (1882) 250, *nomen*.

ANOECTOCHILUS INTERMEDIUM Boxall ex Naves Novis. App. (1882) 250, *nomen*.

ANOECTOCHILUS REINWARDTII Blume apud Naves Novis. App. (1882) 250.

A Javan, Sumatran, and Malayan species misdetermined by Naves.

ANOECTOCHILUS ROXBURGHII Lindl. apud Naves Novis. App. (1882) 250.

A native of India and China, wrongly determined by Naves.

27. **MACODES** Lindley

MACODES PETOLA (Blume) Lindl. Gen. & Sp. Orch. (1840) 497; Reichb. f. Xen. Orch. 1 (1858) 227, t. 96, fig. I, 1-5; Blume Fl. Jav. Orch. (1858) 100 (Orch. Arch. Ind. 119) t. 31, fig. 2, t. 36D; Naves Novis. App. (1882) 249; Ridl. in Journ. Linn. Soc. 32 (1896) 399; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 103, Fig.-Atlas (1908) fig. 73; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 215; Ames Orch. 5 (1915) 36.

Neottia petola Blume Bijdr. (1825) 407, fig. 2.

Luzon (Laguna, Tayabas), Mindanao (Agusan), *B. S. 13705 Ramos, 9464 Robinson, Elmer 14180, 18108, 18436*. Mt. Maquiling and Mt. Binuang, forest slope, the latter at 300 m altitude. In Malay Peninsula and Java.

EXCLUDED OR SYNONYMOUS SPECIES

MACODES SETACEUS Boxall ex Naves Novis. App. (1882) 249, *nomen*.

MACODES STRIATUS Boxall ex Naves Novis. App. (1882) 249, *nomen*.

MACODES VEITCHII Boxall ex Naves Novis. App. (1882) 249, *nomen*.

MACODES XANTHOPHYLLUS Boxall ex Naves Novis. App. (1882) 249, *nomen*.

28. **CHEIROSTYLIS** Blume

CHEIROSTYLIS OCTODACTYLA Ames in Philip. Journ. Sci. 2 (1907) Bot. 314, Orch. 3 (1908) 46, t. 40, 5 (1915) 37.

Luzon (Laguna), Mindoro, *Merrill 5834, 7495*. Mt. Banahao and Mt. Halcon, in forest at 2,500 to 2,700 m altitude.

CHEIROSTYLIS PHILIPPINENSIS Ames Orch. 2 (1908) 59, 5 (1915) 37.

Luzon (Kalinga, Benguet, Tayabas), Panay (Iloilo), Negros (Oriental Negros), *Elmer 6035, 9701, Topping 19, Boettcher s. n., Merrill 4015, B. S. 18196 Robinson*. In mountainous regions at 330 to 830 m altitude.

29. **PHILIPPINAEA** Ames and Schlechter

PHILIPPINAEA WENZELII (Ames) Ames & Schltr. in Ames Orch. 6 (1920) 278, t. 100.

Adenostylis wenzelii Ames Orch. 5 (1915) 41.

Luzon (Sorsogon), Leyte, Mindanao (Agusan), *Elmer 13884, 16668, Wenzel 0119, 0210, 0510, 0562*. Terrestrial at 60 to 500 m altitude.

30. **ADENOSTYLIS** Blume

ADENOSTYLIS BENGUETENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1551, Orch. 5 (1915) 39.

Luzon (Benguet), *Elmer 8856*. Baguio.

ADENOSTYLIS ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1552, in Philip. Journ. Sci. 8 (1913) Bot. 408, Orch. 5 (1915) 39.

Luzon (Bontoc, Benguet, Bataan), Palawan, *Vanoverbergh 69, 1099, 2259, 4054, Elmer 8648, 12634, F. B. 6240 Curran*. Terrestrial, on woody hillocks up to 1,300 m altitude.

ADENOSTYLIS LEYTENSIS Ames Orch. 5 (1915) 39.

Luzon (Sorsogon), Leyte, *Elmer 15672, Wenzel 0349*. Terrestrial.

ADENOSTYLIS LUZONENSIS Ames Orch. 2 (1908) 57, *text cut*, 5 (1915) 39.

Zeuxine luzonensis Ames in Philip. Journ. Sci. 2 (1907) Bot. 314, *nomen*.

Luzon (Bontoc, Lepanto), Mindoro, *Vanoverbergh 787, Merrill 4604, 5841*. Terrestrial on mountains, in the forest, at 930 to 1,300 m altitude.

ADENOSTYLIS MARIVELENSIS Ames Orch. 2 (1908) 58, *text cut*, 5 (1915) 39.

Luzon (Bataan), Negros (Occidental Negros), *Merrill 798, 3963, F. B. 6517 Curran, Weber 17, Merrill 7019, Banks s. n., R. S. Williams 627*.

ADENOSTYLIS PHILIPPINENSIS Ames Sched. Orch. 6 (1923) 9.

Leyte, *Wenzel 0375*. Terrestrial in forest at 60 m altitude.

ADENOSTYLIS STRATEUMATICA (Linn.) Ames Orch. 2 (1908) 59, 5 (1915) 40.

Orchis strateumatica Linn. Sp. Pl. (1753) 943, ed. 2 (1763) 1336.

Spiranthes strateumatica Lindl. in Bot. Reg. 10 (1824) sub *t.* 823.

Adenostylis emarginata Blume Bijdr. (1825) 414.

Adenostylis integerrima Blume Bijdr. (1825) 414, *fig. 17*.

Pterygodium sulcatum Roxb. Fl. Ind. ed. 2, 3 (1832) 452.

Tripleura pallida Lindl. in Wall. Cat. (1832) No. 7391, *nomen*, Gen. & Sp. Orch. (1840) 452.

Zeuxine sulcata Lindl. in Wall. Cat. (1832) No. 7392, *nomen*, Gen. & Sp. Orch. (1840) 485; Griff. Notul. 3 (1851) 396, Icon. Pl. Asiat. 3 (1851) *t.* 349; Blume Fl. Jav. Orch. (1858) 55 (Orch. Arch. Ind. 67); Miq. Fl. Ind. Bat. 3 (1859) 723; Benth. Fl. Hongk. (1861) 360; Thwaites Enum. Pl. Zeyl. (1861) 312; Hook. f. Fl. Brit. Ind. 6 (1890) 106; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 286, *t.* 381; Trim. Handb. Fl. Ceyl. 4 (1898) 215; Matsum. Ind. Pl. Jap. 2 (1905) 264; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 108, Fig.-Atlas (1908) *fig. 76*, Orch. Amb. (1905) 12; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 168; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 217.

Zeuxine emarginata Lindl. Gen. & Sp. Orch. (1840) 485; Reichb. f. in Bonpl. 5 (1857) 36.

Zeuxine integerrima Lindl. Gen. & Sp. Orch. (1840) 486; Blume Fl. Jav. Orch. (1858) 55 (Orch. Arch. Ind. 67), *t.* 19, *fig. 1*, *t.* 23c; Miq. Fl. Ind. Bat. 3 (1859) 723.

Zeuxine membranacea Lindl. Gen. & Sp. Orch. (1840) 486; Blume Fl. Jav. Orch. (1858) 56 (Orch. Arch. Ind. 68), *t.* 22, *fig. 2*, *t.* 23A. ? *Corycium humile* Herb. Ham. ex. Lindl. Gen. & Sp. Orch. (1840) 485, in syn.

Neottia strateumatica R. Br. ex Steud. Nom. Bot. ed. 2, 2 (1841) 190.

Zeuxine bracteata Wight Icon. 5¹ (1852) 16, *t.* 1724bis.

Zeuxine brevifolia Wight Icon. 5¹ (1852) 16, *t.* 1725.

Zeuxine robusta Wight Icon. 5¹ (1852) 16, *t.* 1726.

Zeuxine tripleura Lindl. in Journ. Linn. Soc. 1 (1857) 186.

Zeuxine procumbens Blume Fl. Jav. Orch. (1858) 56 (Orch. Arch. Ind. 68) t. 22, fig. 3, t. 23B.

Zeuxine sp. Vidal Rev. Pl. Vasc. Filip. (1886) 273.

Zeuxine strateumatice Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 77.

Mindanao (Misamis, Agusan), *M. S. Clemens* 6, 679a, *Weber* s. n. Terrestrial in the forest, apparently at low altitude. In Afghanistan, India, Ceylon, Malay Peninsula, China, Assam, Japan, Java, and Amboina.

ADENOSTYLIS VANOVERBERGHII Ames in Philip. Journ. Sci. 8 (1913) Bot. 408, Orch. 5 (1915) 40.

Luzon (Bontoc, Lepanto), *Vanoverbergh* 1490, 3936, *Ramos & Edaño* 67, 84. In wet or mossy forest at 930 to 2,540 m altitude.

ADENOSTYLIS WEBERI Ames Orch. 5 (1915) 40.

Mindanao (Davao), *Weber* 280. Mt. Binatun, in bamboo forest at 580 m altitude.

ADENOSTYLIS ZAMBOANGENSIS Ames Sched. Orch. 6 (1923) 10.

Mindanao (Zamboanga), *Copeland* 1567. Terrestrial, in forest at 130 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

ADENOSTYLIS WENZELII Ames Orch. 5 (1915) 41 = *Philippinaea wenzelii* (Ames) Schltr. & Ames.

31. MYRMECHIS Blume

MYRMECHIS GRACILIS Blume Fl. Jav. Orch. (1858) 64 (Orch. Arch. Ind. 77), t. 21, fig. 2; Franch. & Sav. Enum. Pl. Jap. 2 (1879) 39; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 105, Fig.-Atlas (1908) fig. 74; Matsum. Ind. Fl. Jap. 2 (1905) 254; Ames in Philip. Journ. Sci. 2 (1907) Bot. 315, Orch. 2 (1908) 63, 5 (1915) 42.

Anecochilus gracilis Blume Bijdr. (1825) 413.

Anaectochilus gracilis Blume apud Lindl. Gen. & Sp. Orch. (1840) 500.

Rhamphidia alsinefolia Lindl. in Journ. Linn. Soc. 1 (1857) 182; Miq. Fl. Ind. Bat. 3 (1859) 730.

Rhamphidia japonica Reichb. f. in Bot. Zeit. 36 (1878) 75.

Myrmechis japonica Rolfe in Journ. Linn. Soc. 36 (1903) 44, sub *Myrmechis chinensis* Rolfe.

Luzon (Benguet, Bataan), Mindoro, Mindanao (Misamis, Bukidnon, Davao), *B. S.* 8422 *McGregor*, 38520 *Ramos & Edaño*, *Merrill* 3865, *F. B.* 4432, 4460 *Merritt*, 4622 *Mearns & Hutchinson*, *Elmer* 11372. In mountain regions, up to 2,100 m altitude in Benguet, and 3,300 m altitude in Mindanao. In Java and Japan.

MYRMECHIS PERPUSILLA Ames Sched. Orch. 6 (1923) 15.

Luzon (Benguet), *Copeland* 124.

MYRMECHIS PHILIPPINENSIS Ames Orch. 2 (1908) 64, text cut, 5 (1915) 42.

Luzon (Bataan), *Leiberg* 6066. Mt. Mariveles, terrestrial at 800 m altitude.

32. VRYDAGZYNEA Blume

VRYDAGZYNEA ALBIDA Blume Fl. Jav. Orch. (1858) 62 (Orch. Arch. Ind. 75), t. 19, fig. 2; Hook. f. Fl. Brit. Ind. 6 (1890) 97; Ridl. in Journ. Linn. Soc. 32 (1896) 397; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 89, Fig.-Atlas (1908) fig. 61; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 211; Ames in Philip. Journ. Sci. 2 (1907) Bot. 313; Orch. 2 (1908) 55, 5 (1915) 42.

Etaeria albida Blume Bijdr. (1825) 410.

Aetheria albida Blume apud Lindl. Gen. & Sp. Orch. (1840) 491.

Hetaeria albida Blume apud Miq. Fl. Ind. Bat. 3 (1859) 726; Naves Novis. App. (1882) 249.

Luzon (Rizal, Tayabas, Sorsogon), Mindoro, Samar, Negros (Oriental Negros), Leyte, Mindanao (Zamboanga, Davao), Jolo. A rather common and widely distributed species in the Philippines. Terrestrial in dense forest at 137 to 1,000 m altitude. In Malay Peninsula and Java.

VRYDAGZYNEA WEBERI Ames Orch. 5 (1915) 42.

Luzon (Apayao, Rizal, Laguna, Tayabas, Sorsogon), Polillo, Leyte, Weber 333, Elmer 14616, 14685, 17374, 18028, B. S. 9191, 9439 Robinson, 2978 Ramos & Edaño, 25401 Yates, Wenzel 0244, 0630. Terrestrial, up to 600 m altitude in Leyte.

33. HETAERIA Blume

HETAERIA BLACKII Ames Orch. 5 (1915) 43.

Mindanao (Lanao, Davao), M. S. Clemens 439, Black s. n.

HETAERIA LANCIFOLIA Ames Orch. 5 (1915) 44 (as *lancifolium*).

Luzon (Apayao), Weber 327. Abulug-Narig Trail, terrestrial in forest at 76 m altitude.

HETAERIA LEYTENSIS Ames Sched. Orch. 6 (1923) 14.

Leyte, B. S. 42081b Edaño. At 300 m altitude.

HETAERIA MINDANAENSIS Ames Orch. 5 (1915) 45.

Mindanao (Agusan) Weber 317. Sangaan-Cabadbaran Trail, terrestrial in shady forest at 22 m altitude.

HETAERIA OBLONGIFOLIA Blume Fl. Jav. Orch. (1858) 85 (Orch. Arch. Ind. 102), t. 32, fig. 3; Miq. Fl. Ind. Bat. 3 (1859) 725; Kränzl. in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 241; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 113, Fig.-Atlas (1908) fig. 80; Ames Orch. 1 (1905) 70, 2 (1908) 62, 5 (1915) 46; J. J. Sm. in Nova Guinea 8 (1909) 17, 8^a (1912) 525.

Etaeria oblongifolia Blume Bijdr. (1825) 410, fig. 14.

Aettieria oblongifolia Lindl. Gen. & Sp. Orch. (1840) 491.

Etaeria rubicunda Reichb. f. in Bonpl. 3 (1855) 214.

Rhamphidia tenuis Lindl. in Journ. Linn. Soc. 1 (1857) 182; Miq. Fl. Ind. Bat. 3 (1859) 730; Naves Novis. App. (1882) 248.

Hetaeria tenuis Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 273; Rolfe in Journ. Bot. 24 (1886) 59; Ceron Cat. Pl. Herb. Manila (1892) 169.

Goodyera rubicunda J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 121 (as to syn. *Aetheria rubicunda* Reichb. f.).

Luzon (Ilocos Norte, Cagayan, Benguet, Rizal, Laguna, Tayabas), Ticao, Mindoro, Negros (Oriental Negros), Palawan, Balabac, Mindanao (Zamboanga, Lanao). A common and widely distributed species in the Philippines. Terrestrial in forest at 30 to 90 m altitude. In Java, Molucca Islands, New Guinea, and possibly Samoa.

HETAERIA WENZELII Ames Orch. 5 (1915) 46.

Leyte, *Wenzel 0227*. Dagami, terrestrial in forest at 60 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

HETAERIA ALBIDA Blume apud Miq. Fl. Ind. Bat. 3 (1859) 726; Naves Novis. App. (1882) 240 = *Vrydagzynea albida* Blume.

HETAERIA JAVANICA (Blume) Miq. Fl. Ind. Bat. 3 (1859) 725 = *Cystorchis javanica* Blume.

HETAERIA TENUIS Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 273; Rolfe in Journ. Bot. 24 (1886) 59; Ceron Cat. Pl. Herb. Manila (1892) 169 = *Hetaeria oblongifolia* Blume.

HETAERIA VARIEGATA Miq. apud Naves Novis. App. (1882) 249.

This is probably referable to *Cystorchis javanica* Blume.

34. TROPIDIA Lindley

TROPIDIA CALCARATA Ames in Philip. Journ. Sci. 7 (1912) Bot. 7, Orch. 5 (1915) 47.

Luzon (Rizal, Laguna), *B. S. 13603 Ramos, Merrill 7144, Catalan 223, Elmer 18188*.

TROPIDIA MINDANAENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1553, Orch. 5 (1915) 47.

Mindanao (Davao), *Elmer 11785*. Todaya, Mt. Apo.

TROPIDIA MINDORENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 315, Orch. 5 (1915) 47.

Mindoro, *Merrill 5552*. Binabay River, terrestrial in humid forest at 300 m altitude.

TROPIDIA ROBINSONII Ames Sched. Orch. 6 (1923) 16.

Luzon (Tayabas), *B. S. 9496 Robinson*.

TROPIDIA SEPTEMNERVIS (Schauer) Reichb. f. in Linnaea 25 (1852) 230; Ames Orch. 5 (1915) 47, in textu.

Ptychochilus septemnervis Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 431, t. 12, fig. B.

Tropidia minor Ames Orch. 2 (1908) 65, text cut, 5 (1915) 47.

Luzon (Nueva Ecija, Pampanga, Bataan, Laguna), Mindoro, Leyte, *Merrill 4204, Whitford 1368, B. S. 16558 Ramos, 1542 Bermejos, 26267 Ramos & Edaño, 17927 Brown, Wenzel 0557, F. B. 7501 Curran*. Terrestrial in forest at 500 m in Leyte.

EXCLUDED OR SYNONYMOUS SPECIES

TROPIDIA CURCULIGOIDES Lindl. apud Naves Novis. App. (1882) 249.

A native of Java, Borneo, and Malacca, wrongly attributed to the Philippines. Cf. Ames Orch. 5 (1915) 47, in textu.

TROPIDIA GRAMINEA Blume apud Naves Novis. App. (1882) 249.

TROPIDIA GRANDIS Hance apud Naves Novis. App. (1882) 249.

TROPIDIA MINOR Ames Orch. 2 (1908) 65, *text cut*, 5 (1915) 47 = *Tropidia septemnervis* (Schauer) Reichb. f.

35. CORYMBORCHIS Thouars

CORYMBORCHIS CONFUSA Ames Orch. 5 (1915) 48.

Corymbis disticha Lindl. Fol. Orch. Corymbis (1854) 1, quoad Philip.; Thou. apud Miq. Fl. Ind. Bat. 3 (1859) 736, quoad Philip.

Chloidia confusa Ames Orch. 2 (1908) 67, *text cut*.

Corymbis confusa Schltr. in Engl. Bot. Jahrb. 45 (1911) 395.

Luzon (Ilocos Norte, Apayao), Leyte, Bancalan, Mindanao (Agusan, Zamboanga), Jolo, *B. S. 32723, 32752 Ramos, 37295 Ramos & Edaño, 28227 Fenix, Cuming s. n., Weber 320, 339, Wenzel 0209, Elmer 13433, R. S. Williams 2153, J. Clemens s. n.* Terrestrial, in forest at sea level to 830 m altitude.

36. CHRYSOGLOSSUM Blume

CHRYSOGLOSSUM VILLOSUM Blume Bijdr. (1825) 338, *fig. 7*; Lindl. Gen. & Sp. Orch. (1830) 14; De Vriese Ill. Orch. Ind. Neerl. (1854) *t. 12, fig. 4*; Blume Fl. Jav. Orch. (1858) 137 (Orch. Arch. Ind. 162), *t. 47*; Miq. Fl. Ind. Bat. 3 (1859) 626; Reichb. f. in Walp. Ann. 6 (1861) 206; Hook. f. Fl. Brit. Ind. 5 (1890) 783; Ridl. in Journ. Linn. Soc. 32 (1896) 316; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 173, Fig.-Atlas (1909) *fig. 131*; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 112; Ames in Philip. Journ. Sci. 2 (1907) Bot. 316, Orch. 2 (1908) 67, 5 (1915) 48, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 143.

Mindoro, *Merrill 5838*. Binabay River, terrestrial at about 300 m altitude. In Malay Peninsula, Java, and Borneo.

37. NEPHELAPHYLLUM Blume

NEPHELAPHYLLUM MINDORENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 316, Orch. 3 (1908) 83, *t. 55, 5* (1915) 48.

Luzon (Nueva Ecija), Mindoro, *B. S. 26319, 26494, 26652 Ramos & Edaño, Merrill 5623*. Terrestrial on dry slopes and in humid forest at 200 to 300 m altitude.

38. DIGLYPHOSA Blume

DIGLYPHOSA ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1555, Orch. 5 (1915) 48.

Mindanao (Davao), *Elmer 11815*. Todaya, Mt. Apo.

39. COELOGYNE Lindley

COELOGYNE ASPERATA Lindl. in Journ. Hort. Soc. 4 (July, 1849) 221, Fol. Orch. Coelog. (1854) 3; J. Linden Pescat. 1 (1854) *text to t. 7*; Miq. Fl. Ind. Bat. 3 (1859) 666; Reichb. f. in Walp. Ann. 6 (1861) 224; Warner & Will. Orch. Alb. 7 (1888) *t. 311*; Hook. f. Fl. Brit. Ind. 5 (1890) 835; Veitch. Man. Orch. Pl. pt. 6 (1890) 31; Ridl. in Journ. Linn. Soc. 31 (1896) 287, 32 (1896) 326; L. Linden in Lindenia 13 (1897) 15, *t. 582*; Ridl. in Journ. Bot. 36 (1898) 214, Mat. Fl. Mal. Penin. 1 (1907) 129; Pfütz. in

Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 76; Ames Orch. 2 (1908) 68, 5 (1915) 49; J. J. Sm. in Nova Guinea 8 (1909) 20, 8² (1912) 527, 12 (1913) 10, 12³ (1915) 196; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 143.

Coelogyne lowii Lindl. in Paxt. Mag. Bot. 16 (September, 1849) 225, *t. and text cut*; J. Linden Pescat. 1 (1854) *t. 7*; Boxall ex Naves Novis. App. (1882) 237.

Pleione asperata Ktze. Rev. Gen. Pl. 2 (1891) 680.

Samar, Leyte, Negros (Oriental Negros), Mindanao (Agusan, Surigao, Davao), Basilan, *B. S. 17624 Ramos, 15500, 16257 Reillo, Wenzel s. n., Elmer 10008, Weber 55, 250, Bolster 294, F. B. 6103 Hutchinson*. Epiphyte in forest at 75 to 430 m altitude. In Malay Peninsula, Sumatra, Borneo, and New Guinea.

COELOGYNE BILAMELLATA Lindl. Fol. Orch. Coelog. (1854) 14; Miq. Fl. Ind. Bat. 3 (1859) 668; Naves Novis. App. (1882) 237; Ridl. in Journ. Linn. Soc. 31 (1896) 287; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 24; Ames Orch. 2 (1908) 73, *text cut*, 5 (1915) 94.

Panisea bilamellata Reichb. f. in Walp. Ann. 6 (1861) 240.

Pleione bilamellata Ktze. Rev. Gen. Pl. 2 (1891) 680.

Luzon (Laguna, Tayabas), Leyte, Mindanao (Zamboanga, Agusan, Bukidnon, Surigao), *B. S. 15131 Ramos, 39046 Ramos & Edaño, Elmer 7727, 13900, Wenzel 0412, 0494, 0513, Merrill 8202, Bolster 288*. Epiphyte in forest, up to 1,000 m altitude in Mindanao. Reported from Borneo.

COELOGYNE CANDOONENSIS Ames Sched. Orch. 6 (1923) 18, *fig. 4*.

Mindanao (Bukidnon), *B. S. 38894 Ramos & Edaño*. In forest at 1,830 m altitude.

COELOGYNE CHLOROPTERA Reichb. f. in Gard. Chron. II 19 (1883) 466; Kränz. Xen. Orch. 3 (1890) 69, *t. 235, fig. I. 1-6*; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 55, *fig. 18L*; Ames Orch. 2 (1908) 71, *t., 5* (1915) 49.

Coelogyne sparsa Ames Orch. 1 (1905) 72, *p. p. (as to Elmer 5842)*, non Reichb. f.

Luzon (Cagayan, Kalinga, Bontoc, Benguet, Pampanga, Bataan, Tayabas), Negros (Oriental Negros). A fairly common species in Luzon, on trees and rocks at 1,000 to 1,530 m altitude.

COELOGYNE CONFUSA Ames Orch. 5 (1915) 49.

Mindanao (Camiguin Island), *B. S. 14430 Ramos*. Epiphyte.

COELOGYNE ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1556, Orch. 5 (1915) 50.

Mindanao (Bukidnon, Davao), *B. S. 38583 Ramos & Edaño, Elmer 10694*. Mt. Lipa and Todaya, Mt. Apo, up to 1,300 m altitude.

COELOGYNE INTEGERRIMA Ames in Philip. Journ. Sci. 4 (1909) Bot. 665, 6 (1911) Bot. 40; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340; Ames Orch. 5 (1915) 50.

Luzon (Bontoc, Benguet, Tayabas), *Vanoverbergh 3646, Merrill 6350, B. S. 8454, 8519, 8822 McGregor, 31711 Santos, Elmer 9258*. Mountain regions at 1,750 to 2,450 m altitude.

COELOGYNE LOHERI Rolfe in Kew Bull. (1908) 414; Ames Orch. 5 (1915) 50, 6 (1920) 279.

Luzon (Bontoc, Benguet), *Loher* 549, *Vanoverbergh* 1197, 2867, 3906, 4041, *Boettcher* s. n., *B. S.* 31909 *Santos*. Mountain regions up to 1,700 m altitude.

COELOGYNE LONGIRACHIS Ames Orch. 7 (1922) 88.

Mindanao (Bukidnon), *B. S.* 38671 *Ramos & Edaño*. Mahilucot River, along streams in forest at 1,300 m altitude.

COELOGYNE MARMORATA Reichb. f. in Linnaea 41 (1877) 116; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 85; Ames Orch. 5 (1915) 50, Sched. Orch. 6 (1923) 20.

Coeogyne zahlbrucknerae Kränzl. in Fedde Repert. 17 (1921) 389.

Luzon (Benguet), *Wallis* s. n., *Schultze* s. n., *Loher* s. n. Sablan, May, 1909.

COELOGYNE MERRILLII Ames in Philip. Journ. Sci. 6 (1911) Bot. 40, Orch. 5 (1915) 51.

Luzon (Benguet, Rizal), *Merrill* 6620, *Phil. Pl.* 716, *Haight* 14, *B. S.* 31814 *Santos*, 29764 *Ramos & Edaño*, *Lyon* 164. Mountain regions at 300 to 2,300 m altitude.

COELOGYNE MINUTISSIMA Kränzl. in Fedde Repert. 17 (1921) 390. Philippines, probably Luzon, *Loher* s. n.

No specimen seen.

COELOGYNE PALAWANENSE Ames Orch. 5 (1915) 51.

Palawan, *Merrill* 9499. Mt. Capoas at 1,000 m altitude.

COELOGYNE QUINQUELAMELLATA Ames Orch. 6 (1920) 280.

Mindanao (Surigao), *Wenzel* 01175, 01179, 01182, 01183, 01206. Epiphyte in forest at 150 m altitude.

COELOGYNE RAMOSII Ames Orch. 7 (1922) 90.

Mindanao (Bukidnon, Agusan), *B. S.* 38892, 38944 *Ramos & Edaño*, *Elmer* 14130. Mt. Candoon and Cabadbaran (Mt. Urdaneta) up to 1,850 m altitude.

COELOGYNE ROCHUSSENII De Vriese Ill. Orch. Ind. Neerl. (1854) t. 2, t. 11, fig. 6; Miq. Fl. Ind. Bat. 3 (1859) 666; Reichb. Xen. Orch. 1 (1856) 212, t. 85; Ridl. in Journ. Linn. Soc. 32 (1896) 321; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 144, Fig.-Atlas (1909) fig. 107; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 128; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 70, figs. 22E, 23; Ames in Philip. Journ. Sci. 4 (1909) Bot. 664; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 7; Ames Orch. 5 (1915) 52, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 145.

Coeogyne macrobulbon Hook. f. Fl. Brit. Ind. 5 (1890) 830.

Pleione rochussenii Ktze. Rev. Gen. Pl. 2 (1891) 680.

Pleione macrobulbon Ktze. Rev. Gen. Pl. 2 (1891) 680.

Luzon (Pampanga), Mindanao (Lanao), *M. S. Clemens* s. n., *Lyon* s. n. Camp Keithley, September, 1907. In Malay Peninsula, Sumatra, Java, Celebes, and Borneo.

COELOGYNE SPARSA Reichb. f. in Gard. Chron. II 19 (1883) 306; Veitch Man. Orch. Pl. pt. 6 (1890) 49; Kränzl. Xen. Orch. 3 (1890) 70, *t. 235, fig. II. 7-12*; Wittmack in Gartenfl. 46 (1897) 449, *t. 1442*; Ames Orch. 1 (1905) 71 (excl. *Elmer 5842*), 2 (1908) 70, *text cut*, 5 (1915) 53; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 54, *fig. 18, J. K.*

Pleione sparsa Ktze. Rev. Gen. Pl. 2 (1891) 680.

Luzon (Rizal, Bataan), Panay (Antique), Negros (Oriental Negros), Palawan, *B. S. 2176 Ramos, 647 Foxworthy, Weber 11, 15, 16, Merrill 3961, Whitford 1090, Lyon 133, R. S. Williams 626, 759, Yoder s. n., Elmer 10210, McGregor 5806*. Mountain regions, epiphytic at 900 to 1,050 m altitude.

COELOGYNE SWANIANA Rolfe in Kew Bull. (1894) 183, in Orch. Rev. 2 (1894) 198, in Sander Reichenbachia II 2 (1894) 105, *t. 92*; Hook. f. in Bot. Mag. 124 (1898) *t. 7602*; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 68, *fig. 22C*; Ames Orch. 5 (1915) 53.

Only horticultural specimens of this species have been seen.

COELOGYNE VANOVERBERGHII Ames Orch. 5 (1915) 53.

Bontoc, *Vanoverbergh 2865*. Bauco at 1,700 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

COELOGYNE CILIATA Boxall ex Naves Novis. App. (1882) 237, *nomen*.

COELOGYNE CONCHOIDEA Reichb. f. in Walp. Ann. 6 (1861) 237; Naves Novis. App. (1882) 237 = *Pholidota imbricata* Lindl.

COELOGYNE CUMINGII Lindl. apud Naves Novis. App. (1882) 237.

Cuming's specimen, on which this species was based, was from Singapore, not from the Philippines.

COELOGYNE FIMBRIATA Lindl. apud Naves Novis. App. (1882) 237.

COELOGYNE IMBRICATA (Lindl.) Reichb. f. in Walp. Ann. 6 (1861) 238 = *Pholidota imbricata* Lindl.

COELOGYNE LOWII Lindl. in Paxt. Mag. Bot. 16 (September, 1849) 225, *t. and text cut*; J. Linden Pescat. 1 (1854) *t. 7*; Boxall ex Naves Novis. App. (1882) 237 = *Coelogyne asperata* Lindl.

COELOGYNE PALLIDA (Lindl.) Reichb. f. in Walp. Ann. 6 (1861) 238 = *Pholidota imbricata* Lindl.

COELOGYNE PANDURATA Lindl. apud Naves Novis. App. (1882) 237:

COELOGYNE PUMILA Reichb. f. in Walp. Ann. 6 (1861) 236 = *Acoridium pumilum* (Reichb. f.) Rolfe.

COELOGYNE RHOMBOPHORA Reichb. f. in Linnaea 41 (1877) 116 = *Acoridium rhombophorum* (Reichb. f.) Ames.

COELOGYNE RUMPHII Lindl. apud Naves Novis. App. (1882) 237.

COELOGYNE SPARSA Ames Orch. 1 (1905) 72, p. p. (as to *Elmer 5842*), non Reichb. f. = *Coelogyne chloroptera* Reichb. f.

COELOGYNE SPECIOSA Lindl. apud Naves Novis. App. (1882) 237.

COELOGYNE TRIFIDA Reichb. f. apud Naves Novis. App. (1882) 237.

COELOGYNE TRIOTOS Reichb. f. in Walp. Ann. 6 (1861) 238, in Hamb. Gartenz. 18 (1862) 34; Naves Novis. App. (1882) 237 = *Pholidota imbricata* Lindl.

COELOGYNE VENTRICOSA (Blume)' Reichb. f. in Walp. Ann. 6 (1861) 237 = *Pholidota ventricosa* (Blume) Reichb. f.

COELOGYNE VISCOSA Boxall ex Vidal ex Naves Novis. App. (1882) 237, *nomen*.

COELOGYNE ZAHLBRUCKNERAE Kränzl. in Fedde Repert. 17 (1921) 389 = *Coelogyne marmorata* Reichb. f.

40. PSEUDACORIDIUM Ames

PSEUDACORIDIUM WOODIANUM Ames Orch. 7 (1922) 79.

Dendrochilum woodianum Ames in Philip. Journ. Sci. 2 (1907) Bot. 318, *nomen*, Orch. 2 (1908) 80, *text cut*, 5 (1915) 66.

Mindoro, Merrill 5816. Mt. Halcon, at 2,260 to 2,830 m altitude.

41. ACORIDIUM Nees and Meyen

ACORIDIUM AFFINE Ames Orch. 7 (1922) 80.

Dendrochilum affine Ames Orch. 2 (1908) 95, 5 (1915) 55.

Luzon (Bataan), Merrill 3218, Whitford 1354, B. S. 5184 *Topping*.

ACORIDIUM ANFRACTOIDES Ames Orch. 7 (1922) 80.

Dendrochilum anfractoides Ames Orch. 3 (1908) 13, t. 28, 5 (1915) 55, in Philip. Journ. Sci. 4 (1909) Bot. 665; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340.

Luzon (Benguet), F. B. 4312a Mearns, B. S. 8358, 8834, 8858 McGregor, Haight 10, Ramos & Edaño 11, 48, 48a, 49. In mossy forest at 1,930 to 2,160 m altitude.

ACORIDIUM ANFRACTUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 148.

Dendrochilum anfractum Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 118, fig. 38P; Ames Orch. 2 (1908) 91, t. 21, fig. I, 1, 5 (1915) 55.

Luzon (Lepanto, Benguet), Merrill 4482, Ramos & Edaño 10, 39, 40, 47. In mossy forest at 2,100 to 2,290 m altitude.

ACORIDIUM AURICULARE Ames Orch. 7 (1922) 80.

Dendrochilum auriculare Ames in Philip. Journ. Sci. 4 (1909) Bot. 595, Orch. 5 (1915) 55.

Luzon (Benguet), B. S. 5771, 5818 Ramos, Ramos & Edaño 9. In mossy forest at 2,060 m altitude.

ACORIDIUM BINUANGENSE Ames Orch. 7 (1922) 80.

Dendrochilum binuangense Ames Orch. 6 (1920) 281.

Luzon (Tayabas), B. S. 28620 Ramos & Edaño. Summit of Mt. Binuang at 500 m altitude.

ACORIDIUM CINNABARINUM (Pfitz.) Ames Orch. 7 (1920) 80.

Dendrochilum cinnabarinum Pfitz. in Engl. Pflanzenreich IV. 50.

II. B. 7 (1907) 104, fig. 35F; Ames Orch. 3 (1908) 10, 5 (1915) 56, 6 (1920) 282, in Philip. Journ. Sci. 4 (1909) Bot. 597; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340.

Luzon (Benguet, Rizal), Merrill 6475, Boettcher s. n., Ramos & Edaña 43, B. S. 8844 McGregor, F. B. 16348 Curran, Merritt, & Zschokke. In mossy forest at 2,160 to 2,930 m altitude.

ACORIDIUM CONFUSUM Ames Orch. 7 (1922) 80.

Dendrochilum confusum Ames in Philip. Journ. Sci. 6 (1911) Bot. 41, Orch. 5 (1915) 56.

Luzon (Laguna, Sorsogon), Leyte, F. B. 19124 Tamesis, B. S. 5640 Curran, Elmer 15068, Wenzel 0853. In forest at 700 m altitude.

ACORIDIUM CURRANII Ames Orch. 7 (1922) 80.

Dendrochilum curranii Ames Orch. 3 (1908) 15, t. 29, 5 (1915) 57.

Luzon (Rizal, Laguna), B. S. 3053, 16570, 19561 Ramos, 19152 Reillo, 3077 Curran, 6575, 6580 Robinson, Disdon s. n.

ACORIDIUM ELMERI Ames Orch. 7 (1922) 80.

Dendrochilum elmeri Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1558, Orch. 5 (1915) 57.

Mindanao (Davao), Elmer 10635. Todaya, Mt. Apo, epiphyte in mossy forest at 2,500 m altitude.

ACORIDIUM EXASPERATUM Ames Sched. Orch. 6 (1923) 21, fig. 6.

Luzon (Rizal), B. S. 42338 Ramos. In mossy forest at 1,300 m altitude.

ACORIDIUM EXILE Ames Orch. 7 (1922) 80.

Dendrochilum exile Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, nomen, Orch. 2 (1908) 84, text cut, 5 (1915) 57.

Mindoro, Merrill 5721. Mt. Halcon.

ACORIDIUM FOXWORTHYI Ames Orch. 7 (1922) 80.

Dendrochilum foxworthyi Ames Orch. 3 (1908) 8, t. 26, 5 (1915) 57.

Luzon (Zambales), B. S. 2542 Foxworthy. Mt. Pinatubo at 1,800 m altitude.

ACORIDIUM GRAMINIFOLIUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 144.

Dendrochilum graminifolium Pfitz. in Engl. Pflanzenreich IV. 50.

II. B. 7 (1907) 114, fig. 38 Q, R; Ames Orch. 2 (1908) 12, 83, t. 22, fig. II, 2, 5 (1915) 57.

Luzon (Lepanto, Benguet), Merrill 4594, 4764, F. B. 5734 Klemme, 15907 Bacani, Ramos & Edaña 32, B. S. 8487 McGregor.

ACORIDIUM HASTATUM Ames Orch. 7 (1922) 80.

Dendrochilum hastatum Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, nomen, Orch. 2 (1908) 99, text cut, 5 (1915) 58.

Mindoro, F. B. 4441 Merritt, Merrill 5759. Mt. Halcon at 2,260 m altitude.

ACORIDIUM HUTCHINSONIANUM Ames comb. nov.

Dendrochilum hutchinsonianum Ames in Philip. Journ. Sci. 2 (1907)

Bot. 317, *nomen*, Orch. 2 (1908) 96, *text cut*, 5 (1915) 58.

Acoridium hutchinsonii Ames Orch. 7 (1922) 81, *sphalm.*

Dendrochilum hutchinsonii Ames Orch. 7 (1922) 81, in syn.

Mindoro, Merrill 5813. Mt. Halcon, in mossy thickets at 2,660 m altitude.

ACORIDIUM IRIGENSE Ames Orch. 7 (1922) 81.

Dendrochilum irigense Ames Orch. 5 (1915) 58.

Luzon (Camarines), B. S. 22186 Ramos, Santos s. n. Mt. Iriga at 500 m altitude.

ACORIDIUM LOHERI Ames Orch. 7 (1922) 81.

Dendrochilum loheri Ames Orch. 3 (1908) 12, t. 27, fig. I, A, 5

(1915) 59, 6 (1920) 283; Merr. & Merritt in Philip. Journ. Sci.

5 (1910) Bot. 340.

Luzon (Benguet), Loher 461A, Merrill 6476, Boettcher s. n.

ACORIDIUM LONGIBULBUM Ames Orch. 7 (1922) 81.

Dendrochilum longibulbum Ames in Philip. Journ. Sci. 7 (1912) Bot.

26, Orch. 5 (1915) 59.

Luzon (Bontoc, Lepanto), Vanoverbergh 782, 3830, Ramos & Edaño 25, 37, 46. At high altitudes.

ACORIDIUM LUCBANENSE Ames Orch. 7 (1922) 81.

Dendrochilum lucbanense Ames in Elm. Leaf. Philip. Bot. 5 (1912)

1559, Orch. 5 (1915) 60.

Luzon (Tayabas), Elmer 7571. Lucban, in mossy forest at 530 m altitude.

ACORIDIUM LUZONENSE Ames Orch. 7 (1922) 81.

Dendrochilum luzonense Ames Orch. 2 (1908) 121, *text cut*, 5 (1915)

60.

Luzon (Lepanto, Benguet), F. B. 5735 Klemme, 15729 Curran & Merritt, Ramos & Edaño 18, B. S. 5772 Ramos. At 1,600 to 2,000 m altitude.

ACORIDIUM MACGREGORII Ames Orch. 7 (1922) 81.

Dendrochilum sp. Merr. & Merritt in Philip. Journ. Sci. 5 (1910)

Bot. 341.

Dendrochilum macgregorii Ames in Philip. Journ. Sci. 6 (1911) Bot.

42, Orch. 5 (1915) 60.

Luzon (Benguet), B. S. 8849 McGregor. Mt. Pulog.

ACORIDIUM MALEOLENS (Kränzl.) Ames Orch. 7 (1922) 81.

Dendrochilum maleolens Kränzl. in Orchis 2 (1908) 63; Ames Orch.

5 (1915) 54, 60

Luzon (Rizal), Loher s. n.

Described from a plant collected by Loher at Manila, and said to be near *Acoridium oliganthum* Ames. No specimen of this species has been seen.

ACORIDIUM MARGINATUM Ames Sched. Orch. 6 (1923) 23, fig. 5.

Luzon (Rizal), *Ramos s. n.* In forest at 1,630 m altitude.

ACORIDIUM MERRILLII Ames in Proc. Biol. Soc. Wash. 19 (1906) 151.

Dendrochilum merrillii Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 119, fig. 38A, B; Ames Orch. 2 (1908) 12, 102, t. 23, fig. I, 1, text cut, 5 (1915) 60.

Luzon (Lepanto, Benguet), *Merrill 4585, 4858, B. S. 8353 McGregor, Ramos & Edaño 1, 21, 44.* At 2,100 to 2,800 m altitude.

ACORIDIUM MICROCHILUM (Schltr.) Ames Orch. 7 (1922) 81.

Platyclinis microchila Schltr. in Bull. Herb. Boiss. II 6 (April, 1906) 302.

Acoridium venustulum Ames in Proc. Biol. Soc. Wash. 19 (September, 1906) 147.

Dendrochilum venustulum Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38U; Ames Orch. 2 (1908) 12, 88, t. 22, fig. III, 3, in Philip. Journ. Sci. 4 (1909) Bot. 595; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341.

Dendrochilum microchilum Ames Orch. 2 (1908) 87, in textu, 5 (1915) 60, in Philip. Journ. Sci. 7 (1912) Bot. 26.

Luzon (Bontoc, Lepanto, Benguet, Nueva Vizcaya, Laguna). A rather common species in the above region, especially so in Benguet at about 1,800 to 2,400 m altitude.

ACORIDIUM MINDANAENSE Ames Sched. Orch. 6 (1923) 25, fig. 6.

Mindanao (Agusan), *Elmer 14095.*

ACORIDIUM MINDORENSE Ames Orch. 7 (1922) 81.

Dendrochilum mindorense Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, nomen, Orch. 2 (1908) 91, text cut, 5 (1915) 60.

Mindoro, *Merrill 5729, 5795.* Mt. Halcon at 2,000 m altitude.

ACORIDIUM MONTANUM Ames Sched. Orch. 5 (1923) 1.

Acoridium linearifolium Ames Orch. 7 (1922) 141, non Rolfe.

Luzon (Benguet), *Ramos & Edaño 56.* Mt. Boadan, in mossy forest at about 2,000 m altitude.

ACORIDIUM OCELLATUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 151.

Dendrochilum ocellatum Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 117, fig. 38C; Ames Orch. 2 (1908) 12, 101, t. 23, fig. II, 2, 5 (1915) 61, in Philip. Journ. Sci. 4 (1909) Bot. 596.

Luzon (Lepanto), *Merrill 4481a, Ramos & Edaño 7.* Mt. Data at 2,300 m altitude.

ACORIDIUM OLIGANTHUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 150.

Dendrochilum oliganthum Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 117, fig. 38E; Ames Orch. 2 (1908) 12, 98, t. 23, fig. IV, 4, 5 (1915) 61.

Luzon (Lepanto), *Merrill 4481b, Ramos & Edaño 35.* Mt. Data and Mt. Sinapsapan, at about 2,300 m altitude.

ACORIDIUM PARVULUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 146.

Dendrochilum parvulum Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38N; Ames Orch. 2 (1908) 12, 87, t. 21, fig. II, 2, 5 (1915) 61.

Luzon (Bataan, Rizal, Laguna), *Merrill 3217b, Whitford 141, B. S. 29724, 29732, Ramos & Edaño, F. B. 8024 Curran & Merritt. Lamao River, Mt. Mariveles.*

ACORIDIUM PERPLEXUM Ames Orch. 7 (1922) 142.

Luzon (Lepanto), *Ramos & Edaño 55.* In mossy forest, Mt. Sinapsapan at 2,000 m altitude.

ACORIDIUM PHILIPPINENSE Ames in Proc. Biol. Soc. Wash. 19 (1906) 149.

Dendrochilum philippinense Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 118, fig. 38F; Ames Orch. 2 (1908) 12, 97, t. 23, fig. V, 5, 5 (1915) 61, in Philip. Journ. Sci. 4 (1909) Bot. 595.

Luzon (Bontoc, Lepanto, Ifugao, Benguet). A fairly common species in the above region at about 2,000 to 2,300 m altitude.

ACORIDIUM PULCHERRIMUM Ames Orch. 7 (1922) 144.

Luzon (Benguet), *Ramos & Edaño 45.* In mossy forest, Mt. Natoo, at 1,930 m altitude.

ACORIDIUM PULOGENSE Ames Orch. 7 (1922) 81.

Dendrochilum pulogense Ames in Philip. Journ. Sci. 4 (1909) Bot. 594; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340; Ames Orch. 5 (1915) 61.

Luzon (Benguet), *F. B. 16347A Curran, Merritt, & Zschokke, Merrill 6478, B. S. 8355, 8842, 8843 McGregor, Haight 2, Boettcher s. n., Ramos & Edaño 24, 33.*

ACORIDIUM PUMILUM (Reichb. f.) Rolfe in Orch. Rev. 12 (1904) 220; Ames Orch. 7 (1922) 81.

Dendrochilum pumilum Reichb. f. in Bonpl. 3 (1855) 222, in Walp. Ann. 6 (1864) 927; Naves Novis. App. (1882) 231; Vidal Phan. Cuming. Philip. (1885) 81, 149, Rev. Pl. Vasc. Filip. (1886) 268; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 58; Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, 4 (1909) Bot. 596; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 162; Ames Orch. 2 (1908) 93, text cut, 3 (1908) 6, 5 (1915) 61.

Coelogyne pumila Reichb. f. in Walp. Ann. 6 (1861) 236.

Luzon (Bataan, Laguna, Tayabas). Fairly frequent, especially in Laguna, on the mountains at 950 to 2,000 m altitude.

ACORIDIUM PURPUREUM Ames Orch. 7 (1922) 81.

Dendrochilum purpureum Ames Orch. 6 (1920) 286.

Luzon (Apayao), *Boettcher s. n.* Mt. Pasoan at 2,000 m altitude.

ACORIDIUM QUADRILOBUM Ames Orch. 7 (1922) 81.

Dendrochilum quadrilobum Ames Orch. 5 (1915) 61.

Luzon (Apayao, Ifugao), *Boettcher s. n., B. S. 19894, 19896 McGregor.*

ACORIDIUM RECURVUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 148.

Dendrochilum recurvum Ames in Philip. Journ. Sci. 2 (July, 1907)

Bot. 318; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (November, 1907) 117, fig. 38T; Ames Orch. 2 (1908) 12, 90, t. 22, fig. IV, 4, 5 (1915) 62.

Luzon (Lepanto, Benguet), Mindoro, Merrill 4584, 5509, 5831, Ramos & Edaño 2, 4, 23, 27. In mossy forest, in mountain regions at 2,230 to 2,730 m altitude.

ACORIDIUM RENIFORME Ames Orch. 7 (1922) 81.

Dendrochilum reniforme Ames Orch. 5 (1915) 63.

Luzon (Rizal, Laguna), B. S. 13759, 20440 Ramos, Calica s. n. Mt. Canumay and San Antonio.

ACORIDIUM RHOMBOPHORUM (Reichb. f.) Ames Sched. Orch. 5 (1923) 1.

Coelogyne rhombophora Reichb. f. in Linnaea 41 (1877) 116.

Pholidota rhombophora Reichb. f. in Linnaea 41 (1877) 117, in textu;

Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 153.

Acoridium whitfordii Rolfe ex Ames Orch. 1 (1905) 73.

Dendrochilum whitfordii Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 162; Ames Orch. 2 (1908) 100, text cut, 5 (1915) 64.

Dendrochilum rhombophorum Ames Orch. 3 (1908) 7, 5 (1915) 64.

Luzon (Bataan, Rizal, Laguna, Tayabas), Wallis s. n., Whitford 139, Topping 360, Elmer 7395, 7679, F. B. 6516 Curran, B. S. 27935 Ocampo, 28619, 28620 Ramos & Edaño. On mountains, usually at a fairly high altitude.

ACORIDIUM SACCOLABIUM (Kränzl.) Ames Sched. Orch. 5 (1923) 2.

Dendrochilum saccolabium Kränzl. in Ann. k. k. Hofmus. Wien 30 (1916) 56.

Without locality, *Loher* s. n. A fragment from the living type plant cultivated at Nymphenburg, near Munich, Bavaria, is in the Ames Herbarium.

ACORIDIUM SERRATOI Ames Sched. Orch. 6 (1923) 27, fig. 6.

Luzon (Laguna), *Serrato* s. n. In dense forest.

ACORIDIUM SIMULACRUM Ames Orch. 7 (1922) 82.

Dendrochilum simulacrum Ames Orch. 2 (1908) 92, text cut, 5 (1915) 64.

Luzon (Benguet, Tayabas), F. B. 4981 Curran, Elmer 7586. Mt. Tooglon and Lucban.

ACORIDIUM SPHACELATUM Ames Orch. 1 (1905) 1, t. 1, figs. 1, 4-6, 8, 10, 12, 14, 16, 18, 20.

Dendrochilum sphacelatum Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 112, fig. 37 A, B, D, F, H, K; Ames Orch. 2 (1908) 82, 5 (1915) 64, 6 (1920) 287.

Luzon (Benguet), Elmer 6554, Boettcher s. n., Sandkuhl 97. Mt. Santo Tomas at 2,300 m altitude.

ACORIDIUM STRICTIFORME Ames in Proc. Biol. Soc. Wash. 19 (1906) 147.

Dendrochilum strictiforme Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38L; Ames Orch. 2 (1908) 12, 89, t. 21, fig. III, 3, 5 (1915) 64, in Philip. Journ. Sci. 4 (1909) Bot. 597.

Luzon (Lepanto, Benguet, Tayabas), Leyte. A fairly common species in the mountains at 660 to 2,300 m altitude.

ACORIDIUM TENELLUM Nees & Meyen in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 131; Rolfe in Orch. Rev. 12 (1904) 219, 220; Ames Orch. 1 (1905) 3, t. 1, figs. 2, 3, 7, 11, 13, 15, 17, 19.

Dendrochilum junceum Reichb. f. in Otia Bot. Hamb. 1 (1878) 54; Reichb. f. Xen. Orch. 3 (1881) 30; Naves Novis. App. (1882) 245; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 78.

Dendrochilum tenellum Ames in Philip. Journ. Sci. 2 (July, 1907) Bot. 318; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (November, 1907) 112, fig. 37C, E, G, J, L; Ames Orch. 2 (1908) 81, 5 (1915) 64.

Babuyan Islands (Camiguin), Luzon (Apayao, Kalinga, Bontoc, Ifugao, Benguet, Nueva Vizcaya, Zambales, Bataan, Laguna, Sorsogon), Mindoro, Leyte. Probably the commonest and most widely distributed species of the genus in the Philippines, growing at fairly high altitudes on the mountains.

ACORIDIUM TENUE Ames in Proc. Biol. Soc. Wash. 19 (1906) 145.

Dendrochilum tenue Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38M; Ames Orch. 2 (1908) 12, 86, t. 21, fig. V, 5, 5 (1915) 64.

Luzon (Bontoc, Ifugao, Laguna), Negros, Leyte, Mindanao (Misamis, Bukidnon, Agusan). One of the commoner and widely distributed species of the genus, growing in mossy forest at 500 to 2,160 m altitude.

ACORIDIUM TENUIFOLIUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 145.

Dendrochilum tenuifolium Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 114, fig. 38O; Ames Orch. 2 (1908) 12, 86, t. 21, fig. IV, 4, 5 (1915) 64; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340.

Luzon (Lepanto, Benguet, Albay). Rather common in Lepanto and Benguet, in mossy forest at 2,060 to 2,260 m altitude.

ACORIDIUM TURPE Ames in Proc. Biol. Soc. Wash. 19 (1906) 149.

Dendrochilum turpe Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 118, fig. 38D; Ames Orch. 2 (1908) 12, 98, t. 23, fig. III, 3, 5 (1915) 64.

Luzon (Benguet), Merrill 4758, Ramos & Edaño 30, 42. Mountain regions at 500 to 2,260 m altitude.

ACORIDIUM UNICORNE Ames Orch. 7 (1922) 145.

Luzon (Benguet), Ramos & Edaño 43a, 54. Mt. Simakoko, in mossy forest at 2,160 m altitude.

ACORIDIUM VANOVERBERGHII Ames Orch. 7 (1922) 82.

Dendrochilum vanoverberghii Ames in Philip. Journ. Sci. 7 (1912) Bot. 27, Orch. 5 (1915) 65.

Luzon (Bontoc, Ifugao), Vanoverbergh 1046, B. S. 19893 McGregor.

ACORIDIUM WENZELII Ames Orch. 7 (1922) 82.

Dendrochilum wenzelii Ames Orch. 5 (1915) 65.

Luzon (Kalinga, Camarines), Leyte, B. S. 22161 Ramos, 42088 Edaña, Wenzel 0264, 0338, 0787, 0803, 0862, 0876, 0880, 0886, Boettcher s. n. In forest at 500 to 700 m altitude.

ACORIDIUM WILLIAMSII Ames in Proc. Biol. Soc. Wash. 19 (1906) 143.

Dendrochilum williamsii Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 114, fig. 38S; Ames Orch. 2 (1908) 12, 82, t. 22, fig. I, 1, 5 (1915) 66.

Luzon (Lepanto, Benguet), Ramos & Edaña 5, 38, R. S. Williams 1939. Mountain region at fairly high altitude.

EXCLUDED OR SYNONYMOUS SPECIES

ACORIDIUM ARACHNITES Rolfe in Orch. Rev. 12 (1904) 219 = *Dendrochilum arachnites* Reichb. f.

ACORIDIUM COBBIANUM Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum cobbianum* Reichb. f.

ACORIDIUM CONVALLARIAEFORME Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum convallariaeforme* Schauer.

ACORIDIUM COPELANDII Ames in Proc. Biol. Soc. Wash. 19 (1906) 152 = *Dendrochilum copelandii* Ames.

ACORIDIUM CUCULLATUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 152 = *Dendrochilum cucullatum* (Ames) Pfitz.

ACORIDIUM CUCUMERINUM Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum cucumerinum* Reichb. f., spec. excl.

ACORIDIUM FILIFORME Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum filiforme* Lindl.

ACORIDIUM GLUMACEUM Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum glumaceum* Lindl.

ACORIDIUM GRACILISCAPUM Ames in Proc. Biol. Soc. Wash. 19 (1906) 152 = *Dendrochilum graciliscapum* (Ames) Pfitz.

ACORIDIUM HUTCHINSONII Ames Orch. 7 (1922) 81, sphalm = *Acoridium hutchinsonianum* Ames.

ACORIDIUM LATIFOLIUM Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum latifolium* Lindl.

ACORIDIUM LINEARIFOLIUM Ames Orch. 7 (1922) 141, non Rolfe = *Acoridium montanum* Ames.

ACORIDIUM LONGILABRE Ames in Proc. Biol. Soc. Wash. 19 (1904) 152 = *Dendrochilum longilabre* (Ames) Pfitz.

ACORIDIUM MAGNUM Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum magnum* Reichb. f.

ACORIDIUM PARVULUM Ames in Proc. Biol. Soc. Wash. 19 (1904) 146 = *Dendrochilum parvulum* (Ames) Pfitz.

ACORIDIUM PHILIPPINENSE Ames in Proc. Biol. Soc. Wash. 19 (1904) 149 = *Dendrochilum philippinense* (Ames) Pfitz.

ACORIDIUM UNCATUM Rolfe in Orch. Rev. 12 (1904) 220 = *Dendrochilum uncatum* Reichb. f.

ACORIDIUM VENUSTULUM Ames in Proc. Biol. Soc. Wash. 19 (September, 1906) 147 = *Acoridium microchilum* (Schltr.) Ames.

ACORIDIUM WHITFORDII Rolfe ex Ames Orch. 1 (1905) 73 = *Acoridium rhombophorum* (Reichb. f.) Ames.

42. DENDROCHILUM Blume

DENDROCHILUM ARACHNITES Reichb. f. in Gard. Chron. II 17 (1882) 256; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 65; Ames in Philip. Journ. Sci. 2 (1907) Bot. 316; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 92; Ames Orch. 2 (1908) 119, *text cut*; 5 (1915) 55; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340.

Platyclinis arachnites Rolfe in Kew Hand-list Orch. (1896) 189, *nomen*.

Acoridium arachnites Rolfe in Orch. Rev. 12 (1904) 219.

Luzon (Lepanto, Benguet, Rizal), Mindoro, Leyte, Mindanao (Zamboanga, Bukidnon, Agusan, Davao). A common and rather widely distributed species. Epiphyte in forest at 660 to 2,300 m altitude.

DENDROCHILUM BICALLOSUM Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, *nomen*, Orch. 2 (1908) 117, *text cut*, 5 (1915) 56, in Philip. Journ. Sci. 4 (1909) Bot. 596.

Luzon (Laguna, Camarines), Mindoro, *F. B. 7797 Curran & Merritt, B. S. 21983 Ramos, 26674 Brown, 26438 Catalan, Santos s. n., Merrill 5812*. Mountain regions, epiphyte in trees at 300 to 830 m altitude.

Var. *MINOR* Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, *nomen*, Orch. 2 (1908) 118.

Mindoro, *Merrill 5663*. Mt. Halcon.

DENDROCHILUM CAGAYANENSE Ames in Philip. Journ. Sci. 6 (1911) Bot. 41, Orch. 5 (1915) 56.

Luzon (Cagayan, Apayao, Rizal, Laguna), Leyte, Mindanao (Davao), *F. B. 16722 Curran, B. S. 28425 Fenix, 15392, 16583 Ramos, Reillo s. n., Wenzel 0512, 0532, 0602, Elmer 11844*. Epiphyte, in forest, up to 700 m altitude in Leyte.

DENDROCHILUM CLEMENSIAE Ames Orch. 2 (1908) 109, *text cut*, 5 (1915) 56.

Mindanao (Lanao), *M. S. Clemens s. n., 642*.

DENDROCHILUM COBBIANUM Reichb. f. in Gard. Chron. II 14 (1880) 748; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 75; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 101; Ames Orch. 2 (1908) 112, *text cut*, 5 (1915) 56.

Platyclinis cobbiana Hemsl. in Gard. Chron. II 16 (1881) 656; Veitch Man. Orch. Pl. pt. 5 (1889) 80.

Acoridium cobbianum Rolfe in Orch. Rev. 12 (1904) 220.

Luzon (Bontoc, Benguet), Mindanao (Agusan). A fairly common species in Luzon at 1,400 to 2,400 m altitude.

DENDROCHILUM CONVALLARIAEFORME Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 429; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 97 (excl. syn. *Acoridium copelandii* Ames); Ames Orch. 5 (1915) 56.

Acoridium convallariaeforme Rolfe in Orch. Rev. 12 (1904) 220.

Manila, *Meyen*.

No specimens of this species have been seen.

DENDROCHILUM COPELANDII Ames Orch. 2 (1908) 12, 115, t. 24, fig. I, 1, 5 (1915) 56.

Acoridium copelandii Ames in Proc. Biol. Soc. Wash. 19 (1906) 153.

Mindanao (Zamboanga), *Copeland s. n.*

DENDROCHILUM CUCULLATUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 98, fig. 38J; Ames Orch. 2 (1908) 12, 115, t. 24, fig. III, 3, 5 (1915) 56.

Acoridium cucullatum Ames in Proc. Biol. Soc. Wash. 19 (1906) 153.

Mindanao (Davao), *Copeland s. n.* Mt. Apo, on mossy trees at 2,000 m altitude.

DENDROCHILUM CYMBIFORME Ames in Philip. Journ. Sci. 6 (1911) Bot. 41, Orch. 5 (1915) 57.

Luzon (Abra, Kalinga), *B. S. 7133, 7135 Ramos, Boettcher s. n.* Mt. Bawagan and Mt. Pastan, at 550 to 2,000 m altitude.

DENDROCHILUM ECALLOSUM Ames Orch. 6 (1920) 282.

Catanduanes, *B. S. 30240 Ramos*. Mt. Mariquidon, on tree in forest at 140 m altitude.

DENDROCHILUM FILIFORME Lindl. in Bot. Reg. 26 (1840) Misc. 52; Regel in Gartenfl. 18 (1869) 33, t. 604; Naves Novis. App. (1882) 231; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 68; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 97; Ames Orch. 2 (1908) 113, text cut, 5 (1915) 57.

Dendrochilum glumaceum André in Ill. Hort. 25 (1878) t. 323, non Lindl.

Platyclinis filiformis Benth. ex Hemsl. in Gard. Chron. II 16 (1881) 656; Veitch Man. Orch. Pl. pt. 5 (1889) 80; Cogn. in Dict. Icon. Orch. (1902) *Platyclinis t. 1*.

Acoridium filiforme Rolfe in Orch. Rev. 12 (1904) 220.

Luzon (Rizal), *Cuming s. n., Ramos s. n.* Up to 660 m altitude.

DENDROCHILUM GLUMACEUM Lindl. in Bot. Reg. 27 (1841) Misc. 23; Hook. in Bot. Mag. 81 (1855) t. 4853; Miq. Fl. Ind. Bat. 3 (1859) 627; Reichb. f. in Walp. Ann. 6 (1861) 218; Batem. Second Cent. Orch. Pl. (1867) t. 134; Naves Novis. App. (1882) 231; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 76; Ames in Philip. Journ. Sci. 2 (1907) Bot. 317; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 104; Ames Orch. 2 (1908) 110, text cut, 5 (1915) 57.

Platyclinis glumacea Benth. in Journ. Linn. Soc. 18 (1881) 295, by inference only; ex Hemsl. in Gard. Chron. II 16 (1881) 656; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1888) 128,

fig. 127 B-F; Veitch Man. Orch. Pl. pt. 5 (1889) 81, t.; Cogn. in Dict. Icon. Orch. (1902). *Platyclinis* t. 2; Ames Orch. 1 (1905) 74. *Acoridium glumaceum* Rolfe in Orch. Rev. 12 (1904) 220.

Luzon (Isabela, Bontoc, Lepanto, Benguet, Nueva Vizcaya, Pampanga, Bataan, Rizal, Laguna, Tayabas, Camarines), Catanduanes, Leyte, Mindanao (Davao). The commonest species of the genus in the Philippines, especially so in Luzon. Epiphyte in forest at 30 to 2,300 m altitude.

Var. **VALIDUM** (Rolfe). Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 104; Ames Orch. 2 (1908) 112.

Platyclinis glumacea Benth. ex Hemsl. var. *valida* Rolfe in Orch. Rev. 1 (1893) 115, 12 (1904) 73, fig. 18.

DENDROCHILUM GRACILISCAPUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 99, fig. 38H; Ames Orch. 2 (1908) 12, 118; t. 24, fig. II, 2, 5 (1915) 57.

Acoridium graciliscapum Ames in Proc. Biol. Soc. Wash. 19 (1906) 152.

Mindanao (Davao), *Copeland* 1019, *Elmer* 10633, 10647, 11359. Mt. Apo, epiphyte on mossy trees at 1,800 m altitude.

DENDROCHILUM LATIFOLIUM Lindl. in Bot. Règ. 29 (1843) Misc. 56; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 71; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 102, fig. B, C; Ames Orch. 2 (1908) 104, text cut, 5 (1915) 59.

Platyclinis latifolia Hemsl. in Gard. Chron. II 16 (1881) 656; Rendle in Journ. Bot. 34 (1896) 355; Ames Orch. 1 (1905) 74.

Acoridium latifolium Rolfe in Orch. Rev. 12 (1904) 220.

Luzon (Pampanga, Bataan, Tayabas, Camarines), Panay (Antique, Capiz), Negros (Occidental Negros), Mindanao (Lanao, Davao). A common species, especially in Luzon. Epiphyte in mountain regions at 300 to 1,000 m altitude.

DENDROCHILUM LONGILABRE (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 99, fig. 38K; Ames Orch. 2 (1908) 12, 119, t. 24, fig. IV, 4, 5 (1915) 59.

Acoridium longilabre Ames in Proc. Biol. Soc. Wash. 19 (1906) 152.

Mindanao (Davao), *Copeland* 1025, ? *Elmer* 11419. Mt. Apo, epiphyte on mossy trees at 1,530 m altitude.

DENDROCHILUM LONGISPICATUM Ames Orch. 2 (1908) 105, text cut, 5 (1915) 59.

Luzon (Bontoc, Benguet), Sibuyan, Panay (Capiz), Negros (Oriental Negros), *Vanoverbergh* 899, 1533, 1729, 3871, 3940, 4043, *R. S. Williams* 1942, *Merrill* 4336, *Elmer* 9909, 12436, *B. S.* 35586, 35605 *Martelino & Edaño*, 4462 *Mearns*. Epiphyte, in mountain regions up to 1,700 m altitude in Bontoc.

DENDROCHILUM MAGNUM Reichb. f. in Walp. Ann. 6 (1861) 240; Naves Novis. App. (1882) 231; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 74; Ames in Philip. Journ. Sci. 2 (1907) Bot. 317; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 100, fig. 33 H, J; Ames Orch. 2 (1908) 106, text cut, 5 (1915) 60.

Acoridium magnum Rolfe in Orch. Rev. 12 (1904) 220.

Luzon (Rizal), Mindoro, Panay (Antique); *Cuming s. n.*, Merrill 5611, 5730, Ramos s. n., B. S. 32383 McGregor. Mt. Halcon, in mossy forest at 2,000 m altitude, and Mt. Iriga at 1,630 m altitude.

DENDROCHILUM MALINDANGENSE Ames Orch. 2 (1908) 114, *text cut*, 5 (1915) 60.

Mindanao (Dapitan), F. B. 4597 Mearns & Hutchinson. Mt. Malindang at 1,230 m altitude.

DENDROCHILUM MEARNsii Ames Orch. 2 (1908) 116, *text cut*, 5 (1915) 60.

Camiguin de Misamis, Mindanao (Misamis, Bukidnon); B. S. 14426 Ramos, 38523 Ramos & Edaño, F. B. 4605 Mearns & Hutchinson. Up to 3,060 m altitude on Mt. Malindang.

DENDROCHILUM NIVEUM Ames Orch. 6 (1920) 284.

Luzon (Kalinga), Boettcher s. n. Balbalan at 1,000 m altitude.

DENDROCHILUM PALAWANENSE Ames Orch. 2 (1908) 103, *text cut*, 5 (1915) 61, in Elm. Leaf. Philip. Bot. 5 (1912) 1559.

Palawan, B. S. 553, 557 Foxworthy, Elmer 13195. In mossy scrub at 1,250 m altitude.

DENDROCHILUM PANGASINANENSE Ames Sched. Orch. 6 (1923) 29.

Luzon (Pangasinan), B. S. 8286 Ramos.

DENDROCHILUM PRODIGIOSUM Ames Orch. 7 (1922) 98.

Mindanao (Bukidnon), B. S. 38521 Ramos & Edaño. Mt. Lipa, in mossy forest at 2,160 m altitude.

DENDROCHILUM PROPINQUUM Ames Orch. 6 (1920) 285.

Leyte, Wenzel 0279, 0329, 0376, 0398, 0449, 0497, 0548, 0582, 0682, 0805, 0809, 0857, 0868, 0911, 0920. Jaro, epiphyte in forest at 500 to 700 m altitude.

DENDROCHILUM RAMOSII Ames in Philip. Journ. Sci. 8 (1913) Bot. 410, Orch. 5 (1915) 62.

Luzon (Laguna), B. S. 15003 Ramos, Reillo 4. San Antonio.

DENDROCHILUM UNCATUM Reichb. f. in Bonpl. 3 (1855) 222, in Walp. Ann. 6 (1864) 927, in Gard. Chron. II 16 (1881) 780; Naves Novis. App. (1882) 231; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 72; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 103, fig. 35 G, H; Ames Orch. 2 (1908) 108, *text cut*, 5 (1915) 65; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340.

Dendrochilum uncatum Reichb. f. var. *lancifolium* Reichb. f. in Gard. Chron. II 16 (1881) 780; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 104.

Platyclinis uncata Rolfe ex Vidal Phan. Cuming. Philip. (1885) 148; Vidal Rev. Pl. Vasc. Filip. (1886) 267.

Acoridium uncatum Rolfe in Orch. Rev. 12 (1904) 220.

Luzon (Benguet, Nueva Vizcaya, Pampanga, Bataan, Rizal, Laguna), Panay (Antique). A fairly common species in the mountains of Luzon, especially of Benguet Subprovince, at 1,000 to 2,300 m altitude.

DENDROCHILUM WEBERI Ames in Philip. Journ. Sci. 8 (1913) Bot. 410, Orch. 5 (1915) 65.

Mindanao (Agusan), Basilan, *Weber 59, B. S. 16240 Reillo*. Type from Cabadbaran River at 80 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

DENDROCHILUM AFFINE Ames Orch. 2 (1908) 95, 5 (1915) 55 = *Acoridium affine* Ames.

DENDROCHILUM ANFRACTOIDES Ames Orch. 3 (1908) 13, t. 28, 5 (1915) 55, in Philip. Journ. Sci. 4 (1909) Bot. 665; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340 = *Acoridium anfractoides* Ames.

DENDROCHILUM ANFRACTUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 118, fig. 38P; Ames Orch. 2 (1908) 91, t. 21, fig. I. 1, 5 (1915) 55 = *Acoridium anfractum* Ames.

DENDROCHILUM AURICULARE Ames in Philip. Journ. Sci. 4 (1909) Bot. 595, Orch. 5 (1915) 55 = *Acoridium auriculare* Ames.

DENDROCHILUM AURITUM Reichb. f. apud Naves Novis. App. (1882) 231.

DENDROCHILUM BINUANGENSE Ames Orch. 6 (1920) 281 = *Acoridium binuangense* Ames.

DENDROCHILUM BISTORTUM (Wendl. & Kränzl.) J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 64; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 96, fig. 34 A-D; Ames Orch. 5 (1915) 54.

Platyclinis bistorta Wendl. & Kränzl. Xen. Orch. 3 (1900) 169, t. 299, fig. I. 1-5.

"Indischer Archipel. Insel Maschate (?) leg. W. Micholitz, imp. F. Sander."—Wendland and Kränzl. Merrill writes: "I do not know of any Malayan island by this name and suspect that Masbate of the Philippine Archipelago was intended; Micholitz has carried on extensive explorations in the Philippines." It has not been possible to refer this to any species of Philippine origin.

DENDROCHILUM CINNABARINUM Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 104, fig. 35F; Ames Orch. 3 (1908) 10, 5 (1915) 56, 6 (1920) 282, in Philip. Journ. Sci. 4 (1909) Bot. 597; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340 = *Acoridium cinnabarinum* (Pfitz.) Ames.

DENDROCHILUM CONFUSUM Ames in Philip. Journ. Sci. 6 (1911) Bot. 41, Orch. 5 (1915) 56 = *Acoridium confusum* Ames.

DENDROCHILUM CUCUMERINUM Reichb. f. in Gard. Chron. II 22 (1884) 56, 649; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 71; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 103, fig. 35D, E; Ames Orch. 5 (1915) 54, 57.

Platyclinis cucumerina Rolfe in Kew Hand-list Orch. (1896) 189, nomen.

Acoridium cucumerinum Rolfe in Orch. Rev. 12 (1904) 220.

This was described from cultivated specimens of unknown origin and has been ascribed to the Philippines; it may prove to be a native of some other region.

- DENDROCHILUM CURRANII* Ames Orch. 3 (1908) 15, t. 29, 5 (1915) 57 = *Acoridium curranii* Ames.
- DENDROCHILUM ELMERI* Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1558, Orch. 5 (1915) 57 = *Acoridium elmeri* Ames.
- DENDROCHILUM EXILE* Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, nomen, Orch. 2 (1908) 84, text cut, 5 (1915) 57 = *Acoridium exile* Ames.
- DENDROCHILUM FOXWORTHYI* Ames Orch. 3 (1908) 8, t. 26, 5 (1915) 57 = *Acoridium foxworthyi* Ames.
- DENDROCHILUM GLUMACEUM* André in Ill. Hort. 25 (1878) t. 323, non Lindl. = *Dendrochilum filiforme* Lindl.
- DENDROCHILUM GRAMINIFOLIUM* (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 114, fig. 38 Q, R; Ames Orch. 2 (1908) 12, 83, t. 22, fig. II, 2, 5 (1915) 57 = *Acoridium graminifolium* Ames.
- DENDROCHILUM GRANDIFLORUM* Schltr. in Fedde Repert. 8 (1910) 563; Ames Orch. 5 (1915) 54, 58, non Pfitz.
- The geographic origin of this species is not certainly known; Schlechter stated "Philippinen?," but a drawing received from him is marked "India." This species is very close to *Dendrochilum latifolium* Lindl.
- DENDROCHILUM HASTATUM* Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, nomen, Orch. 2 (1908) 99, text cut, 5 (1915) 58 = *Acoridium hastatum* Ames.
- DENDROCHILUM HUTCHINSONIANUM* Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, nomen, Orch. 2 (1908) 96, text cut, 5 (1915) 58, 7 (1922) 81, in syn. (as *D. hutchinsonii*) = *Acoridium hutchinsonianum* Ames.
- DENDROCHILUM HYMENANTHUM* Reichb. f. apud Vidal Phan. Cuming. Philip. (1885) 149; Vidal Rev. Pl. Vasc. Filip. (1886) 268, sphalm = *Dendrobium hymenanthum* Reichb. f.
- DENDROCHILUM IRIGENSE* Ames Orch. 5 (1915) 58 = *Acoridium irigense* Ames.
- DENDROCHILUM JUNCEUM* Reichb. f. Otia Bot. Hamb. 1 (1878) 54, Xen. Orch. 3 (1881) 30; Naves Novis. App. (1882) 231; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 78 = *Acoridium tenellum* Nees & Meyen.
- DENDROCHILUM LOHERI* Ames Orch. 3 (1908) 12, t. 27, I, A, 5 (1915) 59, 6 (1920) 283; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340 = *Acoridium loheri* Ames.
- DENDROCHILUM LONGIBULBUM* Ames in Philip. Journ. Sci. 7 (1912) Bot. 26, Orch. 5 (1915) 59 = *Acoridium longibulbum* Ames.
- DENDROCHILUM LONGIFOLIUM* Reichb. f. apud Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 108, fig. 36 B (as to *Jagor 680*).
- Jagor 680* is a specimen from Luzon, probably incorrectly identified by Pfitzer who had not seen the type of the species, which is a native of the Malay Peninsula.
- DENDROCHILUM LUCBANENSE* Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1559, Orch. 5 (1915) 60 = *Acoridium lucbanense* Ames.

- DENDROCHILUM LUZONENSE Ames Orch. 2 (1908) 121, *text cut*, 5 (1915) 60 = *Acoridium luzonense* Ames.
- DENDROCHILUM MACGREGORII Ames in Philip. Journ. Sci. 6 (1911) Bot. 42, Orch. 5 (1915) 60 = *Acoridium macgregorii* Ames.
- DENDROCHILUM MALEOLENS Kränzl. in Orchis 2 (1908) 63; Ames Orch. 5 (1915) 54, 60 = *Acoridium maleolens* Ames.
- DENDROCHILUM MERRILLII (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 119, *fig. 38 A, B*; Ames Orch. 2 (1908) 12, 102, *t. 23, fig. I, 1*, 5 (1915) 60 = *Acoridium merrillii* Ames.
- DENDROCHILUM MICROCHILUM (Schltr.) Ames Orch. 2 (1908) 87, in *textu*, 5 (1915) 60, in Philip. Journ. Sci. 7 (1912) Bot. 26 = *Acoridium microchilum* (Schltr.) Ames.
- DENDROCHILUM MINDORENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, *nomen*, Orch. 2 (1908) 91, *text cut*, 5 (1915) 60 = *Acoridium mindorense* Ames.
- DENDROCHILUM OCELLATUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 117, *fig. 38 C*; Ames Orch. 2 (1908) 12, 101, *t. 23, fig. II, 2*, 5 (1915) 61, in Philip. Journ. Sci. 4 (1909) Bot. 596 = *Acoridium ocellatum* Ames.
- DENDROCHILUM OLIGANTHUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 117, *fig. 38 E*; Ames Orch. 2 (1909) 12, 98, *t. 23, fig. IV, 4*, 5 (1915) 61 = *Acoridium oliganthum* Ames.
- DENDROCHILUM PARVULUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, *fig. 38 N*; Ames Orch. 2 (1908) 12, 87, *t. 21, fig. II, 2*, 5 (1915) 61 = *Acoridium parvulum* Ames.
- DENDROCHILUM PHILIPPINENSE (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 118, *fig. 38 F*; Ames Orch. 2 (1908) 12, 97, *t. 23, fig. V, 5*, 5 (1915) 61, in Philip. Journ. Sci. 4 (1909) Bot. 595 = *Acoridium philippinense* Ames.
- DENDROCHILUM PULOGENSE Ames in Philip. Journ. Sci. 4 (1909) 594; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340; Ames Orch. 5 (1915) 61 = *Acoridium pulogense* Ames.
- DENDROCHILUM PUMILUM Reichb. f. in Bonpl. 3 (1855) 222, in Walp. Ann. 6 (1864) 927; Naves Novis. App. (1882) 231; Vidal Phan. Cuming. Philip. (1885) 81, 149, Rev. Pl. Vasc. Filip. (1886) 268; J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 58; Ames in Philip. Journ. Sci. 2 (1907) Bot. 317, 4 (1909) Bot. 596; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 162; Ames Orch. 2 (1908) 93, *text cut*, 3 (1908) 6, 5 (1915) 61 = *Acoridium pumilum* (Reichb. f.) Rolfe.
- DENDROCHILUM PURPUREUM Ames Orch. 6 (1920) 286 = *Acoridium purpureum* Ames.
- DENDROCHILUM QUADRILOBUM Ames Orch. 5 (1915) 61 = *Acoridium quadrilobum* Ames.

DENDROCHILUM RECURVUM Ames in Philip. Journ. Sci. 2 (July, 1907) Bot. 318; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (November, 1907) 117, fig. 38 T; Ames Orch. 2 (1908) 12, 90, t. 22, fig. IV, 4, 5 (1915) 62 = *Acoridium recurvum* Ames.

DENDROCHILUM RENIFORME Ames Orch. 5 (1915) 63 = *Acoridium reniforme* Ames.

DENDROCHILUM RHOMBOPHORUM (Reichb. f.) Ames Orch. 3 (1908) 7, 5 (1915) 64 = *Acoridium rhombophorum* (Reichb. f.) Ames.

DENDROCHILUM SACCOLABIUM Kränzl. in Ann. k. k. Hofmus. Wien 30 (1916) 56 = *Acoridium saccolabium* (Kränzl.) Ames.

DENDROCHILUM SIMULACRUM Ames Orch. 2 (1908) 92, text cut, 5 (1915) 64 = *Acoridium simulacrum* Ames.

DENDROCHILUM SPHACELATUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 112, fig. 37 A, B, D, F, H, K; Ames Orch. 2 (1908) 82, 5 (1915) 64, 6 (1920) 287 = *Acoridium sphacelatum* Ames.

DENDROCHILUM STRICTIFORME (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38 L; Ames Orch. 2 (1908) 12, 89, t. 21, fig. III, 3, 5 (1915) 64, in Philip. Journ. Sci. 4 (1909) Bot. 597 = *Acoridium strictiforme* Ames.

DENDROCHILUM TENELLUM (Nees & Meyen) Ames in Philip. Journ. Sci. 2 (July, 1907) Bot. 318; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (November, 1907) 112, fig. 37 C, E, G, J, L; Ames Orch. 2 (1908) 81, 5 (1915) 64 = *Acoridium tenellum* Nees & Meyen.

DENDROCHILUM TENUE (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38 M; Ames Orch. 2 (1908) 12, 86, t. 21, fig. V, 5, 5 (1915) 64 = *Acoridium tenue* Ames.

DENDROCHILUM TENUIFOLIUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 114, fig. 38 O; Ames Orch. 2 (1908) 12, 85, t. 21, fig. IV, 4, 5 (1915) 64; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 340 = *Acoridium tenuifolium* Ames.

DENDROCHILUM TURPE (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 118, fig. 38 D; Ames Orch. 2 (1908) 12, 98, t. 23, fig. III, 3, 5 (1915) 64 = *Acoridium turpe* Ames.

DENDROCHILUM VANOVERBERGHII Ames in Philip. Journ. Sci. 7 (1912) Bot. 27, Orch. 5 (1915) 65 = *Acoridium vanoverberghii* Ames.

DENDROCHILUM VENUSTULUM (Ames) Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 116, fig. 38 U; Ames Orch. 2 (1908) 12, 88, t. 22, fig. III, 3, in Philip. Journ. Sci. 4 (1909) Bot. 595; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341 = *Acoridium microchilum* (Schltr.) Ames.

DENDROCHILUM WENZELII Ames Orch. 5 (1915) 65 = *Acoridium wenzelii* Ames.

DENDROCHILUM WHITFORDII (Rolfe) Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 162; Ames Orch. 2 (1908) 100, *text cut*, 5 (1915) 66 = *Acoridium rhombophorum* (Reichb. f.) Ames.

DENDROCHILUM WILLIAMSII Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 114, *fig. 38 S*; Ames Orch. 2 (1908) 12, 82, *t. 22, fig. I, 1*, 5 (1915) 66 = *Acoridium williamsii* Ames.

DENDROCHILUM WOODIANUM Ames in Philip. Journ. Sci. 2 (1907) Bot. 318, *nomen*, Orch. 2 (1908) 80, *text cut*, 5 (1915) 66 = *Pseudacoridium woodianum* Ames.

43. PHOLIDOTA Lindley

PHOLIDOTA CARNEA (Blume) Lindl. Gen. & Sp. Orch. (1830) 37; Miq. Fl. Ind. Bat. 3 (1859) 669; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 158, Fig.-Atlas (1909) *fig. 118*; Ames Orch. 6 (1920) 66, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 149, Sched. Orch. 6 (1923) 20.

Crinonia carnea Blume Bijdr. (1825) 339.

Coelogyne carnea Reichb. f. in Walp. Ann. 6 (1861) 237.

Crinonia elmeri Ames Orch. 5 (1915) 67, *nomen*.

Mindanao, *Elmer 13834*.

PHOLIDOTA IMBRICATA Lindl. in Hook. Exot. Fl. 2 (January, 1825) *t. 138*, in Bot. Reg. 14 (1829) *t. 1213*, 21 (1835) *t. 1777*, Gen. & Sp. Orch. (1830) 36; Lodd. Bot. Cab. 20 (1833) *t. 1934*; Wight Icon. 3² (1849) 9, *t. 907*; Dalz. & Gibs. Bomb. Fl. (1861) 262; Thw. Enum. Pl. Zeyl. (1864) 300; Benth. Fl. Austral. 6 (1873) 290; Naves Novis. App. (1882) 237; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 269; Fitzger. Austral. Orch. 2⁴ (1893) *t.*; Hook. f. Fl. Brit. Ind. 5 (1890) 845; Grant Orch. Burma (1895) 161; Ridl. in Journ. Linn. Soc. 31 (1896) 287, 32 (1896) 328; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 144, *t. 201*; Trim. Handb. Fl. Ceyl. 4 (1898) 162; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 151, Fig.-Atlas (1909) *fig. 112*; Ames Orch. 1 (1905) 72, 2 (1908) 75, 5 (1915) 66; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 115; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 154; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 135; J. J. Sm. in Nova Guinea 8 (1909) Bot. 21, 8³ (1912) Bot. 527, 12 (1913) Bot. 11; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 11, in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 188; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 149.

Cymbidium imbricatum Roxb. Hort. Beng. (1814) 63, *nomen*, Fl. Ind. ed. 2, 3 (1832) 460.

Ptilocnema bracteatum D. Don Prodr. Fl. Nepal. (February, 1825) 33.

Ornithidium imbricatum Wall. ex Lindl. in Bot. Reg. 14 (1829) sub *t. 1213*, in syn.

Pholidota pallida Lindl. in Bot. Reg. 21 (1835) sub *t. 1777*; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 153.

Pholidota conchoidea Lindl. in Bot. Reg. 26 (1840) Misc. 84; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 154, fig. 50 K.

Pholidota loricata Reichb. f. in Bonpl. 5 (1857) 43; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 153.

Coelogyne conchoidea Reichb. f. in Walp. Ann. 6 (1861) 237; Naves Novis. App. (1882) 237.

Coelogyne imbricata Reichb. f. in Walp. Ann. 6 (1861) 238.

Coelogyne pallida Reichb. f. in Walp. Ann. 6 (1861) 238.

Coelogyne triotos Reichb. f. in Walp. Ann. 6 (1861) 238, in Hamb. Gartenz. 18 (1862) 34; Naves Novis. App. (1882) 237.

Pholidota triotos Reichb. f. in Walp. Ann. 6 (1861) 238, in syn.; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 154.

Luzon (Bontoc, Benguet, Bulacan, Bataan, Rizal), Polillo, Panay (Capiz), Samar, Leyte, Palawan, Mindanao (Zamboanga, Lanao, Davao), Basilan. A common and widely distributed species in the Philippines at 15 to 1,600 m altitude. In India, Burma, Assam, Siam, Malay Peninsula, China, Sumatra, Java, Celebes, Borneo, and New Guinea.

PHOLIDOTA VENTRICOSA (Blume) Reichb. f. in Bonpl. 5 (1857) 43; Rolfe in Gard. Chron. III 5 (1889) 585; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 154, Fig.-Atlas (1909) fig. 114; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 148, fig. 51; Ames Orch. 2 (1908) 75, 5 (1915) 66, in Philip. Journ. Sci. 4 (1909) Bot. 665; J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 43 (1910) 13; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 150.

Chelonanthera ventricosa Blume Bijdr. (1825) 383, fig. 51; Lindl. Gen. & Sp. Orch. (1833) 178; Miq. Fl. Ind. Bat. 3 (1859) 673.

Coelogyne ventricosa Reichb. f. in Walp. Ann. 6 (1861) 237.

Pholidota sesquitorta Kränzl. Xen. Orch. 3 (1893) 114, t. 266, fig. I. 1-7; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 146, fig. 50 F.

Luzon (Lepanto), Mindanao (Zamboanga), F. B. 5715 Klemme, Copeland 1621. At about 300 m altitude in Mindanao. In Sumatra, Java, and introduced into Borneo.

EXCLUDED OR SYNONYMOUS SPECIES

PHOLIDOTA CONCHOIDEA Lindl. in Bot. Reg. 26 (1840) Misc. 84; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 154, fig. 50 K = *Pholidota imbricata* Lindl.

PHOLIDOTA ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1557 = *Thecostele alata* (Roxb.) Par. & Reichb. f.

PHOLIDOTA GLOBOSA Lindl. apud Naves Novis. App. (1882) 237.

PHOLIDOTA MEMBRANACEA Teijsm. & Binn. apud Naves Novis. App. (1882) 237.

PHOLIDOTA RHOMBOPHORA Reichb. f. in Linnaea 41 (1877) 116, in textu; Pfitz. & Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 153 = *Acoridium rhombophorum* (Reichb. f.) Ames.

PHOLIDOTA TRIOTOS Reichb. f. in Walp. Ann. 6 (1861) 238, in syn.; Pfitz. in Engl. Pflanzenreich IV. 50. II. B. 7 (1907) 154 = *Pholidota imbricata* Lindl.

44. **MALAXIS** Solander ex Swartz

MALAXIS ACUMINATA D. Don Prodr. Fl. Nepal. (1825) 29.

Microstylis wallichii Lindl. in Wall. Cat. (1828) No. 1938, *nomen*. Gen. & Sp. Orch. (1830) 20; Ridl. in Journ. Linn. Soc. 24 (1888) 338.

Var. **BILOBA** (Lindl.) Ames comb. nov.

Microstylis biloba Lindl. in Wall. Cat. (1828) No. 1940, *nomen*. Gen. & Sp. Orch. (1830) 20; Par. & Reichb. f. in Trans. Linn. Soc. 30 (1874) 139.

Dienia carinata Reichb. f. in Bonpl. 3 (1855) 223.

Microstylis carinata Reichb. f. in Walp. Ann. 6 (1861) 207; Naves Novis. App. (1882) 230; Vidal Phan. Cuming. Philip. (1885) 148; Rev. Pl. Vasc. Filip. (1886) 266; Ridl. in Journ. Linn. Soc. 24 (1888) 338.

Microstylis wallichii Lindl. var. *biloba* Hook. f. Fl. Brit. Ind. 5 (1888) 686; Grant Orch. Burma (1895) 18; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 16, t. 19; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 88.

Malaxis biloba Ames Orch. 2 (1908) 122, 5 (1915) 69.

Luzon (Bontoc, Bataan, Rizal, Cavite, Batangas, Laguna), Cuming 2144, Lohr s. n., B. S. 1468, 3244 Ahern's collector, 1472, 4562, 13555, 21337, 22384a Ramos, 29515 Ramos & Edaña, Merrill 4175, Leiberg 6081, Native collector s. n., Vanoverbergh 1349, F. B. 24298, 24298bis Bawan & Borromeo. Terrestrial in forest. In India and Burma. The species proper does not occur in the Philippines.

MALAXIS ALAGENSIS Ames Orch. 2 (1908) 122, *text cut*, 5 (1915) 68.

Microstylis alagensis Ames in Philip. Journ. Sci. 2 (1907) Bot. 318.

Luzon (Tayabas); Mindoro, B. S. 25392, 25432, 25543 Yates, Merrill 5801, 5807, Ramos s. n. Terrestrial in humid forest at 130 to 400 m altitude.

MALAXIS ARIETINA Ames in Philip. Journ. Sci. 8 (1913) Bot. 411, Orch. 5 (1915) 68.

Mindanao (Camiguin Island, Agusan), B. S. 14418, 14443 Ramos, Weber 44, 69. Terrestrial up to 315 m altitude in Agusan.

MALAXIS ATROSANGUINEA Ames Sched. Orch. 6 (1923) 30, fig. 7.

Mindanao (Agusan), Elmer 13815.

MALAXIS BALABACENSIS Ames in Philip. Journ. Sci. 6 (1911) Bot. 42, Orch. 5 (1915) 68.

Luzon (Cavite, Laguna, Sorsogon), Leyte, Palawan, Balabac, Mindanao (Camiguin Island, Lanao, Bukidnon, Agusan, Surigao), Sibutu. A fairly common and rather widely distributed species. Terrestrial, in the forest at relatively low altitude, up to 130 m in Agusan and at 800 m in Cavite.

MALAXIS BANCANOIDES Ames Orch. 2 (1908) 129, *text cut*, 5 (1915) 68.

Sibuyan, *McGregor 14*. Mountains back of San Fernando, terrestrial at about 660 m altitude.

MALAXIS BATAANENSIS Ames in Philip. Journ. Sci. 6 (1911) Bot. 43, Orch. 5 (1915) 69.

Malaxis flavescens Ames Orch. 2 (1908) 127, 5 (1915) 70 (excl. syn.).

Luzon (Bataan), *B. S. 1674 Foxworthy, 6211 Robinson, Merrill 3885*. Lamo Forest Reserve, Mt. Mariveles, at 800 to 1,000 m altitude.

MALAXIS BENGUETENSIS Ames in Philip. Journ. Sci. 6 (1911) Bot. 43, Orch. 5 (1915) 69.

Luzon (Benguet), *B. S. 8362 McGregor, Boettcher s. n.* Pauai at 2,100 m altitude.

MALAXIS BINABAYENSIS Ames Orch. 2 (1908) 124, *text cut*, 5 (1915) 69.

Microstylis binabayensis Ames in Philip. Journ. Sci. 2 (1907) Bot. 318.

Mindoro, *Merrill 5804*. Binabay River, terrestrial in forest at 330 m altitude.

MALAXIS BRACTEOSA Ames Orch. 7 (1922) 111.

Luzon (Sorsogon), Mindanao (Bukidnon, Dapitan), *B. S. 38658, 38673 Ramos & Edaño, F. B. 4575 Mearns & Hutchinson, Elmer 15698*. Mahilucot River, terrestrial along streams, in forest at 100 to 1,400 m altitude.

MALAXIS BULUSANENSIS Ames Sched. Orch. 6 (1923) 33.

Luzon (Sorsogon), *Elmer 14830*.

MALAXIS COPELANDII Ames Orch. 2 (1908) 124, *text cut*, 5 (1915) 69.

Mindanao (Zamboanga), *Copeland s. n.*

MALAXIS CUNEIPETALA Ames Orch. 6 (1920) 288.

Leyte, *Wenzel 0724*. Jaro, terrestrial in forest at 800 m altitude.

MALAXIS CURRANII Ames in Philip. Journ. Sci. 6 (1911) Bot. 44, Orch. 5 (1915) 69.

Luzon (Bontoc, Benguet), *Vanoverbergh 3919, F. B. 5105 Curran, B. S. 12721 Fenix*. Terrestrial, up to 1,300 m altitude in Bontoc.

MALAXIS DAVAENSIS Ames Orch. 5 (1915) 69.

Mindanao (Davao), *Weber 243*. Terrestrial in shade of forest at 1,075 m altitude.

MALAXIS DENTATA Ames Orch. 2 (1908) 125, *text cut*, 5 (1915) 70.

Microstylis dentata Ames in Philip. Journ. Sci. 2 (July, 1907) Bot. 319.

Microstylis philippinensis Kränzl. in Orchis 2 (November, 1907) 16.

Malaxis philippinensis Ames Orch. 5 (1915) 71.

Luzon (Rizal, Sorsogon, Camarines), Polillo, Mindoro, Leyte, Mindanao (Agusan), Jolo. A fairly common species, terrestrial or in trees at 60 to 1,000 m altitude.

MALAXIS ELMERI Ames Orch. 2 (1908) 126, *text cut*, 5 (1915) 70.

Luzon (Benguet), *Elmer 6621, R. S. Williams 1110*.

MALAXIS EPIPHYTICA Ames Sched. Orch. 6 (1923) 34.

Mindanao (Cotabato), *Weber 223*. At 900 m altitude.

MALAXIS HUTCHINSONIANA Ames Orch. 2 (1908) 128, *text cut*, 5 (1915) 70.

Microstylis hutchinsoniana Ames in Philip. Journ. Sci. 2 (1907) Bot. 319.

Mindoro, *Merrill 5809*. Mt. Halcon, terrestrial in forest at 760 m altitude.

MALAXIS LATIFOLIA Sm. in Rees Cycl. 22 (1819) No. 3; Ames in Philip.

Journ. Sci. 6 (1911) Bot. 44, Orch. 5 (1915) 70, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 151.

Malaxis plicata Roxb. Hort. Beng. (1814) 63, *nomen*; Roxb. Fl. Ind. ed. 2, 3 (1832) 456.

Dienia congesta Lindl. in Bot. Reg. 10 (1824) sub t. 825, in Walk. Cat. (1828) No. 1936, Gen. & Sp. Orch. (1830) 22; Reichb. f. in Bonpl. 3 (1855) 250; Benth. Fl. Hongk. (1861) 352.

Gastroglottis montana Blume Bijdr. (1825) 387.

Dienia fusca Lindl. Gen. & Sp. Orch. (1830) 22.

Microstylis congesta Reichb. f. in Walp. Ann. 6 (1861) 206; Hance Suppl. Fl. Hongk. (1872) 39; Naves Novis. App. (1882) 230; Ridl. in Journ. Linn. Soc. 24 (1888) 334, 32 (1896) 223; Hook. f. Fl. Brit. Ind. 5 (1890) 689; Grant Orch. Burma (1895) 16 (excl. syn. in part); King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 19, t. 23; Trim. Handb. Fl. Ceyl. 4 (1898) 141; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 14.

Microstylis trilobulata Kurz Rept. Veg. Andam. Isles App. B (1870) xix, *nomen*; N. E. Br. in Gard. Chron. II 18 (1882) 393.

Microstylis bernaysii F. v. Muell. Fragm. 11 (1878) 21.

Liparis bernaysii F. v. Muell. Fragm. 11 (1878) 21, in syn.; Bernays Fl. Queensl. (?) 507; F. M. Bail. Syn. Queensl. Fl. (1883) 508.

Microstylis congesta Reichb. f. var. *fusca* Ridl. in Journ. Linn. Soc. 24 (1888) 335.

Microstylis latifolia J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 248, Fig.-Atlas (1910) fig. 185; Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 124; J. J. Sm. in Nova Guinea 12 (1913) 15.

Microstylis latifolia J. J. Sm. var. *fusca* J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 249.

Malaxis latifolia Sm. var. *fusca* Ames in Philip. Journ. Sci. 6 (1911) Bot. 45.

Luzon (Bontoc, Benguet, Rizal, Laguna, Sorsogon), Mindanao (Surigao). A fairly common species, though apparently localized. Terrestrial, in forest at 150 to 1,300 m altitude. In India, Ceylon, Burma, Malay Peninsula, Siam, China, Sumatra, Java, Celebes, Borneo, New Guinea, and Australia.

MALAXIS LILACINA Ames Sched. Orch. 6 (1923) 35.

Panay (Antique), Leyte, *McGregor* 6205, *Wenzel* 0641, 0667. At 600 to 900 m altitude.

MALAXIS MACGREGORII Ames in Philip. Journ. Sci. 6 (1911) Bot. 45, Orch. 5 (1915) 71.

Malaxis longipedunculata Ames in Philip. Journ. Sci. 8 (1913) Bot. 411, Orch. 5 (1915) 70.

Luzon (Tayabas), Polillo, Leyte, Mindanao (Agusan, Davao), *B. S.* 10440 *McGregor*, 25539 *Yates*, *Wenzel* 033, 052, 0132, 0237, 0238, 0503, 0847, *Weber* 294, 296, 313. Terrestrial in forest at 60 to 600 m altitude.

MALAXIS MERRILLII Ames Orch. 2 (1908) 128, *text cut*, 5 (1915) 71.

Microstylis merrillii Ames in Philip. Journ. Sci. 2 (1907) Bot. 319.

Mindoro, *Merrill* 5820. Mt. Halcon, terrestrial, at base of cliffs at 1,930 m altitude.

MALAXIS MINDANAENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1560, Orch. 5 (1915) 71.

Mindanao (Davao), *Elmer* 11855. Todaya, Mt. Apo, terrestrial in dense forest at 1,330 m altitude.

MALAXIS MINDORENSIS (Rendle) Ames Orch. 2 (1908) 131, 5 (1915) 71.

Microstylis mindorensis Rendle in Journ. Bot. 34 (1896) 356.

Mindoro, Mt. Dulangan, *Whitehead*. Known only from a photograph and a sketch of the type.

MALAXIS NEGROSIANA Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1561, Orch. 5 (1915) 71, 6 (1920) 288.

Negros (Oriental Negros), Leyte, *Elmer* 9600, *Wenzel* 0641, 0667. Terrestrial or epiphytic in forest at 600 to 1,300 m altitude.

MALAXIS PROPINQUA Ames Orch. 6 (1920) 289.

Luzon (Bontoc), *Vanoverbergh* 3916. Terrestrial, on hills at 1,200 m altitude.

MALAXIS QUADRIDENTATA Ames Orch. 2 (1908) 130, *text cut*, 5 (1915) 72.

Microstylis quadridentata Ames in Philip. Journ. Sci. 2 (1907) Bot. 320.

Mindoro, *Merrill* 5818, 5805. Mt. Halcon, terrestrial in forest at 800 to 1,000 m altitude.

MALAXIS QUADRILOBA Ames Orch. 2 (1908) 131, *text cut*, 5 (1915) 72.

Luzon (Bontoc, Benguet, Bataan), Panay, (Antique), *Vanoverbergh* 786, 1348, 3680, 3944, *R. S. Williams* 1108, *B. S.* 32381 *McGregor*, *Merrill* 3886 in part.

MALAXIS RAMOSII Ames in Philip. Journ. Sci. 6 (1911) Bot. 45, Orch. 5 (1915) 72.

Luzon (Rizal, Laguna), *B. S.* 4567 *Ramos*, 17040 *Robinson*, *Baker* 3573. Up to 700 m altitude in Laguna.

MALAXIS RIZALENSIS Ames in Philip. Journ. Sci. 6 (1911) Bot. 46, Orch. 5 (1915) 72.

Luzon (Rizal), *B. S. 4561 Ramos*. Bosoboso, in thickets.

MALAXIS TAYLORI Ames Orch. 6 (1920) 290.

Mindanao (Agusan), *Elmer 13495, Taylor s. n.*

MALAXIS UNCATA Ames in Philip. Journ. Sci. 6 (1911) Bot. 46, Orch. 5 (1915) 72.

Luzon (Laguna), *F. B. 9552 Curran*. Paete-Piapi.

MALAXIS WENZELII Ames in Philip. Journ. Sci. 8 (1913) Bot. 412, Orch. 5 (1915) 72.

Panay (Capiz), Leyte, *B. S. 31141 Ramos, Wenzel 034, 0130, 0144, 0211, 0239, 0503a*. Terrestrial, in forest at 60 to 500 m altitude.

MALAXIS WILLIAMSII Ames Orch. 2 (1908) 131, *text cut*, 5 (1915) 72.

Luzon (Benguet), *R. S. Williams 1109*. Baguio.

EXCLUDED OR SYNONYMOUS SPECIES

MALAXIS ANCEPS Reichb. f. in Walp. Ann. 6 (1861) 214 = *Oberonia anceps* Lindl.

MALAXIS ANGUSTIFOLIA Blume Bijdr. (1825) 393 = *Liparis caespitosa* (Thou.) Lindl.

MALAXIS APOROPHYLLA Reichb. f. in Walp. Ann. 6 (1861) 214 = *Oberonia aporophylla* Reichb. f.

MALAXIS BILOBA Ames Orch. 2 (1908) 122, 5 (1915) 69 = *Malaxis acuminata* D. Don var. *biloba* Ames.

MALAXIS CAESPITOSA Thou. Orch. Isles Afr. (1822) t. 89 = *Liparis caespitosa* (Thou.) Lindl.

MALAXIS COMPRESSA Blume Bijdr. (1825) 390, fig. 54 = *Liparis compressa* (Blume) Lindl.

MALAXIS CYLINDRICA Reichb. f. in Walp. Ann. 6 (1861) 208 = *Oberonia cylindrica* Lindl.

MALAXIS DISTICHA Thou. Orch. Isles Afr. (1822) t. 88 = *Liparis disticha* (Thou.) Lindl.

MALAXIS FLAVESCENS Ames Orch. 2 (1908) 127, 5 (1915) 70 (excl. syn.) = *Malaxis батаанensis* Ames.

MALAXIS IRIDIFOLIA Reichb. f. in Walp. Ann. 6 (1861) 208 = *Oberonia iridifolia* (Roxb.) Lindl.

MALAXIS LATIFOLIA Blume Bijdr. (1825) 393 = *Liparis latifolia* (Blume) Lindl.

MALAXIS LONGIPEDUNCULATA Ames in Philip. Journ. Sci. 8 (1913) Bot. 411, Orch. 5 (1915) 70 = *Malaxis macgregorii* Ames.

MALAXIS MINIATA Reichb. f. in Walp. Ann. 6 (1861) 215 = *Oberonia miniata* Lindl.

MALAXIS MINIMA Blume Bijdr. (1825) 391 = *Liparis caespitosa* (Thou.) Lindl.

MALAXIS MONSTRUOSA Blume Bijdr. (1825) 395 = *Oberonia monstrosa* (Blume) Lindl.

MALAXIS PALLIDA Blume Bijdr. (1825) 391 = *Liparis pallida* (Blume) Lindl.

MALAXIS PARVIFLORA Blume Bijdr. (1825) 392 = *Liparis parviflora* (Blume) Lindl.

MALAXIS PHILIPPINENSIS Ames Orch. 5 (1915) 71 = *Malaxis dentata* Ames.

MALAXIS THISBE Reichb. f. in Walp. Ann. 6 (1861) 213 = *Oberonia thisbe* Reichb. f.

45. OBERONIA Lindley

OBERONIA ANCEPS Lindl. Sert. Orch. (1838) sub t. 8, Fol. Orch. *Oberonia* (1859) 6; Hook. f. Fl. Brit. Ind. 5 (1888) 685; Ridl. in Trans. Linn. Soc. II 3 (1893) 359; Grant Orch. Burma (1895) 28; Ridl. in Journ. Linn. Soc. 32 (1896) 219; Schltr. in Perk. Frag. Fl. Filip. (1904) 43; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 238, Fig.-Atlas (1910) t. 174; Ames Orch. 1 (1905) 78, 5 (1915) 72; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 18; J. J. Sm. in Bull. Jard. Bot. Buitenz. II 26 (1918) 18.

Oberonia imbricata Wight Icon. 5¹ (1852) 3, t. 1629.

Oberonia griffithii Wight Icon. 5¹ (1852) 4.

Malaxis anceps Reichb. f. in Walp. Ann. 6 (1861) 214.

Oberonia ancipita Naves Novis. App. (1882) 230; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 266; Ceron Cat. Pl. Herb. Manila (1892) 166, sphalm.

Iridorchis anceps Ktze. Rev. Gen. Pl. 2 (1891) 669.

Luzon (Bataan, Laguna, Tayabas), Polillo, Samar, Culion, Mindanao (Agusan), *Serrato s. n.*, B. S. 16580, 24156 Ramos, Merrill 139, 569, *s. n.*, Weber *s. n.*

OBERONIA APOROPHYLLA Reichb. f. in Bonpl. 3 (1855) 223; Lindl. Fol. Orch. *Oberonia* (1859) 6; Naves Novis. App. (1882) 230; Ames Orch. 2 (1908) 140, 5 (1915) 72, in Philip. Journ. Sci. 4 (1909) Bot. 666.

Malaxis aporophylla Reichb. f. in Walp. Ann. 6 (1861) 214.

Oberonia imbricata Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 266, non Lindl.

Iridorchis aporophylla Ktze. Rev. Gen. Pl. 2 (1891) 669.

Luzon (Bontoc, Benguet), Mindanao (Lanao), Cuming 2113, Vanoverbergh 322, 1474, 3827, B. S. 8319, 8508 McGregor, M. S. Clemens 699. Mountainous regions at 1,700 to 2,100 m altitude.

OBERONIA BASILANENSIS Ames Orch. 5 (1915) 72.

Basilan, B. S. 16247 Reillo.

OBERONIA BENGUETENSIS Ames in Philip. Journ. Sci. 7 (1912) Bot. 8, Orch. 5 (1915) 73.

Luzon (Benguet), Merrill 4855. Suyoc to Pauai, epiphyte in mossy forest at 2,200 m altitude.

OBERONIA CYLINDRICA Lindl. in Bot. Reg. 26 (1840) Misc. 20, Fol. Orch. Oberonia (1859) 1; Naves Novis. App. (1882) 230; Ames Orch. 1 (1905) 77, 5 (1915) 73; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Ames in Philip. Journ. Sci. 6 (1911) Bot. 47.

Malaxis cylindrica Reichb. f. in Walp. Ann. 6 (1861) 208.

Iridorchis cylindrica Ktze. Rev. Gen. Pl. 2 (1891) 669.

Luzon (Bontoc, Lepanto, Union, Benguet), Leyte, Bohol, Palawan. A fairly common species in the Philippines. Epiphyte at 6 to 2,650 m altitude.

OBERONIA ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1564, Orch. 5 (1915) 73.

Luzon (Benguet), *Elmer 8434, F. B. 5057 Curran*. Baguio, epiphyte at 1,580 m altitude.

OBERONIA HISPIDULA Ames in Philip. Journ. Sci. 7 (1912) Bot. 8, Orch. 5 (1915) 73.

Luzon (Benguet, Rizal, Tayabas), Mindanao (Agusan), *F. B. 5124 Curran, B. S. 19240 Reillo, Elmer 7406, 9261, 14084*. Epiphytic, at 800 to 1,230 m altitude.

OBERONIA IRIDIFOLIA (Roxb.) Lindl. in Wall. Cat. (1828) No. 1948, p. p., Gen. & Sp. Orch. (1830) 15 (excl. syn. *Malaxis ensiformis* Sm.), Fol. Orch. Oberonia (1859) 1; Benth. Fl. Austral. 6 (1873) 274; Naves Novis. App. (1882) 230; Hook. f. Fl. Brit. Ind. 5 (1888) 675; Grant Orch. Burma (1895) 31; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik-Himal.) (1898) 8, t. 8; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 239, Fig.-Atlas (1910) fig. 175; Ames Orch. 1 (1905) 77, 5 (1915) 73; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 16; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 194; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 152.

Cymbidium iridifolium Roxb. Hort. Beng. (1814) 63, nomen, Fl. Ind. ed. 2, 3 (1832) 458.

? *Cymbidium flavescens* Llanos Fragm. Pl. Filip. (1851) 96; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 74; Merr. Sp. Blancoanae (1918) 113.

• *Malaxis iridifolia* Reichb. f. in Walp. Ann. 6 (1861) 208.

Iridorchis iridifolia Ktze. Rev. Gen. Pl. 2 (1891) 669.

Luzon (Bataan, Rizal, Sorsogon), Palawan, Mindanao (Zamboanga, Davao), Basilan, *F. B. 130 Barnes, 5174 Hutchinson, B. S. 5643, 24100 Ramos, Weber 269, 607*. Epiphytic, up to 400 m in Mindanao. In India, Burma, Malay Peninsula, China, Sumatra, Java, Borneo, and Australia.

OBERONIA LEYTENSIS Ames Orch. 6 (1920) 291, t. 98.

Leyte, *Wenzel 0945*. Epiphyte, in forest at sea level.

OBERONIA LIPENSIS Ames Orch. 7 (1922) 114.

Mindanao (Bukidnon), *B. S. 38535 Ramos*. Mt. Lipa, epiphytic in mossy forest at 2,200 m altitude.

OBERONIA LUZONENSIS Ames Orch. 5 (1915) 74.

Luzon (Laguna, Tayabas), Mindanao (Surigao), *Elmer 17558, B. S. 20309 Escritor, 34489 Ramos & Pascasio.*

OBERONIA MCGREGORII Ames in Philip. Journ. Sci. 2 (1907) Bot. 321, Orch. 3 (1908) 65, t. 48, 5 (1915) 74.

Luzon (Laguna), Mindoro, Mindanao (Agusan), *Reillo 16, B. S. 15009 Ramos, Weber 168.* Epiphytic apparently at low altitudes.

OBERONIA MERRILLII Ames in Philip. Journ. Sci. 7 (1912) Bot. 9, Orch. 5 (1915) 74; Merr. Fl. Manila (1912) 168.

Luzon (Nueva Ecija, Rizal, Tayabas), Busuanga, Mindanao (Zamboanga, Davao), *B. S. 5298 McGregor, Merrill 7348, 11219, Ramos s. n., Weber s. n., ? F. B. 9022 Whitford & Hutchinson, 26587 Reyes, Copeland 1309.* Epiphyte at 18 to 400 m altitude.

OBERONIA MINDORENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, Orch. 3 (1908) 67, t. 49, 5 (1915) 75.

Mindoro, Leyte, Mindanao (Agusan), *Merrill 5613, Wenzel 0420, 0425, 0559, Weber 316.* Epiphyte at 400 to 500 m altitude.

OBERONIA MINIMA Ames Orch. 5 (1915) 75.

Leyte, *Wenzel 046.* Dagami, epiphyte at 60 m altitude.

OBERONIA MINUTISSIMA Ames Orch. 7 (1922) 115.

Mindanao (Surigao) *Wenzel 01030, 01032.* Placer, epiphytic in forest at 150 m altitude.

OBERONIA MONSTRUOSA (Blume) Lindl. Gen. & Sp. Orch. (1830) 17; Miq. Fl. Ind. Bat. 3 (1859) 624; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 235, Fig.-Atlas (1910) fig. 171; Ames Orch. 5 (1915) 76.

Malaxis monstrosa Blume Bijdr. (1825) 395.

Mindanao (Agusan), *Weber 173, 312.* Epiphytic at low altitude. In Java.

OBERONIA OBESA Ames Orch. 5 (1915) 76.

Luzon (Bontoc), *Vanoverbergh 407.*

OBERONIA REILLOI Ames Sched. Orch. 6 (1923) 40.

Luzon (Laguna), *Reillo 20.* Epiphyte.

OBERONIA SETIGERA Ames in Philip. Journ. Sci. 7 (1912) Bot. 10, Orch. 5 (1915) 77.

Luzon (Bataan), *Whitford 1122.* Mt. Mariveles, epiphyte, at 1,200 m altitude.

OBERONIA SURIGAENSIS Ames Orch. 6 (1920) 292.

Mindanao (Surigao), *Wenzel 01159.* Epiphyte, in forest at 150 m altitude.

OBERONIA THISBE Reichb. f. in Bonpl. 3 (1855) 223; Lindl. Fol. Orch. Oberonia (1859) 5; Naves Novis. App. (1882) 230; Ames Orch. 2 (1908) 141, 5 (1915) 77.

Malaxis thisbe Reichb. f. in Walp. Ann. 6 (1861) 213.

Habenaria thisbe Vidal Phan. Cuming. Philip. (1885) 151.

Iridorchis thisbe Ktze. Rev. Gen. Pl. 2 (1891) 669.

Oberonia ruflabris Ames Orch. 1 (1905) 78, non Lindl.

Luzon (Kalinga, Benguet, Laguna), Polillo, Mindanao (Surigao), *Boettcher s. n.*, *Elmer 5839*, *Reillo 52*, *B. S. 15039 Ramos*, *9255 Robinson*, *Cuming 2134*, *Piper 197*. Epiphyte at sea level to 2,000 m altitude.

OBERONIA TOPPINGII Ames in Philip. Journ. Sci. 8 (1913) Bot. 413, Orch. 5 (1915) 77.

Luzon (Rizal), *Topping s. n.* Wawa.

OBERONIA WENZELII Ames Orch. 5 (1915) 77.

Leyte, *Wenzel 0386*, *0388*, *0414*, *0426*, *0436*. Jaro, epiphyte in forest at 500 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

OBERONIA ANCIPITA Naves Novis. App. (1882) 230; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 266; Ceron Cat. Pl. Herb. Manila (1892) 166, sphalm = *Oberonia anceps* Lindl.

OBERONIA IMBRICATA Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 266, non Lindl. = *Oberonia aporophylla* Reichb. f.

OBERONIA MINIATA Lindl. in Bot. Reg. 29 (1843) Misc. 6, Fol. Orch. Oberonia (1859) 6; Naves Novis. App. (1882) 230; Ames Orch. 5 (1915) 75.

Malaxis miniata Reichb. f. in Walp. Ann. 6 (1861) 215.

This species was based on material from Singapore. It has been credited to the Philippines, but there is little reason for considering it Philippine. It has not appeared in our modern collections.

OBERONIA RUFILABRIS Ames Orch. 1 (1905) 78, non Lindl. = *Oberonia thisbe* Reichb. f.

OBERONIA SPATULATA Lindl. apud Naves Novis. App. (1882) 230.

46. HIPPEOPHYLLUM Schlechter

HIPPEOPHYLLUM WENZELII Ames in Philip. Journ. Sci. 8 (1913) Bot. 413, Orch. 5 (1915) 78.

Leyte, *Wenzel 020*, *0158*. Dagami, epiphyte, at 60 m altitude.

47. LIPARIS L. C. Richard

LIPARIS AMESIANA Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 210; Ames Orch. 5 (1915) 78.

Cestichis gracilis Ames Orch. 2 (1908) 136, text cut, non *Liparis gracilis* Hook. f.

Luzon (Bataan, Rizal), Mindanao (Bukidnon), *Copeland 274*, *Merrill s. n.*, *Weber 28*, *B. S. 1782*, *38632 Ramos*. Epiphyte in forest at 930 to 1,460 m altitude.

LIPARIS BENGUETENSIS (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 210; Ames Orch. 5 (1915) 78. *Cestichis benguetensis* Ames Orch. 1 (1905) 9, 75, t. 3, fig. I, 1-6, 2 (1908) 132.

Luzon (Bontoc, Benguet), Negros. Fairly common in Luzon in mountainous regions at 700 to 2,300 m altitude.

LIPARIS BONTOCENSIS Ames Sched. Orch. 6 (1923) 38.

Luzon (Bontoc), *Vanoverbergh 1245, 3947a*. At 1,650 to 1,700 m altitude.

LIPARIS CAESPITOSA (Thou.) Lindl. in Bot. Reg. 11 (1825) sub t. 882; A. Rich. Monogr. Orch. Isles France et Bourb. (1827) [in Mém. Soc. Hist. Nat. Par. 4 (1828) 47]; Lindl. Gen. & Sp. Orch. (1830) 32; Bojer Hort. Maur. (1837) 321; S. Moore in Baker Fl. Maur. & Seych. (1877) 342; Ridl. in Journ. Linn. Soc. 22 (1886) 290; Cordem. Fl. Réunion. (1895) 180; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 266, Fig.-Atlas (1910) fig. 204, in Nova Guinea 8 (1909) 35, 12³ (1915) 221; Schltr. in Engl. Bot. Jahrb. 45 Beibl. 104 (1911) 17; Ames Orch. 5 (1915) 78, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 153.

Malaxis caespitosa Thou. Orch. Isles Afr. (1822) t. 89.

Malaxis minima Blume Bijdr. (1825) 391.

Malaxis angustifolia Blume Bijdr. (1825) 393.

Liparis minima Lindl. Gen. & Sp. Orch. (1830) 32; Miq. Fl. Ind. Bat. 3 (1859) 622; Naves Novis. App. (1882) 230; Ridl. in Journ. Linn. Soc. 22 (1886) 297; Schltr. in Fedde Repert. 10 (1911) 28.

Liparis angustifolia Lindl. Gen. & Sp. Orch. (1830) 31; Miq. Fl. Ind. Bat. 3 (1859) 622; Thw. Enum. Pl. Zeyl. (1864) 296.

Liparis auriculata Reichb. f. in Flora 55 (1872) 277, non Blume.

Liparis pusilla Ridl. in Journ. Linn. Soc. 22 (1886) 294; Hook. f. in Hook. Icon. Pl. 19 (1889) t. 1856 A, Fl. Brit. Ind. 5 (1890) 701; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 32, t. 41.

Liparis prainii Hook. f. in Hook. Icon. Pl. 19 (1889) t. 1857 A, Fl. Brit. Ind. 5 (1890) 700.

Liparis duthiei Hook. f. in Hook. Icon. Pl. 19 (1889) t. 1857 B, Fl. Brit. Ind. 5 (1890) 701; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 32, t. 42; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 93.

Liparis obscura Hook. f. in Hook. Icon. Pl. 19 (1889) t. 1886, Fl. Brit. Ind. 5 (1890) 701; Trim. Handb. Fl. Ceyl. 4 (1898) 147.

Leptorchis caespitosa Ktze. Rev. Gen. Pl. 2 (1891) 671.

Leptorchis duthiei Ktze. Rev. Gen. Pl. 2 (1891) 671.

Leptorchis minima Ktze. Rev. Gen. Pl. 2 (1891) 671.

Leptorchis obscura Ktze. Rev. Gen. Pl. 2 (1891) 671.

Leptorchis prainii Ktze. Rev. Gen. Pl. 2 (1891) 671.

Leptorchis pusilla Ktze. Rev. Gen. Pl. 2 (1891) 671.

Cestichis caespitosa Ames Orch. 2 (1908) 132, t.

Luzon (Benguet, Rizal), Leyte, Mindanao (Cotabato, Davao), *R. S. Williams 428, 1954, Merrill 4684, B. S. 13558 Ramos, 19238 Reillo, Wenzel 0734, Weber 232, Elmer 11858*. Epiphyte in forest at 800 to 1,160 m

altitude. In India, Ceylon, Sumatra, Java, Celebes, Borneo, New Guinea, Bourbon, Mauritius, Reunion, and Madagascar.

LIPARIS CLEMENSIAE Ames Orch. 5 (1915) 79.

Cestichis clemensiae Ames in Philip. Journ. Sci. 4 (1909) Bot. 666.

Mindanao (Lanao), *M. S. Clemens* 129. Camp Keithley.

LIPARIS COMPRESSA (Blume) Lindl. Gen. & Sp. Orch. (1830) 32; Miq.

Fl. Ind. Bat. 3 (1859) 621; Naves Novis. App. (1882) 230; Ridl. in Journ. Linn. Soc. 22 (1886) 291, 32 (1896) 230; Hook. f. Fl. Brit. Ind. 5 (1890) 706; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 280, Fig.-Atlas (1910) fig. 215; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 25; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 17; Ames Orch. 5 (1915) 79.

Malaxis compressa Blume Bijdr. (1825) 390, fig. 54.

Leptorchis compressa Ktze. Rev. Gen. Pl. 2 (1891) 670.

Cestichis compressa Ames Orch. 1 (1905) 76, text cut, 2 (1908) 134.

Luzon (Lepanto, Bataan), *Borden 1816, Merrill 4577*. Mt. Data and Mt. Mariveles, epiphyte in forest at 1,060 to 2,300 m altitude. In Malay Peninsula, Sumatra, Java, and Celebes.

LIPARIS CONFUSA J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 275,

Fig.-Atlas (1910) fig. 211; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 17, in Fedde Repert. 10 (1911) 28, in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 205; J. J. Sm. in Bull. Jard. Buitenz. II 14 (1914) 32; Ames Orch. 5 (1915) 79, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 153.

Liparis flaccida Schltr. in K. Schum. & Lauterb. Nachtr. Fl. Deutsch. Schutzgeb. (1905) 103, non Reichb. f.

Cestichis vestita Ames Orch. 2 (1908) 139, p. p., non *Liparis vestita* Reichb. f.

Luzon (Apayao, Bontoc, Benguet, Nueva Vizcaya, Rizal, Laguna, Tayabas, Sorsogon), Mindoro, Leyte, Samar, Mindanao (Zamboanga, Lanao, Agusan, Butuan, Davao), Basilan. Common and widely distributed in the Philippines. Epiphyte in forest at 15 to 1,430 m altitude. In Sumatra, Java, Celebes, Borneo, and New Guinea.

LIPARIS CUMINGII Ridl. in Journ. Linn. Soc. 22 (1886) 292; Hook. f.

Fl. Brit. Ind. 5 (1890) 707; Ames Orch. 5 (1915) 79.

Cestichis cumingii Ames Orch. 1 (1905) 13, in Philip. Journ. Sci. 7 (1912) Bot. 7.

Luzon (Bontoc, Bataan, Laguna, Sorsogon), Leyte, Bohol,* Mindanao (Zamboanga), *Weber 26, B. S. 17037 Robinson, Elmer 14703, 16296, Wenzel 0680, 0808, Copeland s. n., Vanoverbergh 710*. Epiphyte in forest at 600 to 1,400 m altitude.

LIPARIS DUMAGUETENSIS Ames Orch. 5 (1915) 80.

Liparis elmeri Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1561, nec

Cestichis elmeri Ames, nec *Liparis elmeri* (Ames) Schltr.

Luzon (Tayabas), Panay (Antique), Negros (Oriental Negros), Leyte, Mindanao (Camiguin Island), *Elmer 7566, 7579, 9445, 10236, Wenzel 0298, 0299, McGregor 5805, B. S. 14444 Ramos, 41401 Herre*. Terrestrial in forest at 600 to 1,300 m altitude.

* Bopol, according to Hook. f. in Fl. Brit. Ind. 5 (1890).

LIPARIS ELMERI (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 210; Ames Orch. 5 (1915) 80.

Cestichis elmeri Ames Orch. 1 (1905) 10, '75, t. 3, fig. II, 7-12, 2 (1908) 135.

Luzon (Benguet, Laguna, Tayabas), *Elmer 6553*, *R. S. Williams 1928*, *Ramos & Edaño 107*, *F. B. 5065 Curran*, *B. S. 26528 Ramos & Edaño*, *6085 Robinson*, *27940 Ocampo*. Epiphyte, 260 to 2,230 m altitude.

LIPARIS FORBESII Ridl. in Journ. Linn. Soc. 22 (1886) 283; ? J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 268, Fig.-Atlas (1910) fig. 205; Ames Orch. 5 (1915) 80.

Leptorchis forbesii Ktze. Rev. Gen. Pl. 2 (1891) 671.

Cestichis forbesii Ames Orch. 2 (1908) 135.

Luzon (Bontoc, Benguet), Mindanao (Lanao), *Vanoverbergh 711*, *1387*, *1478*, *3947*, *R. S. Williams 1941*, *M. S. Clemens 446*.

LIPARIS FRAGILIS Ames Orch. 5 (1915) 80.

Cestichis fragilis Ames in Philip. Journ. Sci. 8 (1913) Bot. 412.

Mindanao (Zamboanga), *Merrill 8184*, *8289*. Epiphyte at about 900 m altitude.

LIPARIS HALCONENSIS Ames Orch. 5 (1915) 80.

Cestichis halconensis Ames in Philip. Journ. Sci. 2 (1907) Bot. 321, Orch. 3 (1908) 63, t. 47.

Luzon (Ifugao), Mindoro, Panay (Antique), *Merrill 5799*, *McGregor 6182*, *B. S. 19882 McGregor*. Terrestrial at 400 to 730 m altitude.

LIPARIS JARENSIS Ames Orch. 6 (1920) 293.

Leyte, *Wenzel 0628*, *0708*, *0799*, *0813*. Jaro, epiphyte in forest at 600 to 700 m altitude.

LIPARIS LEYTENSIS Ames Orch. 5 (1915) 80.

Panay (Capiz), Leyte, Mindanao (Agusan), *Wenzel 0216*, *0353*, *0721*, *Weber 160*, *B. S. 35607 Martelino & Edaño*. Epiphyte in forest at 12 to 800 m altitude.

LIPARIS LINEARIFOLIA Ames Orch. 5 (1915) 81.

Cestichis linearifolia Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1563.

Mindanao (Davao), *Elmer 11494*, Todaya, Mt. Apo, epiphyte in dense forest at 2,000 m altitude.

LIPARIS LONGIPES Lindl. in Wall. Pl. Asiat. Rar. 1 (1830) 31, t. 35, Gen. & Sp. Orch. (1830) 30; Wight Icon. Pl. 3² (1849) 9, t. 906; Benth. in Hook. Kew Journ. Bot. 7 (1855) 33; Reichb. f. in Bonpl. 5 (1857) 57; Benth. Fl. Hongk. (1861) 352; Reichb. f. in Walp. Ann. 6 (1861) 218; Thw. Enum. Pl. Zeyl. (1864) 295; Reichb. f. in Seem. Fl. Vit. (1868) 302; Ridl. in Journ. Linn. Soc. 22 (1886) 293 (excl. syn. *L. elegans* Lindl.), 32 (1896) 229; Hook. f. Fl. Brit. Ind. 5 (1890) 703; Grant Orch. Burma (1895) 21; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 29, t. 37; Trim. Handb. Fl. Ceyl. 4 (1898) 147; Forbes & Hemsl. in Journ. Linn. Soc. 36 (1903) 7; Matsum. Ind. Pl. Jap. 2 (1905) 252; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.)

(1906) 93; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 25; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 200; Ames Orch. 6 (1920) 94, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 154.

Liparis pendula Lindl. in Bot. Reg. 24 (1838) Misc. 94; Regel Ind. Sem. Hort. Petrop. (1868) 79; Kränzl. in Orchis 2 (1907) 32.

Liparis spathulata Lindl. in Bot. Reg. 26 (1840) Misc. 81.

Sturmia longipes Reichb. f. in Bonpl. 3 (1855) 250.

? *Liparis boothii* Regel Ind. Sem. Hort. Petrop. (1864) Suppl. 14.

? *Liparis nesophila* Reichb. f. Otia Bot. Hamb. 1 (1878) 56, Xen. Orch. 3 (1881) 31.

Liparis longipes Lindl. var. *spathulata* Ridl. in Journ. Linn. Soc. 22 (1886) 294; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 30, t. 38.

Liparis triloba Ridl. in Journ. Linn. Soc. 22 (1886) 295; Ames Orch. 5 (1915) 84.

Cestichis pendula Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1889) 131, fig. 128 C-E, G.

Leptorchis longipes Ktze. Rev. Gen. Pl. 2 (1891) 670.

Leptorchis triloba Ktze. Rev. Gen. Pl. 2 (1891) 671.

Stichorchis pendula Pfitz. in Engl. & Prantl Nat. Pflanzenfam. Nachtr. 1 (1897) 103.

Cestichis longipes Ames Orch. 1 (1905) 75 (excl. Luzon specimen), 2 (1908) 137.

Luzon (Ilocos Norte, Laguna), Mindoro, Mindanao (Zamboanga), *Cuming s. n.*, *B. S. 27138 Ramos, 9679 Robinson, Ramos & Edaño s. n.*, *R. S. Williams 2176, F. B. 9218 Whitford & Hutchinson*. Epiphyte at 40 to 1,650 m altitude. In India, Ceylon, Burma, Malay Peninsula, China, Formosa, and Samoa.

LIPARIS LYONII Ames Orch. 5 (1915) 81.

Cestichis lyonii Ames in Philip. Journ. Sci. 6 (1911) Bot. 47.

Luzon (Bontoc, Benguet), *Vanoverbergh 3885, Lyon 155, Elmer 8463*.

LIPARIS MAGNICALLOSA Ames Orch. 7 (1922) 108.

Mindanao (Bukidnon), *B. S. 38739 Ramos & Edaño*. Mt. Candoon, epiphyte in forest at 1,600 m altitude.

LIPARIS MERRILLII (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 210; Ames Orch. 5 (1915) 81.

Cestichis merrillii Ames Orch. 1 (1905) 11, 75, t. 3, fig. III, 13-18, 2 (1908) 138, in Philip. Journ. Sci. 2 (1907) Bot. 321.

Luzon (Bontoc, Ifugao, Bataan, Rizal, Tayabas), Mindoro, Leyte, Mindanao (Misamis). A fairly common and widespread species, in mountainous regions, epiphyte in mossy forest at 300 to 3,060 m altitude.

LIPARIS MONOPHYLLA Ames Orch. 6 (1920) 294.

Leyte, *Wenzel 0660, 0673*. Jaro, epiphyte in forest at 600 m altitude.

LIPARIS NEGROSIANA Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1562, Orch. 5 (1915) 81.

Negros (Oriental Negros), *Elmer 9606*. Dumaguete, terrestrial at 1,300 m altitude.

LIPARIS NUTANS Ames Orch. 5 (1915) 81.

Cestichis nutans Ames in Philip. Journ. Sci. 4 (1909) Bot. 597.

Palawan, Mindanao (Lanao, Agusan, Surigao), *B. S. 638 Foxworthy, M. S. Clemens s. n., Weber 70, Bolster 289*. In forest at 60 to 1,050 m altitude.

LIPARIS PALAWANENSIS Ames Sched. Orch. 6 (1923) 39.

Palawan, *Weber 616*. Alfonso XIII at sea level.

LIPARIS PALLIDA (Blume) Lindl. Gen. & Sp. Orch. (1830) 31; Miq. Fl. Ind. Bat. 3 (1859) 622; Naves Novis. App. (1882) 230; Ridl. in Journ. Linn. Soc. 22 (1886) 296; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 272, Fig.-Atlas (1910), *fig. 209*; Ames Orch. 6 (1920) 295.

Malaxis pallida Blume Bijdr. (1825) 391.

Leptorchis pallida Ktze. Rev. Gen. Pl. 2 (1891) 671.

Liparis latifolia Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1562, Orch. 5 (1915) 80, non Lindl.

Leyte, Mindanao (Bukidnon, Davao), *Wenzel 0775, 0908, 0930, B. S. 39120 Ramos & Edaño, Elmer 10648*. Epiphyte in forest at 500 to 2,250 m altitude. In Java.

LIPARIS PARVIFLORA (Blume) Lindl. Gen. & Sp. Orch. (1830) 31; Miq. Fl. Ind. Bat. 3 (1859) 622; J. J. Sm. Fl. Buitenz. 6 (1905) 276, Fig.-Atlas (1910) *fig. 212*; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 17; Ames Orch. 5 (1915) 81, 6 (1920) 95, in Merr. in Journ. Roy. Asiatic Soc. Straits Branch, Special No. (1921) 155.

Malaxis parviflora Blume Bijdr. (1825) 392.

Liparis flaccida Reichb. f. in Linnaea 41 (1876) 45; Ridl. in Journ. Linn. Soc. 22 (1886) 296, 32 (1896) 228; Hook. f. Fl. Brit. Ind. 5 (1890) 703; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 24.

Leptorchis flaccida Ktze. Rev. Gen. Pl. 2 (1891) 670.

Leptorchis parviflora Ktze. Rev. Gen. Pl. 2 (1891) 671.

Cestichis vestita Ames Orch. 2 (1908) 139, p. p., non *Liparis vestita* Reichb. f.

Luzon (Apayao, Bontoc, Rizal, Laguna, Tayabas), Polillo, Leyte, Bu-suanga, Mindanao (Cotabato, Davao), Basilan. A fairly common and widely distributed species in the Philippines. Epiphyte from about sea level to 1,450 m altitude. In Malay Peninsula, Sumatra, Java, Borneo, and Siam.

LIPARIS PECTINATA Ridl. in Journ. Linn. Soc. 22 (1886) 277; Ames Orch. 5 (1915) 82.

Leptorchis pectinata Ktze. Rev. Gen. Pl. 2 (1891) 671.

Luzon (Laguna), Biliran, Leyte, Mindanao (Davao), *Reillo 55, Cuming s. n., B. S. 18934 McGregor, 21450 Escritor, Wenzel 0207, 0223, 0380, 0481, 0578, Black s. n., Weber 279*. Terrestrial in forest at 60 to 500 m altitude.

LIPARIS PHILIPPINENSIS (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 210; Ames Orch. 5 (1915) 83.

Cestichis philippinensis Ames Orch. 1 (1905) 7, 75, t. 2, 2 (1908) 138, in Philip. Journ. Sci. 2 (1907) Bot. 321.

Luzon (Abra, Bontoc, Lepanto, Benguet, Bataan, Pampanga, Rizal, Laguna), Mindoro, Panay (Antique), Negros (Oriental Negros), Leyte, Min-

danao (Agusan, Bukidnon, Davao). A common and widely distributed species, mostly in mountain regions at 300 to 2,900 m altitude.

LIPARIS PRAVA Ames Orch. 7 (1922) 109.

Leyte, Wenzel 0718, 0749, 0804, 0834, 0842. Jaro, epiphyte in forest at 500 to 800 m altitude.

LIPARIS PROPINQUA Ames Orch. 7 (1922) 110.

Liparis gregaria Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 266, non Lindl.

Cestichis disticha Pfitz. in Engl. in Engl. & Prantl Nat. Pflanzenfam. 2 Abt. 6 (1888) 131, quoad Philip.; Ames in Philip. Journ. Sci. 2 (1907) Bot. 320.

Liparis disticha Ames Orch. 5 (1915) 79, non Lindl.

Luzon (Laguna), Mindoro, Palawan, Bancalan, Mindanao (Davao), Cumíng 2099, B. S. 6546 Robinson, Merrill 5638, 5643, F. B. 8598 Merritt, 28185 Elumir, Weber s. n., 011, Elmer 18305. Epiphyte at sea level to 600 m altitude.

LIPARIS RIZALENSIS Ames Orch. 6 (1920) 295.

Luzon (Rizal, Tayabas), B. S. 24070 Ramos, 26892 Edaño, 26533 Ramos & Edaño. Epiphyte on forest slopes at about 300 m altitude.

LIPARIS STRICTA J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 5 (1907) 3; Ames Orch. 6 (1920) 95, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 155.

Basilan, Mindanao (Agusan), B. S. 16250 Reillo, Weber 74. Terrestrial in shaded forest at 400 m altitude in Mindanao. In Borneo.

LIPARIS TRICALLOSA Reichb. f. in Gard. Chron. II 11 (1879) 684; Ridl. in Journ. Linn. Soc. 22 (1886) 275; Hook. f. in Bot. Mag. 127 (1901) t. 7804; Ames Orch. 5 (1915) 83.

Sulu Archipelago (Buut Island), Jolo, Burbidge s. n., J. Clemens s. n. Wooded hillside at 660 m (?) altitude.

LIPARIS TRICHOGLOTTIS (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1911) 203; Ames Orch. 5 (1915) 83.

Cestichis trichoglottis Ames Orch. 2 (1908) 139, text cut.

Luzon (Laguna, Tayabas), Mindanao (Lanao), Basilan, B. S. 16564 Ramos, 16253 Reillo, Elmer 7612, M. S. Clemens 131.

LIPARIS WENZELII Ames Orch. 5 (1915) 84.

Leyte, Wenzel 0403. Epiphyte in forest at 500 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

LIPARIS AFFINIS Lindl. apud Naves Novis. App. (1882) 230.

LIPARIS DIPHYLLOS Nim. in Grah. Cat. Bomb. Pl. (1839) 252 = *Habenaria diphylla* (Nim.) Dalz.

LIPARIS DISTICHA Ames Orch. 5 (1915) 79, non Lindl. = *Liparis propinqua* Ames.

LIPARIS ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1561, nec *Cestichis elmeri* Ames, nec *Liparis elmeri* (Ames) Schltr. = *Liparis dumaguetensis* Ames.

LIPARIS GREGARIA Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 266, non Lindl. = *Liparis propinqua* Ames.

LIPARIS HABENARIA F.-Muell. apud Naves Novis. App. (1882) 230.

LIPARIS LATIFOLIA Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1562, Orch. 5 (1915) 80, non Lindl. = *Liparis pallida* (Blume) Lindl.

LIPARIS MINIMA Lindl. Gen. & Sp. Orch. (1830) 32; Naves Novis. App. (1882) 230 = *Liparis caespitosa* (Thou.) Lindl.

LIPARIS MUCRONATA Lindl. apud Naves Novis. App. (1882) 230.

LIPARIS REFLEXA Lindl. apud Naves Novis. App. (1882) 229.

LIPARIS RHEEDII Lindl. apud Naves Novis. App. (1882) 230.

LIPARIS TRILOBA Ridl. in Journ. Linn. Soc. 22 (1886) 295; Ames Orch. 5 (1915) 84 = *Liparis longipes* Lindl.

LIPARIS VIRIDIFLORA Lindl. apud Naves Novis. App. (1882) 230.

48. **DILOCHIA** Lindley

DILOCHIA ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1554, Orch. 5 (1915) 85.

Luzon (Rizal), Leyte, Mindanao (Davao), *Reillo s. n.*, *Wenzel 0446*, *0579*, *Elmer 10664*. Epiphyte, up to 500 m altitude in Leyte.

49. **CERATOSTYLIS** Blume

CERATOSTYLIS DATAENSIS Ames Orch. 7 (1922) 151.

Luzon (Lepanto), *Ramos & Edaño 76*. Mt. Data, epiphyte in mossy forest at 2,540 m altitude.

CERATOSTYLIS ELMERI Ames Sched. Orch. 6 (1923) 49.

Mindanao (Davao), *Elmer 11358*. Epiphyte at 2,580 m altitude.

CERATOSTYLIS LATIPETALA Ames in Philip. Journ. Sci. 4 (1909) Bot. 671, Orch. 5 (1915) 86.

Luzon (Rizal), Mindanao (Zamboanga, Misamis, Bukidnon, Agusan, Davao), *B. S. 19232 Reillo, 38815, 38953 Ramos & Edaño, F. B. 4661 Mearns & Hutchinson, Elmer 10614, 13801, Merrill 8134, Copeland s. n.* Mountainous regions up to 1,400 m altitude in Misamis.

CERATOSTYLIS MINDANAENSIS Ames Sched. Orch. 6 (1923) 51.

Mindanao (Bukidnon), *B. S. 39015 Ramos & Edaño*. At 1,600 m altitude.

CERATOSTYLIS PHILIPPINENSIS Rolfe ex Ames Orch. 1 (1905) 79, *text cut*, 2 (1908) 149, 5 (1915) 86; Ames in Philip. Journ. Sci. 6 (1911) Bot. 50.

Luzon (Bontoc, Lepanto, Benguet, Bataan, Camarines), Leyte. A rather common species in the mountainous regions of Luzon, at 700 to 2,100 m altitude.

CERATOSTYLIS RAMOSA Ames & Rolfe in Philip. Journ. Sci. 2 (1907) Bot. 323, *nomen*; Ames Orch. 2 (1908) 149, *text cut*, 5 (1915) 86.

Luzon (Bontoc, Bataan), Mindoro, Mindanao (Zamboanga), *Vanoverbergh* 785, 1390, *Whitford* 457, *Leiberg* 6039, *Merrill* 8130, *Merritt* 4418.

CERATOSTYLIS RETISQUAMA Reichb. f. in Bonpl. 5 (1857) 53; Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 269; Ames Orch. 5 (1915) 86.

Philippines, *Cuming* 2152.

CERATOSTYLIS RUBRA Ames in Philip. Journ. Sci. 4 (1909) Bot. 670, Orch. 5 (1915) 86.

Luzon (Ilocos Norte, Rizal, Bataan, Camarines), Mindanao (Lanao), *B. S.* 3042, 26220, 33231 *Ramos*, 3070 *Foxworthy*, 3046, 3072, *M. S.* *Clemens*, *Santos s. n.*, *Lyons J.*

CERATOSTYLIS SENILIS Reichb. f. Otia Bot. Hamb. 1 (1878) 54, Xen. Orch. 3 (1881) 30; Ames Orch. 2 (1908) 150, 5 (1915) 86.

Luzon (Bataan, Rizal, Laguna, Tayabas, Sorsogon), Mindanao (Surigao), *R. S. Williams* 506, *B. S.* 3017, 13761, 19362 *Ramos*, *Elmer* 9262, 14964, *Didson s. n.*, *F. B.* 7788 *Curran & Merritt*, *Bolster* 215, *Taylor* 28. Epiphyte up to 900 m altitude in Tayabas and 1,000 m in Laguna.

CERATOSTYLIS SUBULATA Blume Bijdr. (1825) 306; Lindl. Gen. & Sp. Orch. (1833) 231; Reichb. f. in Bonpl. 5 (1857) 53; Miq. Fl. Ind. Bat. 3 (1859) 706; Reichb. f. Xen. Orch. 2 (1863) t. 127, fig. II, 2, 2 (1865) 93; Naves Novis. App. (1882) 245; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 299, Fig.-Atlas (1910) fig. 229; Ames in Philip. Journ. Sci. 2 (1907) Bot. 323, Orch. 2 (1908) 151, *text cut*, 5 (1915) 86, 6 (1920) 138, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 176.

Appendicula teres Griff. Notul. 3 (1851) 359, Icon. Pl. Asiat. 3 (1851) t. 332.

Ceratostylis teres Reichb. f. in Bonpl. 2 (1854) 89, in Walp. Ann. 6 (1862) 470; Hook. f. Fl. Brit. Ind. 5 (1890) 825; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 247, t. 329.

Ceratostylis cepula Reichb. f. in Bonpl. 5 (1857) 53.

Ceratostylis gracilis Reichb. f. Xen. Orch. 2 (1863) t. 127, fig. I, 1, 2 (1865) 92; Naves Novis. App. (1882) 245; Vidal Rev. Pl. Vasc. Filip. (1886) 269; Ridl. in Journ. Linn. Soc. 32 (1896) 307, p. p., Ames Orch. 1 (1905) 80; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 109, non Blume.

Corymbis gracilis Vidal Rev. Pl. Vasc. Filip. (1886) 272; Ceron Cat. Pl. Herb. Manila (1892) 168, non Blume.

Ceratostylis malaccensis Hook. f. Fl. Brit. Ind. 5 (1890) 825, in Hook. Icon. Fl. 21 (1892) t. 2098; Ridl. in Journ. Linn. Soc. 32 (1896) 308.

Luzon (Ifugao, Benguet, Nueva Vizcaya, Bataan, Pampanga, Rizal), Mindoro, Panay (Capiz), Mindanao (Lanao, Zamboanga, Bukidnon, Davao). A fairly common species in the Philippines. Epiphyte at 1,600

to 2,160 m altitude. In India, Assam, Malay Peninsula, Sumatra, Java, and Borneo.

CERATOSTYLIS WENZELII Ames in Philip. Journ. Sci. 8 (1913) Bot. 421, Orch. 5 (1915) 87.

Leyte, Basilan, *Wenzel 013, 0175, 0661, 0678, 0745, B. S. 16256 Reillo*. Epiphyte at 60 to 800 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

CERATOSTYLIS GIGAS Reichb. f. apud Naves Novis. App. (1882) 245.

CERATOSTYLIS GRACILIS Reichb. f. Xen. Orch. 2 (1863) t. 127, fig. I, 1, 2 (1865) 92; Naves Novis. App. (1882) 245; Vidal Rev. Pl. Vasc. Filip. (1886) 269; Ridl. in Journ. Linn. Soc. 32 (1896) 307, p. p., Ames Orch. 1 (1905) 80; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 109, non Blume = *Ceratostylis subulata* Blume.

CERATOSTYLIS GRAMINEA Blume apud Naves Novis. App. (1882) 245.

50. AGROSTOPHYLLUM Blume

AGROSTOPHYLLUM ELMERI Ames Sched. Orch. 6 (1923) 48.

Luzon (Laguna), *Elmer 18047*.

AGROSTOPHYLLUM HASSELTII (Blume) J. J. Sm. in Icon. Bogor. 2 (1903) 55, in textu, Fl. Buitenz. 6 (Orch. Java) (1905) 288, Fig.-Atlas (1910) fig. 220; Ames Orch. 5 (1915) 87.

Appendicula hasseltii Blume Bijdr. (1825) 304; Lindl. Gen. & Sp. Orch. (1833) 230; Miq. Fl. Ind. Bat. 3 (1859) 704.

Appendicula graminifolia Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 24 (1862) 322.

Appendicula elongata Ridl. in Trans. Linn. Soc. II 3 (1893) 375; Ridl. in Journ. Linn. Soc. 32 (1896) 389.

Podochilus hasseltii Schltr. in Mém. Herb. Boiss. 21 (1900) 61; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 198.

Leyte, Mindanao (Agusan), *Wenzel 0610, 0719, 0742, 0763, 0798, Weber 307*. In the forest at 500 to 800 m altitude. In Malay Peninsula, Sumatra, and Java.

AGROSTOPHYLLUM INOCEPHALUM (Schauer) Ames Orch. 2 (1908) 148, 5 (1915) 87.

Diploconchium inocephalum Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 428, t. 12, fig. A.

Luzon (Benguet, Bataan), Mindanao (Lanao, Agusan), *R. S. Williams 1926, Lyon T., M. S. Clemens s. n., 130, Weber 6, 161, Elmer 13852*. On rocks or in trees from sea level to 500 m altitude.

AGROSTOPHYLLUM LEYTENSE Ames Orch. 6 (1920) 298.

Leyte, *Wenzel 0597, 0632, 0654*. Jaro, in forest at 600 to 700 m altitude.

AGROSTOPHYLLUM LONGIVAGINATUM Ames in Philip. Journ. Sci. 8 (1913) Bot. 420, Orch. 5 (1915) 87.

Luzon (Apayao, Rizal, Laguna), Mindoro, Leyte, *Weber 345, Serrato s. n., B. S. 16584 Ramos, 5656 Merrill, 42061 Lopez, Wenzel 0170, 0171, 0240, 0619, 0646, 0825*. In the forest at 30 to 600 m altitude.

AGROSTOPHYLLUM MALINDANGENSE Ames in Philip. Journ. Sci. 6 (1911) Bot. 50, Orch. 5 (1915) 88.

Mindanao (Misamis, Bukidnon, Davao), *F. B. 4610*, Mearns & Hutchinson, *B. S. 38836* Ramos & Edaña, *Elmer 10868*. On mountains.

AGROSTOPHYLLUM MEARNsii Ames in Philip. Journ. Sci. 8 (1913) Bot. 420, Orch. 5 (1915) 88.

Luzon (Bataan), Mindanao (Misamis, Agusan), with *Whitford 457*, *F. B. 4607* Mearns & Hutchinson, *Elmer 13700*. On mountains at 1,060 to 1,560 m altitude.

AGROSTOPHYLLUM MERRILLii Ames in Philip. Journ. Sci. 2 (1907) Bot. 323, Orch. 5 (1915) 88.

Mindoro, *Merrill 5844*. Mt. Halcon, in forest at 1,000 m altitude.

AGROSTOPHYLLUM MINDANAENSE Ames Orch. 5 (1915) 88.

Agrostophyllum pelorioides Ames in Philip. Journ. Sci. 8 (1913) Bot. 421, non Schltr.

Mindanao (Cotabato, Davao), *Weber 231*, *Copeland 1119*. On mountains at about 860 m altitude.

AGROSTOPHYLLUM PHILIPPINENSE Ames in Philip. Journ. Sci. 4 (1909) Bot. 669, Orch. 5 (1915) 88.

Babuyan Islands (Camiguin Island), Luzon (Bataan, Rizal, Sorsogon), Leyte, Negros. Apparently fairly common, especially in Leyte, and widespread. On mountains, in the forest at 60 to 700 m altitude.

AGROSTOPHYLLUM WENZELii Ames Orch. 5 (1915) 88.

Leyte, Mindanao (Agusan), *Wenzel 028, 0177, 0212, 0752, 0824*, *Weber 104*, *Elmer 13967*. At 60 to 800 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

AGROSTOPHYLLUM PELORIOIDES Ames in Philip. Journ. Sci. 8 (1913) Bot. 421, non Schltr. = *Agrostophyllum mindanaense* Ames.

51. GLOMERA Blume

GLOMERA MERRILLii Ames in Philip. Journ. Sci. 8 (1913) Bot. 419, Orch. 5 (1915) 89.

Mindanao (Zamboanga), *Merrill 8290*. Sax River, epiphyte in damp shaded ravine at about 200 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

GLOMERA ERYTHROSTOMA Blume apud Naves Novis. App. (1882) 245.

52. PODOCHILUS Blume

PODOCHILUS BICAUDATUS Schltr. in Fedde Repert. 3 (1906) 19; Ames in Philip. Journ. Sci. 4 (1909) Bot. 669, 8 (1913) Bot. 415, Orch. 5 (1915) 90.

Palawan, Basilan, *Usteri s. n.*, *Merrill 775*, *B. S. 16255, 16259A* Reillo. Up to 560 m altitude on Palawan.

PODOCHILUS CUMINGII Schltr. in Fedde Repert. 3 (1906) 19; Ames in Philip. Journ. Sci. 4 (1909) Bot. 668, 8 (1913) Bot. 415, Orch. 5 (1915) 89.

Podochilus zollingeri Schltr. in Mém. Herb. Boiss. 21 (1900) 23, non Reichb. f.

Catanduanes, Luzon (Rizal), *B. S. 30310 Ramos, Reillo s. n.* Epiphyte in forest at about 100 m altitude.

PODOCHILUS HYSTRICINUS Ames Sched. Orch. 6 (1923) 42, fig. 8.

Leyte, Wenzel 0262, 0878. At 700 to 750 m altitude.

PODOCHILUS INTRICATUS Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, 416, Orch. 5 (1915) 90.

Luzon (Tayabas), Mindanao (Agusan, Davao), Basilan, *Disdon s. n., Elmer 13704, Copeland 1164, Weber 244, B. S. 16263 Reillo s. n., 19406 Ramos.* Epiphyte at 600 to 1,030 m altitude.

PODOCHILUS LONGILABRIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1565, in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 90.

Luzon (Laguna, Tayabas, Camarines), *Quisumbing s. n., Whitford 939, Elmer 7585, B. S. 22038 Ramos.* Epiphyte at 1,160 to 1,500 m altitude.

PODOCHILUS PLUMOSUS Ames in Philip. Journ. Sci. 4 (1909) Bot. 668, 8 (1913) Bot. 415, Orch. 5 (1915) 90.

Luzon (Nueva Ecija, Zambales, Laguna, Tayabas), Panay (Capiz), Leyte, Mindanao (Zamboanga, Cotabato, Agusan, Surigao). A fairly common species, especially in Luzon. Epiphyte at 80 to 960 m altitude.

PODOCHILUS RAMOSII Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, 417, Orch. 5 (1915) 90.

Mindanao (Camiguin Island), *B. S. 14441 Ramos.*

PODOCHILUS SCIUROIDES Reichb. f. in Bonpl. 5 (1857) 41; Schltr. in Mém. Herb. Boiss. 21 (1900) 20; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 511, Fig.-Atlas (1912) fig. 388; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 192; Ames Orch. 6 (1920) 140, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 177.

Cryptoglottis serpyllifolia Reichb. f. in Bonpl. 5 (1857) 42, non Blume.

Podochilus microphyllus Hook. f. Fl. Brit. Ind. 6 (1890) 81; Grant Orch. Burma (1895) 320; Ridl. in Journ. Linn. Soc. 31 (1896) 301, 32 (1896) 386; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 177, non Lindl.

Podochilus curvunguis Schltr. in Fedde Repert. 10 (1911) 34; Ames Orch. 5 (1915) 89.

Luzon (Pampanga, Bataan, Tayabas), Mindanao (Zamboanga, Lanao), Basilan, Merrill 4260, 8311, *B. S. 1678 Foxworthy, 16259 Reillo, 28652 Ramos & Edaño, M. S. Clemens s. n.* Epiphyte in mossy forest at 450 to 850 m altitude. In Burma, Malay Peninsula, Sumatra, Java, Celebes, and Borneo.

PODOCHILUS SERPYLLIFOLIUS (Blume) Lindl. in Journ. Linn. Soc. 3 (1859) 37, in nota (by inference only); ex Miq. Fl. Ind. Bat. 3 (1859) 688; Naves Novis. App. (1882) 244; Schltr. in Mém. Herb. Boiss. 21 (1900) 20; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.), (1905) 508, Fig.-Atlas (1912) fig. 385; Ames Orch. 5 (1915) 90, 6 (1920) 140, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 177.

Cryptoglottis serpyllifolia Blume Bijdr. (1825) 297, fig. 42; Lindl. Gen. & Sp. Orch. (1833) 230; Reichb. f. in Walp. Ann. 6 (1864) 894.

Hexameria disticha R. Br. in Benn. Pl. Jav. Rar. (1838) 26, t. 7; Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 431; Miq. Fl. Ind. Bat. 3 (1859) 656.

Luzon, *Meyen s. n.*

PODOCHILUS STRICTUS Ames in Philip. Journ. Sci. 4 (1909) Bot. 669, 8 (1913) Bot. 415, Orch. 5 (1915) 90.

Podochilus robinsonii Ames in Philip. Journ. Sci. 6 (1911) Bot. 49, 8 (1913) Bot. 415.

Podochilus lucescens Ames Orch. 5 (1915) 90, non Blume.

Luzon (Cagayan, Rizal, Laguna, Tayabas, Camarines, Sorsogon), Catanduanes, Panay, Negros, Leyte, Samar, Mindanao (Zamboanga, Lanao, Agusan, Surigao, Davao), Basilan. A common, variable, and widely distributed species. Epiphyte at 10 to 1,500 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

PODOCHILUS ANCEPS Schltr. in Mém. Herb. Boiss. 21 (1900) 33; Ames Orch. 2 (1908) 142 = *Appendicula anceps* Blume.

PODOCHILUS CLEMENSIAE Ames in Philip. Journ. Sci. 4 (1909) Bot. 667 = *Appendicula clemensiae* Ames.

PODOCHILUS CORNUTUS Schltr. in Mém. Herb. Boiss. 21 (1900) 34; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 194; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, Orch. 2 (1908) 143 = *Appendicula cornuta* Blume.

PODOCHILUS CROTALINUS Ames in Philip. Journ. Sci. 4 (1909) Bot. 667 = *Appendicula crotalina* (Ames) Schltr.

PODOCHILUS CURVIUNGUIS Schltr. in Fedde Repert. 10 (1911) 34; Ames Orch. 5 (1915) 89 = *Podochilus sciuroides* Reichb. f.

PODOCHILUS ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1565 = *Appendicula elmeri* Ames.

PODOCHILUS FENIXII Ames in Philip. Journ. Sci. 6 (1911) Bot. 48 = *Appendicula fenixii* (Ames) Schltr.

PODOCHILUS FRUTICOSUS Ames in Philip. Journ. Sci. 6 (1911) Bot. 48 = *Appendicula buxifolia* Blume.

PODOCHILUS LONGICALCARATUS Rolfe in Kew Bull. (1904) 186 = *Appendicula undulata* Blume var. *longicalcarata* (Rolfe) Ames.

PODOCHILUS LUCBANENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1566 = *Appendicula lucbanensis* Ames.

- PODOCHILUS LUCESCENS* Ames Orch. 5 (1915) 90, non Blume = *Podochilus strictus* Ames.
- PODOCHILUS LUZONENSIS* Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1567 = *Appendicula luzonensis* Ames.
- PODOCHILUS MALINDANGENSIS* Ames in Philip. Journ. Sci. 6 (1911) Bot. 49 = *Appendicula malindangensis* (Ames) Schltr.
- PODOCHILUS MICRANTHUS* Schltr. in Mém. Herb. Boiss. 21 (1900) 36, 39; Ames Orch. 2 (1908) 144 = *Appendicula micrantha* Lindl.
- PODOCHILUS NEGROSIANUS* Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1568 = *Appendicula negrosiana* Ames.
- PODOCHILUS PENDULUS* Schltr. in Mém. Herb. Boiss. 21 (1900) 48, quoad Philip.; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, 8 (1913) Bot. 415, Orch. 2 (1908) 145 = *Appendicula philippinensis* (Schltr.) J. J. Sm.
- PODOCHILUS PERPLEXUS* Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1569 = *Appendicula perplexa* Ames.
- PODOCHILUS PHILIPPINENSIS* Schltr. in Mém. Herb. Boiss. 21 (1900) 49; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322 = *Appendicula philippinensis* (Schltr.) J. J. Sm.
- PODOCHILUS REFLEXUS* Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, non Schltr. = *Appendicula micrantha* Lindl.
- PODOCHILUS ROBINSONII* Ames in Philip. Journ. Sci. 6 (1911) Bot. 49, 8 (1913) Bot. 415 = *Podochilus strictus* Ames.
- PODOCHILUS SCALPELLIFORMIS* Blume apud Naves Novis. App. (1882) 244.
This is probably *Podochilus strictus* Ames.
- PODOCHILUS TENUIS* Lindl. apud Naves Novis. App. (1882) 244.
- PODOCHILUS UNDULATUS* Schltr. in Mém. Herb. Boiss. 21 (1900) 42 = *Appendicula undulata* Blume.
- PODOCHILUS UNDULATUS* Schltr. var. *LONGICALCARATUS* Schltr. in Mém. Herb. Boiss. 21 (1900) 43; Ames Orch. 2 (1908) 146 = *Appendicula undulata* Blume var. *longicalcarata* (Rolfe) Ames.
- PODOCHILUS XYTRIOPHORUS* Schltr. in Mém. Herb. Boiss. 21 (1900) 47; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, Orch. 2 (1908) 147 = *Appendicula xytriophora* Reichb. f.
- PODOCHILUS ZOLLINGERI* Schltr. in Mém. Herb. Boiss. 21 (1900) 23, non Reichb. f. = *Podochilus cumingii* Schltr.

53. APPENDICULA Blume

- APPENDICULA ANCEPS* Blume Bijdr. (1825) 299; Lindl. Gen. & Sp. Orch. (1833) 228; Miq. Fl. Ind. Bat. 3 (1859) 702; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 518, Fig.-Atlas (1912) fig. 393; Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 91.
Metachilum cyathiferum Lindl. in Wall. Cat. (1829) No. 2022, nomen; Lindl. Gen. & Sp. Orch. (1830) 74.

Appendicula lewisii Griff. in Calc. Journ. Nat. Hist. 4 (1844) 378, t. 19, Notul. 3 (1851) 360, Icon. Pl. Asiat. 3 (1851) t. 337; Reichb. f. in Walp. Ann. 6 (1864) 893; Hook. f. Fl. Brit. Ind. 6 (1890) 83; Ridl. in Journ. Linn. Soc. 32 (1896) 389.

Appendicula cyathifera Reichb. f. MSS ex Hook. f. Fl. Brit. Ind. 6 (1890) 83.

Dendrobium vaginatum Wall. MSS ex Hook. f. Fl. Brit. Ind. 6 (1890) 83.

Appendicula complanata Ridl. in Journ. Linn. Soc. 32 (1896) 389.

Podochilus anceps Schltr. in Mém. Herb. Boiss. 21 (1900) 33; Ames Orch. 2 (1908) 142; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 19.

Mindanao (Zamboanga), Basilan, Merrill 8122, 8128, Copeland s. n., B. S. 16238, 16245 Reillo, 37006, 37361 Ramos & Edaño. Terrestrial in forest at 560 m altitude. In Malay Peninsula, Java, Borneo, and Sumatra.

APPENDICULA BUXIFOLIA Blume Bijdr. (1825) 300; Lindl. Gen. & Sp. Orch. (1833) 228; Miq. Fl. Ind. Bat. 3 (1859) 702; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 532, Fig.-Atlas (1912) fig. 406; Ames Orch. 5 (1915) 93.

Appendicula frutex Ridl. in Journ. Linn. Soc. 31 (1896) 302.

Podochilus buxifolius Schltr. in Mém. Herb. Boiss. 21 (1900) 52.

Podochilus fruticosus Ames in Philip. Journ. Sci. 6 (1911) Bot. 48.

Appendicula fruticosa Ames in Philip. Journ. Sci. 8 (1913) Bot. 415.

Luzon (Laguna, Sorsogon), Mindanao (Misamis, Cotabato), B. S. 16905 Serviñas, Elmer 16138, M. S. Clemens s. n., Weber 216. Apparently in the mountain regions at about 600 m altitude. In Java.

APPENDICULA CLEMENSIAE Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 91.

Podochilus clemensiae Ames in Philip. Journ. Sci. 4 (1909) Bot. 667.

Luzon (Laguna, Tayabas, Sorsogon), Leyte, Samar, Mindanao (Lanao), B. S. 13670, 19514, 24254 Ramos, 24909 Edaño, Disdon s. n., Elmer 15097, 15609, 16593, Wenzel 023, 096, 0187, 0741, M. S. Clemens s. n., Boettcher s. n. Mountain regions usually, but at from 60 to 1,700 m altitude.

APPENDICULA CORNUTA Blume Bijdr. (1825) 302; Lindl. Gen. & Sp. Orch. (1833) 229; Miq. Fl. Ind. Bat. 3 (1859) 704; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 522, Fig.-Atlas (1912) fig. 396; Ames Orch. 5 (1915) 91.

Appendicula congenera Blume Bijdr. (1825) 303; Lindl. Gen. & Sp. Orch. (1833) 229; Miq. Fl. Ind. Bat. 3 (1859) 704.

Dendrobium manillense Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 430.

Appendicula manillensis Reichb. f. ex Miq. Fl. Ind. Bat. 3 (1859) 703; Naves Novis. App. (1882) 245.

Appendicula bifaria Lindl. in Hook. Kew Journ. 7 (1855) 35; Benth. Fl. Hongk. (1861) 358; Reichb. f. in Walp. Ann. 6 (1864) 893; Hook. f. Fl. Brit. Ind. 6 (1890) 82; Ridl. in Journ. Linn. Soc. 31 (1896) 301, 32 (1896) 387; King & Pantl. in Ann. Roy. Bot.

Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 248, *t.* 330; Rolfe in Journ. Linn. Soc. 36 (1903) 39.

Appendicula reduplicata Par. & Reichb. f. *Otia* Bot. Hamb. 1 (1878) 45.

Podochilus cornutus Schltr. in Mém. Herb. Boiss. 21 (1900) 34; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 194; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, Orch. 2 (1908) 143; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1912) 19.

Podochilus brachiatus Schltr. in Mém. Herb. Boiss. 21 (1900) 38.

Podochilus congener Schltr. in Mém. Herb. Boiss. 21 (1900) 40.

Appendicula brachiata Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1912) 337.

Luzon (Laguna), Mindoro, Negros (Oriental Negros), Leyte, Palawan Mindanao (Agusan, Davao), *B. S.* 16573 *Ramos*, 641 *Foxworthy*, 41399 *Herre*, *Merrill* 5541, 5843, *Wenzel* 0106, 0302, 0310, 0311, *Weber* 122, *Elmer* 11329. In humid forest at 210 to 2,000 m altitude. In India, Malay Peninsula, Java, Borneo, and China.

APPENDICULA CROTALINA (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1912) 336, in Philip. Journ. Sci. 8 (1913) Bot. 416, Orch. 5 (1915) 93.

Podochilus crotalinus Ames in Philip. Journ. Sci. 4 (1909) Bot. 667.

Mindanao (Misamis, Cotabato, Davao), *M. S. Clemens s. n.*, *Weber* 230, *Elmer* 10695. Mountainous regions up to 860 m altitude.

APPENDICULA CUNEATA Ames Sched. Orch. 6 (1923) 44, *fig.* 9, habit and flower.

Luzon (Tayabas), Leyte, Samar, Mindanao (Bukidnon, Cotabato, Davao), *B. S.* 28750, 38644 *Ramos & Edaño*, 24236, 24524 *Edaño*, *Wenzel* 0635, 0668, 0685, 0702, *Weber* 237, *Elmer* 10696.

APPENDICULA ELMERI Ames in Philip. Journ. Sci. 8 (1913) Bot. 416, Orch. 5 (1915) 92.

Podochilus elmeri Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1565.

Mindanao (Agusan, Bukidnon, Cotabato, Davao), *Weber* 86, 226, *B. S.* 38630, 39048 *Ramos & Edaño*, *Elmer* 10718. Along streams at 400 to 1,300 m altitude.

APPENDICULA FENIXII (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1912) 336; Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 91.

Podochilus fenixii Ames in Philip. Journ. Sci. 6 (1911) Bot. 48.

Batan Islands (Batan), *B. S.* 3794 *Fenix*. Mt. Iraya, near the summit.

APPENDICULA IRIGENSIS Ames Sched. Orch. 5 (1923) 5.

Luzon (Rizal), *B. S.* 41941C *Ramos*. Mt. Iriga, epiphyte in dense forest at 1,000 m altitude.

APPENDICULA LEYTENSIS Ames Sched. Orch. 6 (1923) 47, *fig.* 9, flower.

Leyte, *Wenzel* 0251, 0283, 0359, 0364, 0417, 0482. Epiphyte in forest at 500 m altitude.

APPENDICULA LUCBANENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 92.

Podochilus lucbanensis Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1566.

Luzon (Apayao, Abra, Bontoc, Rizal, Tayabas), Panay (Capiz), Mindanao (Camiguin Island, Agusan), *B. S.* 28350 *Fenix*, 7143, 14427, 41941b, s. n., *Ramos*, 30778, 37888 *Ramos & Edaño*, *Elmer* 8086, 13688, 13743. Apparently restricted to mountain regions.

APPENDICULA LUZONENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 92.

Podochilus luzonensis Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1567.

Luzon (Tayabas), *Elmer* 7930.

APPENDICULA MALINDANGENSIS (Ames) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) 337; Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 91.

Podochilus malindangensis Ames in Philip. Journ. Sci. 6 (1911) Bot. 49, in Elm. Leaf. Philip. Bot. 5 (1912) 1567.

Negros (Oriental Negros), Mindanao (Misamis, Bukidnon), *Elmer* 9543, *F. B.* 4733 *Mearns & Hutchinson*, *B. S.* 38563 *Ramos & Edaño*. Mountainous regions up to 1,560 m altitude.

APPENDICULA MAQUILINGENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 417, Orch. 5 (1915) 92.

Luzon (Laguna, Camarines, Sorsogon), Catanduanes, Panay (Antique, Capiz), Leyte, Negros (Oriental Negros), Mindanao (Camiguin Island, Bukidnon, Davao), Jolo. A fairly common and widely distributed species at 40 to 2,000 m altitude, on rocks or on trees in mossy forest.

APPENDICULA MERRILLII Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, 418.

Chilopogon merrillii Ames Orch. 5 (1915) 91.

Mindanao (Zamboanga, Cotabato), *Merrill* 8135, *Weber* 219, *B. S.* 36974 *Ramos & Edaño*. Epiphyte at 700 to 1,160 m altitude.

APPENDICULA MICRANTHA Lindl. in Ann. Mag. Nat. Hist. 15 (1845) 386; Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 272; Ames Orch. 5 (1915) 91.

Podochilus micranthus Schltr. in Mém. Herb. Boiss. 21 (1900) 36, 39; Ames Orch. 2 (1908) 144.

Podochilus reflexus Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, non Schltr.

Appendicula wenzelii Ames in Philip. Journ. Sci. 8 (1913) Bot. 419, Orch. 5 (1915) 91.

Luzon (Pampanga, Bataan, Rizal, Laguna, Tayabas), Polillo, Mindoro, Panay, (Capiz, Antique), Negros, Leyte, Palawan, Mindanao (Bukidnon, Agusan, Davao). A common and widely distributed species in the Philippines. Terrestrial or epiphytic at 15 to 600 m altitude.

APPENDICULA NEGROSIANA Ames in Philip. Journ. Sci. 8 (1913) Bot. 416, Orch. 5 (1915) 93.

Podochilus negrosianus Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1568.

Negros (Oriental Negros), *Elmer* 9636. In the Cuernos Mountains.

APPENDICULA PERPLEXA Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 92.

Podochilus perplexus Ames in Elm. Leaf. Philip. Bot. 5 (1921) 1569.

Luzon (Tayabas), *Elmer* 7407. Mt. Banahao, in shaded forest at 800 m altitude.

APPENDICULA PHILIPPINENSIS (Schltr.) J. J. Sm. Orch. Amb. (1905) 91, in textu; Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 92.

Appendicula pendula Blume apud Naves Novis. App. (1882) 245; Ames Orch. 5 (1915) 92.

Podochilus pendulus Schltr. in Mém. Herb. Boiss. 21 (1900) 48, quoad Philip.; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, 8 (1913) Bot. 415, Orch. 2 (1908) 145.

Podochilus philippinensis Schltr. in Mém. Herb. Boiss. 21 (1900) 49; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322.

Luzon (Tayabas, Laguna, Sorsogon), Mindoro, Leyte, Palawan, Mindanao (Zamboanga, Bukidnon). A variable and fairly common species at 115 to 1,600 m altitude.

APPENDICULA POLYANTHA Ames Sched. Orch. 5 (1923) 6.

Luzon (Rizal), *B. S. 42058 Lopez, 29733 Ramos & Edaño*. Mt. Angilog, at 2,200 m altitude.

APPENDICULA TAGALENSIUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 65.

Philippines, *Loher s. n.*

No specimens of this species have been seen.

APPENDICULA UNDULATA Blume Bijdr. (1825) 301; Lindl. Gen. & Sp. Orch. (1833) 229; Miq. Fl. Ind. Bat. 3 (1959) 703; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 527, Fig.-Atlas (1912) t. 402; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 179.

Appendicula purpurascens De Vriese Ill. Orch. Ind. Neerl. (1854) t. 12, fig. 1, non Blume.

Appendicula peruligera Reichb. f. in Bonpl. 5 (1857) 41.

Podochilus unciferus Hook. f. Fl. Brit. Ind. 6 (1890) 81, in Hook. Icon. Pl. 22 (1893) t. 2145; Ridl. in Journ. Linn. Soc. 32 (1896) 385, Mat. Fl. Mal. Penin. 1 (1907) 194.

Podochilus undulatus Schltr. in Mém. Herb. Boiss. 21 (1900) 42.

The species proper is a native of Malay Peninsula, Java, Sumatra, and Borneo, represented in the Philippines by the following variety:

- Var. **LONGICALCARATA** (Rolfe) Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 179.
Podochilus longicalcaratus Rolfe in Kew Bull. (1894) 186.
Podochilus undulatus Schltr. var. *longicalcaratus* Schltr. in Mém. Herb. Boiss. 21 (1900) 43; Ames Orch. 2 (1908) 146.
Appendicula undulata Blume var. *calcarata* Ames in Philip. Journ. Sci. 8 (1913) Bot. 416, Orch. 5 (1915) 93, sphalm.
- Luzon (Rizal), Leyte, Mindanao (Agusan, Lanao, Zamboanga, Cotabato, Surigao, Davao), *Cuming s. n.*, *Reillo s. n.*, *B. S. 15389 Ramos, 37022, 37263 Ramos & Edaño, 42084 Edaño, Elmer 13622, M. S. Clemens 516, R. S. Williams 2331, Copeland 1584, Weber 207, 292, Wenzel 01095*. In mossy forest, apparently in mountainous regions; also in Borneo (Sarawak).
- APPENDICULA WEBERI** Ames in Philip. Journ. Sci. 8 (1913) Bot. 418, Orch. 5 (1915) 93.
 Mindanao (Agusan, Cotabato), *Weber 88, 217, Elmer 13639*. On mountains, in damp shady forest at 430 to 690 m altitude.
- APPENDICULA XYTRIOPHORA** Reichb. f. in Seem. Fl. Vit. (1868) 299; Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 272; Hook. f. Fl. Brit. Ind. 6 (1890) 84; Ridl. in Journ. Linn. Soc. 31 (1896) 301, 32 (1896) 391; Ames in Philip. Journ. Sci. 8 (1913) Bot. 415, Orch. 5 (1915) 92, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 179.
Podochilus xytriophorus Schltr. in Mém. Herb. Boiss. 21 (1900) 47; Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, Orch. 2 (1908) 147.
- Luzon (Benguet, Rizal, Laguna, Tayabas, Sorsogon), Polillo, Mindoro, Panay (Capiz), Leyte, Mindanao (Lanao, Cotabato, Davao). A common and widely distributed species in the Philippines; in mossy forest at 60 to 1,330 m altitude. In Malay Peninsula and Borneo.
- EXCLUDED OR SYNONYMOUS SPECIES
- APPENDICULA ALBA** Blume apud Ames Orch. 5 (1915) 91.
 This species does not occur in the Philippines.
- APPENDICULA FRUTICOSA** Ames in Philip. Journ. Sci. 8 (1913) Bot. 425 = *Appendicula buxifolia* Blume.
- APPENDICULA KUHLLII** Reichb. f. apud Naves Novis. App. (1882) 244.
- APPENDICULA LONGIFOLIA** Blume apud Naves Novis. App. (1882) 245.
- APPENDICULA MANILLENSE** Reichb. f. ex Miq. Fl. Ind. Bat. 3 (1859) 703; Naves Novis. App. (1882) 245 = *Appendicula cornuta* Blume.
- APPENDICULA PENDULA** Blume apud Naves Novis. App. (1882) 245; Ames Orch. 5 (1915) 92 = *Appendicula philippinensis* (Schltr.) J. J. Sm.
- APPENDICULA REFLEXA** Blume ex Ames in Philip. Journ. Sci. 2 (1907) Bot. 322, in syn. = *Appendicula micrantha* Lindl.
- APPENDICULA RHODIOLA** Reichb. f. apud Naves Novis. App. (1882) 245.
- APPENDICULA TORTA** Blume apud Naves Novis. App. (1882) 244.
- APPENDICULA WENZELII** Ames in Philip. Journ. Sci. 8 (1913) Bot. 419, Orch. 5 (1915) 91 = *Appendicula micrantha* Lindl.

54. POLYSTACHYA Hooker

- POLYSTACHYA LUTEOLA** (Sw.) Hook. Exot. Fl. 2 (1824) *t.* 103; Spreng. Syst. Veg. 3 (1826) 742; Lindl. Gen. & Sp. Orch. (1830) 73; Boj. Hort. Maurit. (1837) 320; Lindl. in Bot. Reg. 24 (1838) Misc. 78; Mutel Mém. sur. plus. Orch. (in Mém. Strassb. Soc. Hist. Nat. 3) (1840) 3, *t.* 1, *fig.* 2; Focke in Tijdschr. Nat. Wetensch. 2 (1849) 203; A. Rich. in Sagra Hist. Isla Cuba 11 (Fl. Cub. Faner. 2) (1850) 247 (var. *angustifolia*); Reichb. f. in Nederl. Kruidk. Arch. 4 (1858) 331, in Walp. Ann. 6 (1863) 639; Thw. Enum. Pl. Zeyl. (1864) 308; Griseb. Fl. Brit. W. Ind. (1864) 628, Cat. Pl. Cub. (1866) 265; Sauv. Fl. Cub. (1873) 230; Reichb. f. in Linnaea 41 (1877) 128; Eggers Fl. St. Croix & Virg. Is. (U. S. Nat. Mus. Smithson. Inst. Bull. 13) (1879) 113, No. 970; Hemsl. in Godm. & Salv. Biol. Centr. Am. 3 (1883) 248; Chapm. Fl. So. U. S. ed. 2 (1883) 653, ed. 3 (1897) 482; Reichb. f. in Ber. Deutsch. Bot. Gesell. 3 (1885) 276; Rolfe in Journ. Bot. 23 (1885) 215; Cordem. Fl. l'Île Réun. (1895) 190; Cogn. in Mart. Fl. Bras. 3⁴ (1895) 312; Dur. & Schinz Consp. Fl. Afr. 5 (1895) 35; Duss Fl. Phan. Antill. Franc. (in Ann. Inst. Col. Marseille 3) (1897) 597; Urb. Symb. Antill. 4 (Fl. Portor.) (1903) 170; Ames Contr. Knowl. Orch. Fl. So. Fla. (1904) 16; Cogn. in Bull. Soc. Roy. Bot. Belg. 43 (1906) 304; Pulle Enum. Vasc. Pl. Surinam (1906) 119; Cogn. in Urb. Symb. Antill. 6 (1909) 381; Ames in Philip. Journ. Sci. 4 (1909) Bot. 666; Boldingh Fl. Dutch. W. Ind. 1 (St. Eust., Saba & St. Mart.) (1909) 41.
- Epidendrum minutum* Aubl. Hist. Pl. Guian. Franc. 2 (1775) 824.
- Dendrobium polystachyon* Sw. in Kongl. Svensk. Vet.-Akad. Nya Handl. 21 (1800) 247; Willd. Sp. Pl. 4 (1805) 137; Sw. in Schrad. Neues Journ. Bot. 1 (1806) 95, Fl. Ind. Occ. 3 (1806) 1978, 1998; Pers. Syn. 2 (1807) 523; Lodd. Bot. Cab. 5 (1820) *t.* 458; Lindl. Coll. Bot. (1821) *t.* 20; Thou. Orch. Iles Austr. Afr. (1822) Troisième Tabl. Espec. (as *polystachion*), *t.* 84 (as *polystachys*); A. Rich. Monogr. Orch. Iles Fr. et Bourb. (1827) 50 [in Mém. Soc. Nat. Hist. Par. 4 (1828) 57], *t.* 8, *fig.* 4 (as *polystachyum*).
- Cranichis luteola* Sw. Fl. Ind. Occ. 3 (1806) 1433.
- Onychium flavescens* Blume Bijdr. (1825) 325.
- Polystachya mauritiana* Spreng. Syst. Veg. 3 (1826) 742; Rolfe in Elliot in Journ. Linn. Soc. 29 (1891) 53; Dur. & Schinz. Consp. Fl. Afr. 5 (1895) 35; Palacky Cat. Pl. Madag. Fasc. 1 (1906) 7; Hochreut. Sert. Madag. (in Ann. Conserv. Jard. Bot. Genève 11 & 12) (1908) 55 (21); Schltr. in Beihefte Bot. Centralbl. 33, Abt. 2 (1915) 414.
- Dendrobium flavescens* Lindl. Gen. & Sp. Orch. (1830) 85, in Bot. Reg. 30 (1844) Misc. 57; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 302.
- Epidendrum parviflorum* Pav. ex Lindl. Gen. & Sp. Orch. (1830) 73, in syn.
- Polystachya zeylanica* Lindl. in Bot. Reg. 24 (1838) Misc. 78; Reichb. f. in Walp. Ann. 6 (1863) 639; S. Moore in Baker Fl. Maurit. & Seych. (1877) 362; Hook. f. Fl. Brit. Ind. 6 (1890) 21; Trim. Handb. Fl. Ceyl. 4 (1898) 183.

- Polystachya zollingeri* Reichb. f. in Bonpl. 5 (1857) 39, in Walp. Ann. 6 (1863) 640.
- Polystachya extinctoria* Reichb. f. in Walp. Ann. 6 (1863) 638.
- Callista flavescens* Ktze. Rev. Gen. Pl. 2 (1891) 654.
- Dendrorchis extinctoria* Ktze. Rev. Gen. Pl. 2 (1891) 659.
- Dendrorchis minuta* Ktze. Rev. Gen. Pl. 2 (1891) 658.
- Dendrorchis zollingeri* Ktze. Rev. Gen. Pl. 2 (1891) 659.
- Epidendrum resupinatum* Sessé & Moc. Fl. Mex. ed. 2 (1894) 202, fide Cogn. in Urb. Symb. Antill.
- Polystachya minuta* Frapp. ex Cordem. Fl. l'Île Réunion. (1895) 190, in nota; Britton in Small Fl. SE. U. S. (1903) 328, ed. 2 (1913) 324; Fawc. & Rendle Fl. Jam. 1 (1910) 48, t. 7, figs. 4 & 5; Small Fl. Miami (1913) 55; Ames Orch. 5 (1915) 93; Schltr. in Beihefte Bot. Centralbl. 36, Abt. 2 (1918) 479; Britton & Millsp. Bahama Fl. (1920) 89; Schltr. in Urb. Symb. Antill. 8 (Fl. Doming.) (1920) 122; Britton in Addisonia 7 (1922) 19, t. 234; G. Wilson in Orch. Rev. 30 (1922) 324; Musgrove in Orch. Rev. 31 (1923) 167, in textu, non A. Rich. & Gal.
- Polystachya flavescens* J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 284, Fig.-Atlas (1910) fig. 218; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 24; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 180.

Luzon (Bontoc, Benguet, Pangasinan, Zambales, Rizal, Tayabas), Mindanao (Lanao), *Vanoverbergh* 760, 1332, 3670, *Elmer* 8527, *Fenix* s. n., *Boettcher* s. n., *B. S.* 13550 *Ramos*, 20781 *Escritor*, *M. S. Clemens* s. n. At sea level to 1,200 m altitude. In India, Ceylon, Sumatra, Java, Borneo, Madagascar, Mauritius, Bourbon, and also in Florida, Central America, the West Indies, and northern South America.

The oldest name for the species is *Epidendrum minutum* Aubl. and although the combinations *Polystachya minuta* (Aubl.) Frapp. ex Cordem. and (Aubl.) Britton have been made, it seems inadvisable to use either on account of the already existing *Polystachya minuta* A. Rich. & Gal. in Ann. Sci. Nat. III 3 (1845) 27. The last named is presumably not a *Polystachya*, but its status is at present (and probably will remain) unsettled.

EXCLUDED OR SYNONYMOUS SPECIES

POLYSTACHYA MINUTA Frapp. ex Cordem. Fl. l'Île Réunion. (1895) 190, in nota; Britton in Small Fl. SE. U. S. (1903) 328; Ames Orch. 5 (1915) 93, non A. Rich. & Gal. = *Polystachya luteola* (Sw.) Hook.

55. ACANTHOPHIPPIUM Blume

ACANTHOPHIPPIUM MANTINIANUM L. Linden & Cogn. in Journ. des Orch. 7 (1896) 138; Ames in Philip. Journ. Sci. 6 (1911) Bot. 50, Orch. 5 (1915) 93.

Luzon (Bontoc, Sorsogon), *Vanoverbergh* 1203, *Elmer* 16409. In wet places.

56. PHAJUS Loureiro

PHAJUS FLAVUS (Blume) Lindl. Gen. & Sp. Orch. (1831) 128; Blume Mus. Bot. Lugd.-Bat. 2 (1856) 179; Miq. Fl. Ind. Bat. 3 (1859) 671; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 192, Fig.-Atlas (1909) fig. 144; Ames in Philip. Journ. Sci. 6 (1911) Bot. 50; Schltr. in Engl. Bot. Jahrb. 45 Beibl. 104 (1911) 24; Ames Orch. 5 (1915) 94.

Limodorum flavum Blume Bijdr. (1825) 375.

Bletia woodfordii Hook. in Bot. Mag. 54 (1827) t. 2719.

Phajus maculatus Lindl. in Wall. Cat. (1831) No. 3748, *nomen*, Gen. & Sp. Orch. (1831) 127; Lodd. Bot. Cab. 19 (1832) t. 1803; Hook. in Bot. Mag. 68 (1842) t. 3960; Blume Mus. Bot. Lugd.-Bat. 2 (1856) 180, Fl. Jav. Orch. (1858) 9 (Orch. Arch. Ind. 9) t. 5, E; Reichb. f. in Walp. Ann. 6 (1862) 460; Veitch Man. Orch. Pl. pt. 6 (1890) 12; Hook. f. Fl. Brit. Ind. 5 (1890) 817, 6 (1890) 192; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 107, t. 149.

Bletia flava Wall. in Bibl. Anglo-Indica ic. 1147 ex Lindl. Gen. & Sp. Orch. (1831) 127, in syn.

Phajus flexuosus Blume Mus. Bot. Lugd.-Bat. 2 (1856) 179, Fl. Jav. Orch. (1858) 7 (Orch. Arch. Ind. 6); Miq. Fl. Ind. Bat. 3 (1859) 671.

Phajus crispus Blume Mus. Bot. Lugd.-Bat. 2 (1856) 180, Fl. Jav. Orch. (1858) 8 (Orch. Arch. Ind. 8) t. 4, fig. 2, A, B; Miq. Fl. Ind. Bat. 3 (1859) 671.

Phajus platyphilus Reichb. f. Xen. Orch. 1 (1856) 201, t. 76, fig. III, 4-6, in Bonpl. 5 (1857) 42; Miq. Fl. Ind. Bat. 3 (1859) 740.

Phajus indigoferus Reichb. f. Xen. Orch. 1 (1856) 202, t. 76, fig. IV, 7 & 8 (excl. syn.); Blume Fl. Jav. Orch. (1858) 7 (Orch. Arch. Ind. 7) t. 2, fig. 2; Naves Novis. App. (1882) 246, non Hassk.

Phajus bracteosus Reichb. f. in Bonpl. 5 (1857) 42; Miq. Fl. Ind. Bat. 3 (1859) 740; Naves Novis. App. (1882) 249.

Luzon (Bontoc, Lepanto, Sorsogon), Panay (Antique), *B. S. 5616 Worcester, 32593 McGregor, Elmer 16530*. At 1,200 to 1,500 m altitude in Panay. In India, Java, and Sumatra.

PHAJUS HALCONENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 323, Orch. 5 (1915) 94.

Phajus calanthoides Ames Orch. 2 (1908) 153, *text cut*.

Luzon (Bontoc, Benguet, Nueva Vizcaya, Laguna, Tayabas), Mindoro, Negros. Common in Luzon at fairly high altitudes, up to 2,130 m on Mt. Halcon.

PHAJUS LINEARIFOLIUS Ames in Philip. Journ. Sci. 7 (1912) Bot. 10, Orch. 5 (1915) 94.

Luzon (Bontoc), *Vanoverbergh 1288, 3654*. Terrestrial on hillocks and in brooks at 1,180 to 1,650 m altitude.

PHAJUS LYONII Ames Orch. 5 (1915) 94.

Luzon (Bontoc, Lepanto, Ifugao, Zambales, Laguna, Sorsogon), Leyte, Mindanao (Cotabato, Davao), *B. S.* 5613 Worcester, 22226 Brown, *Lyon s. n.*, *Elmer* 11798, 16524, *Wenzel* 0118, *Weber* 225, *Wester s. n.* Terrestrial in the open or shade at 600 to 850 m altitude.

PHAJUS MINDORENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 324, Orch. 5 (1915) 95.

Luzon (Laguna) Mindoro, *Merrill* 5131, 5612, *B. S.* 17115 Robinson. Epiphyte or terrestrial in mossy forest at 1,000 to 1,430 m altitude.

PHAJUS PHILIPPINENSIS N. E. Br. in Gard. Chron. III 6 (1889) 239; Ames in Philip. Journ. Sci. 8 (1913) Bot. 422, Orch. 5 (1915) 95.

Luzon (Bontoc, Nueva Vizcaya), *Vanoverbergh* 3886, *B. S.* 11139 McGregor. Up to 1,300 m altitude in Bontoc.

PHAJUS RAMOSII Ames Orch. 5 (1915) 96.

Calanthe ramosii Ames in Philip. Journ. Sci. 7 (1912) Bot. 12.

Luzon (Zambales), *B. S.* 4987 Ramos.

EXCLUDED OR SYNONYMOUS SPECIES

PHAJUS AMBOINENSIS Blume apud Naves Novis. App. (1882) 246.

PHAJUS BRACTEOSUS Reichb. f. in Bonpl. 5 (1857) 42; Naves Novis. App. (1882) 249 = *Phajus flavus* (Blume) Lindl.

PHAIUS CALANTHOIDES Ames Orch. 2 (1908) 153, *text cut* = *Phajus halconensis* Ames.

PHAJUS CALLOSUS Lindl. apud Naves Novis. App. (1882) 246.

PHAJUS GRANDIFOLIUS Lour. Fl. Cochinch. (1790) 529; Naves Novis. App. (1882) 246; Ames in Philip. Journ. Sci. 4 (1909) Bot. 671, Orch. 5 (1915) 94.

There seems no real foundation for believing that this species occurs in the Philippines; the material referred to it is presumably *Phajus flavus* (Blume) Lindl.

PHAJUS INDIGOFERUS Reichb. f. Xen. Orch. 1 (1856) 202, *t.* 76, *fig.* IV, 7 & 8 (excl. syn.); Naves Novis. App. (1882) 246, non Hassk. = *Phajus flavus* (Blume) Lindl.

PHAJUS KUHLLII Reichb. f. apud Naves Novis. App. (1882) 246.**PHAJUS WALLICHII** Lindl. apud Naves Novis. App. (1882) 246.57. **CALANTHE** R. Brown

CALANTHE ANGUSTIFOLIA (Blume) Lindl. Gen. & Sp. Orch. (1833) 251, Fol. Orch. *Calanthe* (1854) 5; Miq. Fl. Ind. Bat. 3 (1859) 710; Reichb. f. in Walp. Ann. 6 (1864) 914; Hook. f. Fl. Brit. Ind. 5 (1890) 854; Ridl. in Journ. Linn. Soc. 32 (1896) 330; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 205, Fig.-Atlas (1909) *fig.* 152; Ames in Philip. Journ. Sci. 2 (1907) Bot. 324; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 121; Ames Orch. 2 (1908) 154, 5 (1915) 96, 7 (1922) 150; J. J. Sm. in Bull. Jard. Bot. Buitenz. II 14 (1914) 30.

Ambyglottis angustifolia Blume Bijdr. (1825) 369.

Calanthe phajoides Reichb. f. Xen. Orch. 1 (1856) 207, t. 79, fig. II, 4 & 5, in Bonpl. 5 (1857) 37; Hook. f. in Hook. Ic. Pl. 19 (1839) t. 1864; Naves Novis. App. (1882) 246.

Alismorchis angustifolia Ktze. Rev. Gen. Pl. 2 (1891) 650.

Alismorchis phajoides Ktze. Rev. Gen. Pl. 2 (1891) 650.

Luzon (Benguet), Mindoro, *Ramos & Edaño 82, 109, Merrill 5677*. Terrestrial in mossy forest, at 2,130 to 2,260 m altitude in Mindoro. In Malay Peninsula, Sumatra, and Java.

CALANTHE CONSPICUA Lindl. Fol. Orch. Calanthe (1854) 4; Miq. Fl. Ind. Bat. 3 (1859) 710; Reichb. f. in Walp. Ann. 6 (1864) 913; Naves Novis. App. (1882) 246; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 269; Ames Orch. 5 (1915) 96. *Calanthe lilacina* Loher in Gard. Chron. III 47 (1910) 66, text cut, in Fedde Repert. 11 (1912) 96.

Luzon (Albay, Sorsogon), Palawan, *Cuming 2097, B. S. 6459 Robinson, 23561 Ramos, Foxworthy 576*. At 290 to 485 m altitude.

CALANTHE DAVAENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 422, Orch. 5 (1915) 97.

Mindanao (Davao), *Copeland s. n.* Mt. Apo, terrestrial at 2,000 m altitude.

CALANTHE ELMERI Ames Orch. 2 (1908) 155, text cut, 5 (1915) 97.

Luzon (Abra, Bontoc, Benguet), *B. S. 27003 Ramos, 12936 Fenix, Van-overbergh 2195, 2210, Lyon 153, Elmer 5886, 8382, Topping s. n., Weber 406*. Terrestrial, sea level to 1,800 m altitude.

CALANTHE FURCATA Batem. ex Lindl. in Bot. Reg. 24 (1838) Misc. 28; Knowles & Westc. Floral Cab. 2 (1838) 46; Lindl. in Paxt. Flow. Gard. 3 (1852) 37, Fol. Orch. Calanthe (1854) 25; Moore Ill. Orch. Pl. (1857) Calanthe 5; Miq. Fl. Ind. Bat. 3 (1859) 711; Walp. Ann. 6 (1864) 916; Naves Novis. App. (1882) 246; Vidal Phan Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 269; Ames Orch. 1 (1905) 81.

Calanthe veratrifolia R. Br. apud Naves Novis. App. (1882) 246; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 211, quoad syn. *C. furcata* Batem. et quoad Philip.; J. J. Sm. Orch. Amb. (1905) 23, quoad syn. *C. furcata* Batem. et quoad Philip.

Alismorchis furcata Ktze. Rev. Gen. Pl. 2 (1891) 650.

Calanthe triplicata Ames in Philip. Journ. Sci. 2 (1907) Bot. 326, Orch. 2 (1908) 159, non *Orchis triplicata* Willem.

Luzon (Apayao, Kalinga, Bontoc, Benguet, Nueva Vizcaya, Pampanga, Bataan, Rizal, Laguna, Tayabas, Albay, Sorsogon), Mindoro, Panay (Capiz, Antique), Negros (Oriental Negros), Leyte, Mindanao (Camaguin Island, Zamboanga, Dapitan, Lanao, Bukidnon, Cotabato, Davao), Sibutu. A common and very widely distributed species in the Philippines. Terrestrial at 60 to 3,100 m altitude.

It has been deemed advisable to adopt the name *C. furcata* for the Philippine material as there is no question as to its identity. As pointed out by J. J. Smith, in Merr. Interp. Rumph. Herb. Amb. (1917) 170, the combination *C. triplicata* Ames is untenable as *Orchis triplicata* Willem.,

the name-bringing synonym, is not conspecific. While *C. veratrifolia* R. Br., as to plant figured and described, is very probably conspecific with *C. furcata* Batem., it is open to grave doubt, as pointed out by Smith, l. c., whether *Limodorum veratrifolium* Willd., is the same as the plant that Brown figured and described. Should *Limodorum veratrifolium* Willd. be *Calanthe sylvatica* Lindl., as seems probable, the combination *Calanthe veratrifolia* R. Br., (as to name) would apply to that species. In view of the uncertain status of *Calanthe veratrifolia* R. Br. no attempt has been made to cite its bibliography, synonymy, or distribution.

CALANTHE HALCONENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 324, Orch. 5 (1915) 97.

Luzon (Laguna), Mindoro, *Raymundo s. n.*, Merrill 5489. Terrestrial, in forest at 760 to 1,260 m altitude.

CALANTHE HENNISII Loher in Gard. Chron. III 46 (1909) 34, in Fedde Repert. 8 (1910) 384; Ames in Philip. Journ. Sci. 8 (1913) Bot. 423, Orch. 5 (1915) 97.

Luzon (Rizal, Tayabas), *Loher s. n.*, *Lyon 41*, *B. S. 13111 Foxworthy & Ramos, 41916 Ramos*. At 500 to 1,000 m altitude, according to Loher.

CALANTHE LACERATA Ames in Philip. Journ. Sci. 7 (1912) Bot. 11, Orch. 5 (1915) 97.

Luzon (Lepanto, Benguet, Tayabas, Camarines), *Merrill 4582, 4756, Ramos & Edaño 74, 114, F. B. 11355 Curran, Whitford 960*. Terrestrial or epiphytic ?, in forest at 1,000 to 2,500 m altitude.

CALANTHE LYROGLOSSA Reichb. f. *Otia Bot. Hamb. 1* (1878) 53, *Xen. Orch. 3* (1881) 30; *Naves Novis. App. (1882) 246*; Ames *Orch. 2* (1908) 156, *text cut*, 5 (1915) 97.

Alismorchis lyroglossa Ktze. *Rev. Gen. Pl. 2* (1891) 650.

Luzon (Apayao, Cagayan, Zambales, Bataan, Laguna), Negros (Oriental Negros), Palawan, *Wilkes, s. n.*, *Weber 343, s. n.*, *B. S. 4782, 19558 Ramos, F. B. 6306 Curran, Elmer 10083, Foxworthy 637*. Epiphyte at 50 to 1,100 m altitude.

CALANTHE MAQUILINGENSIS Ames *Orch. 5* (1915) 97.

Luzon (Laguna), Negros (Oriental Negros), Leyte, *B. S. 17246 Brown, Elmer 10063, Wenzel 0658*. Terrestrial at 600 to 1,000 m altitude.

CALANTHE MCGREGORII Ames in Philip. Journ. Sci. 2 (1907) Bot. 325, Orch. 5 (1915) 98.

Polillo, Mindoro, Mindanao (Lanao), *McGregor 177, B. S. 10452 McGregor, M. S. Clemens s. n.* Balete, Rio Baco.

CALANTHE MINDORENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 325, Orch. 5 (1915) 98.

Mindoro, *Merrill 5525*. Terrestrial, ridge forest, Mt. Halcon at 760 m altitude.

CALANTHE PULCHRA (Blume) Lindl. Gen. & Sp. Orch. (1833) 250; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 202, Fig.-Atlas (1909) fig. 150; Ames in Philip. Journ. Sci. 2 (1907) Bot. 326, Orch. 2 (1908) 158, 5 (1915) 98, 6 (1920) 161, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 181.

Amyglottis pulchra Blume Bijdr. (1825) 371.

Styloglossum nervosum Breda Gen. & Sp. Orch. Java (1827) t. 10.

Calanthe curculigoides Lindl. in Wall. Cat. (1832) No. 7340, *nomen*, Gen. & Sp. Orch. (1833) 251, in Bot. Reg. 33 (1847) t. 8; Reichb. f. in Walp. Ann. 1 (1849) 792, 6 (1864) 913; Lindl. Fol. Orch. Calanthe (1854) 4; Miq. Fl. Ind. Bat. 3 (1859) 710; Hook. f. in Bot. Mag. 100 (1874) t. 6104; Hook. f. Fl. Brit. Ind. 5 (1890) 854; Veitch Man. Orch. Pl. pt. 6 (1890) 63; Ridl. in Journ. Linn. Soc. 32 (1896) 330, Mat. Fl. Mal. Penin. 1 (1907) 122.

Alismorchis pulchra Ktze. Rev. Gen. Pl. 2 (1891) 650.

Calanthe curantigiodes Ktze. Rev. Gen. Pl. 2 (1891) 650, *sphalm*, in syn.

Luzon (Sorsogon), Mindoro, Negros (Oriental Negros), Leyte, *Elmer* 9576, 14883, *Merrill* 5849, 5850, *Wenzel* 0648, 0689, mixed with 0704. Terrestrial at 130 to 760 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

CALANTHE AUSTRALIS Ait. ex Loud. Hort. Brit. Suppl. 2 (1839) 615, *nomen* = *Spiranthes sinensis* (Pers.) Ames.

CALANTHE EMARGINATA Lindl. apud Naves Novis. App. (1882) 246.

CALANTHE JUSNERII Boxall ex Naves Novis. App. (1882) 247, *nomen*.

CALANTHE LILACINA Loher in Gard. Chron. III 47 (1910) 66, *text cut*, in Fedde Repert. 11 (1912) 96 = *Calanthe conspicua* Lindl.

CALANTHE MASUCA Boxall ex Naves Novis. App. (1882) 246, *nomen*.

CALANTHE NIVALIS Boxall ex Naves Novis. App. (1882) 246, *nomen*.

CALANTHE PHAJOIDES Reichb. f. Xen. Orch. 1 (1856) 207, t. 79, fig. II, 4 & 5; Naves Novis. App. (1882) 246 = *Calanthe angustifolia* (Blume) Lindl.

CALANTHE RAMOSII Ames in Philip. Journ. Sci. 7 (1912) Bot. 12 = *Phajus ramosii* Ames.

CALANTHE SPECIOSA Lindl. apud Naves Novis. App. (1882) 246.

CALANTHE TRIPPLICATA Ames in Philip. Journ. Sci. 2 (1907) Bot. 326, Orch. 2 (1908) 159, non *Orchis triplicata* Willem. = *Calanthe furcata* Batem.

CALANTHE VERATRIFOLIA R. Br. apud Naves Novis. App. (1882) 246; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 211, quoad syn. *C. furcata* Batem. et quoad Philip.; J. J. Sm. Orch. Amb. (1905) 23, quoad syn. *C. furcata* Batem. et quoad Philip. = *Calanthe furcata* Batem.

58. ASCOTAINIA Ridley

ASCOTAINIA ELMERI Ames Orch. 5 (1915) 99; Schltr. in Fedde Repert. Beihefte 5 (Orch. Sino-Jap. Prodr.) (1919) 246; Ames Sched. Orch. 6 (1923) 79.

Tainia elmeri Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1570.

Tainia inamoena Kränzl. in Fedde Repert. 17 (1921) 387.

Luzon (Bontoc, Benguet), Mindanao (Agusan, Davao), *Vanoverbergh* 1110, 2434, 4052, *Elmer* 8526, *Weber* 042, 061, 075, 0263, *Loher* s. n. Terrestrial in forest, at 3 to 1,300 m altitude.

59. SPATHOGLOTTIS Blume

SPATHOGLOTTIS CHRYSANTHA Ames Orch. 2 (1908) 161, *text cut*, 5 (1915) 99.

Luzon (Bontoc, Lepanto, Ifugao, Benguet, Nueva Vizcaya, Albay). *Vanoverbergh* 378, 1211, 4039, *F. B.* 14494 *Darling*, *Ramos & Edaño* 72, *B. S.* 14387, 19884 *McGregor*, 5374, 5546, 8179 *Ramos*, *Sandkuhl* 295, *Elmer* 5779, 8815, *Lyon* 156, *R. S. Williams* 1064, *Mearns* 2905. At 1,400 to 1,650 m altitude.

SPATHOGLOTTIS ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1572, Orch. 5 (1915) 99.

Spathoglottis aurea Lindl. apud Naves Novis. App. (1882) 246;

Ames in Philip. Journ. Sci. 2 (1907) Bot. 327, Orch. 5 (1915) 99.

Mindoro, Negros (Oriental Negros), *Merritt* 4375, *Elmer* 10080. At 1,415 m altitude on Mt. Halcon and from Dumaguete (Cuernos Mountains).

SPATHOGLOTTIS KIMBALLIANA Hort. Sander ex Gartenflora 41 (1892) 184, *nomen*; Rolfe in Sander Reichenbachia II 2 (1894) 95, t. 88, *text cut*; Hook. f. in Bot. Mag. 121 (1895) t. 7443.

Var. **ANGUSTIFOLIA** Ames Orch. 5 (1915) 99.

Luzon (Ilocos Norte, Isabela, Albay), Sibuyan, *B. S.* 32872 *Ramos*, 21149 *Escritor*, 6284 *Robinson*, *Elmer* 12556. The species proper is from Borneo.

SPATHOGLOTTIS PLICATA Blume Bijdr. (1825) 401, fig. 76; Lindl. Gen. & Sp. Orch. (1831) 119, in Wall. Cat. (1831) No. 3734; Hassk. in Tijdschr. Nat. Gesch. 9 (1842) 139; Lindl. in Bot. Reg. 30 (1844) Misc. 45; Hassk. Pl. Jav. Rar. (1848) 126; De Vriese Ill. Orch. Ind. Neerl. (1854) t.; Moore Ill. Orch. Pl. (1857) *Spathogl.* 2; Miq. Fl. Ind. Bat. 3 (1859) 673; Reichb. f. in Walp. Ann. 6 (1862) 455; Hemsl. in Gard. Chron. II 18 (1882) 532; Naves Novis. App. (1882) 245; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 269; Linden & Rodigas in Lindenia 2 (1886) 15, t. 54; Kränzl. in Engl. Bot. Jahrb. 7 (1886) 438, in K. Schum. & Hollr. Fl. Kais. Wilhelmsland (1889) 33; Hook. f. Fl. Brit. Ind. 5 (1890) 813; Kränzl. Xen. Orch. 3 (1893) 112; Grant Orch. Burma (1895) 192; Ridl. in Journ. Linn. Soc. 31 (1896) 285, 32 (1896) 311; Kränzl. in Reinecke in Engl. Bot. Jahrb. 25 (1898) 603, in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 244; Volkens in Engl. Bot. Jahrb. 31 (1902) 461; Schltr. in Perk. Frag. Fl. Philip. (1904) 43; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 219, Fig.-Atlas (1909) fig. 162, Orch. Amb. (1905) 24; Ames Orch.

1 (1905) 83, 2 (1908) 61, 5 (1915) 100, 6 (1920) 164; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 117; Ames in Philip. Journ. Sci. 2 (1907) Bot. 327; J. J. Sm. in Nova Guinea 8 (1909) 25, 8³ (1912) 529, 12 (1913) 14; Schltr. in Fedde Repert. 10 (1911) 70, in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 27; J. J. Sm. in Philip. Journ. Sci. 12 (1917) Bot. 255, in Merr. Interp. Rumph. Herb. Amb. (1917) 172; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 245; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 182.

Bletia angustata Gaudich. in Freyc. Voy. Uran. et Physic. Bot. (1826) 421.

Bletia angustifolia Gaudich. in Freyc. Voy. Uran. et Physic. Bot. (1826) t. 32.

Paxtonia rosea Lindl. in Bot. Reg. 24 (1838) Misc. 61, t. 60; Moore Ill. Orch. Pl. (1857) Paxtonia 4, t. 1; Reichb. f. in Walp. Ann. 6 (1862) 456; Naves Novis. App. (1882) 237.

Spathoglottis lilacina Griff. Notul. 3 (1851) 323 and Icon. Pl. Asiat. 3 (1851) t. 311, fig. III.

Phajus rhumphii Blume Mus. Bot. Lugd.-Bat. 2 (1856) 179.

Spathoglottis spicata Lindl. in Journ. Linn. Soc. 3 (1859) 9.

Batan Islands, Luzon (Bontoc, Lepanto, Ifugao, Benguet, Nueva Vizcaya, Tarlac, Pampanga, Rizal, Laguna, Tayabas, Camarines, Sorsogon), Mindoro, Panay (Capiz), Negros (Occidental Negros), Samar, Leyte, Palawan, Dinagat, Mindanao (Zamboanga, Lanao, Cotabato, Bukidnon, Agusan, Surigao). A common and widely distributed species in the Philippines. Terrestrial at 160 to 1,500 m altitude. In Malay Peninsula, Formosa, Sumatra, Java, Amboina, Celebes, Borneo, New Guinea, and the Caroline Islands.

SPATHOGLOTTIS TOMENTOSA Lindl. in Bot. Reg. 31 (1843) sub t. 19; Reichb. f. in Walp. Ann. 6 (1862) 455; Naves Novis. App. (1882) 246; Vidal Phan. Cuming. Philip. (1885) 149; Kränzl. Xen. Orch. 3 (1893) 112; Ames Orch. 2 (1908) 163 (excl. Luzon spec.), 5 (1915) 100.

Mindanao (Misamis, Lanao, Agusan, Surigao), Cuming 2108, M. S. Clemens 803, Weber 50, Bolster 256, (?) F. B. 7590 Hutchinson. Terrestrial at 200 to 430 m altitude.

SPATHOGLOTTIS VANOVERBERGHII Ames Orch. 6 (1920) 299.

Spathoglottis tomentosa Ames Orch. 2 (1908) 163, non Lindl., as to specimens from Luzon.

Luzon (Bontoc, Lepanto, Benguet), Vanoverbergh 163, 1109, 1112, 2106, 3943, 4046, Merrill 4287, Elmer 8478. At 1,000 to 1,300 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

SPATHOGLOTTIS AFFINIS De Vriese apud Naves Novis. App. (1882) 245.

SPATHOGLOTTIS AUREA Lindl. apud Naves Novis. App. (1882) 246; Ames in Philip. Journ. Sci. 2 (1907) Bot. 327, Orch. 5 (1915) 99 = *Spathoglottis elmeri* Ames.

SPATHOGLOTTIS TOMENTOSA Ames Orch. 2 (1908) 163, non Lindl., as to specimens from Luzon = *Spathoglottis vanoverberghii* Ames.

60. PLOCOGLOTTIS Blume

PLOCOGLOTTIS BICALLOSA Ames in Elm. Leaf. Philip. Bot. 5 (1915) 1571, Orch. 5 (1915) 100.

Luzon (Sorsogon), Negros (Oriental Negros), Mindanao (Camiguin Island), *Elmer* 9577, 17117, *B. S.* 14437 *Ramos*. Terrestrial at 1,300 m altitude in Negros.

PLOCOGLOTTIS COPELANDII Ames in Philip. Journ. Sci. 2 (1907) Bot. 326, Orch. 5 (1915) 100.

Plocoglottis acuminata Ames Orch. 1 (1905) 82, non Blume.

Luzon (Laguna, Tayabas, Camarines, Sorsogon), Polillo, Mindoro, Panay (Capiz), Biliran, Leyte, Palawan, Mindanao (Camiguin Island, Zamboanga, Cotabato, Agusan, Davao). A common and widely distributed species. Terrestrial in forest at 10 to 580 m altitude.

PLOCOGLOTTIS LUCBANENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1572, Orch. 5 (1915) 100.

Babuyan Islands (Camiguin Island), Luzon (Laguna, Tayabas, Sorsogon), *B. S.* 4147 *Fenix*, *F. B.* 28938 *Willie & Salvosa*, *Elmer* 7707, 17123. Terrestrial in damp forest at 915 m altitude in Tayabas.

PLOCOGLOTTIS MCGREGORII Ames Orch. 7 (1922) 121.

Panay (Antique), *B. S.* 32189 *McGregor*. East of Culasi, in mossy forest at 1,000 m altitude.

PLOCOGLOTTIS MINDORENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 327, Orch. 5 (1915) 101.

Mindoro, Sibuyan, Mindanao (Surigao), *Merrill* 5837, *Elmer* 12147, *Bolster* 278. Terrestrial in forest at 10 to 330 m altitude.

PLOCOGLOTTIS WENZELII Ames Orch. 5 (1915) 101.

Leyte, *Wenzel* 067, 0385, 0453. Ormoc and Jaro, terrestrial in forest at 200 to 500 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

PLOCOGLOTTIS ACUMINATA Ames Orch. 1 (1905) 82, non Blume = *Plocoglottis copelandii* Ames.

PLOCOGLOTTIS LATIFOLIA Blume apud Naves Novis. App. (1882) 247.

PLOCOGLOTTIS LOWII Reichb. f. apud Naves Novis. App. (1882) 247.

PLOCOGLOTTIS MOLUCCANA Blume apud Naves Novis. App. (1882) 247.

61. PACHYSTOMA Blume

PACHYSTOMA PUBESCENS Blume Bijdr. (1825) 376, fig. 29; Reichb. f. in Bonpl. 3 (1855) 251; Miq. Fl. Ind. Bat. 3 (1859) 675; Reichb. f. in Walp. Ann. 6 (1862) 463; Naves Novis. App. (1882) 247; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 186, Fig.-Atlas (1909) fig. 140; Ames in Philip. Journ. Sci. 6 (1911) Bot. 51, Orch. 5 (1915) 102.

Pachychilus pubescens Blume Mus. Bot. Lugd.-Bat. 2 (1856) 173, Fl. Jav. Orch. (1858) 24 (Orch. Arch. Ind. 27) t. 10, fig. 1, t. 11E.

Luzon (Cagayan, Bontoc, Lepanto, Benguet, Nueva Vizcaya, Rizal), Mindoro, *Vanoverbergh* 1102, 2435, 4040, *B. S.* 7050, 7969 *Ramos*, 11135

McGregor, 6774 Robinson, F. B. 11009 Merritt, 10942 Curran, Ramos s. n. Grassy hillsides up to 1,200 m altitude in Bontoc.

62. EULOPHIA R. Brown

EULOPHIA DENTATA Ames in Philip. Journ. Sci. 6 (1911) Bot. 51, Orch. 5 (1915) 102.

Eulophia graminea Naves Novis. App. (1882) 241 (?); Vidal Phan. Cuming. Philip. (1885) 150 (?), Rev. Pl. Vasc. Filip. (1886) 269 (?); Ames Orch. 2 (1908) 166, non Lindl.

Luzon (Bontoc, Isabela, Pangasinan), F. B. 17035 Curran, B. S. 7977 Ramos, Alberto 36, Cuming 2059.

EULOPHIA MACRORHIZA Blume Fl. Jav. Orch. (1858) 155 (Orch. Arch. Ind. 183) t. 63, fig. 2, t. 66B; Miq. Fl. Ind. Bat. 3 (1859) 742; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 227, Fig.-Atlas (1909) fig. 166; Ames Orch. 2 (1908) 167, 5 (1915) 104.

Graphorchis macrorhiza Ktze. Rev. Gen. Pl. 2 (1891) 662.

Luzon (Bataan, Tayabas, Sorsogon), Leyte, Whitford 621, Lyon 165, Elmer 16118, Wenzel 0121. Terrestrial at 60 to 1,000 m altitude. In Java.

EULOPHIA MACROSTACHYA Lindl. Gen. & Sp. Orch. (1833) 183, in Bot. Reg. 23 (1837) t. 1972; Wight Icon. 5¹ (1852) 8, t. 1667-68; Reichb. f. in Walp. Ann. 6 (1863) 646; Thw. Enum. Pl. Zeyl. (1864) 301; Hook. f. in Bot. Mag. 102 (1876) t. 6246, Fl. Brit. Ind. 6 (1890) 4; Trim. Handb. Fl. Ceyl. 4 (1898) 176; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 224, Fig.-Atlas (1909) fig. 164; Ames Orch. 2 (1908) 169, 5 (1915) 104; J. J. Sm. in Nova Guinea 8 (1909) 25, 12³ (1915) 215; Rolfe in Gibbs in Journ. Linn. Soc. 42 (1914) 157; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191.

Eulophia emarginata Blume Fl. Jav. Orch. (1858) 152 (Orch. Arch. Ind. 180); Miq. Fl. Ind. Bat. 3 (1859) 741; Miq. Sumatra (1862) 274, non Lindl.

Graphorchis macrostachya Ktze. Rev. Gen. Pl. 2 (1891) 662.

Graphorchis blumeana Ktze. Rev. Gen. Pl. 2 (1891) 663.

Luzon (Laguna, Camarines, Sorsogon), Leyte, Mindanao (Camiguin Island, Zamboanga, Lanao). Fairly common in the Philippines. Terrestrial in forest at low altitudes. In India, Ceylon, Sumatra, Java, Ternate, Borneo, and New Guinea.

EULOPHIA MERRILLII Ames Orch. 5 (1915) 104.

Mindanao (Surigao), Merrill 5452. Caraga, open grassy slopes at about 100 m altitude.

EULOPHIA SQUALIDA Lindl. in Bot. Reg. 27 (1841) Misc. 77; Hook. f. Fl. Brit. Ind. 6 (1890) 6; Ridl. in Journ. Linn. Soc. 31 (1896) 289, 32 (1896) 332, 36 (1898) 214; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 225, Fig.-Atlas (1909) fig. 165; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 137; Ames in Philip. Journ. Sci. 4 (1909) Bot. 671, Orch. 5 (1915) 105, 6 (1920) 210; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Rolfe in Gibbs

- in Journ. Linn. Soc. 42 (1914) 157; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 192.
- Cyrtopera squalida* Reichb. f. in Bonpl. 5 (1857) 38; Lindl. in Journ. Linn. Soc. 3 (1859) 31; Miq. Fl. Ind. Bat. 3 (1859) 686; Reichb. f. in Walp. Ann. 6 (1863) 668; Naves Novis. App. (1882) 242.
- ? *Eulophia lutea* Blume Fl. Jav. Orch. (1858) 153 (Orch. Arch. Ind. 181); Miq. Fl. Ind. Bat. 3 (1859) 742.
- ? *Eulophia celebica* Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Miq. Fl. Ind. Bat. 3 (1859) 742.
- Eulophia elongata* Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Miq. Fl. Ind. Bat. 3 (1859) 742; Naves Novis. App. (1882) 241.
- Eulophia mucronata* Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Miq. Fl. Ind. Bat. 3 (1859) 742; Naves Novis. App. (1882) 242.
- Cyrtopodium squalidum* Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270.
- Eulophia elata* Hook. f. Fl. Brit. Ind. 6 (1890) 3, fide Ridl.
- Graphorchis elata* Ktze. Rev. Gen. Pl. 2 (1891) 662.
- ? *Graphorchis lutea* Ktze. Rev. Gen. Pl. 2 (1891) 662.
- Graphorchis squalida* Ktze. Rev. Gen. Pl. 2 (1891) 662.
- ? *Eulophia squalida* Lindl. var. *celebica* Schltr. in Fedde Repert. 10 (1911).
- Luzon (Bontoc, Lepanto, Benguet, Zambales), Leyte, Bancalan, Palawan, Mindanao (Davao). A fairly common and rather widespread species in the Philippines. Terrestrial at 5 to 1,300 m altitude. In Malay Peninsula, ? Sumatra, Java, Celebes, and Borneo.
- EULOPHIA STRICTA** (Presl) Ames Orch. 5 (1915) 103.
- Bletia stricta* Presl Rel. Haenk. 1 (1827) 98; Naves Novis. App. (1882) 247.
- Eulophia exaltata* Reichb. f. in Bonpl. 5 (1857) 38; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 228, Fig.-Atlas (1910) fig. 167; Ames Orch. 1 (1905) 85, 2 (1908) 166, 5 (1915) 103; J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 13 (1907) 22.
- Eulophia leschenaultii* Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Naves Novis. App. (1882) 242.
- Cyrtopera ensiformis* Lindl. in Journ. Linn. Soc. 3 (1859) 31; Miq. Fl. Ind. Bat. 3 (1859) 686; Naves Novis. App. (1882) 242.
- Cyrtopodium ensiforme* Vidal Phan. Cuming. Philip. (1885) 80, 150, Rev. Pl. Vasc. Filip. (1886) 270; Schltr. in Perk. Frag. Fl. Philip. (1904) 43.
- Graphorchis exaltata* Ktze. Rev. Gen. Pl. 2 (1891) 662.
- Eulophia warburgiana* Kränzl. Xen. Orch. 3 (1893) 115, t. 266, fig. II, 8-12.
- Eulophia ensiformis* Schltr. in Perk. Frag. Fl. Philip. (1904) 202.
- Luzon (Bontoc, Nueva Vizcaya, Rizal, Camarines), Mindanao (Zamboanga, Bukidnon, Lanao, Butuan, Davao). Not uncommon in the Philippines, terrestrial at 80 to 1,400 m altitude. In Java and Celebes.

EULOPHIA VANOVERBERGHII Ames in Philip. Journ. Sci. 7 (1912) Bot. 13, Orch. 5 (1915) 105.

Luzon (Bontoc), *Vanoverbergh* 336, 1188, 3876, 4051. Terrestrial at about 1,300 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

EULOPHIA BICOLOR Blume apud Naves Novis. App. (1882) 241.

EULOPHIA ELONGATA Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Naves Novis. App. (1882) 242 = *Eulophia squalida* Lindl.

EULOPHIA ENSIFORMIS Schltr. in Perk. Frag. Fl. Philip. (1904) 202 = *Eulophia stricta* (Presl) Ames.

EULOPHIA EXALTATA Reichb. f. in Bonpl. 5 (1857) 38; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 228, Fig.-Atlas (1910) fig. 167; Ames Orch. 1 (1905) 85, 2 (1908) 166, 5 (1915) 103; J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 13 (1907) 22 = *Eulophia stricta* (Presl) Ames.

EULOPHIA GRAMINEA Naves Novis. App. (1882) 241; (?); Vidal Phan. Cuming. Philip. (1885) 150 (?), Rev. Pl. Vasc. Filip. (1886) 269 (?); Ames Orch. 2 (1908) 166, non Lindl. = *Eulophia dentata* Ames.

EULOPHIA LESCHENAULTII Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Naves Novis. App. (1882) 242 = *Eulophia stricta* (Presl) Ames.

EULOPHIA MUCRONATA Blume Fl. Jav. Orch. (1858) 154 (Orch. Arch. Ind. 182); Naves Novis. App. (1882) 242 = *Eulophia squalida* Lindl.

EULOPHIA SAUNDERSIANA Reichb. f. apud Naves Novis. App. (1882) 242.

EULOPHIA SUMATRANA Blume apud Naves Novis. App. (1882) 242.

Closely allied to *Eulophia squalida* Lindl. The Philippine material is presumably referable to that species.

EULOPHIA VENOSA Reichb. f. apud Naves Novis. App. (1882) 242.

EULOPHIA WARBURGIANA Kränzl. Xen. Orch. 3 (1893) 115, t. 266, fig. II. 8-12 = *Eulophia stricta* (Presl) Ames.

63. **GEODORUM** Jackson

GEODORUM NUTANS (Presl) Ames Orch. 2 (1908) 164, 5 (1915) 106, 6 (1920) 306; Merr. Fl. Manila (1912) 166, Sp. Blancoanae (1918) 113; Wells in Philip. Journ. Sci. 14 (1919) 4.

Dendrobium nutans Presl Rel. Haenk. 1 (1827) 102, Symb. Bot. 1 (1832) 34, t. 23; Naves Novis. App. (1882) 234.

Arethusa glutinosa Blanco Fl. Filip. (1837) 641, ed. 2 (1845) 446, ed. 3, 3 (1879) 38, t. 429bis.

Dendrobium haenkeanum Steud. Nomencl. Bot. ed. 2, 1 (1840) 490.

Geodorum semicristatum Lindl. Fol. Orch. Geodorum (1854) 2; Miq. Fl. Ind. Bat. 3 (1859) 712; Reichb. f. in Walp. Ann. 6 (1863)

- 633; Naves Novis. App. (1882) 241; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Merr. in Govt. Lab. Publ. (Philip.) 8 (1903) 45, 52, 61, 75, 88, 152, 27 (1905) 84.
 ? *Geodorum pallidum* Llanos in Mem. Real. Acad. Cienc. Madrid 4 (1859) 507; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 105.

Babuyan Islands (Dalupiri Island), Luzon (Nueva Vizcaya, Bataan, Rizal, Cavite, Laguna, Tayabas, Camarines, Sorsogon), Polillo, Masbate, Panay (Antique), Negros (Oriental Negros), Biliran, Samar, Palawan, Bancalan, Mindanao (Bukidnon, Davao). A common and widely spread, endemic species; terrestrial at sea level to 300 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

- GEODORUM CITRINUM* Jacks. apud Naves Novis. App. (1882) 241.
GEODORUM DILATATUM R. Br. apud Naves Novis. App. (1882) 241.
GEODORUM PALLIDUM Llanos in Mem. Real. Acad. Cienc. Madrid 4 (1859) 507; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 105 = ? *Geodorum nutans* (Presl) Ames.
GEODORUM PICTUM Lindl. apud Naves Novis. App. (1882) 241.
GEODORUM PURPUREUM R. Br. apud Naves Novis. App. (1882) 241.
GEODORUM SEMICRISTATUM Lindl. Fol. Orch. *Geodorum* (1854) 2; Miq. Fl. Ind. Bat. 3 (1859) 712; Reichb. f. in Walp. Ann. 6 (1863) 633; Naves Novis. App. (1882) 241; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Merr. in Govt. Lab. Publ. (Philip.) 8 (1903) 45, 52, 61, 75, 88, 152, 27 (1905) 84 = *Geodorum nutans* (Presl) Ames.

64. *DENDROBIUM* Swartz

- DENDROBIUM ACLINIA* Reichb. f. in Bonpl. 4 (1856) 329.
Dendrobium pseudacclinia Lindl. in Journ. Linn. Soc. 3 (1859) 9; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 297; Ames Orch. 5 (1915) 134.

Luzon (Isabela, Zambales, Rizal), *B. S.* 5618, 19065 (cult.) Curran, *Lyon* 63, 148, 150, *B. S.* 19151 Reillo 7. Epiphyte up to 330 m altitude in Rizal.

Reichenbach states that his specimen came from "Ostindien," a very indefinite localization. Lindley in 1859 renamed the species, without justification, *Dendrobium pseudacclinia*, citing *D. acclinia* Reichb. f. as a synonym, and then made a new *D. acclinia* based on the genus *Aclinia* of Griffith. This proceeding is certainly unjustified and, moreover, the *D. acclinia* of Lindley is considered by J. D. Hooker, Fl. Brit. Ind. 5 (1890) 718, to be a peloric form of *D. incurvum* Lindl. Consequently *D. acclinia* Reichb. f. is reinstated in place of *D. pseudacclinia* Lindl.

- DENDROBIUM ACUMINATISSIMUM* (Blume) Lindl. Gen. & Sp. Orch. (1830) 86; Lindl. in Bot. Reg. 30 (1844) Misc. 50; Miq. Fl. Ind. Bat. 3 (1859) 636; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 350, Fig.-Atlas (1910) fig. 265, in Nova Guinea 8 (1909) 71;

Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 170;
Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 35; Ames Orch.
5 (1915) 114; J. J. Sm. in Philip. Journ. Sci. 12 (1917) Bot. 256.

Grastidium acuminatissimum Bl. Bijdr. (1825) 333.

Dendrobium caudatum Teijsm. & Binn. var. *javanica* Teijsm. & Binn.
in Nat. Tijdschr. Ned. Ind. 24 (1862) 315.

Callista acuminatissima Ktze. Rev. Gen. Pl. 2 (1891) 654.

Leyte, Wenzel 047. Dagami, epiphyte at 60 m altitude. In Sumatra,
Java, Amboina, and New Guinea.

DENDROBIUM ACUMINATUM Rolfe ex Ames Orch. 1 (1905) 86, 2
(1908) 1, 170, t. 17, 5 (1915) 114.

? *Dendrobium cymbidioides* Blume apud Naves Novis. App. (1882) 231.

Sarcopodium cymbidioides Kränzl. in Engl. Pflanzenreich IV. 50.
II. B. 21 (1910) 325, quoad Philip.

Sarcopodium acuminatum Rolfe in Orch. Rev. 18 (August, 1910)
239; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (November,
1910) 329, fig. 32 D, E.

Luzon (Kalinga, Bataan), Lyon 47, Boettcher s. n., Whitford 223, F. B.
738 Borden, Topping 461, Weber 23, Copeland s. n. Epiphyte at 1,000 to
2,000 m altitude.

DENDROBIUM AGUSANENSE Ames Sched. Orch. 6 (1923) 79.

Mindanao (Agusan), Elmer 12462. At 330 m altitude.

DENDROBIUM ALBAYENSE Ames in Philip. Journ. Sci. 7 (1912) Bot.
14, Orch. 5 (1915) 115.

Luzon (Laguna, Tayabas, Albay), Negros (Occidental Negros), Leyte,
Mindanao (Butuan, Agusan, Davao). A fairly common species; epiphytic
at sea level to 900 m altitude.

DENDROBIUM AMETHYSTOGLOSSUM Reichb. f. in Gard. Chron. (1872)
109; Hook. f. in Bot. Mag. 98 (1872) t. 5968; Veitch Man. Orch.
Pl. pt. 3 (1888) 16; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21
(1910) 115; Ames Orch. 5 (1915) 115.

Callista amethystoglossa Ktze. Rev. Gen. Pl. 2 (1891) 654.

Luzon (Benguet), Lyon 157.

DENDROBIUM ANOSMUM Lindl. in Bot. Reg. 31 (1845) Misc. 32; Naves
Novis. App. (1882) 234; J. J. Sm. in Winkler in Engl. Bot. Jahrb.
48 (1912) 101, in Nova Guinea 12 (1913) Bot. 74, 12⁴ (1916)
Bot. 351, in Merr. Interp. Rumph. Herb. Amb. (1917) 175; Merr.
Sp. Blancoanae (1918) 114; Ames in Merr. in Journ. Roy. Asiat.
Soc. Straits Branch, Special No. (1921) 156.

Dendrobium macrophyllum Lindl. in Bot. Reg. 25 (1839) Misc. 36,
28 (1842) Misc. 84, 30 (1844) Misc. 47, Sert. Orch. (1840) t. 35;
Paxt. Mag. Bot. 8 (1841) 97, t.; Hook. Cent. Orch. Pl. (1851)
t. 12; Planch. Hort. Donat. (1855) 156, Atlas t. 1; Reichb.
f. in Linden Pescat. 1 (1860) 40, t. 40, non A. Rich.

Dendrobium macranthum Hook. in Bot. Mag. 69 (1842) t. 3970;
Paxt. Mag. Bot. 9 (1842) 238; Planch. in Fl. des Serres 8 (1852)
21, t. 757; Miq. Fl. Ind. Bat. 3 (1859) 642; Naves Novis. App.
(1882) 233; Ceron Cat. Pl. Herb. Manila (1892) 166, non A. Rich.

- Epidendrum ruibarbarum redolens* Blanco Fl. Filip. ed. 2 (1845) 593.
- Dendrobium macrophyllum* Lindl. var. *giganteum* Lindl. in Gard. Chron. (1854) 219; Warner Sel. Orch. Pl. I (1862-65) t. 26.
- Dendrobium macranthum* Hook. var. *purpureo-marginatum* De Vriese Ill. Orch. Ind. Neerl. (1854) t. 17.
- Dendrobium macrophyllum* Lindl. var. *pallidum* Planch. Hort. Donat. (1855) 156.
- ? *Dendrobium retusum* Llanos in Mem. Real Acad. Cienc. Madr. 4 (1857) 498; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100.
- Dendrobium macranthum* Hook. var. *marginatum* De Vriese ex Miq. Fl. Ind. Bat. 3 (1859) 643; Naves Novis. App. (1882) 233.
- Dendrobium superbum* Reichb. f. in Walp. Ann. 6 (1861) 282, in Gard. Chron. II 17 (1882) 776; Warner & Will. Orch. Alb. 1 (1882) t. 42; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 267; Veitch Man. Orch. Pl. pt. 3 (1888) 77; Du Buys. in l'Orchidoph. 12 (1892) 70; Ridl. in Journ. Bot. 36 (1898) 211; Cogn. in Dict. Icon. Orch. (1899) Dendrob. t. 20; Finet in Bull. Mus. Hist. Nat. Par. 6 (1903) 300; J. J. Sm. Orch. Amb. (1905) 67; Ames Orch. 1 (1905) 90, 2 (1908) 186, 5 (1915) 139; Rolfe in Orch. Rev. 14 (1906) 177, fig. 21; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 56; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 33; O'Brien in Gard. Chron. III 72 (1922) 9.
- Dendrobium superbum* Reichb. f. var. *anosmum* Reichb. f. in Walp. Ann. 6 (1861) 283; Veitch Man. Orch. Pl. pt. 3 (1889) 77; Rolfe in Lindenia 6 (1890) 51, t. 264; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 33.
- Dendrobium superbum* Reichb. f. var. *giganteum* Reichb. f. in Walp. Ann. 6 (1861) 283; Ames Orch. 2 (1908) 187; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 33.
- Dendrobium superbum* Reichb. f. var. *purpureo-marginatum* Reichb. f. in Walp. Ann. 6 (1861) 283; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 33.
- Dendrobium macrophyllum* Lindl. var. *purpureo-marginatum* De Vriese ex Reichb. f. in Walp. Ann. 6 (1861) 283, in syn.
- Dendrobium superbum* Reichb. f. var. *huttonii* Reichb. f. in Gard. Chron. (1869) 206; Rolfe in Lindenia 6 (1890) 52, sub t. 264; Ames Orch. 2 (1905) 187; O'Brien in Gard. Chron. III 72 (1922) 9, fig. 6.
- Epidendrum rhabarbarum redolens* Blanco Fl. Filip. ed. 3, 3 (1879) 45, t. 389.
- Dendrobium macranthum* Hook. var. *giganteum* Naves Novis. App. (1882) 233; Roman in Journ. des Orch. 6 (1895) 284.
- ? *Dendrobium macranthum* Hook. var. *album* Naves Novis. App. (1882) 233.
- Dendrobium superbum* Reichb. f. var. *burkei* Reichb. f. in Gard. Chron. II 21 (1884) 306; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 34; O'Brien in Gard. Chron. III 72 (1922) 9.
- Dendrobium scorteckinii* Hook. f. Fl. Brit. Ind. 5 (1890) 741; Ridl. in Journ. Linn. Soc. 32 (1896) 264.
- Callista anosma* Ktze. Rev. Gen. Pl. 2 (1891) 653.

Callista scortechinii Ktze. Rev. Gen. Pl. 2 (1891) 655.

Dendrobium superbum Reichb. f. var. *dearei* Day ex Rolfe in *Lindenia* 6 (1890) 52, sub t. 264; Rolfe in *Orch. Rev.* 14 (1906) 177, fig. 22; O'Brien in *Gard. Chron.* III 72 (1922) 9.

Dendrobium leucorhodum Schltr. in *Fedde Repert. Beihefte* 1 (*Orch. Deutsch-Neu-Guin.*) (1912) 499.

Luzon (Abra, Bontoc, Lepanto, Benguet, Nueva Vizcaya, Rizal). Fairly frequent in Luzon, up to 760 m altitude in Bontoc. In Malay Peninsula, Indo-China (fide Finet), Celebes, Amboina, Borneo, and New Guinea.

DENDROBIUM APOROIDES (Lindl.) Merr. Sp. Blancoanae (1918) 113.

Aporum incrassatum Brongn. in *Duperr. Voy. Coquille* (1834) 204, t. 42B, non Blume.

Epidendrum equitans Blanco Fl. Filip. (1837) 645, ed. 2 (1845) 449, ed. 3, 3 (1879) 44, non *Dendrobium equitans* Kränzl.

Eria aporoides Lindl. in *Journ. Linn. Soc.* 3 (1859) 60; Miq. Fl. Ind. Bat. 3 (1859) 657; Naves Novis. App. (1882) 236; Merr. in *Govt. Lab. Publ. (Philip.)* 27 (1905) 84.

Pinalia aporodes Ktze. Rev. Gen. Pl. 2 (1891) 678.

Dendrobium brongniartii Kränzl. in *Engl. Pflanzenreich* IV. 50. II. B. 21 (1910) 210; Ames in *Philip. Journ. Sci.* 7 (1912) Bot. 15, *Orch.* 5 (1915) 116.

Dendrobium equitans Kränzl. in *Engl. Pflanzenreich* IV. 50. II. B. 21 (1910) 228, quoad Philip.

Luzon (Ilocos Norte, Cagayan, Nueva Vizcaya, Zambales, Rizal, Tayabas, Camarines, Sorsogon), Leyte, Basilan, Mindanao (Camiguin Island, Agusan, Surigao, Davao). A fairly common and widely distributed species in the Philippines at sea level to 1,000 m altitude.

DENDROBIUM APPENDICULOIDES Ames *Orch.* 7 (1922) 93.

Mindanao (Bukidnon), *B. S.* 38940 *Ramos & Edaño*. Mt. Candoon; epiphyte in forest at 1,650 m altitude.

DENDROBIUM BASILANENSE Ames in *Philip. Journ. Sci.* 7 (1912) Bot. 14, *Orch.* 5 (1915) 116.

Panay (Iloilo), Basilan, Mindanao (Zamboanga, Agusan), *B. S.* 18187 *Robinson, F. B.* 3968 *Hutchinson, Merrill* 8068, *Weber s. n.*, 45, 48, 132. Epiphyte at sea level to 400 m altitude.

DENDROBIUM BULLENIANUM Reichb. f. in *Bot. Zeit.* 20 (1862) 214; Kränzl. in *Engl. Pflanzenreich* IV. 50. II. B. 21 (1910) 133; Ames *Orch.* 5 (1915) 117.

Dendrobium erythroxanthum Reichb. f. in *Gard. Chron.* II 2 (1874) 162; Boxall ex Naves Novis. App. (1882) 234; Kränzl. in *Engl. Bot. Jahrb.* 25 (1898) 604.

No specimens seen, but drawings have been studied.

Regarding *Dendrobium bullenianum* Reichenbach states: "Sie soll aus Manilla stammen." Kränzlin gives in distribution "Luzon?... Samoa-Inseln." Reichenbach's specimen may have come from Polynesia rather than from the Philippines. *Dendrobium erythroxanthum* is supposed to have come from the Philippines.

It seems probable that *Dendrobium chameleon* Ames is conspecific, but sufficient evidence of this fact is at present lacking.

DENDROBIUM BUSUANGENSE Ames Orch. 6 (1920) 296, *t.* 101.

Busuanga, *Taylor s. n.* (cult. in Manila).

DENDROBIUM CABADBARENSE Ames Orch. 5 (1915) 117.

Mindanao (Agusan, Davao), *Weber 157, 192, 297*, with 310, *Black s. n.* Epiphyte at sea-level to 100 m altitude.

DENDROBIUM CANDOONENSE Ames Orch. 7 (1922) 94.

Mindanao (Bukidnon), *B. S. 38716 Ramos & Edaño*. Epiphyte on forest slope, Mt. Candoon, at 1,650 m altitude.

DENDROBIUM CARINATUM (Linn.) Willd. Sp. Pl. 4 (1805) 133; Lindl.

Gen. & Sp. Orch. (1830) 92; Kränzl. in Engl. Pflanzenreich IV. 50.

II. B. 21 (1910) 298; Ames Orch. 5 (1915) 118.

Bontia luzonica, *geniculis inferioribus carinulatis* Petiver Gazophyll.

Decas 5 (1702) 70, *t.* 44, *fig.* 10.

Epidendrum carinatum Linn. Sp. Pl. (1753) 953, ed. 2 (1763) 1350.

Dendrobium robinsonii Ames in Philip. Journ. Sci. 8 (1913) Bot. 425, Orch. 5 (1915) 138.

Luzon (Zambales) Mindoro, *F. B. 6315 Curran*, *B. S. 3059 Foxworthy*, 5642 *Robinson*. Epiphyte in open sandy spots.

DENDROBIUM CERAULA Reichb. f. in Linnaea 41 (1876) 40; Kränzl.

in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 104; Ames Orch.

5 (1915) 118.

Luzon, *Jagor 895, 896* (in Herb. Berol.).

DENDROBIUM CHAMELEON Ames Orch. 2 (1908) 174, *text cut*, 5

(1915) 119; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21

(1910) 134.

Luzon (Benguet), *R. S. Williams 1940*. Baguio; epiphyte, hanging from the trees.

DENDROBIUM CHRYSOGRAPHATUM Ames Orch. 5 (1915) 119.

Dendrobium angustifolium Ames in Philip. Journ. Sci. 7 (1912) Bot. 14, non Lindl.

Luzon (Bontoc, Benguet, Zambales, Bataan, Rizal), *Vanoverbergh 708, 709, 1346, 1522, Ramos & Edaño 95, F. B. 6972 Curran, R. S. Williams 764, B. S. 29588 Ramos & Edaño*. Epiphyte at 1,450 to 2,000 m altitude.

DENDROBIUM CLEMENSIAE Ames in Philip. Journ. Sci. 7 (1912) Bot.

16, Orch. 5 (1915) 120.

Negros (Occidental Negros), Leyte, Mindanao (Lanao), *Lyon 40, Wenzel 0530, 0591, 0599, 0906, M. S. Clemens 508*. Epiphyte at 500 to 700 m altitude in Leyte.

DENDROBIUM CRUMENATUM Sw. in Schrad. Journ. Bot. 2 (1799) 237,

in Kongl. Svensk. Vet.-Akad. Nya Handl. 21 (1800) 246; Willd.

Sp. Pl. 4 (1805) 137; Lindl. in Trans. Hort. Soc. 7 (1830) 40,

Gen. & Sp. Orch. (1830) 88; Roxb. Fl. Ind. ed. 2, 3 (1832) 480;

Lindl. in Bot. Reg. 25 (1839) t. 22; Paxt. Mag. Bot. 6 (1839) 90, 10 (1843) 189; Edit. in Allg. Berl. Gartenz. 7 (1839) 254, 11 (1843) 175; Knowles & Westc. Floral Cab. 3 (1840) 47; Hook. in Bot. Mag. 69 (1843) t. 4013; Griff. Notul. 3 (1851) 315; De Vriese Ill. Orch. Ind. Neerl. (1852) t. 3, fig. VI; Miq. Fl. Ind. Bat. 3 (1859) 638; Reichb. f. in Walp. Ann. 6 (1861) 291; Miq. Sumatra (1862) 274, Choix des Pl. (1863) sub t. 22, p. p. (excl. fig.); Thw. Enum. Pl. Zeyl. (1864) 297; Naves Novis. App. (1882) 232; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Rolfe in Lindenia 5 (1889) 33, t. 207; Hook. f. Fl. Brit. Ind. 5 (1890) 729; Ridl. in Trans. Linn. Soc. II 3 (1893) 362; Grant Orch. Burma (1895) 80; Ridl. in Journ. Linn. Soc. 31 (1896) 270, 32 (1896) 254; Trim. Handb. Fl. Ceyl. 4 (1898) 151; Finet in Bull. Mus. Hist. Nat. Par. 6 (1903) 297; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 330, Fig.-Atlas (1910) fig. 249, Orch. Amb. (1905) 42; Ames Orch. 1 (1905) 88, 2 (1908) 175, 5 (1915) 121, 7 (1922) 95; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 48; J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 43 (1910) 45; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 235, fig. 19 C, D, 6-8; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 34; Merr. Fl. Manila (1912) 168; J. J. Sm. in Winkler in Engl. Bot. Jahrb. 48 (1912) 98; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 158.

Onychium crumenatum Blume Bijdr. (1825) 326.

Callista crumenata Ktze. Rev. Gen. Pl. 2 (1891) 653.

Dendrobium cumulatum Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 115, quoad Philip.; Ames Orch. 5 (1915) 121.

Dendrobium caninum Merr. in Philip. Journ. Sci. 19 (1921) 344; Ridl. in Journ. Bot. 60 (1922) 274; Merr. in Journ. Bot. 61 (1923) 173, non *Epidendrum caninum* Burm. f.

Luzon (Benguet, Rizal, Zambales, Tayabas), Polillo, Samar (Basiao Island), Biliran, Leyte, Palawan, Mindanao (Camiguin Island, Zamboanga, Lanao, Agusan, Surigao, Davao). A common and widely distributed species in the Philippines, at sea level to 500 m altitude. In Ceylon, Burma, Malay Peninsula, Andaman Islands, Cochinchina, Sumatra, Java, Amboina, Timor, and Borneo.

DENDROBIUM CULTRATUM Schltr. in Fedde Repert. 10 (1911) 71; Ames Orch. 5 (1915) 121.

Dendrobium sinuatum Schltr. in Perk. Frag. Fl. Philip. (1904) 43; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 216, quoad Philip., non Lindl.

Dendrobium atropurpureum Ames Orch. 1 (1905) 86, 2 (1908) 172; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 217, quoad Philip., non Miq.

Luzon (Apayao, Ifugao, Pangasinan, Rizal, Tayabas), Polillo, Mindoro, Panay (Capiz), Negros (Occidental Negros), Samar, Palawan, Mindanao (Agusan, Surigao, Davao), Sibutu. A fairly common and widely distributed species in the Philippines, at sea level to 150 m altitude. In Celebes.

DENDROBIUM DEAREI Reichb. f. in Gard. Chron. II 18 (1882) 361; Warner & Will. Orch. Alb. 3 (1884) t. 120; Veitch Man. Orch. Pl. pt. 3 (1888) 37, *text cut*; Cogn. in Dict. Icon. Orch. (1905) Dendrob. t. 36; Ames in Philip. Journ. Sci. 4 (1909) Bot. 599; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 89; Rolfe in Orch. Rev. 19 (1911) 261; Ames Orch. 5 (1915) 122.

Mindoro, Dinagat, Mindanao, *F. B. 8525 Merritt, Lyon 121.*

DENDROBIUM DISTICHUM (Presl) Reichb. f. in Linnaea 41 (1876) 39; Naves Novis. App. (1882) 232; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 211 (excl. syn. in part); Ames Orch. 5 (1915) 122.

Schizmoceras disticha Presl Rel. Haenk. 1 (1827) 96, t. 13, fig. 2. *Dendrobium indivisum* Naves Novis. App. (1882) 231; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 217, quoad Philip.; Ames in Philip. Journ. Sci. 6 (1911) Bot. 52, non Miq.

Luzon (Benguet, Zambales, Bataan, Rizal, Camarines, Sorsogon), Polillo, Sulu Archipelago. A fairly common species in Luzon, at sea level to 900 m altitude.

DENDROBIUM ELMERI Ames in Elm. Leaff. Philip. Bot. 5 (1912) 1573, Orch. 5 (1915) 123.

Luzon (Laguna, Tayabas), *B. S. 16568, 20458 Ramos, 20851 Escritor, Serrato s. n., Calica s. n., Elmer 9249.* Epiphyte, in forest at 830 m altitude in Tayabas.

DENDROBIUM EPIDENDROPSIS Kränzl. in Orchis 2 (1908) 79, fig. 12; Ames in Philip. Journ. Sci. 4 (1909) Bot. 598; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 106; Ames Orch. 5 (1915) 123.

Luzon (Benguet, Bataan, Rizal), *Loher s. n., Lyon 118, 161, F. B. 2109 Borden, Elmer 6839, B. S. 3060 Ramos.*

DENDROBIUM ESCRITORII Ames Orch. 5 (1915) 123.

Luzon (Tayabas, Camarines), Mindanao (Surigao), *B. S. 20683 Escritor, 22019 Ramos, 27405 Alambra, Wenzel 01091.* Epiphyte at 35 to 150 m altitude.

DENDROBIUM EURORUM Ames Orch. 5 (1915) 124.

Leyte, *Wenzel 065, 0612.* Epiphyte in forest at 120 to 500 m altitude.

DENDROBIUM GERLANDIANUM Kränzl. in Fedde Repert. 6 (1909) 317, in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 238; Ames Orch. 5 (1915) 125.

The species was described from a *Loher* plant from Manila. The type has not been seen, but from the description it would seem to approach *D. philippinense* Ames.

DENDROBIUM HERCOGLOSSUM Reichb. f. in Gard. Chron. II 26 (1886) 487; Edit. in Hamb. Gartenz. 42 (1886) 558; Hook. f. Fl. Brit. Ind. 6 (1890) 185; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 75.

Dendrobium aduncum Hook. f. in Bot. Mag. 110 (1884) t. 6784, non Lindl.

Mindanao (Lanao) (so reported), *Lyon* 138.

In view of the above cited (reported Philippine) specimen it does not seem improbable that the material referred to *D. aduncum* Lindl. by Naves and by Ceron is in reality this species.

DENDROBIUM HETEROCARPUM Wall. Cat. (1828) No. 20, *nomen*, ex Lindl. Gen. & Sp. Orch. (1830) 78, Pl. Asiat. Rar. 2 (1831) 84, t. 196; Lindl. in Bot. Reg. 30 (1844) Misc. 49; Paxt. Mag. Bot. 14 (1847) 68; Hook. in Bot. Mag. 79 (1853) t. 4708; Planch. in Fl. des Serres 8 (1854?) 253, t. 842; Lem. Jard. Fleur. 4 (1854) t. 386; Batem. Second Cent. Orch. Pl. (1867) t. 150; Douglas in Gard. Chron. II 25 (1886) 397; Hook. f. Fl. Brit. Ind. 5 (1890) 737; Berkl. in Orch. Rev. 1 (1893) 82; Grant Orch. Burma (1895) 97; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik-Himal.) (1898) 53, t. 74; Trim. Handb. Fl. Ceyl. 4 (1898) 154; Cogn. in Dict. Icon. Orch. (1898) Dendrob. t. 10; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 368, Fig.-Atlas (1911) fig. 281; Ames Orch. 1 (1905) 89, 2 (1908) 176, *text cut*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 45; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Schltr. in Fedde Repert. 10 (1911) 37, Orchideen (1915) 260.

Dendrobium aureum Lindl. Gen. & Sp. Orch. (1830) 77, in Bot. Reg. 30 (1844) Misc. 51; Wight Icon. 5¹ (1852) 6, t. 1646; Lindl. in Fl. des Serres 7 (1852) 210; Reichb. f. in Walp. Ann. 6 (1861) 287; Thw. Enum. Pl. Zeyl. (1864) 297; Hemsl. in Gard. Chron. II 16 (1881) 688; Veitch Man. Orch. Pl. pt. 3 (1888) 19, *text cut*; Reichb. f. in Sander Reichenbachia I 2 (1890) 31, t. 63; Stein Orchideenb. (1892) 186; Du Buyss. in l'Orchidoph. 12 (1892) 184; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 56; Ames Orch. 5 (1915) 115.

Dendrobium aureum Lindl. var. *pallidum* Lindl. in Bot. Reg. 25 (1839) t. 20; Knowles & Westc. Floral Cab. 3 (1840) 48; Reichb. f. in Walp. Ann. 6 (1861) 287; Veitch Man. Orch. Pl. pt. 3 (1888) 19; Stein Orchideenb. (1892) 186.

Dendrobium rhombeum Lindl. in Bot. Reg. 29 (1843) t. 17, 30 (1844) Misc. 56; Reichb. f. in Bot. Zeit. 1 (1843) 333; Lindl. in Paxt. Flow. Gard. 3 (1852-53) 52; Naves Novis. App. (1882) 233; Reichb. f. in Sander Reichenbachia I 2 (1890) 31, in obs.

Dendrobium heterocarpum Wall. var. *henshallii* Hook. in Bot. Mag. 83 (1857) t. 4970; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 45.

Dendrobium aureum Lindl. var. *philippinense* Reichb. f. in Gard. Chron. II 13 (1880) 72; Veitch Man. Orch. Pl. pt. 3 (1888) 19; Stein Orchideenb. (1892) 187.

Dendrobium atractodes Ridl. in Journ. Linn. Soc. 23 (1885) 123, fide Kränzl.

Dendrobium aureum Lindl. var. *henshallii* Veitch Man. Orch. Pl. pt. 3 (1882) 19; Stein Orchideenb. (1892) 186.

Callista aurea Ktze. Rev. Gen. Pl. 2 (1891) 653.

Callista heterocarpa Ktze. Rev. Gen. Pl. 2 (1891) 654.

Dendrobium aureum Lindl. var. *rhombeum* Stein Orchideenb. (1892) 187.

Dendrobium heterocarpum Wall. var. *pallidum* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 45.

Dendrobium heterocarpum Wall. var. *philippinense* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 45.

Dendrobium hildebrandii Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 34, p. p., non Rolfe, fide Schltr.

Dendrobium minahassae Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 107, fide Schltr.

Luzon (Bontoc, Benguet, Bataan, Rizal, Albay, Sorsogon), Mindanao (Davao). Fairly common in the Philippines, especially in Luzon at 1,300 to 1,500 m altitude. In India, Ceylon, Burma, Malay Peninsula, Java, and Celebes.

It has seemed advisable to adopt the name *D. heterocarpum* in place of *D. aureum* Lindl., because the first definite union of the species occurs in Bot. Mag. 79 (1853) t. 4708 (cf. International Rules adopted at Vienna, Sect. 6, Art. 46) and also because it is the earliest name, although a *nomen*, and still further because it has been more commonly accepted by reputable authors.

DENDROBIUM HYMENANTHUM Reichb. f. in Bonpl. 3 (1855) 222, in Walp. Ann. 6 (1861) 302; Naves Novis. App. (1882) 234; Ames in Philip. Journ. Sci. 4 (1909) Bot. 598; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 288; Ames Orch. 5 (1915) 125.

Dendrochilum hymenanthum Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268, sphalm.

Luzon (Rizal, Tayabas), Polillo, Cuming 2135, Lyon 147, B. S. 3035 Ramos, 5647, 10453 McGregor, F. B. 28368 Mabesa.

DENDROBIUM INTERJECTUM Ames Orch. 5 (1915) 125.

Leyte, Wenzel 065a, 0534, 0574, 0594, 0677, 0691. Epiphyte in forest at 120 to 700 m altitude.

DENDROBIUM LEYTENSE Ames Orch. 5 (1915) 126.

Leyte, Wenzel 082, 0203, 0415. Epiphyte in forest at 60 to 500 m altitude.

DENDROBIUM LOHERIANUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 62.

Philippines, probably Luzon, *Loher s. n.*

Known only from the type collection.

DENDROBIUM LUNATUM Lindl. in Journ. Linn. Soc. 3 (1859) 4; Miq. Fl. Ind. Bat. 3 (1859) 630; Naves Novis. App. (1882) 231; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 218; Ames in Philip. Journ. Sci. 7 (1912) Bot. 17, in Elm. Leaf. Philip. Bot. 5 (1912) 1574, Orch. 5 (1915) 127.

Palawan, *Cuming s. n.*, 2136, B. S. 733, 907 Foxworthy, Elmer 12764, Weber 606, Merrill 9337, 9623, *Ledesma s. n.* Epiphyte at sea level to 500 m altitude.

DENDROBIUM LUZONENSE Lindl. in Bot. Reg. 30 (1844) Misc. 54; Reichb. f. in Walp. Ann. 6 (1861) 290; Naves Novis. App. (1882) 234; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 172, fig. 12 A-D; Ames Orch. 5 (1915) 127.

Dendrobium alagense Ames in Philip. Journ. Sci. 2 (1907) Bot. 328.

Luzon (Kalinga, Benguet, Sorsogon), Catanduanes, Mindoro, Leyte, Mindanao (Lanao, Agusan, Davao). A fairly common species; epiphyte at sea level to 1,300 m altitude.

DENDROBIUM LYONII Ames Orch. 2 (1908) 177, 5 (1915) 127, in Philip. Journ. Sci. 4 (1909) Bot. 672.

Dendrobium acuminatum Lyon in Gard. Chron. III 42 (1907) 210, fig. 88; Rolfe in Orch. Rev. 15 (1907) 312, as to plant spoken of; Kränzl. in Orchis 2 (1908) 73, t. 16, non Rolfe ex Ames; Edit. in Gard. Chron. III 46 (1909) 150, fig. 64, in Gardners' Mag. 52 (1909) 659, 660, text cut, non Rolfe ex Ames.

Sarcopodium lyonii Rolfe in Orch. Rev. 18 (1910) 240; Ames in Philip. Journ. Sci. 6 (1911) Bot. 53, in Addisonia 8 (1923) 17, t. 265.

Sarcopodium acuminatum Rolfe var. *lyonii* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 329, fig. 32 A-C.

Luzon (Bataan), *Lyon s. n.*, 10, F. B. 7153 Curran. Epiphyte at 660 to 1,100 m altitude.

DENDROBIUM MACGREGORII Ames in Philip. Journ. Sci. 7 (1912) Bot. 17, Orch. 5 (1915) 128.

Polillo, B. S. 10457 McGregor.

DENDROBIUM MARIVELENSE Ames Orch. 2 (1908) 180, text cut, 5 (1915) 128.

Dendrobium parviflorum Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 222, quoad Philip.; Ames in Philip. Journ. Sci. 7 (1912) Bot. 18, non Reichb. f.

Luzon (Bataan), Mindanao (Lanao), *Merrill s. n.*, Whitford 1127, M. S. Clemens s. n.

DENDROBIUM MERRILLII Ames Orch. 2 (1908) 181, *text cut*, 5 (1915) 126, in Philip. Journ. Sci. 4 (1909) Bot. 599; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 299.

Dendrobium aloifolium Reichb. f. apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 206, *quoad syn. D. merrillii*; Ames Orch. 5 (1915) 115 (*D. aloefolium*).

Luzon (Ifugao, Pangasinan, Rizal, Tayabas), Samar, *B. S. 19989 McGregor, 29863 Fenix, 24355 Ramos, Loher 6017, Merrill 3357*. Epiphyte at apparently rather low altitude (350 m).

DENDROBIUM MINDANAENSE Ames in Philip. Journ. Sci. 8 (1913) Bot. 423, Orch. 5 (1915) 128.

Luzon (Rizal), Polillo, Negros (Occidental Negros), Leyte, Culion, Palawan, Mindanao (Zamboanga, Agusan, Davao). A fairly common species of rather wide distribution. Terrestrial ? or epiphyte at low altitudes.

DENDROBIUM MODESTUM Reichb. f. in Bonpl. 3 (1855) 222, in Walp. Ann. 6 (1861) 281; Naves Novis. App. (1882) 234; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 220; Ames Orch. 5 (1915) 128.

Llanos s. n.

Reichenbach described this species from a *Llanos* specimen from Manila and ascribed to it affinity with *D. aciculare* Lindl. Kränzl. in his monograph cites a *Cuming* specimen in the Lindley Herbarium. This species has not been recognized in any of the material collected during the past twenty years.

DENDROBIUM MULTIRAMOSUM Ames Orch. 5 (1915) 129.

Luzon (Benguet, Rizal), *Williams 1934, Lyon 42*.

DENDROBIUM O'BRIENIANUM Kränzl. in Gard. Chron. III 11 (1892) 266; Ames Orch. 2 (1908) 182, *text cut*, 5 (1915) 130; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 105, *fig. 5 N-O*.

Dendrobium striatum Hort. ex Gard. Chron. III 38 (1905) 382; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 298.

Known only from a cultivated specimen sent from Luzon by Micholitz to F. Sander Co.

DENDROBIUM ORNITHOFLORUM Ames Orch. 3 (1908) 60, *t. 46*, 5 (1915) 130.

Mindanao (Misamis), *F. B. 4627 Mearns & Hutchinson*. Mt. Malindang at 1,900 m altitude.

DENDROBIUM PAPILIO Loher in Gard. Chron. III 21 (1897) 416; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 231, *excl. Williams 1934*; Rolfe in Orch. Rev. 22 (1914) 272; Ames Orch. 5 (1915) 130.

Dendrobium vanoverberghii Ames in Philip. Journ. Sci. 8 (1913) Bot. 425, *t. 13*.

Luzon (Bontoc, Benguet), Negros (Oriental Negros), Mindanao (Davao), *Vanoverbergh 704, 1345, B. S. 31846 Santos, Weber 246, Herre s. n., Boettcher s. n.* Epiphyte at 1,450 to 2,200 m altitude.

DENDROBIUM PERGRACILE Ames in Philip. Journ. Sci. 8 (1913) Bot. 423, Orch. 5 (1915) 131.

Mindanao (Lanao), *M. S. Clemens* 999. Camp Keithley, Lake Lanao.

DENDROBIUM PHILIPPINENSE Ames in Philip. Journ. Sci. 8 (1913) Bot. 424, Orch. 5 (1915) 131.

Dendrobium aciculare Lindl. apud Vidal Phan. Cuming. Philip. (1885) 148; Vidal Rev. Pl. Vasc. Filip. (1886) 267; Hook. f. Fl. Brit. Ind. 5 (1890) 725, quoad Philip.; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 241, quoad Philip.; Ames Orch. 5 (1915) 113.

Luzon (Laguna, Tayabas), Mindoro, Leyte, *Cuming* 2112, *Loher* 487, 6022, *Lyon* 67, *Reillo* s. n., *B. S.* 5641 *Griffin*, 20684 *Escritor*, 5658 *Merrill*, *Wenzel* 06, 077, 086, 0142. Epiphyte, apparently at low altitudes.

DENDROBIUM PLANIBULBE Lindl. in Bot. Reg. 29 (1843) Misc. 54; Reichb. f. in Walp. Ann. 6 (1861) 301; Naves Novis. App. (1882) 232; Ames Orch. 5 (1915) 132.

Dendrobium blumei Lindl. in Journ. Linn. Soc. 3 (1859) 14; Miq. Fl. Ind. Bat. 3 (1859) 638; Naves Novis. App. (1882) 232; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 234, quoad syn. *D. planibulbe* Lindl. et quoad Philip.

Samar, Mindanao (Zamboanga, Agusan), *Cuming* s. n., *B. S.* 24434 *Ramos*, *Reillo* s. n., *Sablaya* s. n., *F. B.* 5176 *Hutchinson*, *Weber* 174. Epiphyte at low altitude.

DENDROBIUM PLATYCAULON Rolfe in Kew Bull. (1892) 139, in Gard. Chron. III 12 (1892) 29, fig. 212; Kränzl. in Engl. Pflanzenreich IV. 50 II. B. 21 (1910) 267; Ames in Philip. Journ. Sci. 6 (1911) Bot. 52, Orch. 5 (1915) 132.

? *Dendrobium lamellatum* Lindl. apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 269, quoad Philip.; Ames Orch. 5 (1915) ix.

Merrill (cult.).

Known only from cultivated specimens.

DENDROBIUM PLICATILE Lindl. in Bot. Reg. 26 (1840) Misc. 10; Reichb. f. in Walp. Ann. 6 (1861) 307; Naves Novis. App. (1882) 234; Ames Orch. 5 (1915) 132.

Dendrobium macraei Ames Orch. 1 (1905) 89, 2 (1908) 176, non Lindl.

Desmotrichum fimbriatum Blume apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 354, quoad Philip.

Luzon (Tarlac, Pampanga, Bataan, Rizal, Laguna, Tayabas), Panay (Capiz), Leyte, Samar, Mindanao (Lanao, Cotabato, Agusan). A common and rather widely distributed species; epiphyte at 60 to 800 m altitude.

DENDROBIUM POLYTRICHUM Ames in Philip. Journ. Sci. 2 (1907) Bot. 328, *nomen*, Orch. 2 (1908) 183, 5 (1915) 133, 6 (1920) 297; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 240.

Luzon (Cagayan, Isabela, Bataan, Rizal, Laguna, Tayabas), Mindoro. A fairly common species, especially in Luzon; up to 260 m altitude in Mindoro.

DENDROBIUM PRISTINUM Ames Orch. 5 (1915) 133.

Luzon (Rizal), *B. S.* 3041 Ramos.

DENDROBIUM PROFUSUM Reichb. f. in Gard. Chron. II 21 (1884) 510; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 259; Ames Orch. 5 (1915) 134.

Roebelen s. n.

Known only from the description and a tracing of the analytical drawings in the Reichenbach Herbarium.

DENDROBIUM PSEUDOCONVEXUM Ames Orch. 5 (1915) 135.

Luzon (Bataan, Rizal, Laguna, Sorsogon). A number of collections have been made in Luzon. Epiphyte in forest at 600 to 700 m altitude.

DENDROBIUM PTEROCARPUM Ames Sched. Orch. 6 (1923) 80.

Mindanao (Davao), *Elmer* 11528. Epiphyte at 2,080 m altitude.

DENDROBIUM PURPUREOSTELIDIUM Ames Orch. 5 (1915) 136.

Luzon (Laguna, Sorsogon), Leyte, Mindanao (Agusan), Basilan, *B. S.* 15103 Ramos, 16258 Reillo, *Elmer* 16203, *Wenzel* 0193, 0228, *Weber* 191. Epiphyte at sea level and low altitudes.

DENDROBIUM RAMOSII Ames in Philip. Journ. Sci. 7 (1912) Bot. 18, Orch. 5 (1915) 138.

Luzon (Bontoc, Lepanto, Benguet), *Vanoverbergh* 9, 1715, *B. S.* 7049 Ramos, *Lyon* 154, *Merrill* 4682, *Ramos & Edaña* 115. Apparently terrestrial in rather open ground in mountainous regions at 1,200 to 1,300 m altitude.

DENDROBIUM SANDERAE Rolfe in Gard. Chron. III 45 (1909) 374, fig. 163, in Orch. Rev. 17 (1909) 209, fig. 17; Ames in Philip. Journ. Sci. 4 (1909) Bot. 673; Fedde Repert. 8 (1910) 382; Rolfe in Bot. Mag. 136 (1910) t. 8351; Ames Orch. 5 (1915) 138.

Luzon (Bontoc, Lepanto, Benguet), *Vanoverbergh* 560, 1263, 3112, *B. S.* 5614 Worcester, 3497 Mearns, *Lyon* 114. Epiphyte at 1,000 to 1,650 m altitude.

DENDROBIUM SCHUETZEI Rolfe in Orch. Rev. 19 (1911) 224, 20 (1912) 289, 308, 337, fig. 47, in Gard. Chron. III 50 (1911) 42, III 52 (1912) 229, fig. 102; G. Wilson in Orch. World 3 (1912) 19, t.; Rolfe in Bot. Mag. 139 (1913) t. 8495; Ames Orch. 5 (1915) 138, 7 (1922) 96.

Mindanao (Surigao), *Lyon s. n.*

DENDROBIUM SCOPA Lindl. in Bot. Reg. 28 (1842) Misc. 55; Reichb. f. in Walp. Ann. 6 (1861) 303; Naves Novis. App. (1882) 234; Ames in Philip. Journ. Sci. 6 (1911) Bot. 52, 9 (1914) Bot. 13, Orch. 5 (1915) 139.

Dendrobium thysanochilum Schltr. in K. Schum. & Lauterb. Nachtr. Fl. Deutsch. Schutzgeb. (1905) 152, fide Kränzl.

Desmotrichum scopa Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 349.

Luzon (Tarlac, Rizal, Laguna), Mindoro, Leyte, Mindanao (Agusan), *Loddiges* 273, *Lyon s. n.*, 110, *Reillo* 53, *Merrill* 9167, *Wenzel* 064, *Weber*

163, *Ramos s. n.* Epiphyte at sea level to 350 m altitude. In Guam and also probably in Formosa, Celebes (fide Kränzlin), New Guinea, and Samoa.

DENDROBIUM SECUNDUM Wall. Cat. (1828) No. 1996, *nomen*; Lindl. in Bot. Reg. 15 (1829) t. 1291, 30 (1844) Misc. 65, Gen. & Sp. Orch. (1830) 81; Hook. in Bot. Mag. 74 (1848) t. 4352; De Vriese Ill. Orch. Ind. Neerl. (1854) t. 6; Miq. Fl. Ind. Bat. 3 (1859) 641; Reichb. f. in Walp. Ann. 6 (1861) 296; Miq. Sumatra (1862) 274; Naves Novis. App. (1882) 233; Veitch Man. Orch. Pl. pt. 3 (1888) 73; Hook. f. Fl. Brit. Ind. 5 (1890) 730; Stein Orchideenb. (1892) 207; Grant Orch. Burma (1895) 120; Ridl. in Journ. Linn. Soc. 31 (1896) 271, 32 (1896) 259; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 358, Fig.-Atlas (1910) fig. 272; Kränzlin in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 98; Ames Orch. 5 (1915) 139, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 165.

Pedilonum secundum Blume Bijdr. (1825) 322:

Dendrobium bursigerum Lindl. in Journ. Linn. Soc. 3 (1859) 17; Miq. Fl. Ind. Bat. 3 (1859) 642; Edit. in l'Orchidoph. 2 (1882) 421; Reichb. f. in Gard. Chron. II 17 (1882) 424; Naves Novis. App. (1882) 233; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Ames Orch. 2 (1908) 173, 5 (1915) 117; Kränzlin in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 119, fig. 10 G-O; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 157.

Callista bursigera Ktze. Rev. Gen. Pl. 2 (1891) 654.

Callista secunda Ktze. Rev. Gen. Pl. 2 (1891) 653.

Dendrobium secundum Wall. var. *bursigerum* Ridl. in Journ. Linn. Soc. 31 (1896) 271.

Luzon (Bataan, Rizal, Camarines), Mindanao (Lanao, Agusan), *R. S. Williams* 765, *B. S.* 3097, 22155 *Ramos*, *Weber* 4, 144, *F. B.* 12287 *Curran*, *Merrill* 9821 (cult.). Epiphyte at 15 to 230 m altitude. In Burma, Malay Peninsula, Sumatra, Java, and Borneo.

DENDROBIUM SINUOSUM Ames Orch. 7 (1922) 96.

Leyte, *Wenzel* 0996. Tacloban, Tigbao, at sea level.

DENDROBIUM SPURIUM (Blume) J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 343, Fig.-Atlas (1910) fig. 260; Ames Orch. 2 (1908) 185, *text cut*, 5 (1915) 139; Kränzlin in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 264, figs. 34 A, B, 35 E-H; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 165. *Dendrocolla spuria* Blume Bijdr. (1825) 290; Miq. Fl. Ind. Bat. 3 (1859) 697.

Aërides spurium Lindl. Gen. & Sp. Orch. (1833) 241.

Dendrobium unguiculatum Teijsm. & Binn. in Tijdschr. Ned. Ind. 5 (1853) 491; Miq. Fl. Ind. Bat. 3 (1859) 634; Reichb. f. in Walp. Ann. 6 (1861) 303; Naves Novis. App. (1882) 231.

Dendrobium euphlebiium Reichb. f. ex Lindl. in Journ. Linn. Soc. 3 (1859) 7; Miq. Fl. Ind. Bat. 3 (1859) 633; Reichb. f. Xen. Orch. 2 (1862) 26, t. 110, II 6; Naves Novis. App. (1882) 232;

Ridl. in Journ. Linn. Soc. 31 (1896) 268, 32 (1896) 258, Mat. Fl. Mal. Penin. 1 (1907) 49.

Sarcochilus spurius Reichb. f. in Walp. Ann. 6 (1863) 500.

Thrixspermum spurium Reichb. f. Xen. Orch. 2 (1867) 122.

Callista euphlebia Kzte. Rev. Gen. Pl. 2 (1891) 654.

Callista unguiculata Ktze. Rev. Gen. Pl. 2 (1891) 655.

Luzon (Cagayan, Bataan, Sorsogon), *F. B. 16727 Curran, Merrill 3692, Whitford 1118, Lyon 43, Weber 31, B. S. 23427 Ramos, Elmer 16384, 17311*. Epiphyte at 100 to 930 m altitude. In Malay Peninsula, Java, and Borneo.

DENDROBIUM STELLA-SILVAE (Loher & Kränzl.) Ames Orch. 5 (1915) 139.

Sarcopodium stella-silvae Loher & Kränzl. in Fedde Repert. 7 (1909) 40; Rolfe in Orch. Rev. 18 (August, 1910) 240; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 326, fig. 31 A-C; Ames in Philip. Journ. Sci. 8 (1913) Bot. 426.

Luzon (Abra, Bontoc, Rizal), *B. S. 7134 Ramos s. n., Vanoverbergh 2418, 2868, Loher s. n.* On forest slope at 1,000 m altitude.

DENDROBIUM TAURINUM Lindl. in Bot. Reg. 29 (1843) t. 28, 30 (1844) Misc. 62; Paxt. Mag. Bot. 10 (1843) 143, 217, t. & text cut; G. K. in Bot. Zeit. 1 (1843) 511; Reichb. f. in Walp. Ann. 6 (1861) 298; Van Houtte in Fl. des Serres 18 (1870) 133, t. 1904; Anderson in Gard. Chron. (1872) 974; Blanco Fl. Filip. ed. 3, Laminas (1877-83) t. 429; Naves Novis. App. (1882) 234; Hemsl. in Gard. Chron. II 17 (1882) 776; Veitch Man. Orch. Pl. pt. 3 (1888) 78; Du Buyss. in l'Orchidoph. 12 (1892) 37; Stein Orchideenb. (1892) 210; L. Linden in Lindenia 13 (1898) 93, t. 621; Ames Orch. 1 (1905) 90, 2 (1908) 187, 5 (1915) 139; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 145, fig. 10A.

? *Dendrobium veratrifolium* Naves Novis. App. (1882) 233, non Lindl.

Callista taurina Ktze. Rev. Gen. Pl. 2 (1891) 655.

Luzon (Rizal, Laguna), Guimaras, Mindanao (Davao), *Lyon s. n., 24, B. S. 2018, 21333, 22245 Ramos, Reillo s. n., 41, Foxworthy s. n., Loher 480, Everett 4206, Black s. n., 3, Copeland 505*.

DENDROBIUM TERNATENSE J. J. Sm. in Bull. Dept. Agr. Ind. Néerl. 22 (1909) 26; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 246, fig. 20 A-C; Ames Orch. 5 (1915) 139.

Dendrobium macrophyllum Ames Orch. 2 (1908) 179, non A. Rich.; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 244, quoad Philip.

Luzon (Bataan), *Whitford 1083*. Mt. Mariveles, Lamao River. Apparently rare in the Philippines. Epiphyte at 900 m altitude. In Ternate Island.

DENDROBIUM TOPAZIACUM Ames Orch. 5 (1915) 140.

Leyte, *Wenzel 0202*. Dagami, epiphyte at 60 m altitude.

DENDROBIUM UNICORNE Ames Orch. 5 (1915) 140.

Luzon (Laguna), Leyte, *B. S. 14987 Ramos, Wenzel 0292, 0527, 0563, 0872*. Epiphyte at 500 to 700 m altitude in Leyte.

DENDROBIUM UNIFLORUM Griff. Notul. 3 (1851) 305, Icon. Pl. Asiat. 3 (1851) t. 303; Ridl. in Trans. Linn. Soc. II 3 (1893) 363, in Journ. Linn. Soc. 31 (1896) 271, 32 (1896) 256, Mat. Fl. Mal. Penin. 1 (1907) 46; Ames Orch. 1 (1908) 91; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 185, quoad bibliog. et spec. (excl. descr.); Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 166.

Dendrobium revolutum Hook. f. Fl. Brit. Ind. 5 (1890) 732, quoad syn. *D. uniflorum* Griff. et spec. Malacca; Grant Orch. Burma (1895) 118, quoad syn. *D. uniflorum* Griff. et spec. Malacca; Ames Orch. 2 (1908) 184, quoad syn. *D. uniflorum* Griff. et spec., 5 (1915) 138; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 185, quoad descr. (excl. bibliog. et spec.), non Lindl.

Luzon (Ilocos Norte, Bataan, Laguna, Sorsogon), Sibuyan, Negros (Occidental Negros), Leyte, Dinagat, Mindanao (Lanao, Agusan, Surigao). Fairly common and of wide distribution in the Philippines. Epiphyte at 60 to 700 m altitude. In Malay Peninsula and Borneo (Sarawak).

Further study of the Philippine material has proved that the species concerned is *D. uniflorum* Griff. and not its near ally *D. revolutum* Lindl. with which it has been much confused.

DENDROBIUM USTERI Schltr. in Usteri Viert. Naturf. Gesell. Zurich 56 (1906) 450, in Bull. Herb. Boiss. II 6 (1906) 458; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 300; Ames Orch. 5 (1915) 141.

Negros, Guimaras, *Usteri s. n., Lyon 137*.

DENDROBIUM USTERIODES Ames Orch. 5 (1915) 142.

Luzon, Mindoro, *Lyon 22, F. B. 3693 Merritt 75*.

DENDROBIUM VENTRICOSUM Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 209; Ames in Philip. Journ. Sci. 7 (1912) Bot. 19, Orch. 5 (1915) 144.

Luzon (Zambales, Sorsogon), Mindanao (Agusan, Surigao, Davao), *Lohr s. n., 6004, Elmer 14441, 14779, 16423, Weber 119, 176, 179, Wenzel 01156, Black s. n.* Epiphyte at low altitudes in Mindanao.

DENDROBIUM VERRUCULOSUM Ames in Philip. Journ. Sci. 8 (1913) Bot. 426, Orch. 5 (1915) 144.

Luzon (Ilocos Norte), Leyte, *Lyon s. n., 143, B. S. 33236 Ramos, Wenzel 0140*. Epiphyte.

DENDROBIUM VICTORIAE REGINAE Loher in Gard. Chron. III 21 (1897) 399, 22 (1897) 121, *fig. 34*; Cogn. Dict. Icon. Orch. (1899) Dendrob. t. 21; Ames Orch. 1 (1905) 92, 5 (1915) 144; Kränzl. in Engl. Pflanzenreich IV. 50 II. B. 21 (1910) 109, *fig. 6 A, B*; Rolfe in Orch. Rev. 20 (1912) 17, *fig. 6*.

Dendrobium coeleste Loher in Gard. Chron. III 22 (1897) 1; Ames Orch. 5 (1915) 120.

Dendrobium victoriae reginae Loher var. *exile* Ames in Philip. Journ. Sci. 2 (1907) Bot. 328.

Luzon (Benguet), Mindoro, Negros, Mindanao (Camiguin Island), *Elmer 6272, B. S. 4326 Mearns, 31797 Santos, 14429 Ramos, Lyon 122, Boettcher s. n., Merrill 5503, 7011, Loher 483*. Epiphyte at 1,300 to 2,650 m altitude.

DENDROBIUM WENZELII Ames Orch. 5 (1915) 144.

Leyte, Mindanao (Lanao, Surigao), *Wenzel 039, 0139, M. S. Clemens 43, Hutchinson 7551*. Epiphyte at 30 to 60 m altitude.

DENDROBIUM ZAMBOANGENSE Ames Orch. 5 (1915) 145.

Mindanao (Zamboanga, Agusan), *Merrill 8174, Elmer 13697*. Epiphyte at about 1,100 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

DENDROBIUM ACICULARE Lindl. apud Vidal Phan. Cuming. Philip. (1885) 148; Vidal Rev. Pl. Vasc. Filip. (1886) 267; Hook. f. Fl. Brit. Ind. 5 (1890) 725, quoad Philip.; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 241, quoad Philip.; Ames Orch. 5 (1915) 113 = *Dendrobium philippinense* Ames.

DENDROBIUM ACUMINATUM Lyon in Gard. Chron. III 42 (1907) 210, *fig. 88*; Rolfe in Orch. Rev. 15 (1907) 312, as to plant spoken of; Kränzl. in Orchis 2 (1908) 73, t. 16; Edit. in Gard. Chron. III 46 (1909) 150, *fig. 64*, in Gardners' Mag. 52 (1909) 659, 660, *text cut*, non Rolfe ex Ames = *Dendrobium lyonii* Ames.

DENDROBIUM ADUNCUM Wall. ex Lindl. apud Boxall ex Naves Novis. App. (1882) 234; Ceron Cat. Pl. Herb. Manila (1892) 166; Ames Orch. 5 (1915) 114, in nota.

This is a Himalayan species, and there is no sufficient reason for considering that it grows in the Philippines; cf. Ames l. c. for a critical note. It is not improbable that the plant reported as growing there may in reality be the closely related species *Dendrobium hercoglossum* Reichb. f.

DENDROBIUM ALAGENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 328 = *Dendrobium luzonense* Lindl.

DENDROBIUM ALBUM Boxall ex Naves Novis. App. (1882) 234, *nomen*; Ceron Cat. Pl. Herb. Manila (1892) 166, *nomen*.

Boxall probably intended *Dendrobium album* Wight, which is a synonym of *D. aqueum* Lindl., an Indian species.

DENDROBIUM ALOIFOLIUM Reichb. f. apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 206, quoad syn. *D. merrillii*; Ames Orch. 5 (1915) 115 (*D. aloefolium*) = *Dendrobium merrillii* Ames.

- DENDROBIUM ANGUSTIFOLIUM** Ames in Philip. Journ. Sci. 7 (1912) Bot. 14, non Lindl. = *Dendrobium chrysographatum* Ames.
- DENDROBIUM ANTENNATUM** Lindl. apud Naves Novis. App. (1882) 233.
- DENDROBIUM ATROPURPUREUM** Ames Orch. 1 (1905) 86, 2 (1908) 172; Kränz. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 217, quoad Philip., non Miq. = *Dendrobium cultratum* Schltr.
- DENDROBIUM AUREOROSEUM** Reichb. f. apud Kränz. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 125, p. p.; Ames Orch. 5 (1915) 115.
- This is a Javan species to which Kränzlin has referred a Philippine specimen collected by Cuming. J. J. Smith considers it to be a synonym of *Dendrobium nudum* Lindl.
- DENDROBIUM AUREUM** Lindl. Gen. & Sp. Orch. (1830) 77; Ames Orch. 5 (1915) 115 = *Dendrobium heterocarpum* Lindl.
- DENDROBIUM BIFARIUM** Lindl. apud Naves Novis. App. (1882) 233.
- DENDROBIUM BINNENDIJKII** Reichb. f. apud Naves Novis. App. (1882) 232.
- DENDROBIUM BLUMEI** Lindl. in Journ. Linn. Soc. 3 (1859) 14; Miq. Fl. Ind. Bat. 3 (1859) 638; Naves Novis. App. (1882) 232; Kränz. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 234, quoad syn. *D. planibulbe* Lindl. et quoad Philip. = *Dendrobium planibulbe* Lindl.
- DENDROBIUM BRONGNIARTII** Kränz. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 210; Ames in Philip. Journ. Sci. 7 (1912) Bot. 15, Orch. 5 (1915) 116 = *Dendrobium aporoides* (Lindl.) Merr.
- DENDROBIUM BURSIGERUM** Lindl. in Journ. Linn. Soc. 3 (1859) 17; Miq. Fl. Ind. Bat. 3 (1859) 642; Naves Novis. App. (1882) 233; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Ames Orch. 2 (1908) 173, 5 (1915) 117; Kränz. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 119, fig. 10 G-O = *Dendrobium secundum* Wall.
- DENDROBIUM CALCEOLUM** Roxb. apud Naves Novis. App. (1882) 231.
- DENDROBIUM CALOPHYLLUM** Reichb. f. apud Naves Novis. App. (1882) 232.
- DENDROBIUM CALOPOGON** Reichb. f. apud Naves Novis. App. (1882) 232.
- DENDROBIUM CANALICULATUM** R. Br. apud Naves Novis. App. (1882) 232.
- DENDROBIUM CANINUM** Merr. in Philip. Journ. Sci. 19 (1921) 344, non *Epidendrum caninum* Burm. f. = *Dendrobium crumenatum* Sw.
- DENDROBIUM CAPILLIPES** Reichb. f. apud Naves Novis. App. (1882) 233.
- DENDROBIUM CARNOSUM** Teijsm. & Binn. apud Naves Novis. App. (1882) 234.
- DENDROBIUM CERINUM** Reichb. f. in Gard. Chron. II 12 (1879) 554; Ridl. in Journ. Linn. Soc. 32 (1896) 263, in textu sub *D. sanguinolentum* Lindl.; Kränz. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 121; Ames Orch. 5 (1915) 119.

Reichenbach merely states that the type material came from the Malay Archipelago; Ridley, under *D. sanguinolentum* Lindl., doubts its being a

distinct species and states that Burbidge's specimen came from the Sulu Archipelago. No material comparable with this species has been seen from the Philippines; for discussion cf. Ames l. c.

DENDROBIUM COBBIANUM Reichb. f. in Gard. Chron. II 14 (1881) 748; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 297.

The origin of this plant is unknown; it was sent to Reichenbach without localization by the firm of Hugh Low & Co. Why it is included among the Philippine species is unknown, as there seems no ground upon which to assume that it came from the Philippines.

DENDROBIUM COELESTE Loher in Gard. Chron. III 22 (1897) 1; Ames Orch. 5 (1915) 120 = *Dendrobium victorinae reginae* Loher.

DENDROBIUM CONCINNUM Miq. apud Naves Novis. App. (1882) 234.

DENDROBIUM CONNATUM Lindl. apud Naves Novis. App. (1882) 231.

DENDROBIUM CONOSTALIX Reichb. f. in Walp. Ann. 6 (1861) 292; Usteri Beitr. Ken. Philip. Veg. (1905) 128; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 162; Ames Orch. 5 (1915) 120.

This is a Malayan species which has been reported from the Philippines. No material from the Philippines matches it and the reports seem erroneous. Cf. Ames l. c. for discussion.

DENDROBIUM CONVEXUM Lindl. apud Naves Novis. App. (1882) 231.

DENDROBIUM CREPIDATUM Lindl. apud Naves Novis. App. (1882) 232.

DENDROBIUM CUMULATUM Lindl. apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 115, quoad Philip.; Ames Orch. 5 (1915) 121.

The Philippine specimens referred here by Kränzl. represent *Dendrobium crumenatum* Sw. Cf. Ames l. c.

DENDROBIUM CYMBIDIODES Blume apud Naves Novis. App. (1882) 231 = ? *Dendrobium acuminatum* Rolfe ex Ames.

DENDROBIUM DAYANUM Boxall ex Naves Novis. App. (1882) 234, *nomen*; Ceron Cat. Pl. Herb. Manila (1892) 166, *nomen*.

DENDROBIUM DRACONIS Reichb. f. apud Naves Novis. App. (1882) 233.

DENDROBIUM EQUITANS Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 228, quoad Philip. = *Dendrobium aporoides* (Lindl.) Merr.

This species is known only from Formosa.

DENDROBIUM EROSUM Lindl. apud Naves Novis. App. (1882) 233.

DENDROBIUM ERYTHROXANTHUM Reichb. f. in Gard. Chron. II 2 (1874) 162; Boxall ex Naves Novis. App. (1882) 234 = *Dendrobium bullenianum* Reichb. f.

DENDROBIUM EUPHLEBIUM Reichb. f. ex Lindl. in Journ. Linn. Soc. 3 (1859) 7; Naves Novis. App. (1882) 232 = *Dendrobium spurium* (Blume) J. J. Sm.

DENDROBIUM FIMBRIATUM Hook. apud Naves Novis. App. (1882) 232.

DENDROBIUM FLABELLUM Reichb. f. apud Naves Novis. App. (1882) 232.

DENDROBIUM GEMELLUM Lindl. apud Naves Novis. App. (1882) 233.

DENDROBIUM GOLDSCHMIDTIANUM Kränzl. in Fedde Repert. 7 (1909) 40,
in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 116.

"Provinz der Philippinen und Formosa."—Kränzlin. The specimen probably originated in Formosa, not in the Philippines.

DENDROBIUM GOULDII Reichb. f. apud Naves Novis. App. (1882) 234.

DENDROBIUM GRACILE Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 230, quoad Philip.; Ames Orch. 5 (1915) 125.

Kränzlin, in his monograph of the genus, refers *Loher 496* to this species. What the Loher plant is is open to question, but no material referable to *D. gracile* has been seen in the collections received.

DENDROBIUM GRANDIFLORUM Lindl. apud Naves Novis. App. (1882) 232.

DENDROBIUM HAENKEANUM Steud. Nomencl. Bot. ed. 2, 1 (1840) 490 =
Geodorum nutans (Presl) Ames.

DENDROBIUM HASSELTII Blume apud Naves Novis. App. (1882) 232.

DENDROBIUM HETEROIDEUM Blume apud Naves Novis. App. (1882) 231.

DENDROBIUM HYMENOPHYLLUM Lindl. apud Naves Novis. App. (1882) 233.

DENDROBIUM INDIVISUM Naves Novis. App. (1882) 231; Vidal Phan.
Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267;
Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 217, quoad
Philip.; Ames in Philip. Journ. Sci. 6 (1911) Bot. 52, non Miq. =
Dendrobium distichum (Presl) Reichb. f.

DENDROBIUM INTERMEDIUM Teijsm. & Binn. apud Naves Novis. App. (1882) 232.

DENDROBIUM JAVANICUM Lindl. apud Naves Novis. App. (1882) 234.

DENDROBIUM JOHANNIS Reichb. f. apud Naves Novis. App. (1882) 233.

DENDROBIUM KÜHLII Lindl. apud Naves Novis. App. (1882) 233.

DENDROBIUM LABUANUM Lindl. apud Naves Novis. App. (1882) 231.

DENDROBIUM LAMELLATUM Lindl. apud Kränzl. in Engl. Pflanzenreich IV.
50. II. B. 21 (1910) 269, quoad Philip.; Ames Orch. 5 (1915) ix.

Cuming 2074, referred here by Kränzlin, is probably *Dendrobium platycaulon* Rolfe; cf. Ames l. c.

DENDROBIUM LILACINUM Reichb. f. apud Naves Novis. App. (1882) 233.

DENDROBIUM LOBBII Lindl. apud Naves Novis. App. (1882) 231.

DENDROBIUM LONGICOLLE Lindl. in Bot. Reg. 26 (1840) Misc. 74; Ames
Orch. 5 (1915) 127.

Diplocaulobium longicolle Kränzl. in Engl. Pflanzenreich IV. 50. II.
B. 21 (1910) 340.

The type was a specimen sent to Loddiges by Cuming from Singapore, where the species still occurs. There is no reason to consider it as Philippine.

- DENDROBIUM LUCENS Reichb. f. apud Naves Novis. App. (1882) 233.
- DENDROBIUM MACCARTHIAE Thw. apud Naves Novis. App. (1882) 233.
- DENDROBIUM MACRAEI Ames Orch. 1 (1905) 89, 2 (1908) 176, non Lindl. = *Dendrobium plicatile* Lindl.
- DENDROBIUM MACRANTHUM Hook. in Bot. Mag. 69 (1842) t. 3970; Naves Novis. App. (1882) 233; Ceron Cat. Pl. Herb. Manila (1892) 166, non A. Rich. = *Dendrobium anosmum* Lindl.
- DENDROBIUM MACROPHYLLUM Ames Orch. 2 (1908) 179, non A. Rich.; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 244, quoad Philip. = *Dendrobium ternatense* J. J. Sm.
- DENDROBIUM MACROPHYLLUM Lindl. in Bot. Reg. 25 (1839) Misc. 36, 28 (1842) Misc. 84, 30 (1844) Misc. 47, Sert. Orch. (1840) t. 35; Paxt. Mag. Bot. 8 (1841) 97, t.; Hook. Cent. Orch. Pl. (1851) t. 12; Planch. Hort. Donat. (1855) 156, Atlas t. 1; Reichb. f. in Linden Pescat. 1 (1860) 40, t. 40, non A. Rich. = *Dendrobium anosmum* Lindl.
- DENDROBIUM MANILLENSE Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 430 = *Appendicula cornuta* Blume.
- DENDROBIUM METACHILINUM Reichb. f. in Bonpl. 3 (1855) 222, in Walp. Ann. 6 (1861) 290; Naves Novis. App. (1882) 234; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Ames Orch. 5 (1915) 128.
- The type of this species is *Cuming 2057*, which was unquestionably from the Malay Peninsula, not from the Philippines. The species is known only from the Malay Peninsula. Cf. Ames l. c.
- DENDROBIUM MINAX Reichb. f. apud Naves Novis. App. (1882) 233.
- DENDROBIUM MONILIFORME Sw. apud Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100; Naves Novis. App. (1882) 233.
- DENDROBIUM MONOPHYLLUM F. v. Muell. apud Naves Novis. App. (1882) 232.
- DENDROBIUM NUDUM Lindl. apud Naves Novis. App. (1882) 232.
- DENDROBIUM NUTANS Presl Rel. Haenk. 1 (1827) 102; Presl Symb. Bot. 1 (1832) 34, t. 23; Naves Novis. App. (1882) 234 = *Geodorum nutans* (Presl) Ames.
- DENDROBIUM PARCIFLORUM Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 222, quoad Philip.; Ames in Philip. Journ. Sci. 7 (1912) Bot. 18, non Reichb. f. = *Dendrobium marivelense* Ames.
- DENDROBIUM PARCUM Reichb. f. apud Naves Novis. App. (1882) 233.
- DENDROBIUM PAUCILACINIATUM J. J. Sm. apud Prain in Ind. Kew. Suppl. 3 (1908) 58.

Index Kewensis cites "Ins. Philip." as the locality; this is an error as the type was from Ternate.

DENDROBIUM PIERARDII Roxb. apud Naves Novis. App. (1882) 232.

DENDROBIUM PSEUDACLINIA Lindl. in Journ. Linn. Soc. 3 (1859) 9; Ames Orch. 5 (1915) 134 = *Dendrobium acclinia* Reichb. f.

DENDROBIUM PUGIONIFORME A. Cunn. apud Naves Novis. App. (1882) 232.

DENDROBIUM PUMILUM Reichb. f. apud Vidal Phan. Cuming. Philip. (1885) 148.

Probably *Dendrobium pumilum* Roxb. was intended, as Reichenbach apparently described no such species. Vidal referred here *Cuming 2102* as Philippine; the specimen in all probability came from the Malay Peninsula where the species is seemingly not uncommon.

DENDROBIUM PURPUREUM Blume apud Naves Novis. App. (1880) 233.

DENDROBIUM PUSILLUM Lindl. apud Naves Novis. App. (1882) 231.

DENDROBIUM RADIANIS Reichb. f. apud Naves Novis. App. (1882) 233.

DENDROBIUM RAMOSUM Lindl. apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 53, quoad Philip.

This is an Indian species, Kränzlin giving its other range as "Philippinen (?)." There is no reason for considering it as a Philippine species.

DENDROBIUM RETUSUM Llanos in Mem. Real. Acad. Cienc. Madr. 4 (1857) 498; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100 = ? *Dendrobium anosmum* Lindl.

DENDROBIUM REVOLUTUM Hook. f. Fl. Brit. Ind. 5 (1890) 732, quoad syn. *D. uniflorum* Griff. et spec. Malacca; Grant Orch. Burma (1895) 118, quoad syn. *D. uniflorum* Griff. et spec. Malacca; Ames Orch. 2 (1908) 184, quoad syn. *D. uniflorum* Griff. et spec., 5 (1915) 138; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 185, quoad descr. (excl. bibliog. et spec.), non Lindl. = *Dendrobium uniflorum* Griff.

DENDROBIUM RHOMBEUM Lindl. in Bot. Reg. 29 (1843) t. 17, 30 (1844) Misc. 56; Reichb. f. in Bot. Zeit. 1 (1843) 333; Lindl. in Paxt. Flow. Gard. 3 (1852-53) 52; Naves Novis. App. (1882) 233 = *Dendrobium heterocarpum* Wall.

DENDROBIUM ROBINSONII Ames in Philip. Journ. Sci. 8 (1913) Bot. 425, Orch. 5 (1915) 138 = *Dendrobium carinatum* (Linn.) Willd.

DENDROBIUM RUCKERI Lindl. in Bot. Reg. 29 (1843) Misc. 25, t. 60, 30 (1844) Misc. 52; Reichb. f. in Walp. Ann. 6 (1861) 288; Naves Novis. App. (1882) 234; Ames Orch. 5 (1915) 138.

When first described this was thought to be of Philippine origin. R. A. Rolfe, in Orch. Rev. 4 (1896) 87, has shown it to be an Indian species. Cf. Ames l. c.

DENDROBIUM SCABRILINGUE Lindl. apud Naves Novis. App. (1882) 232.

DENDROBIUM SCULPTUM Reichb. f. apud Naves Novis. App. (1882) 233.

DENDROBIUM SERRA Lindl. apud Naves Novis. App. (1882) 231.

DENDROBIUM SINUATUM Schltr. in Perk. Frag. Fl. Philip. (1904) 43; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 216, quoad Philip., non Lindl. = *Dendrobium cultratum* Schltr.

- DENDROBIUM STRIATUM Hort. ex Gard. Chron. III 38 (1905) 382; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 298 = *Dendrobium o'brienianum* Kränzl.
- DENDROBIUM STRIOLATUM Reichb. f. apud Naves Novis. App. (1882) 232.
- DENDROBIUM SUBTERES Lindl. apud Naves Novis. App. (1882) 231.
- DENDROBIUM SUBULATUM Blume apud Naves Novis. App. (1882) 232.
- DENDROBIUM SUPERBUM Reichb. f. in Walp. Ann. 6 (1861) 282; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 267; Ames Orch. 1 (1905) 90, 2 (1908) 186, 5 (1915) 139 = *Dendrobium anosmum* Lindl.
- DENDROBIUM TENELLUM Blume apud Naves Novis. App. (1882) 231.
- DENDROBIUM TERES Blanco Fl. Filip. (1837) 638, non Roxb. = *Luisia teretifolia* Gaudich.
- DENDROBIUM TERETIFOLIUM Blanco Fl. Filip. ed. 2 (1845) 444, ed. 3, 3 (1879) 34, non R. Br. = *Luisia teretifolia* Gaudich.
- DENDROBIUM TERETIFOLIUM R. Br. apud Naves Novis. App. (1882) 232.
- DENDROBIUM TETRAEDRE Blume apud Naves Novis. App. (1882) 232.
- DENDROBIUM TETRAGONUM A. Cunn. apud Naves Novis. App. (1882) 232.
- DENDROBIUM TMESIPTERIS Lindl. apud Naves Novis. App. (1882) 232.
- DENDROBIUM TUADENUM Boxall ex Naves Novis. App. (1882) 234, *nomen.*
- DENDROBIUM UNGUICULATUM Teijsm. & Binn. in Tijdschr. Ned. Ind. 5 (1853) 941; Naves Novis. App. (1882) 231 = *Dendrobium spurium* (Blume) J. J. Sm.
- DENDROBIUM VANOVERBERGHII Ames in Philip. Journ. Sci. 8 (1913) Bot. 425, t. 13 = *Dendrobium papilio* Loher.
- DENDROBIUM VEITCHIANUM Lindl. apud Naves Novis. App. (1882) 231.
- DENDROBIUM VERATRIFOLIUM Naves Novis. App. (1882) 233, non Lindl. = ? *Dendrobium taurinum* Lindl.
- DENDROBIUM XANTHOLEUCUM Reichb. f. apud Naves Novis. App. (1882) 231.

65. ERIA Lindley

ERIA ALBOLUTEA Rolfe in Kew Bull. (1917) 80.

Described from cultivated specimens originating in the Philippines.

ERIA ANCEPS Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 219, 237, fig. 18; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 129; Ames Orch. 5 (1915) 146.

Negros (Occidental Negros), Palawan, *Whitford 1601*, F. B. 7275 *Everett*, B. S. 636 *Foxworthy*.

ERIA BARBIFRONS Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 52.

"Provinz der Philippinen und Süd-Formosa: Philippinen, vermuthlich Manila, von A. Loher gesandt." Known only from the description.

ERIA BINABAYENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 329; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 223, 242; Ames Orch. 5 (1915) 146.

Trichotosia binabayensis Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21². (1911) 141.

Luzon (Laguna), Mindoro, B. S. 16554 Ramos, Merrill 5661. Epiphyte in humid forest at 215 m altitude in Mindoro.

ERIA BONTOCENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 427, Orch. 5 (1915) 146.

Luzon (Bontoc), Vanoverbergh 1386.

ERIA BRACHYSTACHYA Reichb. f. in Bonpl. 3 (1855) 223; Lindl. in Journ. Linn. Soc. 3 (1859) 59; Miq. Fl. Ind. Bat. 3 (1859) 657; Reichb. f. in Walp. Ann. 6 (1861) 276; Naves Novis. App. (1882) 236; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 221, 241; Ames Orch. 2 (1908) 188, 5 (1915) 146; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 100.

Pinalia brachystachya Ktze. Rev. Gen. Pl. 2 (1891) 678.

Eria wenzelii Ames in Philip. Journ. Sci. 8 (1913) Bot. 428.

Babuyan Islands (Camiguin Island), Luzon (Kalinga, Rizal, Camarines, Sorsogon), Leyte, Mindanao (Agusan). A fairly common species of wide distribution. Epiphyte at sea level to 1,000 m altitude.

ERIA BRACDESCENS Lindl. in Bot. Reg. 27 (1841) Misc. 18, 30 (1844) t. 29; Paxt. Mag. Bot. 8 (1841) 118, 11 (1844) 116; Edit. in Gard. Chron. (1841) 231; Reichb. f. in Walp. Ann. 6 (1861) 277; Hemsl. in Gard. Chron. II 18 (1882) 468; Hook. f. Fl. Brit. Ind. 5 (1890) 796; Ridl. in Trans. Linn. Soc. II 3 (1893) 366; Grant Orch. Burma (1895) 135; Ridl. in Journ. Linn. Soc. 32 (1896) 293; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 120, t. 166; Williams in Bull. Herb. Boiss. II 4 (1904) 368; J. J. Sm. Orch. Amb. (1905) 74; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 101; Ames Orch. 2 (1908) 189, 5 (1915) 147; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 209, 231, fig. 11; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 88; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 168.

Eria dillwynii Hook. in Bot. Mag. 71 (1845) t. 4163; Lindl. in Bot. Reg. 31 (1845) Misc. 63; F. Kl. in Bot. Zeit. 3 (1845) 550; Paxt. Mag. Bot. 12 (1846) 140; Lindl. in Journ. Linn. Soc. 3 (1859) 55; Naves Novis. App. (1882) 236; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 87.

Eria litoralis Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 24 (1862) 312.

Pinalia bractescens Ktze. Rev. Gen. Pl. 2 (1891) 679.

Luzon (Bataan, Rizal), Basilan, Samar, Mindanao (Zamboanga), Lyon 123, 150, F. B. 4017, 6562 Hutchinson, Whitford 1082, B. S. 3000, 6562, 17641 Ramos. Epiphyte, apparently at low altitudes.

ERIA CANDOONENSIS Ames Orch. 7 (1922) 101.

Mindanao (Bukidnon), *B. S. 38941 Ramos & Edaño*. Mt. Candoon, epiphyte at 1,650 m altitude.

ERIA CARNICOLOR Ames Orch. 7 (1922) 152.

Luzon (Bontoc), *B. S. 37937 Ramos & Edaño*. Mt. Masapilid, epiphyte in mossy forest at 1,600 m altitude.

ERIA CLEMENSIAE Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 220, 238, fig. 20; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 126; Ames Orch. 5 (1915) 147.

Luzon (Rizal), Panay (Antique), Leyte, Mindanao (Lanao, Bukidnon), *Reillo s. n.*, *B. S. 32192 McGregor, 15390 Ramos, 38891 Ramos & Edaño, Wenzel 0393, 0495, M. S. Clemens 602*. Epiphyte in forest at 500 to 1,000 m altitude.

ERIA COCHLEATA Lindl. in Bot. Reg. 30 (1844) Misc. 14; Reichb. f. in Walp. Ann. 6 (1861) 269; Naves Novis. App. (1882) 236.

Imported from the Philippines by Loddiges. Known only from the original description in which Lindley states that it has the appearance of *E. stellata* which is a synonym of *E. javanica* (Sw.) Blume. Kränzl. does not treat *E. cochleata* in his monograph of the genus in Das Pflanzenreich.

ERIA COMPACTA Ames in Philip. Journ. Sci. 2 (1907) Bot. 329; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 208, 231; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 77, fig. 15 A-D; Ames Orch. 5 (1915) 147.

Mindoro, *Merritt 4420*. Mt. Halcon, at 1,415 m altitude.

ERIA COPELANDII Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 203, 226, fig. 3; Ames Orch. 5 (1915) 147; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 217 in textu.

Eria formosana Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 47, non Rolfe.

Mindanao (Misamis, Bukidnon, Davao), *F. B. 4596, 4600 Mearns & Hutchinson, B. S. 38524 Ramos & Edaño, Copeland 1423*. Epiphyte at 1,560 to 3,060 m altitude.

Var. FUSIFORMIS Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 204, 227.

Eria formosana Kränzl. var. *fusiformis* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 47.

Luzon (Lepanto, Benguet), *Merrill 4590, Ramos & Edaño 85, 106, 111, F. B. 5067 Curran*. Epiphyte at 2,100 to 2,530 m altitude.

ERIA CURRANII Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 210, 233, fig. 13; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 79; Ames Orch. 5 (1915) 147.

Palawan, *F. B. 3904 Curran*.

ERIA CYLINDROSTACHYA Ames Orch. 2 (1908) 190, text cut, 5 (1915) 147; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 211, 234, fig. 15; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 69, fig. 51 L, M.

Luzon (Benguet, Rizal), Mindanao (Misamis, Bukidnon, Davao), *Boettcher s. n.*, B. S. 29371, 38648, 39079 Ramos & Edaña, F. B. 4612 Mearns & Hutchinson, Copeland 1424, Elmer 10632, 10632a. Epiphyte at 100 to 1,560 m altitude.

ERIA CYMBIFORMIS J. J. Sm. in Rec. Trav. Bot. Néerl. 1 (1904) 152, *text cut*; Ames in Philip. Journ. Sci. 2 (1907) Bot. 330; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 224, 243, *fig. 26*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 104; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 40; Ames Orch. 5 (1915) 147.

Luzon (Cagayan, Bataan, Rizal, Laguna, Tayabas, Mindoro, Negros (Oriental Negros), Leyte, Mindanao (Davao). A fairly common and widely distributed species. Epiphyte at 500 to 1,015 m altitude, in Sumatra.

ERIA DAGAMENSIS Ames Orch. 5 (1915) 147.

Luzon (Ifugao), Leyte, *Sandkuhl 296*, *Wenzel 0123, 0305, 0308, 0314, 0315, 0317, 0318, 0320, 0321, 0322, 0323, 0345, 0346*, B. S. 42087 Edaña Epiphyte in forest at 500 to 600 m altitude in Leyte.

ERIA DAVAENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1575, Orch. 5 (1915) 149.

Mindanao (Davao), *Elmer 11321*. Todaya, Mt. Apo; epiphyte in humid forest at 1,300 m altitude.

ERIA FASTIGATIFOLIA Ames Orch. 2 (1908) 191, *text cut*, 5 (1915) 149; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 221, 240, *fig. 22*. *Trichotosia fastigiatifolia* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 162.

Luzon (Bontoc, Ifugao, Benguet, Rizal, Laguna, Tayabas, Sorsogon), Negros (Oriental Negros), Leyte. A fairly common species, especially in Luzon. Epiphyte at 300 to 1,300 m altitude.

ERIA FLORIBUNDA Lindl. in Wall. Cat. (1832) No. 7408, *nomen*, in Bot. Reg. 29 (1843) Misc. 43, 30 (1844) t. 20; Paxt. Mag. Bot. 11 (1844) 91; Lindl. in Journ. Hort. Soc. 3 (1848) 15, in Journ. Linn. Soc. 3 (1859) 60; Miq. Fl. Ind. Bat. 3 (1859) 661; Reichb. f. in Walp. Ann. 6 (1861) 276; Hemsl. in Gard. Chron. II 18 (1882) 468; Hook. f. Fl. Brit. Ind. 5 (1890) 792; Ridl. in Trans. Linn. Soc. II 3 (1893) 366; Grant Orch. Burma (1895) 140; Ridl. in Journ. Linn. Soc. 32 (1896) 290; Williams in Bull. Herb. Boiss. II 4 (1904) 368; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 400, Fig.-Atlas (1911) *fig. 303*; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 96; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 209, 231, *fig. 10* (excl. syn. *E. densa* Ridl.); Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 120, *fig. 25 A-D*; Winkler in Engl. Bot. Jahrb. 48 (1912) 101; Ames Orch. 5 (1915) 149, 6 (1920) 124, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 170.

Eria leucostachya Lindl. in Journ. Hort. Soc. 3 (1848) 15.

Eria floribunda Lindl. var. *leucostachya* Lindl. in Paxt. Flow. Gard. 3 (1853) 116, *fig. 288*.

Pinalia floribunda Ktze. Rev. Gen. Pl. 2 (1891) 678.

Palawan, Mindanao (Cotabato, Agusan, Bukidnon), *B. S.* 550, 633 *Foxworthy*, 38911, 39014, 39107 *Ramos & Edaña*, *Merrill* 9617, *Weber* 152, 228. Epiphyte at 15 to 1,260 m altitude. In Burma, Malay Peninsula, Sumatra, Java, and Borneo.

ERIA FUSCA Blume Mus. Bot. Lugd.-Bat. 2 (1856) 183; Miq. Fl. Ind. Bat. 3 (1859) 658; Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 388; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 222, 241; Ames Orch. 5 (1915) 149, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 170.

? *Eria hirta* Blume Mus. Bot. Lugd.-Bat. 2 (1856) 184; Miq. Fl. Ind. Bat. 3 (1859) 658, fide Kränzl.

Trichotosia ciliata Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 5 (1853) 492.

Trichotosia fusca Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 5 (1853) 492; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 145.

Eria ciliata Miq. Fl. Ind. Bat. 3 (1859) 659; Reichb. f. in Walp. Ann. 6 (1861) 273.

Pinalia ciliata Ktze. Rev. Gen. Pl. 2 (1891) 678.

Pinalia fusca Ktze. Rev. Gen. Pl. 2 (1891) 678.

? *Pinalia hirta* Ktze. Rev. Gen. Pl. 2 (1891) 678.

? *Trichotosia hirta* Blume in Herb. Mus. Lugd.-Bat. ex Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 145, in syn.

Negros (Occidental Negros), Leyte, *F. B.* 5582 *Everett*, *Copeland* 138, *Wenzel* 045, 0143, *B. S.* 40147 *Lopez*. Epiphyte in forest at 60 m altitude in Leyte. In Java and Borneo and, perhaps, Celebes.

ERIA GIGANTEA Ames Orch. 2 (1908) 192, *text cut*, 5 (1915) 149; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 218, 239, *fig. 21*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 130; Ames in Addisonia 6 (1921) 41, t. 213.

Luzon (Isabela, Bataan, Rizal, Laguna), Mindanao (Misamis, Lanao), *Lyon* 146, *Topping* 472, *Merrill* 4110, *B. S.* 3430 *Ramos*, 18995 *Brown*, *F. B.* 4606 *Mearns & Hutchinson*, *M. S. Clemens s. n.*, 437. Terrestrial (?) at 660 to 1,300 m altitude.

ERIA GRACILISCAPA Rolfe ex Ames Orch. 1 (1905) 93, 2 (1908) 193, 3 (1908) 88, t. 57, 5 (1915) 149; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 205, 228; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 108.

Luzon (Kalinga, Bataan, Tayabas), *Boettcher s. n.*, *Whitford* 307, *Topping* 474, *F. B.* 795 *Borden*, *Elmer* 7405. Epiphyte at 600 to 1,000 m altitude.

ERIA HALCONENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 330; Ames Orch. 3 (1908) 85, t. 56, 5 (1915) 149; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 224, 241.

Trichotosia halconensis Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 149.

Mindoro, Panay (Capiz), *Merrill 5510, 5742, B. S. 30716 Ramos & Edaño*. Terrestrial and epiphytic at 330 to 2,000 m altitude.

ERIA HIRSUTIPETALA Ames Orch. 7 (1922) 102.

Mindanao (Bukidnon), *B. S. 39124 Ramos & Edaño*. Mt. Candoon, epiphyte in mossy forest at 1,830 m altitude.

ERIA HUTCHINSONIANA Ames in Philip. Journ. Sci. 2 (1907) Bot. 330; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 204, 227, *fig. 4*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 80, *fig. 17 A-D*; Ames Orch. 5 (1915) 150.

Mindoro, *Merrill 5514*. Mt. Halcon, terrestrial on mossy ridge at 2,300 m altitude.

ERIA JARENSIS Ames Orch. 5 (1915) 150.

Luzon (Nueva Vizcaya, Laguna), Leyte, Mindanao (Surigao), *Wenzel 0356, 0357, 0358, 0365, 0377, Lyon 130, B. S. 20436 Ramos, 20090 McGregor*. Epiphyte in forest at 500 to 750 m altitude.

ERIA JAVANICA (Sw.) Blume Rumphia 2 (1836) 23; Ames Orch. 2 (1908) 193; J. J. Sm. in Nova Guinea 8 (1909) 86, 8³ (1912) 577, 12⁴ (1916) 352; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 26; Schltr. in Fedde Repert. 10 (1911) 91; J. J. Sm. Orch. Jav. Fig.-Atlas (1911) *fig. 287*; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 171.

Dendrobium javanicum Sw. in Kongl. Svensk. Vet.-Akad. Nya Handl. 21 (1800) 247; Willd. Sp. Pl. 4 (1805) 137.

Dendrolirium rugosum Blume Bijdr. (July, 1825) 345.

Eria stellata Lindl. in Bot. Reg. 11 (August, 1825) *t. 904*; Lindl. Gen. & Sp. Orch. (1830) 67; Paxt. Mag. Bot. 2 (1836) 137, 4 (1838) 234; Hook. in Bot. Mag. 64 (1837) *t. 3605*; Reichb. f. in Bonpl. 5 (1857) 54; Miq. Fl. Ind. Bat. 3 (1859) 661; Reichb. f. in Walp. Ann. 6 (1861) 269; Hemsl. in Gard. Chron. II 18 (1882) 500; Naves Novis. App. (1882) 236; Ridl. in Trans. Linn. Soc. II 3 (1893) 366, in Journ. Linn. Soc. 31 (1896) 285, 32 (1896) 304; Kränzl. in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 248; Usteri Beitr. Ken. Philip. Veg. (1905) 128; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 98.

Octomeria stellata Spreng. Syst. Veg. 4 Cur. Post (1827) 310.

Octomeria vaginata Breda Gen. et Sp. Orch. Jav. (1827) *t. 5*.

Eria rugosa Lindl. Gen. & Sp. Orch. (1830) 66; Miq. Fl. Ind. Bat. 3 (1859) 661; Hemsl. in Gard. Chron. II 18 (1882) 500; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 378, Orch. Amb. (1905) 74; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 212, 235; Ames Orch. 5 (1915) 158.

Eria fragrans Reichb. f. in Bot. Zeit. 22 (1864) 415, in Trans. Linn. Soc. 30 (1874) 137; Hook. f. Fl. Brit. Ind. 5 (1890) 794; Grant Orch. Burma (1895) 140; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 122, *t. 168*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 24.

Eria striolata Reichb. f. in Gard. Chron. III 3 (May 5, 1888) 554, in Ill. Hort. 35 (May, 1888) 35, *t. 48*; Ridl. in Journ. Linn. Soc.

- 31 (1896) 285, 36 (1896) 304, Mat. Fl. Mal. Penin. 1 (1907) 99; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 25.
Tainia stellata Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2 Abt. 6 (1888) 153.
Dendrobium perakense Hook. f. Fl. Brit. Ind. 5 (1890) 712, in Hook. Icon. Pl. 21 (1890) t. 2019.
Pinalia rugosa Ktze. Rev. Gen. Pl. 2 (1891) 679.
Pinalia fragrans Ktze. Rev. Gen. Pl. 2 (1891) 679.
Pinalia stellata Ktze. Rev. Gen. Pl. 2 (1891) 679.
Pinalia striolata Ktze. Rev. Gen. Pl. 2 (1891) 679.
Eria vaginata Benth. ex Jacks. Ind. Kew. (1893) 864.
Sarcopodium perakense Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 328.
Eria javanica Blume var. *philippinensis* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) fig. 14 E-J.

Luzon (Albay, Bontoc, Benguet, Rizal, Laguna), Leyte, Mindanao (Zamboanga, Lanao), *Loher* 6003, *Vanoverbergh* 3911, *Lyon* s. n., *B. S.* 15135 *Ramos*, 17905 *Brown*, *Wenzel* 0100, *F. B.* 12364 *Hutchinson*, *M. S. Clemens* s. n. Up to 1,200 m altitude in Bontoc. In India, Burma, Malay Peninsula, Sumatra, Celebes, Amboina, Borneo, and New Guinea.

It seems probable that *Eria nicholitzii* Kränzl. in Oesterr. Bot. Zeitschr. (1894) 459; Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1912) 653, *Eria pseudostellata* Schltr. in K. Schum. & Lauterb. Nachtr. Fl. Deutsch. Schutzgeb. (1905) 182, and *Eria papuana* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 25 are conspecific. Schlechter reduces his *E. pseudostellata* and *E. papuana* Kränzl. (to which Kränzl. reduces his *E. nicholitzii*) to *E. nicholitzii* Kränzl. J. J. Smith states that *E. pseudostellata* Schltr. is probably the same as *E. javanica* Blume, and Kränzl. in Das Pflanzenreich, cites it as a synonym of that species.

ERIA LAGUNENSIS Ames Orch. 5 (1915) 150.

Luzon (Laguna), *Serrato* s. n. Epiphyte in thicket forest.

ERIA LAMELLATA Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1576, Orch. 5 (1915) 152.

Biliran, Mindanao (Davao), *B. S.* 18942 *McGregor*, *Elmer* 11715. Epiphyte in humid forest at 660 to 1,300 m altitude.

ERIA LEAVITTII Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 64, fig. 17 G, H.

Eria ventricosa Leavitt var. *benguetensis* Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 212, 234.

Luzon (Benguet), *Merrill* 4849, *B. S.* 4986 *Curran*. Terrestrial and epiphytic in mossy forest up to 1,950 m altitude.

ERIA LEYTENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 427, Orch. 5 (1915) 152.

Leyte, Negros, *Wenzel* 017, 0169, 0674, *F. B.* 19072 *Curran*. Epiphyte at 60 to 600 m altitude.

ERIA LOHERIANA Kränzl. in Fedde Repert. 17 (1921) 387.

Loher s. n. Known only from the original description; no material has been seen.

ERIA LONGIBRACTEATA Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 219, 238, fig. 19; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 126; Ames Orch. 5 (1915) 152.

Luzon (Cagayan), Mindanao (Zamboanga, Misamis), *F. B. 16735 Curran, 4728 Mearns & Hutchinson, Merrill 8142*. Up to 1,900 m altitude in Misamis.

ERIA LONGICRURIS Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 205, 228, fig. 6; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 109; Ames Orch. 5 (1915) 152.

Mindanao (Misamis, Davao), *F. B. 4609 Mearns & Hutchinson, Elmer 11351*. At 2,800 m in Misamis.

ERIA LONGILABRIS Lindl. in Bot. Reg. 27 (1841) Misc. 38, 30 (1844) sub t. 29, fig. 3; Reichb. f. in Walp. Ann. 6 (1861) 277; Naves Novis. App. (1882) 236; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 210, 232; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 86; Ames Orch. 5 (1915) 152.

Panay, *Cuming s. n.* Known only from the description and the Cuming specimen.

As pointed out by Leavitt and by Ames this may be only a variety of *E. bractescens* Lindl., but for the present it is deemed best to maintain it as a separate species.

ERIA LYONII Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 212, 234, fig. 17; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 87; Ames Orch. 5 (1915) 152.

Luzon (Tarlac), *Lyon B, 116* in part (Luzon material). Terrestrial (?) near sea level.

ERIA MACERA Ames Orch. 7 (1922) 103.

Mindanao (Bukidnon), *B. S. 38512 Ramos & Edaño*. Epiphyte on mossy forest slope at 2,130 m altitude.

ERIA MAJOR Ridl. ex Stapf in Trans. Linn. Soc. II 4 (1894) 237, in syn.; Ridl. in Journ. Linn. Soc. 31 (1896) 280, 32 (1896) 288, Mat. Fl. Mal. Penin. 1 (1907) 90; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 218, 239; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 128; Ames Orch. 5 (1915) 155, 6 (1920) 131, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 172.

Eria kingii Hook. f. Fl. Brit. Ind. 5 (1890) 790, in Hook. Icon. Pl. 21 (1891) t. 2066, non F. v. Muell.

Eria scortechinii Stapf in Trans. Linn. Soc. II 4 (1894) 237, non Hook. f.

Mindanao (Misamis), *F. B. 4595 Mearns & Hutchinson*. Mt. Malindang at 1,750 m altitude.

ERIA MAQUILINGENSIS Ames Orch. 5 (1915) 152.

Luzon (Laguna), *B. S. 17909 Brown*. Mt. Maquiling, epiphyte in forest at 700 m altitude.

ERIA MCGREGORII Ames Orch. 5 (1915) 153.

Luzon (Ifugao), *B. S. 19876 McGregor*. Mt. Polis.

ERIA MEARNSII Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 216, 236; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 37; Ames Orch. 5 (1915) 155.

Mindanao (Misamis, Agusan, Davao), *F. B. 4729 Mearns & Hutchinson, Elmer 11850, 13802*. Epiphyte up to 1,700 m altitude in Misamis.

ERIA MERRILLII Ames in Philip. Journ. Sci. 2 (1907) Bot. 331; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 213, 235; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 56; Ames Orch. 5 (1915) 155.

Luzon (Rizal, Sorsogon), Mindoro, Leyte, Mindanao (Surigao), *Lyon 116 in part, 5626, Elmer 15552, Merrill 5519, Wenzel 0104, 0717*. Epiphyte at 60 to 800 m altitude.

ERIA MERRITTII Ames in Philip. Journ. Sci. 2 (1907) Bot. 331; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 207, 230; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 107; Ames Orch. 5 (1915) 155.

Luzon (Rizal, Laguna), Mindoro, Leyte, *B. S. 19205 Reillo, 15122 Ramos, F. B. 4537 Merritt*. Epiphyte at 500 to 1,300 m altitude.

ERIA MICROCHILA Ames Orch. 7 (1922) 104.

Luzon (Bontoc), *B. S. 37997 Ramos & Edaña*. Mt. Causal. Epiphyte in mossy forest at 1,530 m altitude.

ERIA MINDANAENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1577, Orch. 5 (1915) 155, 6 (1920) 207.

Sibuyan, Panay (Capiz), *Elmer 12271, B. S. 35379, 35633 Martelino & Edaña*. Epiphyte at 250 to 660 m altitude.

ERIA ODORIFERA Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 223, 242, fig. 24; Ames Orch. 5 (1915) 156.

Trichotosia odorifera Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 153.

Negros (Oriental Negros), Mindanao (Zamboanga, Davao), *Elmer 9521, 9658, Copeland 1182, Merrill 8153*. Epiphyte up to 1,600 m altitude in Davao.

ERIA ORNATA (Blume) Lindl. Gen. & Sp. Orch. (1830) 66; De Vriese Ill. Orch. Ind. Néerl. (1854) t.; Reichb. f. in Bonpl. 5 (1857) 54; Lindl. in Journ. Linn. Soc. 3 (1859) 48 (excl. Indian and Burman plants); Miq. Fl. Ind. Bat. 3 (1859) 660 (excl. Indian and Burman plants); Reichb. f. in Walp. Ann. 6 (1861) 270; Naves Novis. App. (1882) 236; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268; Ridl. in Journ. Linn. Soc. 31 (1896) 285, ? 32 (1896) 302; Schltr. in Perk. Frag. Fl. Philip. (1904) 43; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 380, Fig.-Atlas (1911) fig. 288; Ames Orch. 1 (1905) 94, 2 (1908) 195, 5 (1915)

156; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 217, 237; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 50, fig. 11 A, B; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 173.

Dendrolirium ornatum Blume Bijdr. (1825) 345.

Eria armeniaca Lindl. in Bot. Reg. 27 (1841) Misc. 38, t. 42.

Pinalia ornata Ktze. Rev. Gen. Pl. 2 (1891) 678.

Luzon (Bontoc, Benguet, Nueva Vizcaya, Zambales, Rizal), Negros (Occidental Negros), Mindanao (Zamboanga, Agusan). A fairly common species in the Philippines at 10 to 1,450 m altitude. In ? Siam, Sumatra, Java, and Borneo.

ERIA OVATA Lindl. in Bot. Reg. 30 (1844) sub t. 29; G. K. in Bot. Zeit. 3 (1845) 95; Reichb. f. in Walp. Ann. 6 (1861) 277; Naves Novis. App. (1882) 236; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268; Ames Orch. 2 (1908) 195, 5 (1915) 156; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 207, 230, fig. 8; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 76.

Eria elmeri Ames Orch. 1 (1905) 93, 2 (1908) 196, 5 (1915) 149; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 208, 230, fig. 10; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 74.

Luzon (Bontoc, Lepanto, Benguet, Bataan, Laguna), Negros (Oriental Negros), Leyte. A fairly common species, epiphyte at 500 to 2,550 m altitude.

ERIA PALAWANENSIS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1578, Orch. 5 (1915) 156.

Palawan, *Elmer 13209*. Mt. Pulgar.

ERIA PERSPICABILIS Ames Orch. 5 (1915) 156.

Luzon (Ifugao), *B. S. 19733 McGregor*. Mt. Polis.

ERIA PHILIPPINENSIS Ames Orch. 1 (1905) 94, 5 (1915) 157; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 204, 227, fig. 5; Ames in Philip. Journ. Sci. 4 (1909) Bot. 673; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 66.

Luzon (Abra, Bontoc, Lepanto, Benguet, Bataan, Laguna). A common species in Luzon. Epiphyte in forest at relatively high altitudes.

ERIA POLYURA Lindl. in Bot. Reg. 27 (1841) Misc. 55, 28 (1842) t. 32; Paxt. Mag. Bot. 9 (1842) 142; Reichb. f. in Walp. Ann. 6 (1861) 279; Naves Novis. App. (1882) 236; Hemsl. in Gard. Chron. II 18 (1882) 500; Ames Orch. 1 (1905) 95, 2 (1908) 196, 5 (1915) 157; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 203, 226, fig. 2; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 115.

Pinalia polyura Ktze. Rev. Gen. Pl. 2 (1891) 679.

Luzon (Cagayan, Bontoc, Benguet, Bataan, Rizal, Laguna, Tayabas), Sibuyan, Panay (Antique), Leyte, Mindanao (Lanao, Surigao). A rather common and widely distributed species in the Philippines. Epiphyte at 60 to 2,050 m altitude.

ERIA PROFUSA Lindl. in Bot. Reg. 28 (1842) Misc. 3; Paxt. Mag. Bot. 12 (1845) 46; Reichb. f. in Walp. Ann. 6 (1861) 271; Hemsl. in Gard. Chron. II 18 (1882) 500; Trim. Handb. Fl. Ceyl. 4 (1898) 168; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 210, 233, fig. 14; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 105.

Luzon (Isabela, Rizal), Negros (Oriental Negros), Mindanao (Agusan), *Lyon F.* 132, *B. S.* 3079, 3083 *Ramos*, *F. B.* 11237 *Everett*, *Weber* 111. Epiphyte at 100 to 700 m altitude.

ERIA PROPINQUA Ames Orch. 7 (1922) 105.

Mindanao (Bukidnon), *B. S.* 38939 *Ramos & Edaño*. Mt. Candoon, on trees of forested slopes, at 1,650 m altitude.

ERIA PUGUAHAANENSIS Ames Orch. 5 (1915) 157.

Leyte, *Wenzel* 0221. Dagami, epiphyte in forest at 60 m altitude.

ERIA RACEMOSA Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 206, 229; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 115; Ames Orch. 5 (1915) 158.

Luzon (Benguet), Mindanao (Misamis), *Curran s. n.*, *Sandkuhl* 225, *F. B.* 4609 *Mearns & Hutchinson* in part.

ERIA RAMOSA Ames nom. nov.

Eria vagans Ames Orch. 7 (1922) 106, non Schltr.

Luzon (Bontoc, Benguet), *Vanoverbergh* 2223, 3896, *Elmer* 8801.

ERIA RAMOSII Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 224, 243, fig. 25; Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1579, Orch. 5 (1915) 158.

Trichotosia ramosii Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 156.

Luzon (Rizal), Negros (Oriental Negros), Mindanao (Agusan), *B. S.* 1775 *Ramos*, 29731 *Ramos & Edaño*, *Elmer* 9803, 13755, 15098, 16136. Epiphyte, forming dense mats, at 1,330 m altitude in Negros.

ERIA RETROFLEXA Lindl. in Journ. Linn. Soc. 3 (1859) 60; Miq. Fl. Ind. Bat. 3 (1859) 662; Naves Novis. App. (1882) 236; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268; Ames Orch. 2 (1908) 197, 5 (1915) 158; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 206, 229, fig. 7; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 76.

Pinalia retroflexa Ktze. Rev. Gen. Pl. 2 (1891) 679.

Luzon (Benguet, Bataan, Rizal, Tayabas), Negros (Oriental Negros), Leyte, Mindanao (Cotabato, Agusan). A not uncommon species, epiphyte at 60 to 860 m altitude.

ERIA RHODOPTERA Reichb. f. in Gard. Chron. II 18 (1882) 586; Rolfe in Orch. Rev. 17 (1909) 220, in Bot. Mag. 136 (1910), t. 8296; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 84; Ames Orch. 5 (1915) 158.

Eria laucheana Kränzl. in Gard. Chron. III 11 (1892) 809.

Eria bractescens Lindl. var. *latipetala* Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 210, 232, fig. 12; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 88.

Luzon (Rizal), Palawan, *B. S. 3000 Ramos, Weber 404*. Epiphyte at sea level to 600 m altitude.

ERIA RINGENS Reichb. f. in Bonpl. 3 (1885) 222; Lindl. in Journ. Linn. Soc. 4 (1859) 55 (erroneously referred to *E. ovata* R. Br.); Reichb. f. in Walp. Ann. 6 (1861) 275, in Trans. Linn. Soc. 30 (1874) 137; Naves Novis. App. (1882) 236; Hook. f. Fl. Brit. Ind. 5 (1890) 793; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 207, 229; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 74; Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1579, Orch. 5 (1915) 158.

Eria polyura Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268, non Lindl.

Pinalia ringens Ktze. Rev. Gen. Pl. 2 (1891) 679.

Negros (Oriental Negros), *Cuming 2128, Elmer 10192*. Cuernos Mountains. In Burma.

ERIA SENILIS Ames Orch. 5 (1915) 159.

Luzon (Sorsogon), Panay (Antique), Leyte, Mindanao (Lanao), *Elmer 16375, B. S. 32575 McGregor, Wenzel 0333, 0335, 0348, 0455, 0852, 0890, 0892, 0894, 0897, 0898, 0899, 0900, 0901, 0902, 0903, 0932, 0933, M. S. Ciemens-s. n.* Epiphyte in forest at 500 to 900 m altitude.

ERIA TAYLORII Ames Sched. Orch. 5 (1923) 31.

Luzon (Laguna), *Taylor 25*. Mt. Maquiling, epiphyte in mossy forest at 1,000 m altitude.

ERIA TRIDENS Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1580, Orch. 5 (1915) 159.

Mindanao (Davao), *Elmer 11850a*. Todaya, Mt. Apo.

ERIA VANOVERBERGHII Ames in Philip. Journ. Sci. 7 (1912) Bot. 20, Orch. 5 (1915) 159.

Luzon (Kalinga, Bontoc, Tayabas), *Boettcher s. n., Vanoverbergh 789, 1347, B. S. 26523 Ramos & Edaño*. Epiphyte at 300 to 1,550 m altitude.

ERIA VENTRICOSA Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 211, 234, fig. 16; Merr. & Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 63; Ames Orch. 5 (1915) 159.

Luzon (Bontoc, Ifugao, Benguet, Bataan, Tayabas), Mindoro, and possibly from Mindanao. A fairly common species in Luzon, epiphyte in forest at 1,650 to 2,400 m altitude.

ERIA VULPINA Reichb. f. in Bonpl. 3 (1855) 222; Lindl. in Journ. Linn. Soc. 3 (1859) 57; Miq. Fl. Ind. Bat. 3 (1859) 660; Reichb. f. in Walp. Ann. 6 (1861) 273; Naves Novis. App. (1882) 236; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268; Ames in Philip. Journ. Sci. 2 (1907) Bot. 332, Orch. 2 (1908) 198, 5 (1915) 160; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 222 241, fig. 23.

Eria vestita Lindl. in Bot. Reg. 30 (1844) Misc. 76, 31 (1845) sub t. 2 in textu, quoad spec. *Cuming*; Reichb. f. in Walp. Ann. 6 (1861) 274, quoad Philip.; Hook. f. in Bot. Mag. 95 (1868) sub t. 5807 in textu, quoad Philip.; Naves Novis. App. (1882) 236.

Pinalia vulpina Ktze. Rev. Gen. Pl. 2 (1891) 679.

Trichotosia vulpina Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 141.

Luzon (Bontoc, Pampanga, Rizal, Laguna, Tayabas, Sorsogon), Mindoro, Sibuyan, Negros (Oriental Negros), Leyte, Mindanao (Camiguin Island, Zamboanga), Basilan. A common and widely distributed species in the Philippines. Epiphyte at 60 to 1,600 m altitude.

ERIA WHITFORDII Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 215; 236; Ames Orch. 5 (1915) 160.

Eria aeridostachya Reichb. f. apud Ames in Philip. Journ. Sci. 2 (1907) Bot. 329; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 213, nota; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 35, quoad syn. *E. whitfordii* et quoad Philip.

Luzon (Isabela, Bataan, Laguna, Sorsogon), Mindoro, Negros (Oriental Negros), Leyte, *Lyon 135, G, Whitford 1119, Merrill s. n., 5518, Reillo 10, Elmer 9731, 16457, Wenzel 0434, 0580*. Epiphyte at 500 to 2,150 m altitude.

ERIA WOODIANA Ames in Philip. Journ. Sci. 2 (1907) Bot. 332; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 205, 228; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 105, fig. 14 J, K.

Luzon (Laguna, Camarines), Mindoro, Leyte, *B. S. 17072 Robinson, 42082 Edaño, Taylor 1, Merrill 5490, Wenzel 0324, 0864, 0924*. Epiphyte at 500 to 1,500 m altitude.

ERIA ZAMBOANGENSIS Ames Orch. 5 (1915) 160.

Mindanao (Zamboanga, Bukidnon), *Merrill 8143, B. S. 38686, 39123 Ramos & Edaño*. Epiphyte in mossy forest at 1,300 to 1,830 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

ERIA AERIDOSTACHYA Reichb. f. apud Ames in Philip. Journ. Sci. 2 (1907) Bot. 239; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 213, nota; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 35, quoad syn. *E. whitfordii* et quoad Philip. = *Eria whitfordii* Leavitt.

For a discussion of the differences and identity of *Eria aeridostachya* Reichb. f. ex Lindl. cf. Leavitt. l. c.

ERIA ALBIDO-TOMENTOSA Lindl. apud Naves Novis. App. (1882) 236.

ERIA AMICA Reichb. f. apud Naves Novis. App. (1882) 236.

ERIA APOROIDES Lindl. in Journ. Linn. Soc. 3 (1859) 60; Miq. Fl. Ind. Bat. 3 (1859) 657; Naves Novis. App. (1882) 236; Merr. in Govt. Lab. Publ. (Philip.) 27 (1905) 84 = *Dendrobium aporoides* (Lindl.) Merr.

ERIA ARMENIACA Lindl. in Bot. Reg. 27 (1841) Misc. 38, t. 42 = *Eria ornata* (Blume) Lindl.

ERIA BRACTESCENS Lindl. var. **LATIPETALA** Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 210, 232, fig. 12; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 88 = *Eria rhodoptera* Reichb. f.

ERIA COMPRESSA Blume apud Naves Novis. App. (1882) 236.

ERIA DASYPUS Reichb. f. apud Naves Novis. App. (1882) 236.

ERIA DILLWYNII Hook. in Bot. Mag. 71 (1845) t. 4163; Lindl. in Bot. Reg. 31 (1845) Misc. 63; F. Kl. in Bot. Zeit. 3 (1845) 550; Paxt. Mag. Bot. 12 (1846) 140; Lindl. in Journ. Linn. Soc. 3 (1859) 55; Naves Novis. App. (1882) 236; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 87 = *Eria bractescens* Lindl.

ERIA ELMERI Ames Orch. 1 (1905) 93, 2 (1908) 196, 5 (1915) 149; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 208, 230, fig. 10; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 74 = *Eria ovata* Lindl.

ERIA ERECTA Lindl. apud Naves Novis. App. (1882) 236.

ERIA FLAVESCENS Lindl. apud Naves Novis. App. (1882) 236; Vidal Rev. Pl. Vasc. Filip. (1886) 268; Ceron Cat. Pl. Herb. Manila (1892) 167.

This species is known only from Java. Vidal's identification of his 1959 was made with expressed doubt.

ERIA FORMOSANA Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 47, non Rolfe = *Eria copelandii* Leavitt.

ERIA FORMOSANA Kränzl. var. *FUSIFORMIS* Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 47 = *Eria copelandii* Leavitt var. *fusiformis* Leavitt.

ERIA GOLDSCHMIDTIANA Schltr. apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 83.

"Provinz der Philippinen und Süd-Formosa."—Kränzl. The type was from Formosa; the species has not been found in the Philippines.

ERIA LANICEPS Reichb. f. apud Naves Novis. App. (1882) 236.

ERIA LAUCHEANA Kränzl. in Gard. Chron. III 11 (1892) 809 = *Eria rhotoptera* Reichb. f.

ERIA MONOSTACHYA Lindl. apud Naves Novis. App. (1882) 236.

ERIA MUCRONATA Lindl. in Bot. Reg. 28 (1842) Misc. 27, in Journ. Linn. Soc. 3 (1859) 59; Miq. Fl. Ind. Bat. 3 (1859) 657; Naves Novis. App. (1882) 236.

Trichotomia mucronata Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 157.

The type of this may or may not have been a Philippine plant. However, the collections of twenty years have not produced a specimen referable to it.

ERIA POLYURA Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 268, non Lindl. = *Eria ringens* Reichb. f.

ERIA PROREPENS Reichb. f. Otia Bot. Hamb. 1 (1878) 54, Xen. Orch. 3 (1878) 31; Naves Novis. App. (1882) 236 = *Phreatia prorepens* Reichb. f.

ERIA PUSILLA Lindl. apud Naves Novis. App. (1882) 236.

ERIA RETUSA Lindl. apud Naves Novis. App. (1882) 236.

ERIA RUGOSA Lindl. Gen. & Sp. Orch. (1830) 66; Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 212, 235; Ames Orch. 5 (1915) 158 = *Eria javanica* (Sw.) Blume.

ERIA STELLATA Lindl. in Bot. Reg. 11 (August, 1825) t. 904; Naves Novis. App. (1882) 236; Usteri Beitr. Ken. Philip. Veg. (1905) 128 = *Eria javanica* (Sw.) Blume.

ERIA VAGANS Ames Orch. 7 (1922) 106, non Schltr. = *Eria ramosa* Ames.

ERIA VENTRICOSA Leavitt var. **BENGUETENSIS** Leavitt in Philip. Journ. Sci. 4 (1909) Bot. 212, 234 = *Eria leavittii* Kränzl.

ERIA VESTITA Lindl. in Bot. Reg. 30 (1844) Misc. 76, 31 (1845) sub t. 2 in textu, quoad spec. *Cuming*; Reichb. f. in Walp. Ann. 6 (1861) 274, quoad Philip.; Hook. f. in Bot. Mag. 95 (1869) sub t. 5807; in textu, quoad Philip.; Naves Novis. App. (1882) 236 = *Eria vulpina* Reichb. f.

Lindley originally placed the Cuming specimen from Manila under *Eria vestita*, but in 1859 (in Journ. Linn. Soc. 3: 57) he corrected this error and gave the points of difference between the two species.

ERIA WENZELII Ames in Philip. Journ. Sci. 8 (1913) Bot. 428 = *Eria brachystachya* Reichb. f.

66. BULBOPHYLLUM Thouars

BULBOPHYLLUM ADENOPETALUM Lindl. in Bot. Reg. 28 (1842) Misc. 85; Reichb. f. in Walp. Ann. 6 (1861) 255; Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 267; Hook. f. Fl. Brit. Ind. 5 (1890) 770, 6 (1890) 188; Ridl. in Journ. Linn. Soc. 32 (1896) 273, Mat. Fl. Mal. Penin. 1 (1907) 75; Ames in Philip. Journ. Sci. 2 (1907) Bot. 333, 7 (1912) Bot. 126, Orch. 2 (1908) 206, 5 (1915) 162.

Bulbophyllum penicillium Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 267; Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, non Par. & Reichb. f.

Bulbophyllum ramosii Ames in Philip. Journ. Sci. 7 (1912) Bot. 25, 128, Orch. 5 (1915) 186.

Phyllorchis adenopetala Ktze. Rev. Gen. Pl. 2 (1891) 677.

Luzon (Apayao, Bontoc, Bataan, Rizal, Laguna, Tayabas), Mindoro, Leyte, Mindanao (Dapitan, Bukidnon). A fairly common species in the Philippines, epiphyte at 500 to 3,000 m altitude. In Malay Peninsula.

BULBOPHYLLUM AEOLIUM Ames in Philip. Journ. Sci. 8 (1913) Bot. 429, Orch. 5 (1915) 164.

Luzon (Rizal), Leyte, Mindanao (Bukidnon), *Reillo s. n.*, *B. S. 33678* Ramos & Edaña, 15391 Ramos, Wenzel 0644, 0652, 0671, 0835. Epiphyte, in forest at 600 to about 1,300 m altitude.

BULBOPHYLLUM AESTIVALE Ames Orch. 5 (1915) 164.

Leyte, Wenzel 0395, 0505, 0506, 0568, 0626. Epiphyte in forest at 500 to 600 m altitude.

BULBOPHYLLUM ALAGENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 333, 7 (1912) Bot. 126, Orch. 3 (1908) 54, t. 44, fig. I. A, a, 5 (1915) 165.

Luzon (Rizal, Laguna), Mindanao, Leyte, *Serrato s. n.*, B. S. 29736 Ramos & Edaña, 16579 Ramos, Merrill 5495, Wenzel 0777. Epiphyte in forest at 400 to 500 m altitude.

BULBOPHYLLUM ALBOROSEUM Ames Orch. 7 (1922) 147.

Luzon (Benguet), Ramos & Edaña 105. Mt. Lusong, epiphyte at 2,000 m altitude.

BULBOPHYLLUM ALSIOSUM Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1583, Orch. 5 (1915) 165.

Luzon (Rizal), Negros (Oriental Negros), Leyte, *Reillo s. n.*, Elmer 9817, Wenzel 0642, 0643, 0666, 0672. Epiphyte in forest at 600 to 1,300 m altitude.

BULBOPHYLLUM ANTENNIFERUM (Lindl.) Reichb. f. in Walp. Ann. 6 (1861) 248; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 138, Orch. 5 (1915) 165.

Cirrhopetalum antenniferum Lindl. in Bot. Reg. 29 (1843) sub t. 49. *Phyllorchis antennifera* Ktze. Rev. Gen. Pl. 2 (1891) 677.

Leyte, Mindanao (Lanao). *Cuming s. n.*, Wenzel 0791, M. S. Clemens 445.

BULBOPHYLLUM ARRECTUM Kränzl. in Fedde Repert. 17 (1921) 384; Ames Sched. Orch. 6 (1923) 82.

Philippines, probably Luzon, *Loher s. n.*

Very close to *Bulbophyllum stellatum* Ames, but material at hand is inadequate to determine its status more definitely.

BULBOPHYLLUM AURATUM (Lindl.) Reichb. f. in Walp. Ann. 6 (1861) 261; Naves Novis. App. (1882) 235; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 82; Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 139, Orch. 5 (1915) 165.

Cirrhopetalum auratum Lindl. in Bot. Reg. 26 (1840) Misc. 50, 29 (1843) t. 61; G. K. in Bot. Zeit. 2 (1844) 103; Moore Ill. Orch. Pl. (1857) Cirrhpet. 5, t. 4; Ridl. in Journ. Linn. Soc. 32 (1896) 284.

Phyllorchis aurata Ktze. Rev. Gen. Pl. 2 (1891) 677.

Leyte, Mindanao, Butuan, Agusan, Davao), *Cuming s. n.*, Lyon 139, Wenzel 0200, 0511, B. S. 12128 Celestino, Weber 106, 137, Black 8. Epiphyte at 5 to 500 m altitude. In Malay Peninsula.

BULBOPHYLLUM BATAANENSE Ames Orch. 1 (1905) 96, 2 (1908) 207, 5 (1915) 166, in Philip. Journ. Sci. 4 (1909) Bot. 599, 7 (1912) Bot. 126.

Luzon (Zambales, Bataan, Laguna), Mindanao (Lanao, Davao), B. S. 3028 Mrs. Lyon, 5001, 16576 Ramos, F. B. 8995, 9504 Merritt & Curran, 796 Borden, Whitford 121, 163, Topping 473, Merrill 5182, Black 10.

- BULBOPHYLLUM BLUMEI** (Lindl.) J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 459, Fig.-Atlas (1911) fig. 351; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, Orch. 5 (1915) 167.
- Ephippium ciliatum* Blume Bijdr. (1825) 309, fig. 65, non *Bulbophyllum ciliatum* Lindl.
- Cirrhopetalum blumei* Lindl. Gen. & Sp. Orch. (1830) 59, in Bot. Reg. 29 (1843) sub t. 49; Miq. Fl. Ind. Bat. 3 (1859) 654; Ridl. in Journ. Linn. Soc. 32 (1896) 285.
- Cirrhopetalum maxillare* Lindl. in Bot. Reg. 29 (1843) sub t. 49.
- Bulbophyllum cheiri* Lindl. in Bot. Reg. 30 (1844) Misc. 44; Reichb. f. in Walp. Ann. 6 (1861) 246; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, Orch. 5 (1915) 169.
- Sarcopodium cheiri* Lindl. Fol. Orch. Sarcopod. (1853) 4; Moore Ill. Orch. Pl. (1857) Sarcopod. 4.
- Bulbophyllum maxillare* Reichb. f. in Walp. Ann. 6 (1861) 248; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, Orch. 5 (1915) 181.
- Bulbophyllum cuspidilingue* Reichb. f. in Walp. Ann. 6 (1861) 264.
- Phyllorchis blumei* Ktze. Rev. Gen. Pl. 2 (1891) 676.
- Phyllorchis cheiri* Ktze. Rev. Gen. Pl. 2 (1891) 677.
- Phyllorchis maxillaris* Ktze. Rev. Gen. Pl. 2 (1891) 677.
- Bulbophyllum ephippium* Ridl. Mat. Fl. Mal. Penin. 1 (1907) 78.
- Leyte, *Cuming s. n.*, Wenzel 050, 0107, 0736. Epiphyte at 60 m altitude. In Malay Peninsula and Java.
- BULBOPHYLLUM BOLSTERI** Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 139, Orch. 5 (1915) 168.
- Bulbophyllum chryseum* Ames in Philip. Journ. Sci. 6 (1911) Bot. 54, quoad spec.
- Luzon (Rizal), Mindanao (Lanao, Surigao), B. S. 3076 Ramos, M. S. Clemens s. n., Bolster 349. Epiphyte.
- BULBOPHYLLUM BONTOCENSE** Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 135, Orch. 5 (1915) 168.
- Luzon (Bontoc, Zambales), Vanoverbergh 705, 1366, 1382, 3681, F. B. 8135 Curran & Merritt. Epiphyte at 1,450 to 1,700 m altitude.
- BULBOPHYLLUM BRACCATUM** Reichb. f. in Linnaea 41 (1877) 117; Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, Orch. 5 (1915) 168.
- Philippines, probably Luzon, Wallis s. n. Probably conspecific with *Bulbophyllum hortensoides* Ames.
- BULBOPHYLLUM CANLAONENSE** Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 136, Orch. 5 (1915) 168.
- Luzon (Benguet), Negros (Oriental Negros), Mindanao (Bukidnon), B. S. 38717 Ramos & Edaña, 4314 Mearns, Elmer 9936, Merrill 7015. Epiphyte at 1,300 to 2,300 m altitude.
- BULBOPHYLLUM CATENULATUM** Kränzl. in Fedde Repert. 17 (1921) 382.
- Philippines, probably Luzon, Loher s. n.

BULBOPHYLLUM CHRYSSENDTUM Ames Orch. 5 (1915) 169.

Leyte, *Wenzel* 0219, 0621, 0657. Dagami and Jaro, epiphyte in forest at 60 to 600 m altitude.

BULBOPHYLLUM CHRYSSEUM (Kränzl.) Ames in Philip. Journ. Sci. 6 (1911) Bot. 54, (excl. spec.), 7 (1912) Bot. 127, Orch. 5 (1915) 170; J. J. Sm. in Bull. Jard. Bot. Buitenz. 8 (1912) 23.

Cirrhopetalum chryseum Kränzl. in Fedde Repert. 8 (1910) 97.

Luzon (Rizal), Leyte, *Loher s. n.*, *Lyon* 152, *Wenzel* 093, 0138, 0214. Epiphyte at low altitudes, 60 m in Leyte.

BULBOPHYLLUM CLEMENSIAE Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 129, Orch. 5 (1915) 170.

Mindanao (Lanao), *M. S. Clemens* 440. Camp Keithley, Lake Lanao.

BULBOPHYLLUM COLUBRIMODUM Ames Sched. Orch. 6 (1923) 82, fig. 12.

Leyte, *Wenzel* 0698. Epiphyte in forest at 600 m altitude.

BULBOPHYLLUM COPELANDII Ames Orch. 1 (1905) 97, 2 (1908) 5, t. 19, 5 (1915) 170, in Philip. Journ. Sci. 7 (1912) Bot. 127.

Luzon (Laguna), Mindanao (Davao), *B. S.* 20434 *Ramos*, *Calica s. n.*, *Copeland* 1316, *Weber* 265.

BULBOPHYLLUM COSTATUM Ames Orch. 6 (1920) 299.

Leyte, *Wenzel* 0665. Epiphyte in forest at 600 m altitude.

BULBOPHYLLUM CUBICUM Ames Orch. 7 (1922) 83.

Mindanao (Bukidnon), *B. S.* 38633 *Ramos & Edaño*. Mahilucot River, epiphyte in forest at 1,400 m altitude.

BULBOPHYLLUM CUMINGII (Lindl.) Reichb. f. in Walp. Ann. 6 (1861) 261; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 6 (1911) Bot. 55, 7 (1912) Bot. 127, Orch. 5 (1915) 170.

Cirrhopetalum cumingii Lindl. in Paxt. Mag. Bot. 8 (1841) 165, in Bot. Reg. 29 (1843) sub t. 49; G. K. in Bot. Zeit. 1 (1843) 791; Moore Ill. Orch. Pl. (1857) *Cirrhopet.* 5; Hook. in Bot. Mag. 83 (1857) t. 4996; Miq. Fl. Ind. Bat. 3 (1859) 653; Veitch Man. Orch. Pl. pt. 3 (1888) 100; Rodigas in Ill. Hort. 39 (1892) 62, in textu.

Phyllorchis cumingii Ktze. Rev. Gen. Pl. 2 (1891) 677.

Luzon (Rizal, Laguna, Tayabas), Panay (Capiz), Negros (Occidental Negros), *Cuming s. n.*, *B. S.* 13763 *Ramos*, 31273 *Ramos & Edaño*, 7334 *Celestino*, *Serrato s. n.*, *F. B.* 13070, 13071 *Curran*, 5227 *Danao & Aspillera*. Apparently at low altitudes.

BULBOPHYLLUM CUNEATUM Rolfe ex Ames Orch. 1 (1905) 98, 2 (1908) 207, 5 (1915) 127, in Philip. Journ. Sci. 4 (1909) Bot. 674, 7 (1912) Bot. 127.

Luzon (Bataan), *Merrill* 3730, *B. S.* 5183 *Topping*, *R. S. Williams* 431. Mt. Mariveles, at about 1,000 m altitude.

BULBOPHYLLUM CURRANII Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 130, Orch. 5 (1915) 170.

Luzon (Benguet), *F. B. 4953 Curran, Haight 7, Ramos & Edaña 62*. Epiphytic vine at about 3,000 m altitude.

BULBOPHYLLUM DAGAMENSE Ames Orch. 5 (1915) 170.

Leyte, *Wenzel 057, 0912*. Dagami and Jaro, epiphyte in forest at 60 to 500 m altitude.

BULBOPHYLLUM DASYPETALUM Rolfe ex Ames Orch. 1 (1905) 98, *text cut*, 2 (1908) 207, 5 (1915) 171; Ames in Philip. Journ. Sci. 2 (1907) Bot. 333, 7 (1912) Bot. 127.

Luzon (Bataan), *Merrill 3720*. Mt. Mariveles.

BULBOPHYLLUM DEAREI (Hort.) Reichb. f. in Flora 71 (April, 1888) 156; Veitch Man. Orch. Pl. pt. 3 (May or June, 1888) 95; Rolfe in *Lindenia* 8 (1892) t. 345; Cogn. in Dict. Icon. Orch. (1902) Bulbo. t. 2; Rolfe in Orch. Rev. 13 (1905) 197; Ames in Philip. Journ. Sci. 4 (1909) Bot. 674, 7 (1912) Bot. 127, Orch. 5 (1915) 171, 6 (1920) 174, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 185.

Sarcopodium dearei Hort. ex Gard. Chron. II 20 (1883) 108, fig. 17; Flor. & Pomol. 22 (1883) 108; Journ. Hort. III 24 (1892) 237; Kew Bull. Add. Ser. 4 (1900) 357.

Sarcopodium godseffianum Hort. ex Gard. Chron. III 7 (1890) 800, *nomen*.

Bulbophyllum godseffianum Weathers in Gardeners' Mag. 33 (1890) 540, *text cut*.

Phyllorchis dearei Ktze. Rev. Gen. Pl. 2 (1891) 677.

Palawan, Mindanao (Lanao), *Weber 7, M. S. Clemens s. n.* Apparently at low altitudes. In Borneo.

BULBOPHYLLUM DECEPTUM Ames Orch. 5 (1915) 171.

Leyte, *Wenzel 0148*. Dagami, epiphyte in forest.

BULBOPHYLLUM DISSOLUTUM Ames in Philip. Journ. Sci. 8 (1913) Bot. 429, Orch. 5 (1915) 172.

Luzon (Laguna), *B. S. 15069 Ramos, Reillo 5*.

BULBOPHYLLUM DORYPHOROIDE Ames Orch. 5 (1915) 172.

Luzon (Ifugao), *B. S. 19873 McGregor*. Mt. Polis.

BULBOPHYLLUM EBRACTEOLATUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 58.

Philippines, probably Luzon, *Loher s. n.* Known only from the original description; no material of this species has been seen.

BULBOPHYLLUM ELMERI Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 136, Orch. 5 (1915) 173.

Luzon (Benguet), *Elmer 6618*. Baguio.

BULBOPHYLLUM ELONGATUM (Blume) Hassk. Cat. Hort. Bogor. (1844) 39; Miq. Fl. Ind. Bat. 3 (1859) 648; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 426, Fig.-Atlas (1911) fig. 321; Ames Orch. 5 (1915) 173, non Kränzl.

Ephippium elongatum Blume Bijdr. (1825) 309.

Cirrhopetalum elongatum Lindl. in Bot. Reg. 29 (1843) sub t. 49;

Reichb. f. in Walp. Ann. 6 (1861) 264.

Bulbophyllum sceptrum Reichb. f. in Walp. Ann. 6 (1861) 254.

Phyllorchis elongata Ktze. Rev. Gen. Pl. 2 (1891) 677.

Phyllorchis sceptrum Ktze. Rev. Gen. Pl. 2 (1891) 678.

Bulbophyllum gigas Ridl. in Journ. Linn. Soc. 32 (1896) 277, Mat.

Fl. Mal. Penin. 1 (1907) 70, fide J. J. Sm.

Mindanao (Cotabato, Bukidnon), *Weber 227*, *B. S. 38616 Ramos & Edaño*. Epiphyte in forest at 100 to 900 m altitude. In Malay Peninsula and Java.

BULBOPHYLLUM ERRATUM Ames Orch. 7 (1922) 85.

Mindanao (Bukidnon), *B. S. 38685 Ramos & Edaño*. Mahilucot River, along stream in forest at about 1,400 m altitude.

BULBOPHYLLUM ESCRITORII Ames Orch. 5 (1915) 174.

Luzon (Benguet, Tayabas), *Ramos & Edaño 104*, *B. S. 20734 Escritor*, *F. B. 13088 Curran*. Epiphyte up to 2,000 m in Benguet.

BULBOPHYLLUM EXILE Ames Orch. 2 (1908) 208, *text cut*, 5 (1915) 175, in Philip. Journ. Sci. 7 (1912) Bot. 127.

Mindanao (Davao), *Copeland 1123*, *R. S. Williams 2563*. Mt. Apo, epiphyte at 3,000 to 3,150 m altitude.

BULBOPHYLLUM EXQUISITUM Ames Sched. Orch. 6 (1923) 84.

Bulbophyllum macgregorii Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 137, Orch. 5 (1915) 180, non Schltr. (1911).

Luzon (Benguet), *B. S. 8322 McGregor*, *31900 Santos*, *Merrill 6624*, *Haight 1*. Epiphyte in mossy forest at 2,100 to 2,400 m altitude.

BULBOPHYLLUM FENIXII Ames in Philip. Journ. Sci. 8 (1913) Bot. 430, Orch. 5 (1915) 175.

Babuyan Islands (Camiguin Island), *B. S. 4138 Fenix*.

BULBOPHYLLUM FILICOIDES Ames Sched. Orch. 6 (1923) 84, *fig. 11*.

Mindanao (Agusan), *Elmer 14200*. Epiphyte at 1,500 m altitude.

BULBOPHYLLUM GIMAGAANENSE Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 130, Orch. 5 (1915) 175.

Luzon (Laguna), Panay (Capiz), Negros (Occidental Negros), Leyte, *Reillo 14*, *B. S. 22885 McGregor*, *35462*, *35518 Martelino & Edaño*, *Whitford 1610*, *Wenzel 0538*, *0541*, *0544*, *0556*, *0613*, *0615*, *0656*. Epiphyte at 100 to 700 m altitude.

BULBOPHYLLUM GLANDULOSUM Ames Sched. Orch. 6 (1923) 87, *fig. 12*.

Luzon (Tayabas), *B. S. 28787 Ramos & Edaño*. Epiphyte at 1,000 m altitude.

BULBOPHYLLUM GNOMONIFERUM Ames Orch. 2 (1908) 209, *text cut*, 5 (1915) 175, in Philip. Journ. Sci. 7 (1912) Bot. 127.

Mindanao (Davao), *Copeland 1023*. Mt. Apo, epiphyte at 1,500 m altitude.

BULBOPHYLLUM GOEBELIANUM Kränzl. in Fedde Repert. 17 (1921) 385.

Philippines, probably Luzon, *Loher s. n.*

Allied to *Bulbophyllum dearei* (Hort.) Reichb. f.

BULBOPHYLLUM GRACILISCAPUM Ames & Rolfe in Ames Orch. 5 (1915) 175.

Mindanao (Davao), *Copeland 1127*. Mt. Apo, epiphyte at 2,000 m altitude.

BULBOPHYLLUM HALCONENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 127, Orch. 3 (1908) 54, t. 44, fig. III. C, c, 5 (1915) 176.

Mindoro, *Merrill 5832*. Mt. Halcon, epiphyte in ridge forest at 1,500 to 2,250 m altitude.

BULBOPHYLLUM HORTENSOIDES Ames Orch. 6 (1920) 300.

Leyte, *Wenzel 0649, 0771*. Epiphyte in forest at 600 m altitude.

Probably conspecific with *Bulbophyllum braccatum* Reichb. f.

BULBOPHYLLUM INVISUM Ames Orch. 7 (1922) 148.

Luzon (Benguet), *Ramos & Edaño 58*. Mt. Pulogloco, epiphyte in mossy forest at 2,200 m altitude.

BULBOPHYLLUM JARENSE Ames Orch. 6 (1920) 301.

Leyte, *Wenzel 0761*. Epiphyte in forest at 500 m altitude.

BULBOPHYLLUM LANCIFOLIUM Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1583, Orch. 5 (1915) 176.

Negros, Mindanao (Bukidnon, Davao), *F. B. 13662 Curran, Elmer 11352, B. S. 38954 Ramos & Edaño*. Epiphyte at about 2,600 m altitude in Davao.

BULBOPHYLLUM LANCILABIUM Ames Orch. 5 (1915) 176.

Luzon (Laguna), *B. S. 20439 Ramos*. Epiphyte at 1,000 m altitude.

BULBOPHYLLUM LANCIPETALUM Ames in Philip. Journ. Sci. 7 (1912) Bot. 23, 127, Orch. 5 (1915) 177.

Mindanao (Dapitan, Davao), *F. B. 4608 Mearns & Hutchinson, Copeland X*. At 2,000 to 2,400 m altitude.

BULBOPHYLLUM LASIOGLOSSUM Rolfe ex Ames Orch. 1 (1905) 100, 2 (1908) 3, 210, t. 18, 5 (1915) 177; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127.

Luzon (Bataan), *F. B. 797 Borden, Whitford s. n, 138, 1094, Copeland s. n.* Epiphyte at about 1,000 m altitude.

BULBOPHYLLUM LASIOPETALUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 57.

Philippines, probably Luzon, *Loher s. n.*

BULBOPHYLLUM LEIBERGII Ames & Rolfe in Ames Orch. 5 (1915) 177

Luzon (Bataan), *Leiberg 6034, B. S. 702 Foxworthy*. Mt. Mariveles a 900 m altitude.

BULBOPHYLLUM LEPANTENSE Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 131, Orch. 5 (1915) 178.

Luzon (Bontoc, Lepanto, Benguet), *Vanoverbergh 1133*, *Merrill 4586*, *B. S. 31778 Santos*. Epiphyte at 2,200 to 2,450 m altitude.

BULBOPHYLLUM LEPTOCAULON Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 59.

Philippines, probably Luzon, *Loher s. n.*

Known only from the original description and the fragment of the type.

BULBOPHYLLUM LEVANAEE Ames Orch. 5 (1915) 178, Sched. Orch. 6 (1923) 89.

Bulbophyllum nymphopolitanum Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 60.

Luzon (Zambales, Sorsogon), Sibuyan, Leyte, *B. S. 4985 Ramos*, *Elmer 12320*, *15836*, *Wenzel 0387*. Epiphyte at 500 m altitude in Leyte.

BULBOPHYLLUM LEYTENSE Ames Orch. 5 (1915) 178.

Leyte, Mindanao (Agusan), *Wenzel 098*, *Elmer 13805*. Epiphyte.

BULBOPHYLLUM LIPENSE Ames Sched. Orch. 3 (1923) 22.

Mindanao (Bukidnon), *B. S. 38532 Ramos & Edaña*. Epiphyte in damp forest at 1,300 m altitude.

BULBOPHYLLUM LOHERIANUM (Kränzl.) Ames comb. nov.

Cirrhopetalum loherianum Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 61.

Philippines, probably Luzon, *Loher*.

Known only from the original description and the fragmentary type specimen.

BULBOPHYLLUM LONGIPETIOLATUM Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1584, Orch. 5 (1915) 180.

Mindanao (Davao), *Elmer 10883*. Todaya, Mt. Apo, epiphyte in humid forest at about 1,300 m altitude.

BULBOPHYLLUM LUZONENSE Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 141, Orch. 5 (1915) 180, Sched. Orch. 6 (1923) 89.

Luzon (Cagayan, Nueva Vizcaya, Laguna), Leyte, Mindanao (Agusan), *F. B. 16787 Curran*, *B. S. 20100 McGregor*, *16557 Ramos*, *Serrato s. n.*, *Wenzel 084*, *0167*, *0405*, *0765*, *Elmer 13623*, *13908*. Epiphyte, at 120 to 500 m altitude in Leyte.

BULBOPHYLLUM MACULOSUM Ames Orch. 5 (1915) 180.

Leyte, *Wenzel 0151*, *0199*. Dagami, epiphyte in forest at 60 m altitude.

BULBOPHYLLUM MAKOYANUM Reichb. f. in Gard. Chron. II 11 (1879) 234; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 81; Ames in Philip. Journ. Sci. 6 (1911) Bot. 55, 7 (1912) Bot. 127, Orch. 5 (1915) 181.

Cirrhopetalum makoyanum Reichb. f. in Gard. Chron. II 11 (1879) 234; Hook. f. in Bot. Mag. 118 (1892) t. 7259; Ridl. in Journ. Linn. Soc. 32 (1896) 284.

Luzon (Ifugao, Rizal), Polillo, Busuanga, Mindanao (Agusan), *B. S. 10438, 19990 McGregor, 19185 Reillo, Weber 37, 407* in part. Epiphyte, apparently at low altitudes. In Malay Peninsula.

BULBOPHYLLUM MARCIDUM Ames Sched. Orch. 6 (1923) 90, *fig. 12*.

Mindanao (Bukidnon), *B. S. 38573 Ramos & Edaño*. Mt. Lipa, epiphyte at 1,630 m altitude.

BULBOPHYLLUM MARIVELENSE Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 132, Orch. 5 (1915) 181.

Luzon (Bataan), *Merrill 3738*. Mt. Mariveles, epiphyte on exposed ridge at 1,100 m altitude.

BULBOPHYLLUM MASAGANAPENSE Ames Orch. 6 (1920) 302.

Luzon (Rizal), Leyte, *B. S. 29714 Ramos & Edaño, 41632, 41639 Ramos, Wenzel 0681*. Epiphyte at 200 to 1,300 m altitude.

BULBOPHYLLUM MEARNsii Ames in Philip. Journ. Sci. 8 (1913) Bot. 430, Orch. 5 (1915) 181.

Bulbophyllum carinatum Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 140, nec Naves nec Cogn.

Luzon (Sorsogon), Mindanao (Dapitan), *Elmer 17039, F. B. 4593 Mearns & Hutchinson*. At 900 m altitude in Mindanao.

BULBOPHYLLUM MELANOGLOSSUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 60.

Philippines, probably Luzon, *Loher s. n.*

BULBOPHYLLUM MERRITTII Ames in Philip. Journ. Sci. 2 (1907) Bot. 334, 7 (1912) Bot. 127, Orch. 5 (1915) 181.

Mindoro, *F. B. 4338 Merritt*. Mt. Halcon, at about 500 m altitude.

BULBOPHYLLUM MINDANAENSE Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1585, Orch. 5 (1915) 181.

Leyte, Mindanao (Davao), *Wenzel 0295, 0397, 0447, 0743, 0744, 0811, 0815, Elmer 11307, 11625, 14160*. Epiphyte in humid forest at 500 to 1,300 m altitude.

BULBOPHYLLUM MINDORENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 334, 7 (1912) Bot. 127, Orch. 3 (1908) 58, *t. 45*, 5 (1915) 182.

Luzon (Camarines), Mindoro, Leyte, *B. S. 22185 Ramos, Merrill 5796, Wenzel 0758*. Epiphyte in damp forest at 500 to 1,000 m altitude.

BULBOPHYLLUM MONSTRABILE Ames Orch. 5 (1915) 182.

Leyte, *Wenzel 0443*. Jaro, epiphyte in forest at 500 m altitude.

BULBOPHYLLUM NEGROSIANUM Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1586, Orch. 5 (1915) 182.

Negros (Oriental Negros), *Elmer 9821*. Cuernos Mountains, epiphyte in mossy woods at about 1,100 m altitude.

BULBOPHYLLUM ORTHOGLOSSUM Wendl. & Kränzl. in Gard. Chron. III 19 (1896) 326; Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, Orch. 5 (1915) 183.

"In Sarangni insula det. W. Micholitz, Freder, Sander Verulamensis collecta indefessus."—Kränzlin. Probably Sarangani Island, off the southern end of Mindanao, is intended. Only a single flower and a sketch of the inflorescence have been seen.

BULBOPHYLLUM OTHONIS (Ktze.) J. J. Sm. in Bull. Gard. Bot. Buitenz. II 8 (1912) 26; Ames Orch. 5 (1915) 183.

Cirrhopetalum nutans Lindl. in Bot. Reg. 25 (1839) Misc. 71, 29 (1843) sub t. 49; Hook. in Bot. Mag. 75 (1849) t. 4418; Miq. Fl. Ind. Bat. 3 (1859) 653.

Bulbophyllum nutans Reichb. f. in Walp. Ann. 6 (1861) 260; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 24, 127, non Thou.

Phyllorchis othonis Ktze. Rev. Gen. Pl. 2 (1891) 677.

Luzon (Sorsogon), Polillo, Leyte, *Elmer 16933*, *B. S. 9077 Robinson, Wenzel 0141, 0192*. Epiphyte, apparently at low altitudes.

BULBOPHYLLUM PAMPANGENSIS Ames Sched. Orch. 6 (1923) 92.

Luzon (Pampanga), *Merrill 11214*.

BULBOPHYLLUM PAPILLIPETALUM Ames Orch. 7 (1922) 149.

Luzon (Benguet), *Ramos & Edaño 79*. Mt. Boadan, epiphyte, on mossy forest slope at about 2,300 m altitude.

BULBOPHYLLUM PAUCIFLORUM Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 132, Orch. 5 (1915) 184.

Luzon (Bontoc), *Vanoverbergh 1381*. Epiphyte at 1,650 m altitude.

BULBOPHYLLUM PERAMOENUM Ames in Philip. Journ. Sci. 8 (1913) Bot. 430, Orch. 5 (1915) 184.

Mindoro, Mindanao (Zamboanga), *Merrill 5832, 8271*.

BULBOPHYLLUM PHILIPPINENSE Ames Orch. 6 (1920) 303.

Leyte, *Wenzel 0659*. Jaro, epiphyte in forest at 500 m altitude.

BULBOPHYLLUM PLEUROTHALLOIDES Ames in Philip. Journ. Sci. 2 (1907) Bot. 335, 7 (1912) Bot. 128, Orch. 3 (1908) 55, t. 44, fig. II B, b, 5 (1915) 184.

Mindoro, *Merrill 6128*. Mt. Halcon, epiphyte in mossy forest at 1,500 m altitude.

BULBOPHYLLUM PLUMATUM Ames Orch. 5 (1915) 184.

Mindanao (Bukidnon), *B. S. 21479 Escritor*.

BULBOPHYLLUM PROFUSUM Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, 137, Orch. 5 (1915) 185.

Luzon (Cagayan, Rizal, Sorsogon), Sibuyan, Leyte, Palawan, Basilan, Mindanao (Agusan, Butuan, Surigao). A not uncommon and widely distributed species. Epiphyte apparently at rather low altitudes.

BULBOPHYLLUM PUGUAHAANENSE Ames Orch. 5 (1915) 185.

Leyte, *Wenzel 0128*. Dagami, Puguahaan, epiphyte at 120 m altitude.

BULBOPHYLLUM REILLOI Ames in Philip. Journ. Sci. 8 (1913) Bot. 431, Orch. 5 (1915) 186.

Luzon (Laguna), *Reillo 24*.

BULBOPHYLLUM SANTOSII Ames Orch. 5 (1915) 186.

Luzon (Laguna, Camarines), Leyte, *B. S.* 20461 *Ramos, Santos s. n., Wenzel 0269.* Epiphyte, up to 750 m altitude in Leyte.

BULBOPHYLLUM SAPPHIRINUM Ames Orch. 5 (1915) 187.

Leyte, *Wenzel 092, 0125, 0136, 0231, 0232.* Dagami, epiphyte in forest at 60 m altitude.

BULBOPHYLLUM SAUROCEPHALUM Reichb. f. in Gard. Chron. II 26 (1886) 262; Rolfe in Orch. Rev. 11 (1903) 190, in textu, 13 (1905) 244; Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, Orch. 5 (1915) 187.

No specimens of this species have been seen.

BULBOPHYLLUM SEMPITERNUM Ames Orch. 6 (1920) 304.

Leyte, *Wenzel 0260, 0287, 0442, 0472, 0524, 0546, 0555, 0589, 0595, 0607, 0610, 0675, 0751, 0810, 0820, 0841, 0848, 0849, 0873, 0893, 0913.* Epiphyte in forest at 500 to 750 m altitude.

BULBOPHYLLUM SENSILE Ames Orch. 5 (1915) 188.

Luzon (Bontoc), *Vanoverbergh 1134, 2816, 3918.* At 1,600 to 1,700 m altitude.

BULBOPHYLLUM SIBUYANENSE Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1587, Orch. 5 (1915) 189.

Sibuyan, *Elmer 12140.* Mt. Giting-giting, epiphyte in humid forest at about 400 m altitude.

BULBOPHYLLUM SIMULACRUM Ames Orch. 5 (1915) 189.

Leyte, *Wenzel s. n., 0268, 0297, 0409, 0692, 0699, 0773, 0786, 0812.* Epiphyte in forest at 500 to 750 m altitude.

BULBOPHYLLUM STELLATUM Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, 141, Orch. 5 (1915) 190.

Luzon (Bontoc), *Vanoverbergh 1360, 3914.* Epiphyte at 1,650 to 1,750 m altitude.

BULBOPHYLLUM STRAMINEUM Ames Orch. 5 (1915) 190.

Palawan, *Merrill 9567, F. B. 21577* *Danao.* Epiphyte at 75 to 500 m altitude.

BULBOPHYLLUM SUBAEQUALE Ames Sched. Orch. 6 (1923) 93.

Luzon (Rizal), *B. S. 29296* *Ramos & Edaña.* Epiphyte at 300 m altitude.

BULBOPHYLLUM SYLLECTUM Kränzl. in Fedde Repert. 17 (1921) 383.

Philippines, probably Luzon, *Loher s. n.*

Owing to the scanty material at hand for study, this is maintained as a species, though it is apparently at best hardly more than a variety of *Bulbophyllum luzonense* Ames; the slight differences are presumably the result of cultivation.

BULBOPHYLLUM TOPPINGII Ames in Philip. Journ. Sci. 8 (1913) Bot. 432, Orch. 5 (1915) 191.

Luzon (Benguet), *Topping s. n.*, Merrill 7753. Epiphyte at about 1,500 m altitude.

BULBOPHYLLUM TRIGONOSEPALUM Kränzl. in Fedde Repert. 17 (1921) 384.

Philippines, probably Luzon, *Loher s. n.*

From the material examined this appears to be a large variety of *Bulbophyllum levanae* Ames.

BULBOPHYLLUM TRISETUM Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, 142, Orch. 5 (1915) 191.

Luzon (Benguet), *F. B. 15899 Bacani*. Itogan to Baguio, epiphyte.

BULBOPHYLLUM VAGANS Ames & Rolfe in Philip. Journ. Sci. 2 (1907) Bot. 335, 7 (1912) Bot. 128; Ames Orch. 2 (1908) 210, *text cut*, 5 (1915) 191.

Luzon (Benguet, Bataan, Laguna, Camarines), Mindanao (Camiguin Island, Agusan), *F. B. 2205 Meyer, 15739 Curran & Merritt, Merrill s. n.* (cult.), *Whitford 1120, B. S. 14423 Ramos, 1636 Foxworthy, Santos s. n., Elmer 13699, Weber 30*. Vinelike epiphyte at 800 to 1,000 m altitude.

Var. **ANGUSTUM** Ames in Philip. Journ. Sci. 2 (1907) Bot. 210, 7 (1912) Bot. 128.

Bulbophyllum vagans Ames & Rolfe var. *linearifolium* Ames Orch. 5 (1915) 191, *nomen*.

Mindoro, *Merrill 6218*. Mt. Halcon, epiphyte in mossy forest at 210 m altitude.

BULBOPHYLLUM VANOVERBERGHII Ames in Philip. Journ. Sci. 7 (1912) Bot. 133, Orch. 5 (1915) 191.

Luzon (Bontoc), *Vanoverbergh 1367*.

BULBOPHYLLUM WEBERI Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, 142, Orch. 5 (1915) 192.

Cirrhopetalum thouarsii Lindl. in Bot. Reg. 24 (1838) t. 11, non *C. thouarsii* Lindl. Gen. & Sp. Orch. (1830) 58; Hook. in Bot. Mag. 72 (1846) t. 4237 (excl. syn.).

Bulbophyllum longiflorum Reichb. f. in Walp. Ann. 6 (1861) 260, quoad Philip.; Naves Novis. App. (1880) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, non Thou.

Bulbophyllum baucoense Ames Orch. 5 (1915) 166.

Luzon (Bontoc, Rizal), Mindanao (Bukidnon, Davao), *Vanoverbergh 2613, B. S. 19239 Reillo, 22253, 41917 Ramos, Weber 8, 194, 289*. On rocks and in trees at 122 to 1,500 m altitude.

BULBOPHYLLUM WENZELII Ames in Philip. Journ. Sci. 8 (1913) Bot. 432, Orch. 5 (1915) 192.

Leyte, *Wenzel s. n.*, 0162, 0178. Dagami, epiphyte in forest at about 60 m altitude.

BULBOPHYLLUM WHITFORDII Rolfe ex Ames Orch. 1 (1905) 100, 2 (1908) 211, 5 (1915) 192, in Philip. Journ. Sci. 7 (1912) Bot. 128.

Luzon (Bataan), Busuanga, Basilan, Mindanao (Davao), *Whitford 61*, *Weber 403*, s. n., *F. B. 4021 Hutchinson*, *R. S. Williams 2915*. Epiphyte.

BULBOPHYLLUM ZAMBALENSE Ames in Philip. Journ. Sci. 7 (1912) Bot. 134, 8 (1913) Bot. 433, Orch. 5 (1915) 192.

Luzon (Zambales), *B. S. 4996 Ramos*.

BULBOPHYLLUM ZAMBOANGENSE Ames in Philip. Journ. Sci. 8 (1913) Bot. 433, Orch. 5 (1915) 192.

Mindanao (Zamboanga), *F. B. 9165 Whitford & Hutchinson*. Epiphyte at 20 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

BULBOPHYLLUM AURECEMUM Boxall ex Naves Novis. App. (1882) 235, *nomen*.

BULBOPHYLLUM BALAENICEPS Reichb. f. in Hamb. Gartenz. 19 (1863) 280, Xen. Orch. 2 (1870) 147, t. 157, fig. 1-6; Naves Novis. App. (1882) 234; Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, Orch. 5 (1915) 165.

This species was described from material in the garden of Consul Schiller. The origin of the plant was unknown and it was doubtfully ascribed to the Philippines by Reichenbach. There seems no evidence that it occurs in the Philippines.

BULBOPHYLLUM BAUCOENSE Ames Orch. 5 (1915) 166 = *Bulbophyllum weberi* Ames.

BULBOPHYLLUM BIFLORUM Teijsm. & Binn. apud Naves Novis. App. (1882) 234.

BULBOPHYLLUM CAREYANUM Spreng. apud Ames in Philip. Journ. Sci. 7 (1912) Bot. 126.

The type of the species was from Nepal, and there is no evidence to show that it occurs in the Philippines.

BULBOPHYLLUM CARINATUM Ames in Philip. Journ. Sci. 7 (1912) Bot. 126, 140, nec Naves nec Cogn. = *Bulbophyllum mearnsii* Ames.

BULBOPHYLLUM CARINATUM (Teijsm. & Binn.) Naves Novis. App. (1882) 235.

BULBOPHYLLUM CHEIRI Lindl. in Bot. Reg. 30 (1844) Misc. 44; Reichb. f. in Walp. Ann. 6 (1861) 246; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, Orch. 5 (1915) 169 = *Bulbophyllum blumei* (Lindl.) J. J. Sm.

BULBOPHYLLUM CHRYSSEUM Ames in Philip. Journ. Sci. 6 (1911) 54, quoad spec. = *Bulbophyllum bolsteri* Ames.

BULBOPHYLLUM COREYANUM Spreng. apud Llanos ex Naves & F.-Vill. in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100; Naves Novis. App. (1882) 235.

BULBOPHYLLUM CUPREUM Lindl. in Bot. Reg. 24 (1838) Misc. 95; Reichb. f. in Walp. Ann. 6 (1861) 257; Naves Novis. App. (1882) 235.

Lindley's statements in the original description that this species came from Manila and was collected by Cuming seem doubtful, as no specimens referable to it from the Philippines have been seen.

BULBOPHYLLUM DAYANUM Reichb. f. apud Naves Novis. App. (1882) 234.

BULBOPHYLLUM DENSUM Spreng. apud Llanos ex Naves & F.-Vill. in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100; Naves Novis. App. (1882) 235.

BULBOPHYLLUM ECHINOCHILUM Kränzlin. in Fedde Repert. 17 (1921) 385; Ames Sched. Orch. 6 (1923) 84.

Kränzlin, l. c., states: "Philippinen. Manila (Loher). Blüthe in kgl. botan. Garten zu München-Nymphenburg in Dezember 1916.

"Durch die Form und Behaarung des Labellums erinnert diese Art sehr stark an *B. Calamaria* Lindl. westafrikanischer Herkunft, * * *."

The record of the Philippine origin of this species is doubted; nothing of its alliance has been seen from the Philippines and its affinities are African, as pointed out by Kränzlin. *Bulbophyllum calamaria* Lindl. has a similar labellum, but its habit is very different.

BULBOPHYLLUM GODSEFFIANUM Weathers in Gardeners' Mag. 33 (1890) 540, text cut = *Bulbophyllum dearei* (Hort.) Reichb. f.

BULBOPHYLLUM HENSCHALLII Boxall ex Naves Novis. App. (1882) 235, *nomen*.

BULBOPHYLLUM LASIANTHUM Lindl. in Gard. Chron. (1855) 53; Reichb. f. in Walp. Ann. 6 (1861) 255; Naves Novis. App. (1882) 234; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127.

The type of this was Sumatran; the living specimen examined by Lindley, said to have come from Manila, imported by Loddiges, probably came from the Malay Peninsula or Sumatra. No record for this species as Philippine is known other than Lindley's original statement.

BULBOPHYLLUM LOBBII Lindl. apud Naves Novis. App. (1882) 234.

BULBOPHYLLUM LONGIFLORUM Reichb. f. in Walp. Ann. 6 (1861) 260, quoad Philip.; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, non Thou. = *Bulbophyllum weberi* Ames.

BULBOPHYLLUM MACGREGORII Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, 137, Orch. 5 (1915) 180, non Schltr. (1911) = *Bulbophyllum exquisitum* Ames.

BULBOPHYLLUM MACULATUM Boxall ex Naves Novis. App. (1882) 235, *nomen*.

BULBOPHYLLUM MAXILLARE Reichb. f. in Walp. Ann. 6 (1861) 248; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 127, Orch. 5 (1915) 181 = *Bulbophyllum blumei* (Lindl.) J. J. Sm.

BULBOPHYLLUM MEDUSAE (Lindl.) Naves Novis. App. (1882) 235.

BULBOPHYLLUM MEGALANTHUM Griff. apud Naves Novis. App. (1882) 235.

BULBOPHYLLUM MUCRONATUM Lindl. apud Naves Novis. App. (1882) 234.

- BULBOPHYLLUM NUTANS** Reichb. f. in Walp. Ann. 6 (1861) 260; Naves Novis. App. (1882) 235; Ames in Philip. Journ. Sci. 7 (1912) Bot. 24, 127, non Thou. = *Bulbophyllum othonis* (Ktze.) J. J. Sm.
- BULBOPHYLLUM NYMPHOPOLITANUM** Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 60 = *Bulbophyllum levinae* Ames.
- BULBOPHYLLUM ODORATUM** Lindl. apud Naves Novis. App. (1882) 234.
- BULBOPHYLLUM PENICILLIUM** Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 149, Rev. Pl. Vasc. Filip. (1886) 267; Ames in Philip. Journ. Sci. 7 (1912) Bot. 128, non Par. & Reichb. f. = *Bulbophyllum adenopetalum* Lindl.
- BULBOPHYLLUM PURPUREUM** (Reichb. f.) Naves Novis. App. (1882) 234.
- BULBOPHYLLUM RAMOSII** Ames in Philip. Journ. Sci. 7 (1912) Bot. 25, 128, Orch. 5 (1915) 186 = *Bulbophyllum adenopetalum* Lindl.
- BULBOPHYLLUM REGNELLII** Reichb. f. apud Naves Novis. App. (1882) 234.
- BULBOPHYLLUM RETICULATUM** Boxall ex Naves Novis. App. (1882) 234, *nomen*.
- BULBOPHYLLUM TENELLUM** Lindl. apud Naves Novis. App. (1882) 234.
- BULBOPHYLLUM VAGANS** Ames & Rolfe var. **LINEARIFOLIUM** Ames Orch. 5 (1915) 191, *nomen* = *Bulbophyllum vagans* Ames & Rolfe var. *angustum* Ames.

67. OCTARRHENA Thwaites

- OCTARRHENA AMESIANA** Schltr. in Fedde Repert. 9 (1911) 439; Ames Orch. 5 (1915) 192.
Phreatia parvula Ames Orch. 2 (1908) 204, non Benth. ex Hook. f. Mindanao (Davao), Copeland 1124, 1427, Elmer 10568, 11413, 11457. Mt. Apo, epiphyte at about 2,000 m altitude.
- OCTARRHENA CAULESCENS** Ames Orch. 5 (1915) 192.
Phreatia caulescens Ames Orch. 2 (1908) 200, *text cut*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 12.
 Luzon (Bontoc, Lepanto, Ifugao, Benguet). A fairly common species in the provinces mentioned. Epiphyte in mossy forest at 1,650 to 2,300 m altitude.
- OCTARRHENA ELMERI** Ames Orch. 5 (1915) 192.
Phreatia elmeri Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1581.
 Luzon (Camarines), Mindanao (Davao), B. S. 22184 Ramos, Elmer 10622. Epiphyte in forest at 2,000 m altitude in Mindanao.
- OCTARRHENA ENSIFOLIA** (Ames) Schltr. in Fedde Repert. 9 (1911) 217; Ames Orch. 5 (1915) 192.
Phreatia ensifolia Ames Orch. 2 (1908) 202, *text cut*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 15.
 Luzon (Bataan), Negros (Oriental Negros), Mindanao (Agusan), Whitford 153, Elmer 9733, 13851. Epiphyte at about 1,250 m altitude in Luzon.

OCTARRHENA GEMMIFERA Ames Orch. 5 (1915) 193.

Leyte, Wenzel 0254, 0293, 0404, 0526, 0547, 0592. Jaro, epiphyte at 500 to 700 m altitude.

68. THELASIS Blume

THELASIS CAPITATA Blume Bijdr. (1825) 386; Lindl. Gen. & Sp. Orch. (1833) 253; Blume Mus. Bot. Lugd.-Bat. 2 (1856) 186 (excl. syn. *Euphroboscis pygmaea* Griff.); Reichb. f. in Bonpl. 5 (1857) 37; Blume Fl. Jav. Orch. (1858) 19 (Orch. Arch. Ind. 20) t. 7, fig. 1; Miq. Fl. Ind. Bat. 3 (1859) 700, Sumatra (1862) 275; Naves Novis. App. (1882) 244; Hook. f. Fl. Brit. Ind. 6 (1890) 86; Ridl. in Trans. Linn. Soc. II 3 (1893) 376, in Journ. Linn. Soc. 32 (1896) 393; Schltr. in Mém. Herb. Boiss. 21 (1900) 69; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 493, Fig.-Atlas (1911) fig. 372; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 199; Ames Orch. 2 (1908) 212, 5 (1915) 194; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 54; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 41; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 190.

Thelasis capitata Blume var. *major* Blume Fl. Jav. Orch. (1858) 19 (Orch. Arch. Ind. 20).

Thelasis ochreatea Lindl. in Journ. Linn. Soc. 3 (1859) 63; Miq. Fl. Ind. Bat. 3 (1859) 701; Naves Novis. App. (1882) 244; Ridl. in Journ. Linn. Soc. 31 (1896) 303.

Thelasis elongata Blume var. *major* Ridl. in Journ. Linn. Soc. 32 (1896) 393, fide Ridl. in Mat. Fl. Mal. Penin. 1. c.

Mindoro, F. B. 12057 Merritt. In Malay Peninsula, Sumatra, Java, and Borneo.

THELASIS CARINATA Blume Bijdr. (1825) 386; Lindl. Gen. & Sp. Orch. (1833) 253, Mus. Bot. Lugd.-Bat. 2 (1856) 187, Fl. Jav. Orch. (1858) 21 (Orch. Arch. Ind. 24) t. 7, fig. 3; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 497, Fig.-Atlas (1912) fig. 375; Ames in Philip. Journ. Sci. 2 (1907) Bot. 336; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 43; Ames Orch. 5 (1915) 194, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 190.

Phreatia carinata Miq. Fl. Ind. Bat. 3 (1859) 655, Sumatra (1862) 274.

Eria carinata Reichb. f. in Seem. Fl. Vit. (1868) 301, in obs.

Thelasis elata Hook. f. Fl. Brit. Ind. 6 (1890) 87; Ridl. in Trans. Linn. Soc. II 3 (1893) 376; Hook. f. in Hook. Icon. Pl. 22 (1893) t. 2156.

Oxyanthera elata Hook. f. in Hook. Icon. Pl. 22 (1893) sub t. 2156; Ridl. in Journ. Linn. Soc. 31 (1896) 303, 32 (1896) 394, Mat. Fl. Mal. Penin. 1 (1907) 200.

Oxyanthera carinata Schltr. in Mém. Herb. Boiss. 21 (1900) 76, in Fedde Repert. 10 (1911) 186.

Babuyan Islands (Camiguin Island), Luzon (Bataan, Laguna, Sorsogon), Mindoro, Negros, Leyte, Basilan, Mindanao (Zamboanga, Agusan). A fairly common and widely distributed species in the Philippines. Epi-

phyte, apparently at low altitudes. In Malay Peninsula, Sumatra, Java, Celebes, and Borneo.

THELASIS MICRANTHA (Brongn.) J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 495, Fig.-Atlas (1911) fig. 374; Ames Orch. 2 (1908) 213, 5 (1915) 194; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 44; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191.

Oxyanthera micrantha Brongn. in Duperr. Voy. Coquille Phan. (1834) 198, t. 37 B; Schltr. in Mém. Herb. Boiss. 21 (1900) 77; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 200.

Thelasis contracta Blume Mus. Bot. Lugd.-Bat. 2 (1856) 187, Fl. Jav. Orch. (1858) 21 (Orch. Arch. Ind. 24) t. 5 B; Miq. Fl. Ind. Bat. 3 (1859) 701.

Thelasis carinata Reichb. f. in Trans. Linn. Soc. 30 (1874) 136; Grant. Orch. Burma (1895) 314, non Blume.

Thelasis decurva Hook. f. Fl. Brit. Ind. 6 (1890) 87; Ridl. in Trans. Linn. Soc. II 3 (1893) 376; Hook. f. in Hook. Icon. Pl. 22 (1893) t. 2157.

Oxyanthera decurva Hook. f. in Hook. Icon. Pl. 22 (1893) sub t. 2157; Ridl. in Journ. Linn. Soc. 31 (1896) 303, 32 (1896) 394.

Luzon (Rizal, Laguna, Tayabas), Leyte, Mindanao (Zamboanga), B. S. 974, 15393, 23514 Ramos, Reillo s. n., 17083 Robinson, Weber 5, Wenzel 048, R. S. Williams 2421, Elmer 7403. Epiphyte at 60 to 800 m altitude. In Malay Peninsula, Sumatra, Java, and Borneo.

THELASIS OBTUSA Blume Bijdr. (1825) 386; Lindl. Gen. & Sp. Orch. (1833) 253; Blume Mus. Bot. Lugd.-Bat. 2 (1856) 186, Fl. Jav. Orch. (1858) 18 (Orch. Arch. Ind. 20); Schltr. in Mém. Herb. Boiss. 21 (1900) 72; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 492, Fig.-Atlas (1911) fig. 371; Ames in Philip. Journ. Sci. 4 (1909) Bot. 675; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 43; Ames Orch. 5 (1915) 194.

Luzon (Bataan), Leiberg 6046. Mt. Mariveles, epiphyte at 900 m altitude. In Java.

THELASIS TRIPTERA Reichb. f. in Bonpl. 3 (1855) 219, in Walp. Ann. 6 (1864) 923; Naves Novis. App. (1882) 244; Hemsl. Bot. Chal. Exped. 1^a (1885) 248; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 272; Kränzl. in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 249; Rolfe in Journ. Linn. Soc. 36 (1903) 39.

Thelasis elongata Blume Mus. Bot. Lugd.-Bat. 2 (1856) 187, Fl. Jav. Orch. (1858) 20 (Orch. Arch. Ind. 23) t. 7, fig. 2, t. 5 C; Miq. Fl. Ind. Bat. 3 (1859) 701; Hook. f. Fl. Brit. Ind. 6 (1890) 86; Ridl. in Trans. Linn. Soc. II 3 (1893) 375, in Journ. Linn. Soc. 32 (1896) 393; Schltr. in Mém. Herb. Boiss. 21 (1900) 70; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 494, Fig.-Atlas (1911) fig. 373; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 199; Ames Orch. 2 (1908) 212, 5 (1915) 194; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 41, fig. 5, T-X; Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch.-Neu-Guin.) (1913) 908; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 190.

Thelasis zollingeri Reichb. f. in Bonpl. 5 (1857) 37.

Thelasis pygmaea Hance in Journ. Linn. Soc. 13 (1872) 127, Fl. Hongk. Suppl. (1872) 39, non Lindl.

Thelasis hongkongensis Rolfe in Kew Bull. (1896) 199, in Journ. Linn. Soc. 36 (1903) 39; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 257.

Luzon (Bataan, Rizal, Sorsogon), Polillo, Panay (Antique), Leyte, Mindanao (Zamboanga, Cotabato, Agusan, Surigao, Davao), Cagayan Sulu. A fairly common and widely distributed species in the Philippines. Epiphyte at 10 to 1,500 m altitude. In Malay Peninsula, China, Sumatra, Java, Borneo, and the Admiralty Islands.

EXCLUDED OR SYNONYMOUS SPECIES

THELASIS ELONGATA Blume Mus. Bot. Lugd.-Bat. 2 (1856) 187; Schltr. in Mém. Herb. Boiss. 21 (1900) 70; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 494, Fig.-Atlas (1911) fig. 373; Ames Orch. 2 (1908) 212, 5 (1915) 194; Kränzl. in Engl. Pflanzenreich IV. 50 II. B. 23 (1911) 41, fig. 5, T-X; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 190 = *Thelasis triptera* Reichb. f.

THELASIS OCHREATA Lindl. in Journ. Linn. Soc. 3 (1859) 63; Naves Novis. App. (1882) 244 = *Thelasis capitata* Blume.

69. PHREATIA Lindley

PHREATIA AMESII Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 16; Ames Orch. 5 (1915) 195, 6 (1920) 207.

Phreatia myosurus Ames Orch. 2 (1908) 203, quoad spec. (excl. syn.) non Lindl.

Luzon (Bontoc, Bataan, Sorsogon), Panay (Antique), Negros (Oriental Negros), Leyte, Mindanao (Agusan), *Vanoverbergh* 707, 1290a, 1477a, 3683, *Merrill* 3854, *Whitford* 1361, *Leiberg* 6061, B. S. 23390 *Ramos*, 32363 *McGregor*, *Elmer* 10354, 13804, *Wenzel* 0713, 0829. Epiphyte, at 300 to 1,600 m altitude. In Borneo.

PHREATIA ARISTULIFERA Ames Orch. 2 (1908) 199, text cut, 5 (1915) 195; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 15.

Luzon (Bontoc, Lepanto, Ifugao, Benguet, Rizal, Tayabas). A reasonably common species in the mountainous regions of Luzon. Epiphyte at 300 to 2,500 m altitude.

PHREATIA DENSIFLORA (Blume) Lindl. Gen. & Sp. Orch. (1830) 64; Miq. Fl. Ind. Bat. 3 (1859) 655; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 502, Fig.-Atlas (1912) fig. 380, Orch. Amb. (1905) 117; Ames Orch. 2 (1908) 201, 5 (1915) 195, 6 (1920) 208; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 18; Rolfe in Gibbs in Journ. Linn. Soc. 42 (1914) 152; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191.

Dendrolirium densiflorum Blume Bijdr. (1825) 350.

Eria myosurus Reichb. f. in Bonpl. 5 (1857) 54, non Reichb. f. in Seem. Fl. Vit. (1868) 300.

Phreatia myosurus Lindl. in Journ. Linn. Soc. 3 (1859) 61; Miq. Fl. Ind. Bat. 3 (1859) 655; Hook. f. Fl. Brit. Ind. 5 (1890) 810; Ridl.

in Journ. Linn. Soc. 31 (1896) 285, 32 (1896) 307, Mat. Fl. Mal. Penin. 1 (1907) 106; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191.

Eria stachyurus Reichb. f. in Seem. Fl. Vit. (1868) 301, in obs.

Luzon (Bataan), *Whitford* 131, 1156. Mt. Mariveles, on rocks at about 900 m altitude. In Malay Peninsula, Sumatra, Java, Amboina, and Borneo.

PHREATIA INFUNDIBULIFORMIS Ames in Philip. Journ. Sci. 7 (1912) Bot. 20, Orch. 5 (1915) 195.

Mindanao (Dapitan), *F. B. 4611 Mearns & Hutchinson*. Mt. Malindang at 1,900 m altitude.

PHREATIA LUZONIENSIS Rolfe ex Ames Orch. 1 (1905) 96, 2 (1908) 202, 5 (1915) 195; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 25; Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1581.

Luzon (Nueva Vizcaya, Bataan, Rizal, Laguna, Tayabas, Sorsogon), Negros (Oriental Negros), Leyte, Mindanao (Zamboanga, Cotabato, Agusan, Davao). A fairly common species of rather wide distribution; epiphyte at 10 to 2,000 m altitude.

PHREATIA MEARNsii Ames in Philip. Journ. Sci. 7 (1912) Bot. 21, Orch. 5 (1915) 195.

Luzon (Pampanga), Mindanao (Dapitan, Davao), *F. B. 19318 Curran, 4592 Mearns & Hutchinson, Elmer 10625*. Epiphyte up to 1,900 m altitude in Dapitan.

PHREATIA NEGROSIANA Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1582, Orch. 5 (1915) 195.

Negros (Oriental Negros), *Elmer 10148*. Cuernos Mountains, epiphyte.

PHREATIA PROREPENS Reichb. f. Otia Bot. Hamb. 1 (1878) 54, in textu; Ames in Philip. Journ. Sci. 2 (1907) Bot. 332, 6 (1911) Bot. 53, Orch. 2 (1908) 205, 3 (1908) 52, *t. 43*, 5 (1915) 195; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 32.

Eria prorepens Reichb. f. Otia Bot. Hamb. 1 (1878) 54, Xen. Orch. 3 (1878) 31; Naves Novis. App. (1882) 236.

Pinalia prorepens Ktze. Rev. Gen. Pl. 2 (1891) 679.

Luzon (Abra, Kalinga, Bontoc, Ifugao, Benguet, Laguna), Mindoro, Negros, Leyte, Mindanao (Cotabato). A not uncommon species of wide distribution, originally collected on Mt. Mahaitai in Luzon by the Wilkes Expedition in 1838-1842. Epiphyte in mossy forest at 600 to 2,450 m altitude.

PHREATIA RAMOSII Ames in Philip. Journ. Sci. 7 (1912) Bot. 22, Orch. 5 (1915) 195.

Luzon (Abra), *B. S. 7140 Ramos*. Epiphyte in forest.

PHREATIA SECUNDA (Blume) Lindl. Gen. & Sp. Orch. (1830) 64; Miq. Fl. Ind. Bat. 3 (1859) 656; Naves Novis. App. (1882) 235; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 500, Fig.-Atlas (1912) fig. 378 (excl. syn. *P. microtidis* Lindl.); Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 53; Kränzl. in Engl. Pflanzenreich IV. 50.

II. B. 23 (1911) 25 (excl. descr. p. p. et syn. *P. microtidis* Lindl.); J. J. Sm. in Winkler in Engl. Bot. Jahrb. 48 (1912) 102; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191.

Dendrolirium secundum Blume Bijdr. (1825) 350.

Eria secunda Reichb. f. in Bonpl. 5 (1857) 54, quoad nom. et syn. sed forsitan non quoad spec.

Phreatia minutiflora Lindl. in Journ. Linn. Soc. 3 (1859) 62; Miq. Fl. Ind. Bat. 3 (1859) 655; Naves Novis. App. (1882) 235; Hook. f. Fl. Brit. Ind. 5 (1890) 811; Ridl. in Journ. Linn. Soc. 31 (1896) 285, 32 (1896) 307; Kränzl. in Reinecke in Engl. Bot. Jahrb. 25 (1898) 607; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 107; Ames Orch. 2 (1908) 205, 5 (1915) 195, 6 (1920) 208; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 23, fig. 3 C, a-g; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191.

Eria minutiflora Reichb. f. in Seem. Fl. Vit. (1868) 301, in obs.

Mindanao (Zamboanga), *Copeland* 1658. San Ramon River, epiphyte at about 200 m altitude. In Malay Peninsula, Sumatra, Java, and Borneo; also stated, by Kränzl., to occur in the Celebes, Pacific Islands, and Samoa.

PHREATIA SULCATA (Blume) J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 505, Fig.-Atlas (1912) fig. 383; J. J. Sm. Orch. Amb. (1905) 118; Ames in Philip. Journ. Sci. 2 (1907) Bot. 332, 7 (1912) Bot. 22; Schltr. in Engl. Bot. Jahrb. 45, Beibl. 104 (1911) 55; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 31; Ames Orch. 5 (1915) 195.

Dendrolirium sulcatum Blume Bijdr. (1825) 347.

Eria sulcata Lindl. Gen. & Sp. Orch. (1830) 69; Miq. Fl. Ind. Bat. 3 (1859) 664.

Pinalia sulcata Ktze. Rev. Gen. Pl. 2 (1891) 679.

Phreatia bracteata Ames in Philip. Journ. Sci. 4 (1909) Bot. 674, non Schltr.

Luzon (Bontoc, Lepanto, Pampanga, Bataan, Rizal, Tayabas), Mindoro, Leyte, Mindanao (Agusan), Basilan. A fairly common species of wide distribution in the Philippines. Epiphyte in forest at 600 to 2,500 m altitude. In Sumatra and Java.

PHREATIA VANOVERBERGHII Ames in Philip. Journ. Sci. 7 (1912) Bot. 22, Orch. 5 (1915) 195.

Luzon (Abra, Kalinga, Bontoc, Ifugao, Benguet), B. S. 7145 Ramos, Boettcher s. n., Vanoverbergh 1115, 1493, 2571, 19385 McGregor, 31848 Santos. Epiphyte at 600 to 2,200 m altitude.

PHREATIA WENZELII Ames Orch. 5 (1915) 196.

Leyte, Wenzel 089. Ormoc, epiphyte in forest at 1,000 m altitude.

PHREATIA XANTHOLEUCA Kränzl. in Fedde Repert. 17 (1921) 386.

Philippines, probably Luzon, *Loher* s. n.

This species was described from material which flowered at Nymphenburg, Munich. No material has been seen.

EXCLUDED OR SYNONYMOUS SPECIES

- PHREATIA BRACTEATA** Ames in Philip. Journ. Sci. 4 (1909) Bot. 674, non Schltr. = *Phreatia sulcata* (Blume) J. J. Sm.
- PHREATIA CAULESCENS** Ames Orch. 2 (1908) 200, *text cut*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 12 = *Octarrhena caulescens* Ames.
- PHREATIA ELMERI** Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1581 = *Octarrhena elmeri* Ames.
- PHREATIA ENSIFOLIA** Ames Orch. 2 (1908) 202, *text cut*; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 15 = *Octarrhena ensifolia* (Ames) Schltr.
- PHREATIA LIMENOPHYLAX** Reichb. f. apud Naves Novis. App. (1882) 235.
- PHREATIA MINUTIFLORA** Lindl. in Journ. Linn. Soc. 3 (1859) 62; Naves Novis. App. (1882) 235; Ames Orch. 2 (1908) 205, 5 (1915) 195; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 23 (1911) 23, *fig. 3 C, a-g*; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 191 = *Phreatia secunda* (Blume) Lindl.
- PHREATIA MYOSURUS** Ames Orch. 2 (1908) 203 (excl. syn.), non Lindl. = *Phreatia amesii* Kränzl.
- PHREATIA PARVULA** Ames Orch. 2 (1908) 204, non Benth. ex Hook. f. = *Octarrhena amesiana* Schltr.
- PHREATIA PUSILLA** Lindl. apud Naves Novis. App. (1882) 235.

70. ACRIOPSIS Reinwardt

ACRIOPSIS FLORIBUNDA Ames Orch. 6 (1920) 306.

Bucas Grande Island, B. S. 35112 Ramos & Pascasio.

ACRIOPSIS INDICA Wight Icon. 5¹ (1852) 20, *t. 1748, fig. I*; Reichb. f. in Walp. Ann. 6 (1863) 492; Hook. f. Fl. Brit. Ind. 6 (1890) 79; Grant Orch. Burma (1895) 313; Schltr. in Oesterr. Bot. Zeitschr. 50 (1900) 287; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 539, Fig.-Atlas (1912) *fig. 411*; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 188; Ames Orch. 6 (1920) 307.

Luzon (Kalinga), *Boettcher s. n.* At 1,000 m altitude. In Burma, Malay Peninsula, and Java.

ACRIOPSIS JAVANICA Reinw. ex Blume Cat. Gew. Buitenz. (1823) 97; Blume Bijdr. (1825) 377, *fig. 71*; Lindl. Gen. & Sp. Orch. (1832) 140; De Vriese Ill. Orch. Ind. Neerl. (1854) *t. 3*; Miq. Fl. Ind. Bat. 3 (1859) 709; Reichb. f. in Walp. Ann. 6 (1863) 492; Hook. f. Fl. Brit. Ind. 6 (1890) 79; Grant Orch. Burma (1895) 313; Ridl. in Journ. Linn. Soc. 32 (1896) 384; Schltr. in Oesterr. Bot. Zeitschr. 50 (1900) 286; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 537, Fig.-Atlas (1912) *fig. 410*; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 187; J. J. Sm. in Nova Guinea 8 (1909) Bot. 119, 8³ (1912) Bot. 604; Schltr. in Fedde Repert. 10 (1911) 190; Ames Orch. 5 (1915) 197; J. J. Sm. in Philip. Journ. Sci. 12 (1917) Bot. 238.

? *Spathoglottis trivalvis* Wall. Cat. (1831) No. 3742, *nomen*; Lindl.

Gen. & Sp. Orch. (1831) 120.

Acriopsis picta Lindl. in Bot. Reg. 29 (1843) Misc. 69.

Acriopsis crispa Griff. Notul. 3 (1851) 333 and Icon. Pl. Asiat. 3 (1851) t. 318.

Acriopsis griffithii Reichb. f. in Bonpl. 2 (1854) 92.

Acriopsis philippinensis Ames Orch. 2 (1908) 215, *text cut*.

Luzon (Sorsogon), Leyte, Mindanao (Misamis, Lanao, Davao), *Elmer* 16957, *Wenzel* 0818, 0845, *Weber* 270, 276, *M. S. Clemens* s. n., *R. F. Black* 15, *R. S. Williams* 2778. At 400 to 580 m altitude. In Burma, Malay Peninsula, Sumatra, Java, Borneo, Celebes, and New Guinea.

EXCLUDED OR SYNONYMOUS SPECIES

ACRIOPSIS PHILIPPINENSIS Ames Orch. 2 (1908) 215, *text cut* = *Acriopsis javanica* Reinw. ex Blume.

71. THECOSTELE Reichenbach f.

THECOSTELE ALATA (Roxb.) Par. & Reichb. f. in Trans. Linn. Soc. 30 (1874) 135, 144, t. 29; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 541, Fig.-Atlas (1912) fig. 412; Rolfe in Orch. Rev. 20 (1912) 237; Ames Orch. 5 (1915) 197, 6 (1920) 215, in Merr in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 193. *Cymbidium alatum* Roxb. Hort. Beng. (1814) 63, *nomen*, Fl. Ind. ed. 2, 3 (1832) 459.

Thecostele zollingeri Reichb. f. in Bonpl. 5 (1857) 37, Xen. Orch. 2 (1868) 133, t. 147; Hook. f. Fl. Brit. Ind. 6 (1890) 19; Grant Orch. Burma (1895) 232; Ridl. in Journ. Linn. Soc. 31 (1896) 299, 32 (1896) 385, Mat. Fl. Mal. Penin. 1 (1907) 190.

Pholidota elmeri Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1557.

Thecostele elmeri Ames in Philip. Journ. Sci. 8 (1913) Bot. 434.

Leyte, Palawan, *Wenzel* 036, 0149, 0215, *Elmer* 12971. Epiphyte in forest at 60 to 80 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

THECOSTELE ELMERI Ames in Philip. Journ. Sci. 8 (1913) Bot. 434 = *Thecostele alata* (Roxb.) Par. & Reichb. f.

72. GRAMMATOPHYLLUM Blume

GRAMMATOPHYLLUM SCRIPTUM (Linn.) Blume Rumphia 4 (1848) 48, Mus. Bot. Lugd.-Bat. 1 (1849) 47; Lindl. in Paxt. Flow. Gard. 2 (1852) 159; Reichb. f. in Walp. Ann. 3 (1852) 550, 6 (1863) 629; Miq. Fl. Ind. Bat. 3 (1859) 708; Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 27 (1864) 21; Naves Novis. App. (1882) 245; J. J. Sm. Orch. Amb. (1905) 84, in Nova Guinea 8 (1909) Bot. 103, 12⁴ (1916) Bot. 428; Schltr. Orchideen (1914) 356, in Orchis 9 (1915) 105, fig. 15; J. J. Sm. in Philip. Journ. Sci. 12 (1917) Bot. 258, in Merr. Interp. Rumph. Herb. Amb. (1917) 177. *Epidendrum scriptum* Linn. Sp. Pl. ed. 2 (1763) 1351; Lam. Encycl. 1 (1783) 185.

- Cymbidium scriptum* Sw. in Schrad. Journ. Bot. 2 (1799) 218, in Kongl. Svensk. Vet.-Akad. Nya Handl. 21 (1800) 237; Willd. Sp. Pl. 4 (1805) 100; Pers. Syn. Pl. 2 (1807) 515.
- Vanda scripta* Spreng. Syst. Veg. 3 (1826) 719.
- Gabertia scripta* Gaudich. in Freyc. Voy. Uranie et Physic. (1829) Bot. 425.
- Grammatophyllum speciosum* Lindl. Gen. & Sp. Orch. (1833) 173, p. p. et quoad syn.; Veitch Man. Orch. Pl. pt. 9 (1893) 33, p. p. non Blume.
- Ophrys cernua* Blanco Fl. Filip. (1837) 639 (as *Ophiris*), ed. 2 (1845) 445, ed. 3, 3 (1879) 35, t. 276, non Linn.
- Grammatophyllum multiflorum* Lindl. in Bot. Reg. 24 (1838) Misc. 46, 25 (1839) t. 65, in Paxt. Mag. 6 (1839) 217, t. & text cut; Edit. in Allg. Berl. Gartenz. 8 (1840) 32; Lindl. in Paxt. Flow. Gard. 2 (1852) 160; Guidon in Journ. Soc. Imp. et Centr. Hort. Par. 4 (1858) 539; Reichb. f. Xen. Orch. 2 (1862) 16, in Walp. Ann. 6 (1863) 629; Naves Novis. App. (1882) 245; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Veitch Man. Orch. Pl. pt. 9 (1893) 32; Ames Orch. 2 (1908) 217, 5 (1915) 198; Schltr. Orchideen (1914) 356, in Orchis 9 (1915) 104; Merr. Sp. Blancoanae (1918) 14.
- Grammatophyllum multiflorum* Lindl. var. *tigrinum* Lindl. in Bot. Reg. 28 (1842) t. 69, in Paxt. Flow. Gard. 2 (1852) 160; Guidon in Journ. Soc. Imp. et Centr. Hort. Par. 4 (1858) 539; Veitch Man. Orch. Pl. pt. 9 (1893) 32; Schltr. in Orchis 9 (1915) 105.
- Grammatophyllum fenzlianum* Reichb. f. Xen. Orch. 2 (1862) 16; Naves Novis. App. (1882) 245; Veitch Man. Orch. Pl. pt. 9 (1893) 31; Ames Orch. 5 (1915) 197, in nota.
- Grammatophyllum rumphianum* Miq. Ann. Mus. Bot. Lugd.-Bat. 4 (1869) 219, tt. 8, 9; Rolfe in Orch. Rev. 3 (1895) 208; Hook. f. in Bot. Mag. 122 (1896) t. 7507; Cogn. in Dict. Icon. Orch. (1899) Grammat. t. 1; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 192.
- Grammatophyllum boweri* F. v. Muell. in Wing South. Sci. Rec. (1883).
- Grammatophyllum leopardinum* Reichb. f. in Flora 71 (1888) 151; Rolfe in Orch. Rev. 23 (1915) 7.
- Grammatophyllum measuresianum* Weathers in Gard. & For. 2 (1889) 524; Edit. in Journ. Hort. III 23 (1891) 342, fig. 69; Stein Orchideenb. (1892) 267; Ames Orch. 2 (1908) 216, 5 (1915) 197; Schltr. in Orchis 9 (1915) 108.
- Grammatophyllum seegerianum* Hort. ex Gard. Chron. III 10 (1891) 49, nomen; Stein Orchideenb. (1892) 267; Rolfe in Orch. Rev. 3 (1895) 208, in textu.
- Grammatophyllum fenzlianum* Reichb. f. var. *measuresianum* Veitch Man. Orch. Pl. pt. 9 (1893) 32, text cut.
- Grammatophyllum guilelmi-secundi* Kränzl. in Gartenfl. 43 (1894) 114; K. Schum. in Notizbl. Bot. Gart. Berl. 1 (Fl. New-Pomm.) (1898) 105; This.-Dyer in Kew Bull. Add. Ser. 4 (1900) 194;

Kränzl. in K. Schum. & Lauterb. Fl. Deutsch. Schutzgeb. (1901) 250.

Grammatophyllum scriptum (Linn.) Blume var. *boweri* Schltr. in Fedde Rept. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1913) 951.

Luzon (Cavite, Tayabas, Camarines, Sorsogon), Catanduanes, Mindoro, Leyte, Biliran, Palawan, Lumbacan. A very variable, common, and widely distributed species in the Philippines. Epiphyte, at sea level to 500 m altitude. In Amboina, Ternate, Celebes (fide J. J. Sm.), Borneo, and New Guinea.

GRAMMATOPHYLLUM SPECIOSUM Blume Bijdr. (1825) 378, fig. 20; Lindl. Gen. & Sp. Orch. (1833) 173, ex parte; Blume Rumphia 4 (1848) 47, t. 191, Mus. Bot. Lugd.-Bat. 1 (1849) 47; Lindl. in Paxt. Flow. Gard. 2 (1852) 157, 160, t. 69; Reichb. f. in Walp. Ann. 3 (1852) 550, 6 (1863) 628; Guidon in Journ. Soc. Imp. et Centr. Hort. Par. 4 (1858) 539 (excl. syn.); Miq. Fl. Ind. Bat. 3 (1859) 708; Hook. in Bot. Mag. 86 (1860) t. 5157; Van Houtte in Fl. des Serres 13 (1861) 177, t. 1386-7; Reichb. f. Xen. Orch. 2 (1862) 16; Batem. Second Cent. Orch. Pl. (1867) t. 181; Edit. in Gard. Chron. II 10 (1878) 181, 183, fig. 36; Naves Novis. App. (1882) 245; Hook. f. Fl. Brit. Ind. 6 (1890) 18 (excl. syn. scripta et scriptum); Burb. in Gard. Chron. III 7 (1890) 287, 297, fig. 46; Stein Orchideenb. (1892) 267, figs. 86, 87; Ceron Cat. Pl. Herb. Manila (1892) 167; Veitch Man. Orch. Pl. pt. 9 (1893) 33 (excl. syn. scripta et scriptum); Grant Orch. Burma (1895) 263 (excl. syn. scripta et scriptum); Ridl. in Journ. Linn. Soc. 31 (1896) 290, 32 (1896) 335; Rolfe in Orch. Rev. 5 (1897) 268, 23 (1915) 6; Rendle in Journ. Bot. 39 (1901) 175; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 485; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 142 (excl. syn. scripta et scriptum); Edit. in Gard. Chron. III 42 (1907) 168, figs. 69, 70; Schltr. Orchideen (1914) 356, in Orchis 9 (1915) 101; Ames Orch. 5 (1915) 198; W. W. in Gard. Chron. III 64 (1918) 55, in textu; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 192; G. Wilson in Orch. Rev. 29 (1921) 83, 98.

Grammatophyllum fastuosum Lindl. in Paxt. Flow. Gard. 2 (1852) 159.

Pattonia macrantha Wight Icon. 5¹ (1852) 21, t. 1750.

Grammatophyllum macranthum Reichb. f. Xen. Orch. 2 (1862) 16, in Walp. Ann. 6 (1863) 628.

Grammatophyllum wallisii Reichb. f. in Linnaea 41 (1876) 107; Rolfe in Orch. Rev. 23 (1915) 7; Ames Orch. 5 (1915) 198.

Grammatophyllum giganteum Blume apud Reichb. f. in Linnaea 41 (1876) 107, in textu.

Grammatophyllum sanderianum Hort. ex Edit. in Gard. Chron. III 14 (1893) 15, sphalm.

Mindanao (Surigao), Wallis s. n., de Veyra s. n. In Burma, Malay Peninsula, Cochin China (fide J. J. Sm.), Sumatra (fide J. J. Sm.), Java, and Borneo.

EXCLUDED OR SYNONYMOUS SPECIES

GRAMMATOPHYLLUM FENZLIANUM Reichb. f. Xen. Orch. 2 (1862) 16; Naves Novis. App. (1882) 245; Veitch Man. Orch. Pl. pt. 9 (1893) 31; Ames Orch. 5 (1915) 197, in nota = *Grammatophyllum scriptum* (Linn.) Blume.

GRAMMATOPHYLLUM MEASURESIANUM Weathers in Gard. & For. 2 (1889) 524; Edit. in Journ. Hort. III 23 (1891) 342, fig. 69; Stein Orchideenb. (1892) 267; Ames Orch. 2 (1908) 216, 5 (1915) 197; Schltr. in Orchis 9 (1915) 108 = *Grammatophyllum scriptum* (Linn.) Blume.

GRAMMATOPHYLLUM MULTIFLORUM Lindl. in Bot. Reg. 24 (1838) Misc. 46, 25 (1839) t. 65; Lindl. in Paxt. Mag. Bot. 6 (1839) 217, t. & text cut; Edit. in Allg. Berl. Gartenz. 8 (1840) 32; Lindl. in Paxt. Flow. Gard. 2 (1852) 160; Guidon in Journ. Soc. Imp. et Centr. Hort. Par. 4 (1858) 539; Reichb. f. Xen. Orch. 2 (1862) 16, in Walp. Ann. 6 (1863) 629; Naves Novis. App. (1882) 245; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Veitch Man. Orch. Pl. pt. 9 (1893) 32; Ames Orch. 2 (1908) 217, 5 (1915) 198; Schltr. Orchideen (1914) 356; Schltr. in Orchis 9 (1915) 104; Merr. Sp. Blancoanae (1918) 14 = *Grammatophyllum scriptum* (Linn.) Blume.

GRAMMATOPHYLLUM WALLISII Reichb. f. in Linnaea 41 (1876) 107; Rolfe in Orch. Rev. 23 (1915) 7; Ames Orch. 5 (1915) 198 = *Grammatophyllum speciosum* Blume.

73. DIPODIUM R. Brown

DIPODIUM PALUDOSUM (Griff.) Reichb. f. Xen. Orch. 2 (1862) 15, in Walp. Ann. 6 (1863) 648, in Gard. Chron. III 4 (1888) 91; Hook. f. Fl. Brit. Ind. 6 (1890) 19; Warner & Will. Orch. Alb. 9 (1891) t. 422; Hook. f. in Bot. Mag. 122 (1896) t. 7464; Ridl. in Journ. Linn. Soc. 31 (1896) 290, 32 (1896) 339; This-Dyer in Kew Bull. Add. Ser. 4 (1900) 161; Rendle in Journ. Bot. 39 (1901) 175; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 142; Ames in Philip. Journ. Sci. 6 (1911) Bot. 55, Orch. 5 (1915) 199, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 192.

Grammatophyllum paludosum Griff. Notul. 3 (1851) 344.

Grammatophyllum affine Griff. Icon. Pl. Asiat. 3 (1851) t. 323.

Wailesia paludosa Reichb. f. in Bonpl. 2 (1854) 93.

Sibuyan, Leyte, Mindanao (Surigao), *Elmer 12435, Wenzel 0198, 0400, 0418, Bolster s. n., B. S. 34800 Ramos & Pascasio*. Epiphyte in forest at 60 to 600 m altitude. In Malay Peninsula and Borneo; also reported from Cochin China.

EXCLUDED OR SYNONYMOUS SPECIES

DIPODIUM PICTUM Reichb. f. apud Naves Novis. App. (1882) 242.

DIPODIUM PUNCTATUM R. Br. apud Naves Novis. App. (1882) 242.

DIPODIUM SQUAMATUM R. Br. apud Naves Novis. App. (1882) 242.

74. CYMBIDIUM Swartz

CYMBIDIUM ATROPURPUREUM (Lindl.) Rolfe in Orch. Rev. 11 (1903) 190; Ames Orch. 2 (1908) 218, 5 (1915) 199.

Cymbidium pendulum Sw. var. *atropurpureum* Lindl. in Gard. Chron. (1854) 287; Hook. f. in Bot. Mag. 94 (1868) t. 5710.

? *Cymbidium pendulum* Vidal Phan. Cuming. Philip. (1885) 150, p. p., Rev. Pl. Vasc. Filip. (1886) 269, p. p. non Sw.

Cymbidium finlaysonianum Lindl. var. *atropurpureum* Veitch Man. Orch. Pl. pt. 9 (1893) 16.

Cymbidium pendulum Sw. var. *purpureum* W. Wats. Orchids, ed. 2 (1895) 151, sphalm.

Luzon (Rizal), Leyte, Mindanao (Davao), *Lyon 101*, *Copeland 648*, *Black 16*, *Wenzel 0103*.

CYMBIDIUM FINLAYSONIANUM Lindl. in Wall. Cat. (1832) No. 7358, *nomen*, Gen. & Sp. Orch. (1833) 164; Hook. f. Fl. Brit. Ind. 6 (1890) 11; Veitch Man. Orch. Pl. pt. 9 (1893) 16; Rendle in Journ. Bot. 39 (1901) 175; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 481, Fig.-Atlas (1911) fig. 366; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 139; Ames Orch. 2 (1908) 218, 5 (1915) 199; Schltr. in Fedde Repert. 10 (1911) 190; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 192.

Cymbidium pendulum Blume Bijdr. (1825) 379; Miq. Fl. Ind. Bat. 3 (1859) 707; Lindl. in Bot. Reg. 26 (1840) t. 25 (excl. syn.); Reichb. f. in Walp. Ann. 6 (1863) 624, quoad spec.; Naves Novis. App. (1882) 245; Vidal Phan. Cuming. Philip. (1885) 150, p. p., Rev. Pl. Vasc. Filip. (1886) 269, p. p.; Warner & Will. Orch. Alb. 10 (1893) t. 437 (excl. syn. in part) non Sw.

Cymbidium aloifolium Lindl. in Wall. Cat. (1832) No. 7352, A & B; Naves Novis. App. (1882) 245; Ridl. in Trans. Linn. Soc. II 3 (1893) 368, in Journ. Linn. Soc. 32 (1896) 333; Ames Orch. 1 (1905) 101; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 39, non Sw.

Cymbidium wallichii Lindl. Gen. & Sp. Orch. (1833) 165; Miq. Fl. Ind. Bat. 3 (1859) 707; Naves Novis. App. (1882) 245.

Cymbidium pendulum Sw. var. *brevilabre* Lindl. in Bot. Reg. 28 (1842) Misc. 60, 30 (1844) t. 24; Paxt. Mag. Bot. 11 (1844) 115; Reichb. f. in Walp. Ann. 6 (1863) 624.

Cymbidium tricolor Miq. Choix des Pl. (1864) t. 19.

Luzon (Benguet, Bataan, Rizal, Tayabas), Polillo, Bohol, Palawan, Mindanao (Zamboanga, Surigao), Dinagat. Epiphyte at sea level to 250 m altitude. In Burma (fide Hook. f.), Malay Peninsula, Java, Celebes, and Borneo; also reported from Sumatra and Cochin China.

EXCLUDED OR SYNONYMOUS SPECIES

CYMBIDIUM ALOIFOLIUM Lindl. in Wall. Cat. (1832) No. 7352, A. & B; Naves Novis. App. (1882) 245; Ames Orch. 1 (1905) 101; Merr. in Philip. Journ. Sci. 1 (1906) Suppl. 39, non Sw. = *Cymbidium finlaysonianum* Lindl.

- CYMBIDIUM FINLAYSONIANUM Lindl. var. ATROPURPUREUM Veitch Man. Orch. Pl. pt. 9 (1893) 16 = *Cymbidium atropurpureum* (Lindl.) Rolfe.
- CYMBIDIUM FLAVESCENS Llanos Fragm. Pl. Filip. (1851) 96; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 74; Merr. Sp. Blancoanae (1918) 113 = ? *Oberonia iridifolia* (Roxb.) Lindl.
- CYMBIDIUM HUTTONII Boxall ex Naves Novis. App. (1882) 245, *nomen*.
- CYMBIDIUM IMBRICATUM Roxb. Hort. Beng. (1814) 63, *nomen*, Fl. Ind. ed. 2, 3 (1832) 460 = *Pholidota imbricata* Lindl.
- CYMBIDIUM IRIDIFOLIUM Roxb. Hort. Beng. (1814) 63, *nomen*, Fl. Ind. ed. 2, 3 (1832) 458 = *Oberonia iridifolia* (Roxb.) Lindl.
- CYMBIDIUM PENDULUM Blume Bijdr. (1825) 379; Naves Novis. App. (1882) 245; Vidal Phan. Cuming. Philip. (1885) 150, p. p., Rev. Pl. Vasc. Filip. (1886) 269, p. p., non Sw. = *Cymbidium finlaysonianum* Lindl.
- CYMBIDIUM PENDULUM Vidal Phan. Cuming. Philip. (1885) 150, p. p., Rev. Pl. Vasc. Filip. (1886) 269, p. p., non Sw. = ? *Cymbidium atropurpureum* (Lindl.) Rolfe.
- CYMBIDIUM PENDULUM Sw. var. ATROPURPUREUM Lindl. in Gard. Chron. (1854) 287; Hook. f. in Bot. Mag. 94 (1868) t. 5710 = *Cymbidium atropurpureum* (Lindl.) Rolfe.
- CYMBIDIUM PENDULUM Sw. var. PURPUREUM W. Wats. Orchids, ed. 2 (1895) 151, sphalm = *Cymbidium atropurpureum* (Lindl.) Rolfe.
- CYMBIDIUM WALLICHII Lindl. Gen. & Sp. Orch. (1833) 165; Naves Novis. App. (1882) 245 = *Cymbidium finlaysonianum* Lindl.

75. CHAMAEANTHUS Schlechter

CHAMAEANTHUS WENZELII Ames Orch. 5 (1915) 200.

Luzon (Sorsogon), Leyte, Mindanao (Davao), *Elmer 15767, 16877, Wenzel 0110 A, 0190, Weber 260*. Epiphyte at sea level to 60 m altitude.

76. THRIXSPERMUM Loureiro

THRIXSPERMUM ACUMINATISSIMUM (Blume) Reichb. f. Xen. Orch. 2 (1867) 121, in Flora 51 (1868) 53; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 569, Fig.-Atlas (1912) fig. 427; Schltr. in Orchis 5 (1911) 54; Ames Orch. 5 (1915) 201.

Dendrocolla acuminatissima Blume Bijdr. (1825) 288.

Aërides acuminatissimum Lindl. Gen. & Sp. Orch. (1833) 240; Miq. Fl. Ind. Bat. 3 (1859) 696.

Sarcochilus acuminatissimus Reichb. f. in Walp. Ann. 6 (1863) 498.

Sarcochilus notabilis Hook. f. Fl. Brit. Ind. 6 (1890) 42, in Hook. Icon. Pl. 22 (1893) t. 2126.

Thrixspermum notabile Ridl. in Journ. Linn. Soc. 32 (1896) 379, Mat. Fl. Mal. Penin. 1 (1907) 183.

Luzon (Laguna), Leyte, *Reillo 56, B. S. 16560, 20447, 20455 Ramos, Wenzel 060, 0101, 0163*. Epiphyte at 60 m altitude in Leyte. In Malay Peninsula and Java.

THRIXSPERMUM AGUSANENSE Ames Orch. 5 (1915) 201.

Mindanao (Agusan), *Weber 147*. Veruela, Agusan River, epiphyte at 15 m altitude.

THRIXSPERMUM AMPLEXICAULE (Blume) Reichb. f. Xen. Orch. 2 (1867) 121, in *Flora* 51 (1868) 53; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 573, Fig.-Atlas (1912) *fig. 430*, Orch. Amb. (1905) 96; Schltr. in *Orchis* 5 (1911) 54; Ames in *Philip. Journ. Sci.* 8 (1913) Bot. 438, Orch. 5 (1915) 202; J. J. Sm. in *Philip. Journ. Sci.* 12 (1917) Bot. 259; Ames in *Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No.* (1921) 194.

Dendrocolla amplexicaulis Blume *Bijdr.* (1825) 288.

Aërides amplexicaule Lindl. *Gen. & Sp. Orch.* (1833) 239; *Miq. Fl. Ind. Bat.* 3 (1859) 696.

Sarcophilus lilacinus Griff. *Notul.* 3 (1851) 334, *Icon. Pl. Asiat.* 3 (1851) *t. 320, fig. II*; Reichb. f. in *Walp. Ann.* 6 (1863) 499; Hook. f. *Fl. Brit. Ind.* 6 (1890) 40; *Ridl. in Trans. Linn. Soc. II* 3 (1893) 371; Hook. f. in *Bot. Mag.* 127 (1901) *t. 7754*.

Orsidice amplexicaulis Reichb. f. in *Bonpl.* 2 (1954) 93.

Orsidice lilacina Reichb. f. in *Bonpl.* 2 (1854) 93.

Sarcophilus amplexicaulis Reichb. f. in *Walp. Ann.* 6 (1863) 499; Rolfe in *Journ. Bot.* 23 (1885) 215; Vidal Phan. *Cuming. Philip.* (1885) 150, *Rev. Pl. Vasc. Filip.* (1886) 271.

Thrixspermum lilacinum Reichb. f. Xen. Orch. 2 (1867) 121, in *Flora* 51 (1868) 53; *Ridl. in Journ. Linn. Soc.* 31 (1896) 298, 32 (1896) 377, *Mat. Fl. Mal. Penin.* 1 (1907) 181.

Luzon (Nueva Vizcaya), Mindanao (Lanao), *Cuming 2056*, *B. S. 11143* *McGregor, M. S. Clemens 949*. In bogs, twining on grasses. In Malay Peninsula, Sumatra, Java, Amboina, Celebes, and Banda Islands.

THRIXSPERMUM BICRISTATUM Ames Orch. 5 (1915) 202.

Leyte, Mindanao (Agusan), *Wenzel 030*, *Weber 145*. Epiphyte at 15 to 40 m altitude.

THRIXSPERMUM ELONGATUM Ames Orch. 5 (1915) 203.

Luzon (Bataan, Rizal, Laguna, Sorsogon), Polillo, Siquijor, Leyte, Palawan, Mindanao (Butuan, Agusan, Surigao, Davao). A fairly common and fairly widely distributed species. Epiphyte at sea level to 525 m altitude.

THRIXSPERMUM LINEARIFOLIUM Ames Orch. 5 (1915) 205.

Mindanao (Dapitan, Bukidnon), *F. B. 4734* *Mearns & Hutchinson, B. S. 38499* *Ramos & Edaño*. Mt. Malindang and Mt. Lipa. Epiphyte in mossy forest at 1,900 to 2,200 m altitude.

THRIXSPERMUM QUINQUELOBUM Ames Orch. 5 (1915) 206.

Luzon (Laguna), *B. S. 10016*, *16561*, *16562*, *20444* *Ramos, Reillo 19*, *Serrato s. n.* Epiphyte in forest.

THRIXSPERMUM ROBINSONII Ames Orch. 5 (1915) 207.

Luzon (Bataan, Laguna, Tayabas, Sorsogon), Negros (Occidental Negros), Leyte, Mindanao (Agusan), *Whitford 137*, *1508*, *B. S. 9447*, *17096* *Robinson*, *17313* *Robinson & Brown*, *16282* *Brown*, *Elmer 13833*, *17134*, *Wenzel 0408*. Epiphyte at 600 to 2,150 m altitude.

THRIXSPERMUM ROSTRATUM Ames Orch. 5 (1915) 208.

Panay, *Lyon 159*. Patolan, plant climbing.

THRIXSPERMUM SUBULATUM (Blume) Reichb. f. Xen. Orch. 2 (1867) 122, in *Flora* 51 (1868) 53; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 578, Fig.-Atlas (1912) *fig. 434*, Orch. Amb. (1905) 97; Schltr. in *Orchis* 5 (1911) 57; J. J. Sm. in Bull. Jard. Buitenz. II 9 (1913) 102; Ames Orch. 7 (1922) 134.

Dendrocolla subulata Blume Bijdr. (1825) 291.

Aërides subulatum Lindl. Gen. & Sp. Orch. (1833) 241; Miq. Fl. Ind. Bat. 3 (1859) 698.

Sarcochilus subulatus Reichb. f. in Walp. Ann. 6 (1863) 500; Naves Novis. App. (1882) 238.

Thrixspermum falcilobum Schltr. in Bull. Herb. Boiss. II 6 (1906) 469; Schltr. in *Orchis* 5 (1911) 57 (fide J. J. Sm. in Bull. Jard. Buitenz. I. c.).

Luzon (Bontoc), *B. S. 37905 Ramos & Edaño*. Mt. Masapilid, epiphyte at 1,000 m altitude. In Sumatra, Java, and Amboina.

THRIXSPERMUM VANOVERBERGHII Ames in Philip. Journ. Sci. 8 (1913) Bot. 438, Orch. 5 (1915) 209.

Luzon (Bontoc), *Vanoverbergh 632, 1091, 1531, 1792*. Epiphyte at about 1,400 m altitude.

THRIXSPERMUM WEBERI Ames Orch. 7 (1922) 134.

Mindanao (Agusan), *Weber 187, 187X, 187X'*. Cabadbaran, epiphyte in swamp at sea level.

THRIXSPERMUM WENZELII Ames Orch. 5 (1915) 209.

Luzon (Laguna, Sorsogon), Leyte, Palawan, Mindanao (Agusan), Basilan. A not uncommon species of fairly wide distribution. Epiphyte at sea level to 60 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

THRIXSPERMUM MCGREGORII Ames in Philip. Journ. Sci. 2 (1907) Bot. 337 = *Sarcochilus mcgregorii* Ames.

THRIXSPERMUM MINDANAENSE Ames in Philip. Journ. Sci. 8 (1913) Bot. 436 = *Sarcochilus mindanaensis* Ames.

THRIXSPERMUM PALLIDUM Reichb. f. Xen. Orch. 2 (1867) 122; Ames Orch. 2 (1908) 253 = *Sarcochilus pallidus* (Blume) Reichb. f.

THRIXSPERMUM PHILIPPINENSE Ames in Philip. Journ. Sci. 8 (1913) Bot. 437 = *Sarcochilus philippinensis* Ames.

THRIXSPERMUM UNGUICULATUM Reichb. f. Xen. Orch. 2 (1867) 122, in *Flora* 51 (1868) 53; Warner & Will. Orch. Alb. 6 (1887) *t. 266* = *Sarcochilus pallidus* (Blume) Reichb. f.

77. SARCOCHILUS R. Brown

SARCOCHILUS APPENDICULATUS (Blume) J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 564, Fig.-Atlas (1912) *fig. 425*.

Dendrocolla appendiculata Blume Bijdr. (1825) 289.

Aërides pusillum Lindl. Gen. & Sp. Orch. (1833) 241; Miq. Fl. Ind. Bat. 3 (1859) 697.

Sarcochilus pusillus Reichb. f. in Walp. Ann. 6 (1863) 500.

Grosourdyia appendiculata Reichb. f. Xen. Orch. 2 (1867) 123; Naves Novis. App. (1882) 239.

Thrixspermum appendiculatum Ktze. Rev. Gen. Pl. 2 (1891) 682.

Palawan, Weber 605. Alfonso XIII, epiphyte at sea level. In Java.

J. J. Smith, Fl. Buitenz. l. c., states that *Sarcochilus hirtulus* Hook. f. Fl. Brit. Ind. 6 (1890) 39, Hook. f. in Hook. Icon. Pl. 22 (1892) t. 2121: *Ascochilus hirtulus* Ridl. in Journ. Linn. Soc. 31 (1896) 375, Ridl. Mat. Fl. Mal. Penin. 1 (1907) 178 is perhaps conspecific.

SARCOCHILUS LEYTENSIS Ames Orch. 5 (1915) 211.

Leyte, ? Basilan, Wenzel s. n., 0109, ? B. S. 16242 Reillo. Epiphyte at 60 m altitude.

SARCOCHILUS LONGICALCARUS Ames & Rolfe in Ames Orch. 5 (1915) 212.

Luzon (Bontoc, Bulacan, Tayabas, Sorsogon, Camarines), Mindoro, Leyte, Mindanao (Butuan, Davao). A not uncommon species of wide distribution. Epiphyte at sea level to 500 m altitude.

SARCOCHILUS MCGREGORII Ames Orch. 5 (1915) 213.

Thrixspermum mcgregorii Ames in Philip. Journ. Sci. 2 (1907) Bot. 337.

Mindoro, McGregor 288. Balete, Baco River, epiphyte.

SARCOCHILUS MERRILLII Ames Orch. 5 (1915) 213.

Mindanao (Surigao), Merrill 5432. Baganga, epiphyte at near sea level.

SARCOCHILUS MINDANAENSIS Ames Orch. 5 (1915) 214.

Thrixspermum mindanaense Ames in Philip. Journ. Sci. 8 (1913) Bot. 436.

Luzon (Sorsogon), Sibuyan, Leyte, Mindanao (Agusan, Davao), Elmer 12530, 15361 in part, Wenzel 0110B, 0280, 0366, 0390, Weber 99, 139, 255. Epiphyte at sea level to 640 m altitude.

SARCOCHILUS PALAWANENSIS Ames Orch. 5 (1915) 214.

Palawan, Mindanao (Davao), Weber 614, Elmer 10515. Epiphyte at sea level in Palawan.

SARCOCHILUS PALLIDUS (Blume) Reichb. f. in Walp. Ann. 6 (1863) 500; Naves Novis. App. (1882) 238; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 559, Fig.-Atlas (1912) fig. 422, Orch. Amb. (1905) 92, in Winkler in Engl. Bot. Jahrb. 48 (1912) 104; Ames Orch. 5 (1915) 215, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 196.

Dendrocolla pallida Blume Bijdr. (1825) 290.

Aërides pallidum Lindl. Gen. & Sp. Orch. (1833) 241, in Paxt. Flow. Gard. 2 (1839) 143; Miq. Fl. Ind. Bat. 3 (1859) 697, nec Blume nec Roxb.

- Sarcochilus unguiculatus* Lindl. in Bot. Reg. 26 (1840) Misc. 67, 32 (1846) sub t. 19; Edit. in Allg. Berl. Gartenz. 14 (1846) 167; Reichb. f. in Walp. Ann. 6 (1863) 501, in Gard. Chron. II 15 (1881) 562; Naves Novis. App. (1882) 238; Rolfe in Orch. Rev. 2 (1894) 231, 14 (1906) 273, fig. 37; Ridl. in Journ. Linn. Soc. 31 (1896) 298, Mat. Fl. Mal. Penin. 1 (1907) 175; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 196.
- Aërides diurnum* Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 27 (1864) 19.
- Thrixspermum pallidum* Reichb. f. Xen. Orch. 2 (1867) 122, in Flora 51 (1868) 53; Ames Orch. 2 (1908) 253.
- Thrixspermum unguiculatum* Reichb. f. Xen. Orch. 2 (1867) 122, in Flora 51 (1868) 53; Warner & Will. Orch. Alb. 6 (1887) t. 266.
- Phalaenopsis ruckeri* Hort. ex Reichb. f. in Gard. Chron. II 15 (1881) 562, in textu.
- Phalaenopsis ruckeriana* Hort. ex Warner & Will. Orch. Alb. 6 (1887) t. 266, in syn.
- Sarcochilus aureus* Hook. f. Fl. Brit. Ind. 6 (1890) 35, in Ann. Roy. Bot. Gard. Calc. 5 (Cent. Ind. Orch.) (1895) 42, t. 64; Ridl. in Journ. Linn. Soc. 32 (1896) 373.
- Thrixspermum aureum* Ktze. Rev. Gen. Pl. 2 (1891) 682.
- ? *Phalaenopsis fugax* Kränzl. in Gard. Chron. III 14 (1893) 360; Rolfe in Orch. Rev. 13 (1905) 230; cf. infra.
- Sarcochilus unguiculatus* Lindl. var. *aureus* Ridl. Mat. Fl. Mal. Penin. 1 (1907) 175.

Luzon (Cagayan, Benguet, Nueva Vizcaya, Rizal, ? Laguna, Tayabas, Sorsogon), Polillo, Mindoro, Leyte, Mindanao (Zamboanga, Agusan, Butuan, Davao), Basilan. A common and widely distributed species in the Philippines. Epiphyte apparently at low altitudes. In Malay Peninsula, Sumatra, Java, Amboina, Celebes, and Borneo.

J. J. Smith, Fl. Buitenz. 6 and Orch. Amb. II. cc., reduces *Phalaenopsis fugax* Kränzl. to this species, whereas Ridley, Mat. Fl. Mal. Penin. 1 l. c., makes it a synonym of *Sarcochilus caligaris* Ridl. J. J. Smith also considers *Sarcochilus cladostachys* Hook. f. Fl. Brit. Ind. 6 (1890) 35, in Ann. Roy. Bot. Gard. Calc. 5 (1895) 43, t. 65 to be conspecific, but Ridley considers them as distinct species.

SARCOCHILUS PHILIPPINENSIS Ames Orch. 5 (1915) 215, non Vidal.
Thrixspermum philippinense Ames in Philip. Journ. Sci. 8 (1913) Bot. 437.

Luzon (Apayao, Zambales, Rizal, Laguna, Sorsogon), Polillo, Leyte, Mindanao (Agusan). A fairly common species of wide distribution. Epiphyte occurring, as far as records show, at 3 to 60 m altitude.

SARCOCHILUS TRIPERCUS Ames Orch. 7 (1922) 125.

Leyte, Wenzel 0968. Tacloban, Tigbao, epiphyte in forest at sea level.

SARCOCHILUS ZAMBOANGENSIS Ames Sched. Orch. 5 (1923) 39.

Mindanao (Zamboanga), Merrill 11640. Flecha Point, epiphyte in forest at about 10 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

SARCOCHILUS AMPLEXICAULIS Reichb. f. in Walp. Ann. 6 (1863) 499; Rolfe in Journ. Bot. 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271 = *Thrixspermum amplexicaule* (Blume) Reichb. f.

SARCOCHILUS CALCEOLUS Lindl. in Bot. Reg. 32 (1846) t. 19; G. K. in Bot. Zeit. 4 (1846) 798; Paxt. Mag. Bot. 13 (1846) 93; Reichb. f. in Walp. Ann. 1 (1849) 781, 6 (1863) 501; Miq. Fl. Ind. Bat. 3 (1859) 689; Naves Novis. App. (1882) 238; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271; Ridl. in Trans. Linn. Soc. II 3 (1893) 371; Ames Orch. 5 (1915) 210, in textu. *Thrixspermum calceolus* Reichb. f. Xen. Orch. 2 (1867) 122, in Flora 51 (1868) 53; Ridl. in Journ. Linn. Soc. 32 (1896) 378, Mat. Fl. Mal. Penin. 1 (1907) 181.

This was originally described as Philippine, but there is every reason to believe that Cuming's specimen came from Singapore, where the species is common and where Cuming also collected botanical material; it has not appeared in any Philippine collections made since Cuming's visit (1836-1840); cf. Ames l. c.

SARCOCHILUS CENTIPEDA Naves Novis. App. (1882) 238 (excl. syn.).

SARCOCHILUS COMPRESSUS Reichb. f. apud Naves Novis. App. (1882) 238.

SARCOCHILUS CROCEUS Lindl. in Bot. Reg. 32 (1846) sub t. 19; Reichb. f. in Walp. Ann. 1 (1849) 781; Miq. Fl. Ind. Bat. 3 (1859) 689; Naves Novis. App. (1882) 238; Ames Orch. 5 (1915) 210, in nota.

This was described from a single leaf and bud sent to Lindley by Messrs. Loddiges. No material from the Philippines referable to this species has been seen.

SARCOCHILUS FALCATUS R. Br. apud Naves Novis. App. (1882) 238.

SARCOCHILUS NEPALENSIS Spreng. apud Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 99.

SARCOCHILUS OLIVACEUS Lindl. apud Naves Novis. App. (1882) 238.

SARCOCHILUS PARVIFLORUS Lindl. apud Naves Novis. App. (1882) 238.

SARCOCHILUS PHILIPPINENSIS Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271 = *Camarotis philippinensis* Lindl.

SARCOCHILUS RHOPALORRHACHIS Reichb. f. apud Naves Novis. App. (1882) 238.

SARCOCHILUS SUBULATUS Reichb. f. in Walp. Ann. 6 (1863) 500; Naves Novis. App. (1882) 238 = *Thrixspermum subulatum* (Blume) Reichb. f.

SARCOCHILUS TENUIFOLIUS (Linn.) Naves Novis. App. (1882) 238.

SARCOCHILUS TERES Reichb. f. apud Naves Novis. App. (1882) 238.

SARCOCHILUS UNGUICULATUS Lindl. in Bot. Reg. 26 (1840) Misc. 67, 32 (1846) sub t. 19; Edit. in Allg. Berl. Gartenz. 14 (1846) 167; Reichb. f. in Walp. Ann. 6 (1863) 501; Reichb. f. in Gard. Chron.

II 15 (1851) 562; Naves Novis. App. (1882) 238; Rolfe in Orch. Rev. 2 (1894) 231, 14 (1906) 273, *fig. 37*; Ridl. in Journ. Linn. Soc. 31 (1896) 298, Mat. Fl. Mal. Penin. 1 (1907) 175; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 196 = *Sarcochilus pallidus* (Blume) Reichb. f.

SARCOCHILUS USNEOIDES Reichb. f. apud Naves Novis. App. (1882) 238.

78. KINGIELLA Rolfe

KINGIELLA PHILIPPINENSIS (Ames) Rolfe in Orch. Rev. 25 (1917) 197; Ames Orch. 6 (1920) 307.

Doritis philippinensis Ames Orch. 2 (1908) 235, *text cut*, 5 (1915) 215.

Luzon (Cagayan, Bataan, Bulacan, Rizal), Leyte, Palawan, Bancalan, Mindanao (Cotabato, Bukidnon, Agusan, Davao). A common species of wide distribution. Epiphyte at nearly sea level to 500 m altitude.

79. PHALAEOPSIS Blume

PHALAEOPSIS AMABILIS (Linn.) Blume Bijdr. (1825) 294, *fig. 44*; Lindl. Gen. & Sp. Orch. (1833) 213; Bennett Pl. Jav. Rar. (1838) 28, *t. 8*; Blume Rumphia 4 (1848) 52, *tt. 194, fig. 1, 199A* (excl. citat. Lindl.); Reichb. f. in Bot. Zeit. 10 (1852) 672, in Walp. Ann. 3 (1852) 561 (excl. pars Philip.); Miq. Fl. Ind. Bat. 3 (1859) 690 (excl. citat. Bot. Mag. et pars Philip.), Sumatra (1862) 274; Reichb. f. Xen. Orch. 2 (1862) 5; Duchartre in Journ. Soc. Imp. et Centr. Hort. Par. 8 (1862) 730; Lem. in Ill. Hort. 10 (1863) Misc. 13; Linden & Rodigas in Lindenia 2 (1886) 65, *quoad syn. in part*; Rolfe in Gard. Chron. II 26 (1886) 212, *fig. 43A*; Kerchove in Rev. Hort. Belg. 16 (1890) 25, *t. 3*; Veitch Man. Orch. Pl. pt. 7 (1891) 22, *t.*; Stein Orchideenb. (1892) 504, *fig. 153*; Rolfe in l'Orchidoph. 13 (1893) 100; Will. Orch. Grow. Man. ed. 7 (1894) 660, *text cut* (excl. bibliog. in parte et *quoad* Philip.); Ridl. in Journ. Linn. Soc. 31 (1896) 292; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 549, Fig.-Atlas (1912) *fig. 416* (excl. bibliog. in parte et *quoad* Philip.), Orch. Amb. (1905) 92 (excl. citat. Bot. Mag. *t. 4297*); Ames Orch. 1 (1905) 101, *quoad syn. et descr. (excl. spec.)*, 2 (1908) 224; J. J. Sm. in Nova Guinea 8 (1909) Bot. 119 (excl. Philip.), 8^a (1912) Bot. 604, 9 (1913) Bot. 100, 12^a (1916) Bot. 455; Schltr. Orchideen (1914) 543, *fig. 183*; Rolfe in Gibbs in Journ. Linn. Soc. 42 (1914) 157; J. J. Sm. in Philip. Journ. Sci. 12 (1917) Bot. 258, in Merr. Interp. Rumph. Herb. Amb. (1917) 177; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 196.

Epidendrum amabile Linn. Sp. Pl. (1753) 953, ed. 2 (1763) 1351; Sw. in Nov. Act. Ups. 6 (1799) 67, in Schrad. Journ. Bot. 2 (1799) 210; Willd. Sp. Pl. 4 (1805) 115; Pers. Syn. Pl. 2 (1807) 518; Spreng. Syst. Veg. 3 (1826) 737.

Cymbidium amabile Roxb. Hort. Beng. (1814) 63, *nomen*, Fl. Ind. ed. 2, 3 (1832) 457.

Phalaenopsis grandiflora Lindl. in Gard. Chron. (1848) 39, *fig. 1*; Reichb. f. in Walp. Ann. 3 (1852) 561; Moore Ill. Orch. Pl. (1857) Phalaenop. 7; Miq. Fl. Ind. Bat. 3 (1859) 690; Hook. in Bot. Mag. 86 (1860) *t. 5184*; Groenland in Rev. Hort. 32 (1860) 238, *figs.*

- 53, 54; Gedney in Floral World (1876) 323; Puydt Orch. (1880) 307, t. 34; Burb. in The Garden 22 (1882) 118; Hemsl. Bot. Chal. Exped. 1 (1885) 199; Warner & Will. Orch. Alb. 6 (1887) t. 277; Rolfe in Gard. Chron. III 5 (1889) 88; Gower in The Garden 35 (1889) 363; Rolfe in l'Orchidoph. 13 (1893) 100; Will. Orch. Grow. Man. ed. 7 (1894) 664.
- Phalaenopsis aphrodite* Lem. in Ill. Hort. 10 (1863) Misc. 14, quoad citat. Rumph., non Reichb. f.
- Phalaenopsis amabilis* Blume var. *grandiflora* Batem. Second Cent. Orch. Pl. (1867) t. 114.
- Phalaenopsis amabilis* Blume var. *rimestadtiana* L. Linden in Lindenia 16 (1902) 35, t. 736; Edit. in Gard. Chron. III 32 (1902) 306, t.; Pauwels in Rev. Hort. Belg. 29 (1903) 88, figs. 22-24; Rolfe in Orch. Rev. 14 (1906) 233, fig. 27, 23 (1915) 254.
- Phalaenopsis rimestadtiana* "Argus" in Orch. Rev. 13 (1905) 260, in textu, *nomen*; Rolfe in Orch. Rev. 25 (1917) 151, in textu, *nomen*.

Although the species proper does not occur in the Philippines, it has been deemed advisable to cite a full synonymy and bibliography of it on account of the confusion arising from the inclusion in it, by Lindley and subsequently by others, of Philippine material. The species is represented in the Philippines by variety *aphrodite* (Reichb. f.) Ames and its variations.

Var. **APHRODITE** (Reichb. f.) Ames Orch. 2 (1908) 226, 5 (1915) 216.

- Phalaenopsis amabilis* Lindl. in Bot. Reg. 24 (1838) t. 34 (excl. syn.); Paxt. Mag. Bot. 5 (1838) 20, 162, 7 (1840) 49, text cut, t. (excl. syn.); Maund Botanist 3 (1839?) t. 133 (excl. syn.); Lem. in Fl. des Serres 1 (1845) 85, text cut, t. 36 (excl. syn.); Blanco Fl. Filip. ed. 2 (1845) 592, ed. 3, 3 (1879) 41; Edit. in Allg. Berl. Gartenz. 13 (1845) 389, 15 (1847) 215 (excl. syn. in both); Hook. in Bot. Mag. 73 (1847) t. 4297 (excl. syn. in part); Blume Rumphia 4 (1848) 52, quoad citat. Lindl.; Lindl. in Gard. Chron. (1848) 39, fig. 2; Reichb. f. in Walp. Ann. 1 (1849) 790, 3 (1852) 561, quoad Philip.; Moore Ill. Orch. Pl. (1857) Phalaenop. 6, t. 1 (excl. syn.); Miq. Fl. Ind. Bat. 3 (1859) 690, quoad citat. Bot. Mag. et quoad Philip.; Gedney in Floral World (1876) 323; Reichb. f. in Gard. Chron. II 5 (1876) 654; Burb. in The Garden 22 (1882) 119; Naves Novis. App. (1882) 243; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Linden & Rodigas in Lindenia 2 (1886) 65, t. 79 (excl. syn. in parte); Gower in The Garden 35 (1889) 362, fig. 363; Roebelen in Gard. Chron. III 7 (1890) 459, in textu; Will. Orch. Grow. Man. ed. 7 (1894) 660, quoad bibliog. in parte et quoad Philip.; Merr. in Govt. Lab. Publ. (Philip.) 8 (1903) 57, 174, 27 (1905) 84; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 549, quoad bibliog. in parte et quoad Philip.; Ames Orch. 1 (1905) 101, quoad spec. (excl. syn. et descr.); Merr. Sp. Blancoanae (1918) 115, non Blume.
- Phalaenopsis aphrodite* Reichb. f. in Hamb. Gartenz. 18 (1862) 35, Xen. Orch. 2 (1862) 6; Duchartre in Journ. Soc. Imp. et Centr. Hort. Par. 8 (1862) 730; Lem. in Ill. Hort. 10 (1863) Misc. 14 (excl. citat. Rumph.); Rolfe in Gard. Chron. II 26 (1886) 212,

fig. 43 B; Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1889) 211, fig. 230; Veitch Man. Orch. Pl. pt. 7 (1891) 24, t.; Stein Orchideenb. (1892) 505, fig. 155; Rolfe in Orch. Rev. 2 (1894) 209, fig. 30, 8 (1900) 135, 136, fig. 20, 13 (1905) 232, fig. 52, 25 (1917) 151, in textu; Cogn. in Dict. Icon. Orch. (1898) Phalaenop. t. 1; Edit. in Sem. Hort. 4 (1900) 392, fig. 132; Rolfe in Forbes & Hemsl. in Journ. Linn. Soc. 36 (1903) 34; Matsum. Ind. Pl. Jap. 2 (1905) 257; Ames in Philip. Journ. Sci. 2 (1907) Bot. 336; Schltr. Orchideen (1914) 544 (excl. syn. *sanderiana*); Schltr. in Fedde Repert. Beih. hefte 4 (Orch. Sino-Jap. Prodr.) (1919) 277 (excl. syn. *sanderiana*).

Phalaenopsis ambigua Reichb. f. in Hamb. Gartenz. 18 (1862) 35, Xen. Orch. 2 (1862) 6; Duchartre in Journ. Soc. Imp. et Centr. Hort. Par. 8 (1862) 311, 731; Lem. in Ill. Hort. 10 (1863) Misc. 13; Naves Novis. App. (1882) 243; Rolfe in Gard. Chron. II 26 (1886) 212.

Phalaenopsis amabilis Blume var. *ambigua* Burb. in The Garden 22 (1882) 119.

Luzon (Ifugao, Rizal, Tayabas, Sorsogon), Samar, Palawan, Bancalan, Mindanao (Zamboanga, Lanao, Cotabato, Davao), Lumbukan. Fairly common and widely distributed in the Philippines. Epiphyte at sea level to 300 m altitude. No attempt has been made to segregate the varieties or forms in the distribution above given. In China and Japan.

The following variations of variety *aphrodite* have been noted and published as species, varieties, subvarieties, or forms, but are of little scientific value, as they are largely based on color and size:

Subvar. **DAYANA** (Hort. ex Warner & Will.) Ames Orch. 2 (1908) 227.

Phalaenopsis amabilis Lindl. var. *dayana* Hort. ex Warner & Will. Orch. Alb. 1 (1882) t. 11; Gower in The Garden 35 (1889) 363; Will. Orch. Grow. Man. ed. 7 (1894) 661.

Phalaenopsis aphrodite Reichb. f. var. *dayana* Veitch Man. Orch. Pl. pt. 7 (1891) 24; Stein Orchideenb. (1892) 505.

Subvar. **ERUBESCENS** (Burb.) Ames Orch. 2 (1908) 227.

Phalaenopsis amabilis Lindl. var. *erubescens* Burb. in The Garden 22 (1882) 119.

Subvar. **GLORIOSA** (Reichb. f.) Ames Orch. 2 (1908) 227.

Phalaenopsis gloriosa Reichb. f. in Gard. Chron. III 3 (1888) 554; Gower in The Garden 35 (1889) 362, t. 697; Will. Orch. Grow. Man. ed. 7 (1894) 664; Rolfe in Orch. Rev. 8 (1900) 136, in textu, 13 (1905) 228.

Phalaenopsis aphrodite Reichb. f. var. *gloriosa* Veitch Man. Orch. Pl. pt. 7 (1891) 25; Stein Orchideenb. (1892) 505.

Sulu Archipelago.

Subvar. **SANDERIANA** (Reichb. f.) Ames Orch. 2 (1908) 228.

Phalaenopsis sanderiana Reichb. f. in Flora 65 (1882) 466, in Gard. Chron. II 19 (1883) 607, 656, 20 (1883) 110; Burb. in The Garden 24 (1883) 270, t. 407; Edit. in l'Orchidoph. 3 (1833) 661; Godefr.-Lebeuf in l'Orchidoph. 5 (1885) 18, t.; Linden & Rodigas in Lindenia 1 (1885) 51, t. 23; Will. Orch. Grow. Man. ed. 6 (1885) 535, ed. 7 (1894) 670; Rolfe in Gard. Chron. II 26 (1886)

212; Warner & Will. Orch. Alb. 5 (1886) *t.* 209; Gower in The Garden 35 (1889) 363; Roebelen in Gard. Chron. III 7 (1890) 459, in textu; Stein Orchideenb. (1892) 511; Kränzl. in Sander Reichenbachia II 2 (1894) 41, *t.* 68 (upper fig.); Rolfe in Orch. Rev. 8 (1900) 133, 15 (1907) 185, *fig.* 22; Cogn. in Dict. Icon. Orch. (1903) Phalaenop. Hybr. *t.* 2; Rolfe in l'Orchidoph. 13 (1903) 101; G. Wilson in Orch. World 6 (1916) 266.

Forma ALBA (Veitch) Ames Orch. 2 (1908) 228.

Phalaenopsis sanderiana Reichb. f. subvar. *alba* Veitch Man. Orch. Pl. pt. 7 (1891) 35, *text cut.*

Phalaenopsis sanderiana var. *alba* Stein Orchideenb. (1892) 511; G. Wilson in Orch. World 6 (1916) 266, *text cut.*

Phalaenopsis sanderiana Reichb. f. forma *alba* Will. Orch. Grow. Man. ed. 7 (1894) 670.

Forma MARMORATA (Reichb. f.) Ames Orch. 2 (1908) 228.

Phalaenopsis sanderiana Reichb. f. var. *marmorata* Reichb. f. in Gard. Chron. II 20 (1883) 812; Rolfe in Gard. Chron. II 26 (1886) 212; Stein Orchideenb. (1892) 511; Will. Orch. Grow. Man. ed. 7 (1894) 670.

Phalaenopsis sanderiana Reichb. f. subvar. *marmorata* Veitch Man. Orch. Pl. pt. 7 (1891) 35.

Forma PUNCTATA (O'Brien) Ames Orch. 2 (1908) 229.

Phalaenopsis sanderiana Reichb. f. var. *punctata* O'Brien in Gard. Chron. III 7 (1890) 78; Stein Orchideenb. (1892) 511; Will. Orch. Grow. Man. ed. 7 (1894) 670.

Phalaenopsis sanderiana Reichb. f. subvar. *punctata* Veitch Man. Orch. Pl. pt. 7 (1891) 35.

PHALAEOPSIS BOXALLII Reichb. f. in Gard. Chron. II 19 (1883) 274; Rolfe in Gard. Chron. II 26 (1886) 276; Veitch Man. Orch. Pl. pt. 7 (1891) 26; Ames Orch. 5 (1915) 216.

PHALAEOPSIS EQUESTRIS (Schauer) Reichb. f. in Linnaea 22 (1849) 864; Lindl. in Paxt. Flow. Gard. 2 (1852) 174; Reichb. f. in Walp. Ann. 3 (1852) 562, 6 (1864) 860; Miq. Fl. Ind. Bat. 3 (1859) 690; Reichb. f. in Hamb. Gartenz. 16 (1860) 116; Duchartre in Journ. Soc. Imp. et Centr. Hort. Par. 6 (1860) 869, 8 (1862) 727; Reichb. f. Xen. Orch. 2 (1862) 4; Naves Novis. App. (1882) 242; Ames Orch. 2 (1908) 229, 5 (1915) 216.

Stauroglottis equestris Schauer in Nov. Act. Acad. Nat. Cur. 19 Suppl. 1 (1843) 432.

Phalaenopsis rosea Lindl. in Gard. Chron. (1848) 671, *text cut*; Paxt. Mag. Bot. 16 (1849) 60, 189, *text cut*; Lindl. in Paxt. Flow. Gard. 2 (1852) 173, *t.* 72; Reichb. f. in Bot. Zeit. 10 (1852) 673; Moore Ill. Orch. Pl. (1857) Phalaen. 7; Hook. in Bot. Mag. 86 (1860) *t.* 5212; Lem. in Jard. Fleur. 3 (1853) *t.* 233, in Ill. Hort. 10 (1863) Misc. 11; Van Houtte in Fl. des Serres 16 (1866) *t.* 1645; Jennings Orch. (1875) *t.* 27; Burb. in The Garden 22 (1822) 119 (excl. var.); Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Rolfe in Gard. Chron. II 26 (1886) 276; Warner &

Will. Orch. Alb. 6 (1887) t. 268; Veitch Man. Orch. Pl. pt. 7 (1891) 34; Ames Orch. 1 (1905) 102.

Phalaenopsis esmeralda Cogn. in Dict. Icon. Orch. (1898) Phalaen. t. 3, non Reichb. f.

Luzon (Ilocos Norte, Bataan, Rizal, Laguna, Tayabas, Camarines), Panay (Capiz), Bohol, Mindanao (Camiguin Island, Lanao). A common and widely distributed species. Epiphyte at sea level to 300 m altitude.

Var. **LEUCASPIS** Reichb. f. in Gard. Chron. II 15 (1881) 688, in l'Orchidoph. 1 (1881) 50; Ames Orch. 2 (1908) 230.

Phalaenopsis rosea Lindl. var. *leucaspis* Rolfe in Gard. Chron. II 26 (1886) 276; Veitch Man. Orch. Pl. pt. 7 (1891) 34.

Var. **LEUCOTANTHE** Reichb. f. in l'Orchidoph. 3 (1883) 490; Ames Orch. 2 (1908) 230.

PHALAENOPSIS FASCIATA Reichb. f. in Gard. Chron. II 18 (1882) 134; Rolfe in Orch. Rev. 13 (1905) 225; Ames Orch. 5 (1915) 217.

No material of this species has been seen. Reichenbach gave the origin of the plant as Philippines; imported by Messrs. Hugh Low & Co. The origin seems open to some doubt. Reichenbach states that the species is near *Phalaenopsis lueddemanniana* Reichb. f.

PHALAENOPSIS FUSCATA Reichb. f. in Gard. Chron. II 2 (1874) 6; Rolfe in Orch. Rev. 13 (1905) 226; Ames Orch. 5 (1915) 216.

Phalaenopsis denisiana Cogn. in Gard. Chron. III 26 (1899) 82; Cogn. in Dict. Icon. Orch. (1899) Phalaenop. t. 6.

The origin of *P. fuscata* is given as the Malay Peninsula, that of *P. denisiana* as the Philippines. No material of this species has been seen.

PHALAENOPSIS LEUCORRHODA Reichb. f. in Gard. Chron. II 3 (March, 1875) 301, 366; W. G. Sm. in Floral Mag. 14 (June, 1875) 41, t. 166; Naves Novis. App. (1882) 243; Rolfe in Gard. Chron. II 26 (1886) 212; Veitch Man. Orch. Pl. pt. 7 (1891) 46; Stein Orchideenb. (1892) 507; Will. Orch. Grow. Man. ed. 7 (1894) 667; Rolfe in Orch. Rev. 13 (1905) 232, fig. 54.

This is a supposed natural hybrid between *Phalaenopsis amabilis* (Linn.) Blume var. *aphrodite* (Reichb. f.) Ames and *P. schilleriana* Reichb. f. Its origin is Philippine, but no specimens have been seen.

PHALAENOPSIS LINDENII Loher in Journ. des Orch. 6 (1895) 103; Rolfe in Orch. Rev. 13 (1905) 230, 15 (1907) 296; Ames in Philip. Journ. Sci. 4 (1909) Bot. 599, Orch. 5 (1915) 217; G. Wilson in Orch. Rev. 30 (1922) 354.

Luzon (Benguet), *Lyon 39*, *F. B. 5121 Curran*. Epiphyte.

PHALAENOPSIS LUEDEDEMANNIANA Reichb. f. in Bot. Zeit. 23 (1865) 146, in Gard. Chron. (1865) 434; Moore in Flor. & Pomol. (1865) 257, t. 254; Lem. in Ill. Hort. 12 (1865) Misc. 31; Edit. in Proc. Roy. Hort. Soc. 5 (1865) 137; Otto in Hamb. Gartenz. 21 (1865) 470; G. B. in Belg. Hort. 15 (1865) 229; Carr. in Rev. Hort. 44 (1872) 390, t.; Rolfe in Gard. Chron. II 26 (1886) 277, in *Lindenia* 2 (1886) 95, t. 94, 8 (1892) 63, t. 366; Veitch Man. Orch. Pl. pt.

7 (1891) 30, *text cut*; Cogn. in Dict. Icon. Orch. (1899) *Phalaen. t. 9*; Ames Orch. 2 (1908) 230, 5 (1915) 217.

Phalaenopsis luddemaniai Boxall ex Naves Novis. App. (1882) 243, sphalm.

Luzon (Apayao, Nueva Vizcaya, Bataan, Bulacan, Laguna, Tayabas, Sorsogon), Polillo, Leyte, Palawan, Mindanao (Bukidnon). A common and widely distributed species, epiphyte at 30 to 60 m altitude.

A number of varieties of this species have been characterized from cultivated specimens.

Var. **DELICATA** Reichb. f. in Gard. Chron. (1865) 434; Burb. in The Garden 22 (1882) 119; Rolfe in Gard. Chron. II 26 (1886) 277, in Lindenia 8 (1892) 63, sub *t. 366*; Ames Orch. 2 (1908) 231.

Phalaenopsis lueddemanniana Batem. in Bot. Mag. 91 (1865) *t. 5523*; Van Houtte in Fl. des Serres 16 (1865) 53, *t. 1636*; Batem. Second Cent. Orch. Pl. (1867) *t. 133*, non Reichb. f.

Phalaenopsis lueddemanniana subvar. *delicata* Veitch Man. Orch. Pl. pt. 7 (1891) 30.

Var. **HIEROGLYPHICA** Reichb. f. in Gard. Chron. III 2 (1887) 586; Edit. in l'Orchidoph. 9 (1889) 197; Rolfe in Lindenia 8 (1892) 63, sub *t. 366*; Ames Orch. 2 (1908) 231.

Phalaenopsis lueddemanniana Reichb. f. subvar. *hieroglyphica* Veitch Man. Orch. Pl. pt. 7 (1891) 31.

Var. **OCHRACEA** Reichb. f. in Gard. Chron. (1865) 438; Carr. in Rev. Hort. 44 (1872) 391, *fig. A*; Burb. in The Garden 22 (1882) 119; Rolfe in Gard. Chron. II 26 (1886) 277, in Lindenia 8 (1892) 63, sub *t. 366*; Ames Orch. 2 (1908) 232.

Phalaenopsis lueddemanniana subvar. *ochracea* Veitch Man. Orch. Pl. pt. 7 (1891) 31.

Lyon 48.

Var. **PULCHRA** Reichb. f. in Gard. Chron. II 4 (1875) 36; Burb. in The Garden 22 (1882) 119; Rolfe in Gard. Chron. II 26 (1886) 277, in Lindenia 8 (1892) 64, sub *t. 366*; Ames Orch. 2 (1908) 232.

Phalaenopsis lueddemanniana Reichb. f. subvar. *pulchra* Veitch Man. Orch. Pl. pt. 7 (1891) 31.

PHALAEOPSIS MARIAE Burb. in Warner & Will. Orch. Alb. 2 (1883) *t. 80* et sub *t. 87*; Rolfe in Gard. Chron. II 26 (1886) 277; Hook. f. in Bot. Mag. 113 (1887) *t. 6964*; Veitch Man. Orch. Pl. pt. 7 (1891) 32; Ridl. in Journ. Linn. Soc. 31 (1896) 292; Ames in Philip. Journ. Sci. 8 (1913) Bot. 434, Orch. 5 (1915) 217, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 197.

Luzon (Nueva Vizcaya), Mindanao (Lanao, Cotabato, Bukidnon, Davao), B. S. 11136, 11141 McGregor, 21389 Escritor, M. S. Clemens 626, Weber 220, Loher 6011. Epiphyte up to 700 m altitude in Cotabato. In Borneo.

PHALAEOPSIS MICHOLITZII Rolfe in Gard. Chron. III 8 (1890) 187, in Journ. des Orch. 1 (1890) 198, in Orch. Rev. 13 (1905) 229; Ames Orch. 5 (1915) 217.

Without locality indicated, *Micholitz*.

No technical description of this species has been published. From the material at hand it appears scarcely distinct from *Phalaenopsis lueddemaniana* Reichb. f.

PHALAEOPSIS REICHENBACHIANA Reichb. f. & Sander in Gard. Chron. II 18 (1882) 586; Rolfe in Orch. Rev. 13 (1905) 226; Ames Orch. 5 (1915) 218.

According to Rolfe (in Orch. Rev. l. c.) Micholitz stated that this species is a native of Mindanao. No material has been seen.

PHALAEOPSIS SCHILLERIANA Reichb. f. in Hamb. Gartenz. 16 (1860) 115, 116; Lindl. in Gard. Chron. (1860) 216, (1861) 617, (1862) 95; Duchartre in Journ. Soc. Imp. et Centr. Hort. Par. 7 (1861) 845, 8 (1862) 609, 730, *t. 17*; Warner Sel. Orch. Pl. I (1861) *t. 1*; Reichb. f. Xen. Orch. 2 (1862) 1, 5, *t. 101*; Lem. in Ill. Hort. 10 (1863) Misc. 13, *t. 348*; Lescuyer in Horticulteur Fr. (1863) 127, *t. 11*; Van Houtte in Fl. des Serres 15 (1864) 115, *tt. 1559-60*, 21 (1875) 142, *text cut*; Batem. in Bot. Mag. 91 (1865) *t. 5530*, Second Cent. Orch. Pl. (1867) *t. 171*; Regel in Gartenfl. 17 (1868) 161, *t. 581*, 36 (1887) 57, *fig. 21*; Jennings Orch. (1875) *t. 15*; Edit. in Gard. Chron. II 4 (1875) 169, *fig. 34*, II 12 (1879) 301, *fig. 47 D-F*, III 3 (1888) 528, *fig. 72*; Gedney in Floral World (1876) 321, 323, *t.*; Puydt Orch. (1880) 309, *t. 35*; Burb. in The Garden 22 (1882) 118, 119, *t. 348, fig. 2*; Naves Novis. App. (1882) 243; Edit. in l'Orchidoph. 4 (1884) 279, *t. 212*; Rodigas in Ill. Hort. 35 (1888) 57, *t. 56*; Pfütz. in Engl. & Prantl Pflanzenfam. 2, Abt. 6 (1889) 211, *fig. 227 C*; Rolfe in Lindenia 5 (1889) *t. 227*; Roebelen in Gard. Chron. III 7 (1890) 459; Veitch Man. Orch. Pl. pt. 7 (1891) 36, *text cut*; Rivois in Journ. des Orch. 3 (1892) 146, *fig. 18*; Rolfe in Orch. Rev. 2 (1894) 209, *fig. 21*, 8 (1900) 134-136, *fig. 21*, 13 (1905) 232, *fig. 53*; Grignan in Sem. Hort. 2 (1898) 46, *fig. 214*; Cogn. in Dict. Icon. Orch. (1899) *Phalaen. t. 10*; L. Linden in Sem. Hort. 4 (1900) 424, *fig. 143*; Ames Orch. 2 (1908) 232, 5 (1915) 218.

Luzon (Tayabas), *Lyon 54, Merrill 4018, 4024*. This is one of the most highly prized of the Philippine orchids and is extensively exported. A number of forms have been described from cultivated specimens.

Var. **ADVENA** Reichb. f. in Gard. Chron. II 23 (1885) 174; Edit. in l'Orchidoph. 5 (1885) 200; Ames Orch. 2 (1908) 234.

Var. **IMMACULATA** Reichb. f. in Gard. Chron. II 3 (1875) 429; Rolfe in Gard. Chron. II 26 (1888) 212; Ames Orch. 2 (1908) 234.

Phalaenopsis schilleriana Reichb. f. var. *delicata* Dean in Floral Mag. 16 (1877) *t. 257*.

Phalaenopsis schilleriana Reichb. f. subvar. *immaculata* Veitch Man. Orch. Pl. pt. 7 (1891) 37.

Var. **MAJOR** Rolfe in Gard. Chron. II 26 (1886) 212; Ames Orch. 2 (1908) 234.

Var. **PURPUREA** Hort. ex Rolfe in Journ. des Orch. 2 (1892) 358; Garnier in Journ. des Orch. 5 (1894) 9; O'Brien in Gard. Chron. III 15 (1894) 266; Ames Orch. 2 (1908) 234.

Var. **SPLENDENS** Hort. ex Carr. in Rev. Hort. 58 (1886) 396, *t.*; Warner Sel. Orch. Pl. III (?) *t.* 5; Ames Orch. 2 (1908) 234.

Var. **VESTALIS** Reichb. f. in Gard. Chron. II 17 (1882) 330; Edit. in l'Orchidoph. 2 (1882) 291; Rolfe in Gard. Chron. II 26 (1886) 212; Edit. in Gard. Chron. III 3 (1888) 75; Ames Orch. 2 (1908) 235.

Phalaenopsis schilleriana Reichb. f. var. *alba* Roebelen in Gard. Chron. III 7 (1890) 459; Rolfe in l'Orchidoph. 13 (1893) 101, in Orch. Rev. 8 (1900) 134, in textu.

Phalaenopsis schilleriana Reichb. f. subvar. *vestalis* Veitch Man. Orch. Pl. pt. 7 (1891) 37.

PHALAEOPSIS SCHILLERIANA-STUARTIANA Rolfe in Orch. Rev. 8 (1900) 94, 113, *fig. 18*, 13 (1905) 230, *fig. 50*; G. Wilson in Orch. Rev. 30 (1922) 77, *text cut*.

Said to be a natural hybrid of Philippine origin.

PHALAEOPSIS STUARTIANA Reichb. f. in Gard. Chron. II 16 (1881) 748, 753, *fig. 149*; Hook. f. in Bot. Mag. 108 (1882) *t. 6622*; Rodigas in Ill. Hort. 31 (1884) 175, *t. 540*; Rolfe in Gard. Chron. II 26 (1886) 212; Warner & Will. Orch. Alb. 5 (1886) *t. 237*; Reichb. f. in Sander Reichenbachia I (1888) 95, *t. 42*; Veitch Man. Orch. Pl. pt. 7 (1891) 39, *t.* (opp. p. 29); Cogn. in Dict. Icon. Orch. (1898) Phalaen. *t. 2*; Ames Orch. 5 (1915) 218.

Philippines, without locality, *Lyon s. n.*

PHALAEOPSIS VEITCHIANA Reichb. f. in Gard. Chron. (1872) 935; Burb. in Floral Mag. 15 (1876) *t. 213*; Veitch Man. Orch. Pl. pt. 7 (1898) 47; Ames Orch. 5 (1915) 218; G. Wilson in Orch. Rev. 30 (1922) 346.

EXCLUDED OR SYNONYMOUS SPECIES

PHALAEOPSIS AMABILIS Lindl. in Bot. Reg. 24 (1838) *t. 34* (excl. syn.); Paxt. Mag. Bot. 5 (1838) 20, 162, 7 (1840) 49, *text cut* (excl. syn.); Maund Botanist 3 (1839?) *t. 133* (excl. syn.); Lem. in Fl. des Serres 1 (1845) 85, *text cut*, *t. 36* (excl. syn.); Blanco Fl. Filip. ed. 2 (1845) 592, ed. 3, 3 (1879) 41; Edit. in Allg. Berl. Gartenz. 13 (1845) 389, 15 (1847) 215 (excl. syn. in both); Hook. in Bot. Mag. 73 (1847) *t. 4297* (excl. syn. in parte); Blume Rumphia 4 (1848) 52, quoad citat. Lindl.; Lindl. in Gard. Chron. (1848) 39, *fig. 2*; Reichb. f. in Walp. Ann. 1 (1849) 790, 3 (1852) 561, quoad Philip.; Moore Ill. Orch. Pl. (1857) Phalaenop. 6, *t. 1* (excl. syn.); Miq. Fl. Ind. Bat. 3 (1859) 690, quoad citat. Bot. Mag. et quoad Philip.; Gedney in Floral World (1876) 323; Reichb. f. in Gard. Chron. II 5 (1876) 654; Burb. in The Garden 22 (1882) 119; Naves Novis. App. (1882) 243; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Linden &

Rodigas in *Lindenia* 2 (1886) 65, *t.* 79 (excl. syn. in parte); Gower in *The Garden* 35 (1889) 362, *fig.* 363; Roebelen in *Gard. Chron.* III 7 (1890) 459, in textu; Will. *Orch. Grow. Man.* ed. 7 (1894) 660, quoad bibliog. in parte et quoad Philip.; Merr. in *Govt. Lab. Publ. (Philip.)* 8 (1903) 57, 174, 27 (1905) 84; J. J. Sm. *Fl. Buitenz.* 6 (*Orch. Jav.*) (1905) 549, quoad bibliog. in parte et quoad Philip.; Ames *Orch.* 1 (1905) 101, quoad spec. (excl. syn. et descr.); Merr. *Sp. Blancoanae* (1918) 115, non Blume = *Phalaenopsis amabilis* Blume var. *aphrodite* (Reichb. f.) Ames.

PHALAEOPSIS AMBIGUA Reichb. f. in *Hamb. Gartenz.* 18 (1862) 35, *Xen. Orch.* 2 (1862) 6; Duchartre in *Journ. Soc. Imp. et Centr. Hort. Par.* 2 (1862) 311, 731; Lem. in *Ill. Hort.* 10 (1863) Misc. 13; Naves *Novis. App.* (1882) 243; Rolfe in *Gard. Chron.* II 26 (1886) 212 = *Phalaenopsis amabilis* Blume var. *aphrodite* (Reichb. f.) Ames.

PHALAEOPSIS APHRODITE Reichb. f. in *Hamb. Gartenz.* 18 (1862) 35, *Xen. Orch.* 2 (1862) 6; Duchartre in *Journ. Soc. Imp. et Centr. Hort. Par.* 8 (1862) 730; Lem. in *Ill. Hort.* 10 (1863) Misc. 14 (excl. citat. Rumph.); Rolfe in *Gard. Chron.* II 26 (1886) 212, *fig.* 43B; Pfitz. in *Engl. & Prantl Nat. Pflanzenfam.* 2, Abt. 6 (1889) 211, *fig.* 230; Veitch *Man. Orch. Pl.* pt. 7 (1891) 24, *t.*; Stein *Orchideenb.* (1892) 505, *fig.* 155; Rolfe in *Orch. Rev.* 2 (1894) 209, *fig.* 20, 8 (1900) 135, 136, *fig.* 20, 13 (1905) 232, *fig.* 52, 25 (1917) 151, in textu; Cogn. *Dict. Icon. Orch.* (1898) *Phalaenop. t.* 1; Edit. in *Sem. Hort.* 4 (1900) 392, *fig.* 132; Rolfe in *Forbes & Hemsl. in Journ. Linn. Soc.* 36 (1903) 34; Matsum. *Ind. Pl. Jap.* 2 (1905) 257; Ames in *Philip. Journ. Sci.* 2 (1907) Bot. 336; Schltr. *Orchideen* (1914) 544 (excl. syn. *sanderiana*); Schltr. in *Fedde Repert. Beihefte* 4 (*Orch. Sino-Jap. Prodr.*) (1919) 277 (excl. syn. *sanderiana*) = *Phalaenopsis amabilis* Blume var. *aphrodite* (Reichb. f.) Ames.

PHALAEOPSIS CASTA Boxall ex Naves *Novis. App.* (1882) 243, *nomen*.

Boxall probably had in mind *Phalaenopsis casta* Reichb. f., which is now considered a form or variety of *P. leucorrhoda* Reichb. f.

PHALAEOPSIS CORNU-CERVI Blume apud Naves *Novis. App.* (1882) 243.

PHALAEOPSIS DELICIOSA Reichb. f. apud Naves *Novis. App.* (1882) 243.

PHALAEOPSIS DENISIANA Cogn. in *Gard. Chron.* III 26 (1899) 82; Cogn. in *Dict. Icon. Orch.* (1899) *Phalaenop. t.* 6 = *Phalaenopsis fuscata* Reichb. f.

PHALAEOPSIS DEVRIESIANA Reichb. f. apud Naves *Novis. App.* (1882) 243.

PHALAEOPSIS ESMERALDA Cogn. in *Dict. Icon. Orch.* (1898) *Phalaenop. t.* 3, non Reichb. f. = *Phalaenopsis equestris* (Schauer) Reichb. f.

PHALAEOPSIS GLORIOSA Reichb. f. in *Gard. Chron.* III 3 (1888) 554; Gower in *The Garden* 35 (1889) 362, *t.* 697; Will. *Orch. Grow. Man.* ed. 7 (1894) 664; Rolfe in *Orch. Rev.* 8 (1900) 136, in textu,

13 (1905) 228 = *Phalaenopsis amabilis* (Linn.) Blume subvar. *gloriosa* (Reichb. f.) Ames.

PHALAENOPSIS HEBE Reichb. f. apud Naves Novis. App. (1882) 242.

PHALAENOPSIS INTERMEDIA Lindl. apud Naves Novis. App. (1882) 243.

PHALAENOPSIS INTERMEDIA Lindl. var. **PORTEI** Reichb. f. in Bot. Zeit. 21 (1863) 128; W. G. Sm. in Floral Mag. 14 (1875) t. 162; Naves Novis. App. (1882) 243; Warner Sel. Orch. Pl. II (1865-75) t. 2; Veitch Man. Orch. Pl. pt. 7 (1891) 45; Sander Reichenbachia II 2 (1894) 43, t. 68 (lower fig.).

Phalaenopsis intermedia Lindl. var. *brymeriana* Reichb. f. in Gard. Chron. II 5 (1876) 366; Dean in Floral Mag. 16 (1877) t. 263; Veitch Man. Orch. Pl. pt. 7 (1891) 44; Warner & Will. Orch. Alb. 9 (1891) t. 415.

Phalaenopsis portei Reichb. f. in Gard. Chron. II 5 (1876) 370, figs. 70, 71; Outram in Gard. Chron. II 15 (1881) 181.

There seems no valid authority for including this in the list of Philippine species.

PHALAENOPSIS LOWII Reichb. f. apud Naves Novis. App. (1882) 243.

PHALAENOPSIS LUEDDEMANNIANA Batem. in Bot. Mag. 91 (1865) t. 5523; Van Houtte in Fl. des Serres 16 (1865) 53, t. 1636; Batem. Second Cent. Orch. Pl. (1867) t. 133, non Reichb. f. = *Phalaenopsis lueddemannia* Reichb. f. var. *delicata* Reichb. f.

PHALAENOPSIS LUDDEMANII Boxall ex Naves Novis. App. (1882) 243, sphalm = *Phalaenopsis lueddemanniana* Reichb. f.

PHALAENOPSIS PALLENS (Lindl.) Reichb. f. in Walp. Ann. 6 (1864) 932; Rolfe in Gard. Chron. II 26 (1886) 276, in Orch. Rev. 8 (1900) 327, 13 (1905) 226.

Trichoglottis pallens Lindl. in Journ. Hort. Soc. 5 (1850) 34, in Paxt. Flow. Gard. 1 (1850) 15.

Stauroopsis pallens Reichb. f. in Hamb. Gartenz. 16 (1860) 117, Xen. Orch. 2 (1862) 7; Naves Novis. App. (1882) 243.

For many years this species was ascribed to the Philippines. It does not occur in the Archipelago, and Rolfe, l. c., has shown that the type could not have come from the Philippines.

PHALAENOPSIS PARISHII Reichb. f. apud Naves Novis. App. (1882) 243.

PHALAENOPSIS ROSEA Lindl. in Gard. Chron. (1848) 671, text cut; Paxt. Mag. Bot. 16 (1849) 60, 189, text cut; Lindl. in Paxt. Flow. Gard. 2 (1852) 173, t. 72; Reichb. f. in Bot. Zeit. 10 (1852) 673; Moore Ill. Orch. Pl. (1857) Phalaen. 7; Hook. in Bot. Mag. 86 (1860) t. 5212; Lem. in Jard. Fleur. 3 (1853) t. 283, in Ill. Hort. 10 (1863) Misc. 11; Van Houtte in Fl. des Serres 16 (1866) t. 1645; Jennings Orch. (1875) t. 27; Burb. in The Garden 22 (1882) 119 (excl. var.); Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270; Rolfe in Gard. Chron. II 26 (1886) 276; Warner & Will. Orch. Alb. 6 (1887) t. 268; Veitch Man. Orch. Pl. pt. 7 (1891) 34; Ames Orch. 1 (1905) 102 = *Phalaenopsis equestris* (Schauer) Reichb. f.

PHALAENOPSIS SANDERIANA Reichb. f. in *Flora* 65 (1882) 466, in *Gard. Chron.* II 19 (1883) 607, 656, 20 (1883) 110; Burb. in *The Garden* 24 (1883) 270, *t. 407*; Edit. in *l'Orchidoph.* 3 (1883) 661; Godefr.-Lebeuf in *l'Orchidoph.* 5 (1885) 18, *t.*; Linden & Rodigas in *Lindenia* 1 (1885) 51, *t. 23*; Will. *Orch. Grow. Man.* ed. 6 (1885) 535, ed. 7 (1894) 670; Rolfe in *Gard. Chron.* II 26 (1886) 212; Warner & Will. *Orch. Alb.* 5 (1886) *t. 209*; Gower in *The Garden* 35 (1889) 363; Roebelen in *Gard. Chron.* III 7 (1890) 459, in *textu*; Stein *Orchideenb.* (1892) 511; Kränzl. in *Sander Reichenbachia* II 2 (1894) 41, *t. 68* (upper fig.); Rolfe in *Orch. Rev.* 8 (1900) 133, 15 (1907) 185, *fig. 22*; Cogn. in *Dict. Icon. Orch.* (1903) *Phalaenop. hybr. t. 2*; Rolfe in *l'Orchidoph.* 13 (1903) 101; G. Wilson in *Orch. World* 6 (1916) 266 = *Phalaenopsis amabilis* (Linn.) Blume subvar. *sanderiana* (Reichb. f.) Ames.

PHALAENOPSIS SUMATRANA Korth. apud Naves Novis. App. (1882) 242.

PHALAENOPSIS VIOLACEA Teijsm. & Binn. apud Naves Novis. App. (1882) 243.

80. AERIDES Loureiro

AERIDES AUGUSTIANUM Rolfe in *Lindenia* 5 (1889) 39, *t. 210*, in *Gard. Chron.* III 7 (1890) 9; Veitch *Man. Orch. Pl.* pt. 7 (1891) 65; Ames *Orch.* 5 (1915) 218.

Discovered in the Philippines by Auguste Linden; no specimen seen.

AERIDES JARCKIANUM Schltr. in *Orchis* 9 (1915) 53 *t. 5, figs. 15-21*.

Luzon, *Jarck*. No specimen seen. Reported to have been collected near the "Lagune von Manila."

AERIDES LAWRENCIAE Reichb. f. in *Gard. Chron.* II 20 (1883) 307, 368, 460; Warner & Will. *Orch. Alb.* 6 (1887) *t. 270*; Gower in *The Garden* 35 (1889) 484, *t. 702*; Veitch *Man. Orch. Pl.* pt. 7 (1891) 71, *text cut*; Kränzl. *Xen. Orch.* 3 (1892) *t. 252, fig. 5*; Ames *Orch.* 2 (1908) 249, 5 (1915) 218, in *Philip. Journ. Sci.* 4 (1909) Bot. 600; G. Wilson in *Orch. Rev.* 31 (1923) 19.

Mindanao (Davao), *R. F. Black s. n., Elmer 10787, Lyon Y* (without locality).

Var. **AMESIANUM** Sander ex Kränzl. in *Gard. Chron.* III 10 (1891) 393; Kränzl. *Xen. Orch.* 3 (1892) 94, *t. 252, figs. 1-4, 7-10*; Ames *Orch.* 2 (1908) 250; G. Wilson in *Orch. Rev.* 31 (1923) 19, in *textu*.

Var. **SANDERIANUM** (Reichb. f.) Sander ex Kränzl. *Xen. Orch.* 3 (1892) 95, in *textu*, *t. 252, fig. 6* (as var. *sanderiana*); Ames *Orch.* 2 (1908) 250; G. Wilson in *Orch. Rev.* 31 (1923) 19, in *textu*.

Aërides sanderianum Reichb. f. in *Gard. Chron.* II 22 (1884) 134; O'Brien in *Gard. Chron.* II 22 (1884) 460.

Aërides lawrenciae subvar. *sander's* Veitch *Man. Orch. Pl.* pt. 7 (1891) 72, *text cut*.

AERIDES ODORATUM Lour. *Fl. Cochinch.* 2 (1790) 525; Sw. in *Kongl. Svensk. Vet.-Akad. Nya Handl.* 21 (1800) 244; Lindl. *Gen. & Sp. Orch.* (1833) 239; Knowles & Westc. *Floral Cab.* 2 (1838) *t. 75*; Hook. in *Bot. Mag.* 71 (1845) *t. 4139*; Regel in *Gartenfl.* 6

(1857) *t.* 272; Naves Novis. App. (1882) 239; Hook. f. Fl. Brit. Ind. 6 (1890) 47; Veitch Man. Orch. Pl. pt. 7 (1891) 76; Grant Orch. Burma (1895) 272; Ridl. in Journ. Linn. Soc. 31 (1896) 294, 32 (1896) 376; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 212, *t.* 282; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 581, Fig.-Atlas (1912) *fig.* 436; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 143; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 179; Ames Orch. 2 (1908) 250, 5 (1915) 218, 6 (1920) 220, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 197.

Aërides cornutum Roxb. Hort. Beng. (1814) 63, *nomen*; Roxb. Fl. Ind. ed. 2, 3 (1832) 472; Lindl. in Bot. Reg. 18 (1832) *t.* 1485.

Aërides virens Lindl. in Bot. Reg. 29 (1843) Misc. 41, 30 (1844) *t.* 41; Miq. Fl. Ind. Bat. 3 (1859) 695; Naves Novis. App. (1882) 239; Warner & Will. Orch. Alb. 4 (1885) *t.* 160; Veitch Man. Orch. Pl. pt. 7 (1891) 81.

Aërides suaveolens Blume Rumphia 4 (1848) 53, *t.* 193, *fig.* 1, Mus. Bot. Lugd.-Bat. 1 (1849) 112, non Roxb.

Aërides suavisimum Lindl. in Journ. Hort. Soc. 4 (1849) 264, in Paxt. Flow. Gard. 2 (1851-52) 141, *t.* 66; Hook. f. Fl. Brit. Ind. 6 (1890) 47; Veitch Man. Orch. Pl. pt. 7 (1891) 78, *text cut*; Rolfe in Lindenia 7 (1891) 41, *t.* 307; Ceron Cat. Pl. Herb. Manila (1892) 168.

Aërides flavidum Lindl. in Paxt. Flow. Gard. 2 (1851-52) 101.

Aërides reichenbachii J. Linden ex Reichb. f. in Koch. & Fintelm. Wochenschr. 1 (1858) 61; Reichb. f. Xen. Orch. 2 (1862) 11, *t.* 104; Naves Novis. App. (1882) 239; J. Linden in Lindenia 1 (1885) 7, *t.* 1.

Aërides jucundum Reichb. f. in Hamb. Gartenz. 16 (1860) 281; Morr. in Belg. Hort. 26 (1876) 289.

Aërides nobile Warner Sel. Orch. I (1862) *t.* 11; Regel in Gartenfl. 19 (1870) 40; Naves Novis. App. (1882) 239.

Aërides reichenbachii Linden var. *cochinchinense* Hort. ex Reichb. f. in Gard. Chron. II 14 (1880) 620.

Aërides rohanianum Reichb. f. in Gard. Chron. II 21 (1884) 206; Ames Orch. 5 (1915) 219.

Aërides wilsonianum Hort. ex Gard. Chron. II 23 (1885) 705.

Aërides ballantianum Reichb. f. in Gard. Chron. II 24 (1885) 198.

Mindoro, Lyon P. A very variable and widely distributed species.

Many horticultural varieties have been described and some field varieties, usually based on one of the specific names formerly in use. In India from the Himalayas to Sylhet, in Burma, Malay Peninsula, Cochin China, China, Java, and Borneo.

AERIDES QUINQUEVULNERUM Lindl. Sert. Orch. (1838) *t.* 30; Paxt. Mag. Bot. 6 (1839) 167, 260, 8 (1841) 241, *t.* and *text cut*; Reichb. f. in Walp. Ann. 6 (1864) 899; Naves Novis. App. (1882) 239; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271; J. Linden in Lindenia 4 (1888) 15, *t.* 150; Veitch Man. Orch. Pl. pt. 7 (1891) 77; Warner & Will. Orch. Alb. 11 (1897) *t.* 495; Ames Orch. 1 (1905) 104, 2 (1908) 252, 5 (1915) 219; Merr. Sp. Blancoanae (1918) 116.

Aërides maculatum Llanos Frag. Fl. Filip. (1851) 93; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 72, t. 409, non Buch-Ham.

Aërides fenzlianum Reichb. f. in Hamb. Gartenz. 16 (1860) 282.

Aërides thibautianum Reichb. f. in Gard. Chron. (1866) 101.

Luzon (Bataan, Bulacan, Rizal, Cavite, Laguna, Tayabas). A common species in and about Rizal at relatively low altitude.

Var. **FARMERI** (Boxall) Stein Orchideenb. (1892) 63; Ames Orch. 2 (1908) 253.

Aërides farmeri Boxall ex Naves Novis. App. (1882) 239, *nomen*.

Aërides album Sander ex Stein Orchideenb. (1892) 63, in syn.

Var. **MARGINATUM** (Reichb. f.) Stein Orchideenb. (1892) 63; Ames Orch. 2 (1908) 253.

Aërides marginatum Reichb. f. in Gard. Chron. II 23 (1885) 533, 25 (1886) 170.

Var. **PURPURATUM** Reichb. f. in Gard. Chron. II 16 (1881) 230; Ames Orch. 2 (1908) 253.

Var. **SCHADENBERGIANUM** Stein Orchideenb. (1892) 63; Ames Orch. 2 (1908) 253.

Mindanao, *Schadenberg* (fide Stein).

EXCLUDED OR SYNONYMOUS SPECIES

AËRIDES ALBUM Sander ex Stein Orchideenb. (1892) 63, in syn.

Aërides quinquevulnerum Lindl. var. *farmeri* (Boxall) Stein.

AËRIDES CYLINDRICUM Boxall ex Naves Novis. App. (1882) 239, *nomen*.

AËRIDES DIURNUM Teijsm. & Binn. in Nat. Tijdschr. Ned. Ind. 27 (1864) 19 = *Sarcochilus pallidus* (Blume) Reichb. f.

AËRIDES FARMERI Boxall ex Naves Novis. App. (1882) 239, *nomen* = *Aërides quinquevulnerum* Lindl. var. *farmeri* (Boxall) Stein.

AËRIDES FENZLIANUM Reichb. f. in Hamb. Gartenz. 16 (1860) 282 = *Aërides quinquevulnerum* Lindl.

AËRIDES LARPENTA Boxall ex Naves Novis. App. (1882) 239, *nomen*.

AËRIDES MACULATUM Llanos Frag. Pl. Filip. (1851) 93; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 72, t. 409, non Buch-Ham. = *Aërides quinquevulnerum* Lindl.

AËRIDES MARGARITACEUM Boxall ex Naves Novis. App. (1882) 239, *nomen*.

AËRIDES MARGINATUM Reichb. f. in Gard. Chron. II 23 (1885) 533, 25 (1886) 170 = *Aërides quinquevulnerum* Lindl. var. *marginatum* (Reichb. f.) Stein.

AËRIDES MENDELLII Boxall ex Naves Novis. App. (1882) 239, *nomen*.

AËRIDES NOBILE Warner Sel. Orch. Pl. I (1862) t. 11; Regel in Gartenfl. 19 (1870) 40; Naves Novis. App. (1882) 239 = *Aërides odoratum* Lour.

AËRIDES ORTGESIANUM Reichb. f. in Gard. Chron. II 23 (1885) 501; Ames Orch. 5 (1915) 218.

No specimens from the Philippines have been seen. A near ally, probably a form, of *Aërides quinquevulnerum* Lindl.

AËRIDES REICHENBACHII J. Linden ex Reichb. f. in Koch & Fintelm. Wochenschr. 1 (1858) 61; Naves Novis. App. (1882) 239; J. Linden in Lindenia 1 (1885) t. 1 = *Aërides odoratum* Lour.

AËRIDES ROEBELENI Reichb. f. in Gard. Chron. II 21 (1884) 510; Ames Orch. 5 (1915) 219.

No specimens from the Philippines have been seen. A near ally, probably a form, of *Aërides quinquevulnerum* Lindl.

AËRIDES ROHANIANUM Reichb. f. in Gard. Chron. II 21 (1884) 206; Ames Orch. 5 (1915) 219 = *Aërides odoratum* Lour.

AËRIDES SAVAGEANUM Sander ex Veitch Man. Orch. Pl. pt. 7 (1891) 78; Rolfe in Sander Reichenbachia II 2 (1894) 81, t. 81; O'Brien in Gard. Chron. III 16 (1894) 562; Ames Orch. 5 (1915) 219.

AËRIDES SHIBATITIANA Boxall ex Naves Novis. App. (1882) 239, *nomen*.

Perhaps an error for *Aërides thibautianum* Reichb. f., which is equal to *Aërides quinquevulnerum* Lindl.

AËRIDES SPURIUM Lindl. Gen. & Sp. Orch. (1833) 241 = *Dendrobium spurium* (Blume) J. J. Sm.

AËRIDES SUAVISSIMUM Lindl. in Journ. Hort. Soc. 4 (1849) 264; Ceron Cat. Pl. Herb. Manila (1892) 168 = *Aërides odoratum* Lour.

AËRIDES THIBAUTIANUM Reichb. f. in Gard. Chron. (1866) 101 = *Aërides quinquevulnerum* Lindl.

AËRIDES VIRENS Lindl. in Bot. Reg. 29 (1843) Misc. 41; Naves Novis. App. (1882) 239 = *Aërides odoratum* Lour.

81. RHYNCHOSTYLIS Blume

RHYNCHOSTYLIS RETUSA (Linn.) Blume Bijdr. (1825) 286, fig. 49; Reichb. f. in Bonpl. 5 (1857) 40, in Walp. Ann. 6 (1864) 887 (and vars.); Rolfe in Journ. Bot. 23 (1885) 216; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271; Hook. f. Fl. Brit. Ind. 6 (1890) 32; Veitch Man. Orch. Pl. pt. 7 (1891) 53, t. (and vars.); Stein Orchideenb. (1892) 536, fig. 164 (and vars. and forms); Grant Orch. Burma (1895) 290 (and vars.); Ridl. in Journ. Linn. Soc. 31 (1896) 294, 32 (1896) 356; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik-Himal.) (1898) 213, t. 284; Trim. Handb. Fl. Ceyl. 4 (1898) 187; Rolfe in Orch. Rev. 9 (1901) 281, fig. 42 (var. *guttata*); Cogn. in Dict. Icon. Orch. (1902) Rhynchost. t. 1; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 629, Fig.-Atlas (1914) fig. 471; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 143; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 158; Ames Orch. 2 (1908) 255,

- 5 (1915) 219; Merr. Sp. Blancoanae (1918) 117; Ames in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 197.
- Epidendrum retusum* Linn. Sp. Pl. (1753) 953, ed. 2 (1763) 1351; Poir. in Lam. Encycl. Suppl. 1 (1810) 384.
- Aërides retusum* Sw. in Schrad. Journ. Bot. 2 (1799) 233; Willd. Sp. Pl. 4 (1805) 130; Pers. Syn. Pl. 2 (1807) 522; Spreng. Syst. Veg. 3 (1826) 719; Grah. Cat. Pl. Bomb. (1839) 204.
- Limodorum retusum* Sw. in Nov. Act. Ups. 6 (1799) 80.
- Aërides praemorsum* Willd. Sp. Pl. 4 (1805) 130; Spreng. Syst. Veg. 3 (1826) 719; Grah. Cat. Pl. Bomb. (1839) 264.
- Epidendrum indicum* Poir. in Lam. Encycl. Suppl. 1 (1810) 384.
- Aërides undulatum* Sm. in Rees Cyclop. 39 (1819) No. 13.
- Aërides spicatum* D. Don Prodr. Fl. Nepal. (1825) 31.
- Epidendrum hippium* Buch.-Ham. ex D. Don Prod. Fl. Nepal (1825) 32, in syn.
- Rhynchosstylis praemorsa* Blume Bijdr. (1825) 286; Hassk. in Hoev. & De Vriese Tijdschr. Nat. Gesch. 9 (1842) 143; Hassk. Pl. Jav. Rar. (1848) 128.
- Sarcanthus guttatus* Lindl. in Bot. Reg. 17 (1831) *t.* 1443.
- Aërides guttatum* Roxb. Fl. Ind. ed. 2, 3 (1832) 471; Regel in Gartenfl. 12 (1863) 339, *t.* 415, *figs.* 8-12; Morr. in Belg. Hort. 26 (1876) 291.
- Saccolabium guttatum* Lindl. in Wall. Cat. (1832) No. 7308, *nomen*, Gen. & Sp. Orch. (1833) 220; Paxt. Mag. Bot. 2 (1835) 141, 6 (1839) 214, 11 (1844) 211; Lindl. Sert. Orch. (1841) sub *t.* 47; Hook. in Bot. Mag. 70 (1844) *t.* 4108; Griff. Notul. 1 (1847) 22, Icon. Pl. Asiat. 1 (1847) *tt.* 8, 9; Wight Icon. 5¹ (1852) 19, *t.* 1745-6; Harting. Parad. Vindob. 2 (1848-60) *t.* 13; De Vriese Ill. Orch. Ind. Neerl. (1854) *t.* 14; Lindl. in Journ. Linn. Soc. 3 (1859) 32; Miq. Fl. Ind. Bat. 3 (1859) 693; Dalz. & Gibs. Bomb. Fl. (1861) 263; Thw. Enum. Pl. Zeyl. (1864) 303; Drury Handb. Ind. Fl. 3 (1869) 405; Puydt Orch. (1880) 227, *text cut*; Naves Novis. App. (1882) 241; Godefr.-Lebeuf in l'Orchidoph. 8 (1888) 273, *t.* (var. *giganteum*), 11 (1891) 11; Meylhand in Journ. des Orch. 2 (1892) 255; Will. Orch. Grow. Man. ed. 7 (1894) 701 and var.); Meylhand in Sem. Hort. 3 (1899) 489.
- Saccolabium praemorsum* Lindl. Gen. & Sp. Orch. (1833) 221; Paxt. Mag. Bot. 4 (1837) 94; Hassk. in Tijdschr. Nat. Gesch. 9 (1842) 143, Pl. Jav. Rar. (1848) 128; Edit. in Fl. des Serres 7 (1851-52) 92, *text cut*; Miq. Fl. Ind. Bat. 3 (1859) 693; Drury Handb. Ind. Fl. 3 (1869) 405; Meylhand in Journ. des Orch. 2 (1891) 255, in Sem. Hort. 3 (1899) 489, non Hook. f.
- Saccolabium spicatum* Lindl. Gen. & Sp. Orch. (1833) 221.
- Orchis lanigera* Blanco Fl. Filip. (1837) 641, ed. 2 (1845) 446, ed. 3, 3 (1879) 37.
- Saccolabium blumei* Lindl. in Bot. Reg. 27 (1841) Misc. 55; Lindl. Sert. Orch. (1841) *t.* 47; Miq. Fl. Ind. Bat. 3 (1859) 693; Puydt Orch. (1880) 313, *t.* 27; Naves Novis. App. (1882) 241; Edit. in Gard. Chron. II 23 (1885) 572, *fig.* 105; Warner & Will. Orch. Alb. 4 (1885) *t.* 169; Teran in Ill. Hort. 32 (1885) 125, *t.* 566; Meylhand in Journ. des Orch. 2 (1891) 254; Will. Orch. Grow.

- Man. ed. 7 (1894) 699, t.; E. L. in Sem. Hort. 2 (1898) 376, fig. 375; Meylhand in Sem. Hort. 3 (1899) 489, 516.
- Saccolabium macrostachyum* Lindl. Sert. Orch. (1841) sub t. 47.
- Saccolabium retusum* Voigt Hort. Sub. Calc. (1845) 630; Van Houtte in Fl. des Serres 14 (1861) t. 1463-4; Will. Orch. Grow. Man. ed. 7 (1894) 703.
- Saccolabium rheedii* Wight Icon. 5¹ (1852) 19.
- Saccolabium blumei* Lindl. var. *major* Will. Orch. Grow. Man. (1852) 89, ed. 7 (1894) 699; J. Linden Pescat. 1 (1860) t. 21; Lem. in Ill. Hort. 15 (1868) t. 545; Meylhand in Journ. des Orch. 2 (1891) 254.
- Rhynchostylis guttata* Reichb. f. in Bonpl. 2 (1854) 93; Rolfe in Orch. Rev. 22 (1914) 281, fig. 33.
- Saccolabium gurwalicum* Lindl. in Journ. Linn. Soc. 3 (1859) 32; Reichb. f. in Gard. Chron. II 12 (1879) 102.
- Rhynchostylis gurwalica* Reichb. f. in Walp. Ann. 6 (1864) 888.
- Rhynchostylis retusa* (Linn.) Blumę var. *macrostachya* Reichb. f. in Walp. Ann. 6 (1864) 888.
- Saccolabium furcatum* Hort. ex Will. Orch. Grow. Man. ed. 3 (1868) 201, ed. 7 (1894) 700.
- Saccolabium holfordianum* Hort. ex Warner Sel. Orch. Pl. II (1874) t. 18.
- Saccolabium littorale* Reichb. f. in Gard. Chron. II 16 (1881) 198; Hook. f. Fl. Brit. Ind. 6 (1890) 33, cf. Ridl. in Journ. Linn. Soc. 32 (1896) 356.
- ? *Saccolabium berkeleyi* Reichb. f. in Gard. Chron. II 19 (1883) 814, Edit. in l'Orchidoph. 3 (1883) 693; Hook. f. Fl. Brit. Ind. 6 (1890) 33; Will. Orch. Grow. Man. ed. 7 (1894) 699.
- Saccolabium heathii* Hort. ex Gard. Chron. II 24 (1885) 369.
- Saccolabium turneri* Will. Orch. Grow. Man. ed. 6 (1885) 568, ed. 7 (1894) 703.
- Saccolabium blumei* Lindl. var. *russellianum* Will. in Warner & Will. Orch. Alb. 5 (1886) t. 238; Will. Orch. Grow. Man. ed. 6 (1885) 564, ed. 7 (1894) 699.
- Gastrochilus blumei* Ktze. Rev. Gen. Pl. 2 (1891) 661.
- Gastrochilus gorwalicus* Ktze. Rev. Gen. Pl. 2 (1891) 661.
- Gastrochilus retusoides* Ktze. Rev. Gen. Pl. 2 (1891) 661.
- Gastrochilus rheedii* Ktze. Rev. Gen. Pl. 2 (1891) 661.
- Gastrochilus spicatus* Ktze. Rev. Gen. Pl. 2 (1891) 661.
- ? *Rhynchostylis violacea* (Lindl.) Reichb. f. var. *berkeleyi* Stein Orchideenb. (1882) 537.
- Gastrochilus praemorsus* Dur. & Jacks. Ind. Kew. Suppl. 1 (1903) 178, non Ktze.
- Rhynchostylis violacea* Ames Orch. 1 (1905) 105, non Reichb. f.
- Luzon (Bataan, Rizal), Lyon 13, Whitford 36, Merrill 4190, Sp. *Blancoanae* 1030, B. S. 13548 Ramos, Weber 10, Reillo 9. Epiphyte. In India, Ceylon, Burma, Malay Peninsula, Siam, Java, and Borneo.
- A polymorphic species, which has been variously treated either as several distinct species or as one species with several varieties and forms. There are many varieties and forms based on extra-Philippine material

that are not listed in the above synonymy unless as species. Cf. Stein Orchideenb. (1892) 536-537 for some of these variants.

EXCLUDED OR SYNONYMOUS SPECIES

RHYNCHOSTYLIS VIOLACEA Ames Orch. 1 (1905) 105 (excl. syn.) non Reichb. f. = *Rhynchostylis retusa* (Linn.) Blume.

RHYNCHOSTYLIS VIOLACEA Reichb. f. in Bonpl. 2 (1854) 93; in Walp. Ann. 6 (1864) 888; Van Houtte in Fl. des Serres 17 (1869) 91; Stein Orchideenb. (1892) 537; Ames Orch. 5 (1915) 219 = *Anota violacea* (Lindl.) Schltr.

82. **LUISIA** Gaudichaud

LUISIA FOXWORTHII Ames Orch. 2 (1908) 222, *text cut*, 5 (1915) 220. Luzon (Bataan, Bulacan, Sorsogon), Polillo, *B. S. 1679 Foxworthy, 9257 Robinson, F. B. 7191 Curran, Elmer 16251*. Epiphyte at sea level to 830 m altitude.

LUISIA RAMOSII Ames in Philip. Journ. Sci. 6 (1911) Bot. 55, Orch. 5 (1915) 220.

Luzon (Cagayan, Zambales), *B. S. 7970 Ramos, Boettcher s. n.*

LUISIA TERETIFOLIA Gaudich. in Freyc. Voy. Uranie et Physic. (1829) Bot. 427, *t. 37*; Blume Rumphia 4 (1848) 50, *t. 194, fig. 3, t. 197 D* (excl. syn. as to *triste*); Blume Mus. Bot. Lugd.-Bat. 1 (1849) 63 (excl. syn. *Epidendrum triste* Forst.); Reichb. f. in Walp. Ann. 3 (1852) 549, 6 (1863) 621 (excl. syn. *Epidendrum triste* Forst.); Lindl. Fol. Orch. Luisia (1853) 2 (excl. syn. *Epidendrum triste* Forst.); Miq. Fl. Ind. Bat. 3 (1859) 685; Naves Novis. App. (1882) 242; Hook. f. Fl. Brit. Ind. 6 (1890) 22; Grant Orch. Burma (1895) 236; Ridl. in Journ. Linn. Soc. 32 (1896) 349; King & Pantl. in Ann. Roy. Bot. Gard. Calc. 8 (Orch. Sik.-Himal.) (1898) 203, *t. 271*; Trim. Handb. Fl. Ceyl. 4 (1898) 190; Rolfe in Forbes & Hemsl. in Journ. Linn. Soc. 36 (1903) 33; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 545, Fig.-Atlas (1912) *fig. 413*, Orch. Amb. (1905) 91; Duthie in Ann. Roy. Bot. Gard. Calc. 9 (Orch. North-west. Himal.) (1906) 140; Ames in Philip. Journ. Sci. 8 (1913) Bot. 453, 9 (1914) Bot. 15, Orch. 5 (1915) 220; Schltr. in Fedde Repert. Beihefte 4 (Orch. Sino-Jap. Prodr.) (1919) 281.

Cymbium triste Roxb. Fl. Ind. ed. 2, 3 (1832) 461 (excl. syn.); Hook. in Bot. Mag. 65 (1838) *t. 3648* (excl. syn. in part); Wight Icon. 3² (1849) 10 (excl. *t.*); Griff. Notul. 3 (1851) 340, quoad *Mergui planta*, non Willd.

Dendrobium teres Blanco Fl. Filip. (1837) 638, non Roxb.

Dendrobium teretifolium Blanco Fl. Filip. ed. 2 (1845) 444, ed. 3, 3 (1879) 34, non R. Br.

Cymbidium tenuifolium Wight Icon. 5 (1852) *t. 1689* (excl. text), non Willd.

Luisia burmanica Lindl. Fol. Orch. Luisia (1853) 3; Reichb. f. in Walp. Ann. 6 (1863) 621.

Luisia zeylanica Lindl. Fol. Orch. Luisia (1853) 3; Reichb. f. Xen. Orch. 1 (1856) 204, in Walp. Ann. 6 (1863) 621; Thw. Enum. Pl. Zeyl. (1864) 302.

Luisia trichorrhiza Reichb. f. Xen. Orch. 1 (1856) t. 77, figs. 8-18, non Blume.

Luisia platyglossa Reichb. f. in Walp. Ann. 6 (1863) 622, in Trans. Linn. Soc. 30 (1874) 144.

Luisia tristis Ktze. Rev. Gen. Pl. 2 (1891) 672 (excl. syn.), non Hook. f.

Luzon (Bontoc, Bulacan, Laguna, Sorsogon), Negros (Oriental Negros), Leyte, Palawan, Mindanao (Camiguin Island, Agusan, Surigao). A fairly common, variable, and widely distributed species. Epiphyte at 70 to 500 m altitude. In India, Ceylon, Burma, Malay Peninsula, Siam, China, Sumatra, Java, Amboina, Moluccas, Borneo (fide Ridley), and also the Marianne Islands and New Caledonia.

LUISIA VALIDA Reichb. f. in Bonpl. 10 (1862) 335, in Walp. Ann. 6 (1863) 621; Ames Orch. 5 (1915) 220.

According to Reichenbach (Bonpl. l. c.) of Philippine origin. The description is vague, and it seems probable that the species is conspecific with *Luisia teretifolia* Gaudich.

EXCLUDED OR SYNONYMOUS SPECIES

LUISIA ANTENNIFERA Blume apud Naves Novis. App. (1882) 242.

LUISIA BRACHYSTACHYS Blume apud Naves Novis. App. (1882) 242.

83. ARACHNIS Blume

ARACHNIS LYONII Ames Orch. 5 (1915) 221.

Luzon (Isabela), Mindanao (Lanao), *Lyon 126*, *M. S. Clemens s. n.* Terrestrial at 1,000 m altitude in Luzon.

84. VANDOPSIS Pfitzer

VANDOPSIS LEYTENSIS Ames Orch. 5 (1915) 222.

Luzon (Rizal, Sorsogon), Leyte, *Lyon s. n.*, *Elmer 16860*, *Wenzel 014*, *0154*, *0172*, *0900 C.* Epiphyte at 60 to 800 m altitude.

VANDOPSIS LISSOCHILOIDES (Gaudich.) Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1889) 210, fig. 229; Kerchove Livre des Orch. (1894) 138, fig. 114; J. J. Sm. Orch. Amb. (1905) 102; Ames Orch. 2 (1908) 220, 5 (1915) 223; J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1913) 76 (reprint 6), in Merr. Interp. Rumph. Herb. Amb. (1917) 178.

Fieldia lissochiloides Gaudich. in Freyc. Voy. Uranie et Physic. (1829) Bot. 424, t. 36; Blume Rumphia 4 (1848) t. 194, fig. 2; Van Houtte in Fl. des Serres 17 (1869) 91; Reichb. f. Xen. Orch. 2 (1862) 38, in Walp. Ann. 6 (1864) 870.

Vanda lissochiloides Lindl. Gen. & Sp. Orch. (1833) 216; Blume Rumphia 4 (1848) 49, Mus. Bot. Lugd.-Bat. 1 (1849) 63; Reichb. f. in Walp. Ann. 1 (1849) 790, 3 (1852) 565; Miq. Fl. Ind. Bat. 3 (1859) 680; Naves Novis. App. (1882) 240.

Vanda batemanii Lindl. in Bot. Reg. 32 (1846) t. 59 (as *batemanni*); Paxt. Mag. Bot. 14 (1847) 21; Lindl. in Paxt. Flow. Gard. 2 (1851) 22; S. Ch. in Fl. des Serres 6 (1852) 332; Lindl. Fol. Orch. Vanda (1853) 2; Moore Ill. Orch. Pl. (1857) Vanda 10, t. 1; du Buyss in Fl. des Serres 18 (1870) 161, t. 1921-2.

Grammatophyllum pantherinum Herb. Zipp. ex Blume Rumphia 4 (1848) 49, in syn.

Stauroopsis lissochiloides Benth. ex Pfitz. Vergl. Morph. Orch. (1882) 14; Veitch Man. Orch. Pl. pt. 7 (1891) 3; Cogn. in Journ. des Orch. 5 (1894) 291, fig. 80; Cogn. Dict. Icon. Orch. (1898) *Stauroopsis* t. 1; G. Wilson in Orch. World 4 (1914) 227.

Stauroopsis batemanii Nichols. Dict. Gard. 3 (1886) 493 (as *batemanni*).

Panay, Mindanao (Zamboanga, Bukidnon, Sigaboy Island), *Loher* 6025 (cult. in Manila); *B. S.* 11774 *Robinson*, 21434 *Escritor*, 19345 *Wester*, *Merrill* 3317.

EXCLUDED OR SYNONYMOUS SPECIES

VANDOPSIS KUPPERIANA Kränzl. in Fedde Repert. 17 (1921) 390 = *Staurochilus luzonensis* Ames.

85. RENANTHERA Loureiro

RENANTHERA MONACHICA Ames Orch. 5 (1915) 224.

Luzon (Zambales), *B. S.* 3089 *Curran*.

RENANTHERA STORIEI Reichb. f. in Gard. Chron. II 14 (1880) 296, 16 (1881) 364; Warner & Will. Orch. Alb. 11 (1897) t. 513; Cogn. in Dict. Icon. Orch. (1907) *Renanth.* t. 2; Ames Orch. 2 (1908) 219, 5 (1915) 225; Rolfe in Orch. Rev. 23 (1915) 279, 297, fig. 38; W. T. in The Garden 79 (1915) 445, text cut; G. Wilson in Orch. World 6 (1915) 16.

Vanda storiei Storie ex Reichb. f. in Gard. Chron. II 14 (1880) 296, in syn.; Hook. f. in Bot. Mag. 123 (1897) t. 7537.

Luzon (Bataan, Rizal), Bucas Grande, Dinagat, *F. B.* 20110 *Borden*, *Whitford* 1336, *Loher* 6000, *B. S.* 35085, 35228 *Ramos & Pascasio*, *Lyon* 136. Apparently at low altitudes (up to 130 m).

EXCLUDED OR SYNONYMOUS SPECIES

RENANTHERA AMABILIS Boxall ex Naves Novis. App. (1882) 241, *nomen*.

RENANTHERA ARACHNITIS Lindl. apud Naves Novis. App. (1882) 240.

RENANTHERA COCCINEA Lour. apud Naves Novis. App. (1882) 240.

RENANTHERA ELONGATA Ames Orch. 5 (1915) 224, non Lindl.

Renanthera matutina Lindl. in Bot. Reg. 29 (1843) t. 41, quoad Philip.; G. K. in Bot. Zeit. 1 (1843) 760, quoad Philip.

Renanthera micrantha Blume Rumphia 4 (1848) 60, Mus. Bot. Lugd.-Bat. 1 (1849) 60; Reichb. f. in Walp. Ann. 3 (1852) 566; Miq. Fl. Ind. Bat. 3 (1859) 698; Naves Novis. App. (1882) 240.

Although this species, under the name *Renanthera micrantha*, is reported as being of Philippine origin, no material referable to it has been seen and it seems very doubtful that it occurs in the Islands.

RENANTHERA HOOKERIANA Reichb. f. apud Naves Novis. App. (1882) 241.

RENANTHERA LOWEI Reichb. f. apud Naves Novis. App. (1882) 241.

RENANTHERA MATUTINA Lindl. in Bot. Reg. 29 (1843) t. 41, quoad Philip.; G. K. in Bot. Zeit. 1 (1843) 760, quoad Philip. = *Renanthera elongata* Ames, spec. excl.

RENANTHERA MICRANTHA Blume Rumphia 4 (1848) 60, Mus. Bot. Lugd.-Bat. 1 (1849) 60; Reichb. f. in Walp. Ann. 3 (1852) 566; Miq. Fl. Ind. Bat. 3 (1859) 698; Naves Novis. App. (1882) 240 = *Renanthera elongata* Ames, spec. excl.

RENANTHERA MOLUCCANA Blume apud Naves Novis. App. (1882) 240.

RENANTHERA SULINGI Lindl. apud Naves Novis. App. (1882) 240.

86. *ACAMPE* Lindley

ACAMPE LOHERIANA Kränzl. in Fedde Repert. 17 (1921) 386.

Löher s. n.

Kränzlin states: "Philippinen. Ohne genaueren Standort."

The genus *Acampe* has not heretofore been reported from the Philippines, and field specimens are much desired to authenticate its habitat there. The material examined is inadequate and it is of garden origin.

87. *VANDA* Jones

VANDA CLITELLARIA Reichb. f. in Flora 55 (1872) 277; Ames Orch. 5 (1915) addend. opp. p. 258.

Sulu Archipelago (Mangsi Island), *Wilkes Exped.* (in Gray Herb.).

Very closely allied to *Vanda lamellata* Lindl. and perhaps only a form of it.

VANDA LAMELLATA Lindl. in Bot. Reg. 24 (1838) Misc. 66; Paxt. Mag. Bot. 6 (1839) 143; S. Ch. in Fl. des Serres 6 (1851) 332; Lindl. in Paxt. Flow. Gard. 2 (1851) 22, Fol. Orch. Vanda (1853) 9; Moore Ill. Orch. Pl. (1857) Vanda 12; Miq. Fl. Ind. Bat. 3 (1859) 680; Van Houtte in Fl. des Serres 17 (1867-68) 92; Reichb. f. in Walp. Ann. 6 (1864) 869; Naves in Blanco Fl. Filip. ed. 3, Laminas (1877-83) t. 465; Naves Novis. App. (1882) 240; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271; Veitch Man. Orch. Pl. pt. 7 (1891) 99; Stein Orchideenb. (1892) 583; Schltr. in Perk. Frag. Fl. Philip. (1904) 43; Ames Orch. 1 (1905) 103, 2 (1908) 240, 5 (1915) 225.

Vanda cumingii Lindl. ex Paxt. in Loud. Hort. Brit. Suppl. 3 (1850) 654, *nomen*.

Luzon (Abra, Lepanto, Benguet, Zambales, Bataan, Cavite), Mindoro, Panay (Capiz), Palawan, Comiran. A fairly common species of wide distribution. On rocks or on trees at sea level to 400 m altitude.

- Var. **BOXALLI** Reichb. f. in Gard. Chron. II 13 (1880) 743, 15 (1881) 87, fig. 18, 16 (1881) 19, 786; Edit. in The Garden 19 (1881) 574, t. 287, in l'Orchidoph. 5 (1885) 129; Warner & Will. Orch. Alb. 8 (1889) t. 338; Veitch Man. Orch. Pl. pt. 7 (1891) 100, text cut; Stein Orchideenb. (1892) 583; Cogn. in Dict. Icon. Orch. (1898) Vanda t. 9; Ames Orch. 2 (1908) 241, 5 (1915) 225.
- Vanda boxallii* Reichb. f. in Gard. Chron. II 15 (1881) 336; Linden & Rodigas in Lindenia 1 (1885) 69, t. 32.
- Vanda superba* Linden & Rodigas in Lindenia 3 (1887) 83, t. 136.
- Luzon (Zambales), *Merrill s. n.* (cult. in Manila).
- Var. **COBBIANA** (Reichb. f.) Ames Orch. 2 (1908) 241.
- Vanda boxallii* Reichb. f. var. *cobbiana* Reichb. f. in Gard. Chron. II 16 (1881) 780; Edit. in l'Orchidoph. 1 (1882) 194.
- VANDA LUZONICA** Loher ex Rolfe in Orch. Rev. 23 (1915) 137, 371, fig. 12; Edit. in The Garden 79 (1915) 563, text cut; Rolfe in Bot. Mag. 143 (1917) t. 8709; Ames Orch. 6 (1920) 308.
- Vanda tricolor* Ames Orch. 2 (1908) 245, 5 (1915) 225, non Lindl.
- Luzon (Bulacan), *Loher 6006*.
- VANDAROEBLINGIANA** Rolfe in Kew Bull. (1894) 365; Ames Orch. 2 (1908) 241, 5 (1915) 225; G. Wilson in Orch. Rev. 30 (1922) 96.
- Luzon (Bontoc, Benguet), *Vanoverbergh 1257, 3939, F. B. 5059 Curran*. Up to 1,600 m altitude in Bontoc. In Malay Peninsula.
- VANDA SANDERIANA** Reichb. f. in Gard. Chron. II 17 (1882) 588; Edit. in l'Orchidoph. 2 (1882) 318, 4 (1884) 109, t., in Gard. Chron. II 20 (1883) 402, 440, 446, figs. 67, 68, 22 (1884) 236; Warner & Will. Orch. Alb. 3 (1884) t. 124; Moore in Gard. Chron. II 21 (1884) 9; Rodigas in Ill. Hort. 31 (1884) 139, t. 532; Carr. in Rev. Hort. 57 (1885) 372, t.; Hook. f. in Bot. Mag. 114 (1888) t. 6983; Sander Reichenbachia I 2 (1890) 29, t. 62; Veitch Man. Orch. Pl. pt. 7 (1891) 103, t.; Rolfe in Orch. Rev. 4 (1896) 358, frontisp., 19 (1911) 281, fig. 31, 23 (1915) 361, frontisp.; Cogn. in Dict. Icon. Orch. (1900) Vanda t. 12; Ames Orch. 2 (1908) 242; G. Wilson in Orch. World 2 (1911) 36, t., 5 (1915) 52, 201, text cuts, 6 (1916) 97, in Orch. Rev. 29 (1921) 79, text cut.
- Esmeralda sanderiana* Reichb. f. in Gard. Chron. II 17 (1882) 588, in textu; Kerchove Livre des Orch. (1894) 307, 355, figs. 214, 223; Will. Orch. Grow. Man. ed. 7 (1894) 398, text cut; Ames Orch. 5 (1915) 222.
- Vanda sanderiana* Reichb. f. var. *labello-viridi* Linden & Rodigas in Lindenia 1 (1885) 85, t. 40; Ames Orch. 2 (1908) 244.
- Vanda sanderiana* Reichb. f. var. *albata* Reichb. f. in Gard. Chron. III 2 (1887) 9; Edit. in l'Orchidoph. 8 (1888) 197; Reichb. f. in Sander Reichenbachia I 2 (1890) 29; Ames Orch. 2 (1908) 244.
- Esmeralda sanderiana* Reichb. f. var. *labello-viridi* Will. Orch. Grow. Man. ed. 7 (1894) 400.
- Esmeralda sanderiana* Reichb. f. var. *albata* Will. Orch. Grow. Man. ed. 7 (1894) 400.
- Vanda sanderiana* Reichb. f. var. *froebeliana* Cogn. in Dict. Icon. Orch. (1903) Vanda t. 12A; Ames Orch. 2 (1908) 244.

Euanthe sanderiana Schltr. Orchideen (1914) 568, fig. 194; Ames Orch. 6 (1920) 308.

Mindanao (Davao), R. F. Black 20, Loher 6030 (cult. in Manila).

EXCLUDED OR SYNONYMOUS SPECIES

- VANDA BATEMANII** Lindl. in Bot. Reg. 32 (1846) t. 59 (as *batemanni*); Paxt. Mag. Bot. 14 (1847) 21; S. Ch. in Fl. des Serres 6 (1851) 332; Lindl. in Paxt. Flow. Gard. 2 (1851) 22; Lindl. Fol. Orch. Vanda (1853) 2; Moore Ill. Orch. Pl. (1857) Vanda 10, t. 1; du Buyss in Fl. des Serres 18 (1870) 161, t. 1921-2 = *Vandopsis lissochiloides* (Gaudich.) Pfitz.
- VANDA BOXALLII** Reichb. f. in Gard. Chron. II 15 (1881) 336; Linden & Rodigas in Lindenia 1 (1885) 69, t. 32 = *Vanda lamellata* Lindl. var. *boxallii* Reichb. f.
- VANDA CRISTATA** Lindl. apud Boxall ex Naves Novis. App. (1882) 240; Ceron Cat. Pl. Herb. Manila (1892) 168.
- VANDA CUMINGII** Lindl. ex Paxt. in Loud. Hort. Brit. Suppl. 3 (1850) 654, nomen = *Vanda lamellata* Lindl.
- VANDA FURVA** Lindl. apud Naves Novis. App. (1882) 239.
- VANDA GIGANTEA** Lindl. apud Naves Novis. App. (1882) 240.
- VANDA HELVOLA** Bl. apud Naves Novis. App. (1882) 240.
- VANDA HINDSII** Lindl. apud Naves Novis. App. (1882) 239.
- VANDA INSIGNIS** Bl. apud Naves Novis. App. (1882) 239.
- VANDA LIMBATA** Bl. apud Naves Novis. App. (1882) 240.
- VANDA LISSOCHILOIDES** Lindl. Gen. & Sp. Orch. (1833) 216; Naves Novis. App. (1882) 240 = *Vandopsis lissochiloides* (Gaudich.) Pfitz.
- VANDA ROXBURGHII** R. Br. apud Naves Novis. App. (1882) 240.
- VANDA STANGEANA** Reichb. f. apud Naves Novis. App. (1882) 240.
- VANDA STORIEI** Storie ex Reichb. f. in Gard. Chron. II 14 (1880) 296, in syn.; Hook. f. in Bot. Mag. 123 (1897) t. 7537 = *Renanthera storiei* Reichb. f.
- VANDA SUPERBA** Linden & Rodigas in Lindenia 3 (1887) 83, t. 136 = *Vanda lamellata* Lindl. var. *boxallii* Reichb. f.
- VANDA TRICOLOR** Ames Orch. 2 (1908) 245, 5 (1915) 225, non Lindl. = *Vanda luzonica* Loher ex Rolfe.
- VANDA UNDULATA** Lindl. apud Naves Novis. App. (1882) 240.
- VANDA VIDALII** Boxall ex Naves Novis. App. (1882) 240, nomen; Ceron Cat. Pl. Herb. Manila (1892) 168.
- VANDA VIOLACEA** Lindl. in Bot. Reg. 27 (1841) Misc. 12, 33 (1847) t. 30; Paxt. Mag. Bot. 9 (1842) 119; Van Houtte in Fl. des Serres 17 (1868-69) 92 = *Anota violacea* (Lindl.) Schltr.

88. **ANGRAECUM** Bory

ANGRAECUM PHILIPPINENSE Ames in Philip. Journ. Sci. 2 (1907) Bot. 336, 4 (1909) Bot. 597, Orch. 2 (1908) 246, 3 (1908) 69, *t.* 50, 5 (1915) 226.

Luzon (Bontoc, Nueva Vizcaya), Mindoro, *Lyon* (Curran) 121, *F. B.* 10860 Curran, Merrill 5698. On mountains, in the forest at 850 to 1,400 m altitude.

89. **SACCOLABIUM** Blume

SACCOLABIUM CONFUSUM Ames in Philip. Journ. Sci. 8 (1913) Bot. 438, Orch. 5 (1915) 227.

Luzon (Laguna, Sorsogon), Mindanao (Cotabato), *Serrato s. n.*, *Elmer* 14713, *Weber* 221. Epiphyte, at 700 m altitude in Cotabato.

SACCOLABIUM ESCRITORII Ames Orch. 5 (1915) 227.

Luzon (Tayabas), *B. S.* 20733 *Escritor*.

SACCOLABIUM LOHERI Ames Orch. 5 (1915) 228.

Luzon (Zambales), *Loher* 6026.

SACCOLABIUM LUZONENSE Ames in Philip. Journ. Sci. 8 (1913) Bot. 435, Orch. 5 (1915) 229.

Luzon (Laguna, Sorsogon), *B. S.* 16556 *Ramos*, *Serrato s. n.*, *Elmer* 14860, 17076. Epiphyte.

SACCOLABIUM SARCOCHILOIDES Schltr. in Orchis 5 (1911) 61, *t.* 8; Ames Orch. 5 (1915) 230.

Saccolabium epichysiochilum Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 63.

K. W. John (fide Schltr.), *Loher s. n.*

SACCOLABIUM TENELLUM Ames Orch. 6 (1920) 308, *t.* 99.

Leyte, Mindanao (Surigao), *Wenzel* 0988, 01037. Epiphyte at sea level to 150 m altitude.

SACCOLABIUM VANOVERBERGHII Ames Orch. 5 (1915) 230.

Luzon (Bontoc, Lepanto, Ifugao), *Vanoverbergh* 955, 1712, 2214, *Ramos & Edaña* 91, *B. S.* 19634 *McGregor*. Epiphyte in mountain regions at 1,550 to 2,540 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

SACCOLABIUM ASCENDENS Lindl. apud Naves Novis. App. (1882) 241.

SACCOLABIUM BIFIDUM Lindl. in Bot. Reg. 24 (1838) Misc. 3; Paxt. Mag. Bot. 6 (1839) 262; Reichb. f. in Walp. Ann. 6 (1864) 886; Naves Novis. App. (1882) 241 = *Sarcanthus bifidus* (Lindl.) Ames.

SACCOLABIUM BLUMEI Lindl. in Bot. Reg. 27 (1841) Misc. 55, Sert. Orch. (1841) *t.* 47; Naves Novis. App. (1882) 241 = *Rhynchostylis retusa* (Linn.) Blume.

SACCOLABIUM CHIONANTHUM Lindl. in Journ. Linn. Soc. 3 (1859) 35; Naves Novis. App. (1882) 241; Ames in Philip. Journ. Sci. 4 (1909) Bot. 675 = *Schoenorchis micrantha* Blume.

SACCOLABIUM COMPRESSUM Lindl. in Bot. Reg. 26 (1840) Misc. 9, Sert. Orch. (1840) *t.* 32; Edit. in Allg. Berl. Gartenz. 8 (1840) 336; Reichb. f. in Walp. Ann. 6 (1864) 883; Naves Novis. App. (1882) 241; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271; Ames in Philip. Journ. Sci. 2 (1907) Bot. 336, Orch. 2 (1915) 238.

Gastrochilus compressus Ktze. Rev. Gen. Pl. 2 (1891) 661 = *Robiquetia compressa* (Lindl.) Schltr.

SACCOLABIUM DENSIFLORUM Lindl. in Bot. Reg. 24 (1838) Misc. 56, quoad Philip., non Lindl. in Wall. Cat.; Reichb. f. in Walp. Ann. 6 (1864) 885, quoad Philip.; Naves Novis. App. (1882) 241 = *Robiquetia pantherina* (Kränzl.) Ames.

SACCOLABIUM EPICHYSIOSCHILUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 63 = *Saccolabium sarcochiloides* Schltr.

SACCOLABIUM GUTTATUM Lindl. in Wall. Cat. (1832) No. 7308, *nomen*, Gen. & Sp. Orch. (1833) 220; Naves Novis. App. (1882) 241; Ceron Cat. Pl. Herb. Manila (1892) 168 = *Rhynchostylis retusa* (Linn.) Blume.

SACCOLABIUM HANISSONII Boxall ex Naves Novis. App. (1882) 241, *nomen*.

SACCOLABIUM MACROSTACHYUM Lindl. Sert. Orch. (1841) sub *t.* 47 = *Rhynchostylis retusa* (Linn.) Blume.

SACCOLABIUM MINIATUM Lindl. in Bot. Reg. 33 (1847) sub *t.* 26, *t.* 58; Ames Orch. 2 (1908) 239 (incl. var. *citrinum*) = *Ascocentrum miniatum* (Lindl.) Schltr.

SACCOLABIUM PANTHERINUM Kränzl. in Fedde Repert. 17 (1921) 391 = *Robiquetia pantherina* (Kränzl.) Ames.

SACCOLABIUM PUMILIO Reichb. f. in Walp. Ann. 6 (1864) 886; in Gard. Chron. II 4 (1875) 98; Ames Orch. 5 (1915) 229.

Saccolabium discolor Reichb. f. in Otia Bot. Hamb. 1 (1878) 42.

Reichenbach in Gardener's Chronicle (l. c.) states: "I obtained the novelty from Mr. Bull, who introduced the plant from Manila." The species occurs in Sylhet, British India. Manila seems an error. No material from the Philippines referable to this species has been seen.

SACCOLABIUM REFLEXUM Lindl. apud Naves Novis. App. (1882) 241.

SACCOLABIUM SEMICLAUSUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 63.

Said to have been sent from the Philippines by Loher. No material has been seen.

SACCOLABIUM TRICHROMUM Reichb. f. apud Naves Novis. App. (1882) 241.

SACCOLABIUM VIOLACEUM Reichb. f. in Bonpl. 2 (1854) 93, in syn.; Warner Sel. Orch. 1 (1862-65) *t.* 14; Puydt Orch. (1880) 317, *t.* 39; Naves Novis. App. (1882) 241; Veitch Man. Orch. Pl. pt. 7 (1891) 117; Ceron Cat. Pl. Herb. Manila (1892) 168 = *Anota violacea* (Lindl.) Schltr.

90. GASTROCHILUS D. Don

GASTROCHILUS PHILIPPINENSIS Ames Orch. 5 (1915) 231.

Luzon (Benguet), *Elmer* 6517, Baguio.

91. OMOEA Blume

OMOEA PHILIPPINENSIS Ames Orch. 5 (1915) 232.

Luzon (Pampanga, Bataan), *Merrill* 4208, *B. S.* 1669 *Foxworthy*. Epiphyte in mossy forest at 850 to 1,400 m altitude.

92. ASCOCENTRUM Schlechter

ASCOCENTRUM MINIATUM (Lindl.) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1913) 975; *J. J. Sm.* Orch. Jav. Fig.-Atlas (1914) *fig.* 483; Ames Orch. 5 (1915) 233.

Saccolabium miniatum Lindl. in Bot. Reg. 33 (1847) sub *t.* 26, *t.* 58; Reichb. f. in Walp. Ann. 1 (1849) 791; Miq. Fl. Ind. Bat. 3 (1859) 692; Veitch Man. Orch. Pl. pt. 7 (1891) 117; C. K. in Journ. Hort. III 24 (1892) 449, *fig.* 77; Rolfe in Orch. Rev. 4 (1896) 178; *J. J. Sm.* Fl. Buitenz. 6 (Orch. Jav.) (1905) 648; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 170; Ames Orch. 2 (1908) 239, non Hook. in Bot. Mag. 88 (1862) *t.* 5326.

Saccolabium miniatum Lindl. var. *citrinum* Reichb. f. in Gard. Chron. II 21 (1884) 542; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 170; Ames Orch. 2 (1908) 240.

Gastrochilus miniatum Ktze. Rev. Gen. Pl. 2 (1891) 661.

Luzon (Rizal), Mindanao (Lanao, Davao), *B. S.* 5649 *Ramos*, *Weber* 9, 3020 *Mrs. Lyon*, *Copeland* 687. Epiphyte at 180 to 750 m altitude. In India (fide Rolfe), Malay Peninsula, Siam (fide Ridley), and Java.

93. MALLEOLA J. J. Smith and Schlechter

MALLEOLA CONSTRICTA Ames Orch. 5 (1915) 234.

Luzon (Sorsogon), Leyte, Mindanao (Lanao, Agusan), *Elmer* 13953, 14831, 17199, 17342, *Wenzel* 055, 062, 0124, 0195, 0882, *M. S.* *Clemens* 702. Epiphyte at 60 to 700 m altitude in Leyte.

MALLEOLA LYONII Ames Orch. 5 (1915) 235.

Luzon?, *Lyon* s. n.

EXCLUDED OR SYNONYMOUS SPECIES

MALLEOLA MERRILLII Ames Orch. 5 (1915) 236 = *Robiquetia merrillii* Ames.

94. ROBIQUETIA Gaudichaud

ROBIQUETIA COMPRESSA (Lindl.) Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1913) 983; Ames Orch. 5 (1915) 237.

Saccolabium compressum Lindl. in Bot. Reg. 26 (1840) Misc. 9, Sert. Orch. (1840) *t.* 32; Edit. in Allg. Berl. Gartenz. 8 (1840) 336; Reichb. f. in Walp. Ann. 6 (1864) 883; Naves Novis. App. (1882) 241; Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271; Ames in Philip. Journ. Sci. 2 (1907) Bot. 336; Merr. &

Merritt in Philip. Journ. Sci. 5 (1910) Bot. 341; Ames Orch. 5 (1915) 238.

Gastrochilus compressus Ktze. Rev. Gen. Pl. 2 (1891) 661.

Luzon (Camarines, Sorsogon), Mindanao (Dapitan, Davao), B. S. 22081, 23679 Ramos, F. B. 4603 Mearns & Hutchinson, Mearns s. n. At 550 to 3,000 m altitude.

ROBIQUETIA MERRILLII Ames Orch. 7 (1922) 123.

Malleola merrillii Ames Orch. 5 (1915) 236.

Luzon (Bontoc, Lepanto, Benguet, Bataan, Laguna, Camarines), Mindanao (Agusan), B. S. 5603 Worcester, Merrill 7685, 31999 Santos, 17127 Robinson, 22178 Ramos, Taylor 3, Elmer 13838, F. B. 1596 Borden. Epiphyte at 350 to 3,200 m altitude.

ROBIQUETIA PANTHERINA (Kränzl.) Ames Sched. Orch. 6 (1923) 95.

Cleisostoma spatulatum Blume apud Naves Novis. App. (1882) 238.

Saccolabium densiflorum Lindl. in Bot. Reg. 24 (1838) Misc. 56, quoad Philip., non Lindl. in Wall. Cat.; Reichb. f. in Walp. Ann. 6 (1864) 885, quoad Philip.; Naves Novis. App. (1882) 241.

Robiquetia spathulata Ames Orch. 5 (1915) 237, non J. J. Sm.

Saccolabium pantherinum Kränzl. in Fedde Repert. 17 (1921) 391.

Leyte, Loher s. n., Wenzel 0180. Epiphyte in forest at 60 m altitude.

ROBIQUETIA RAMOSII Ames Orch. 7 (1922) 123.

Catanduanes, Panay (Antique), B. S. 30396 Ramos, 32592 McGregor. Epiphyte at 30 to 300 m altitude.

ROBIQUETIA VANOVERBERGHII Ames Orch. 5 (1915) 238.

Luzon (Bontoc, Benguet), Mindanao (Zamboanga), Vanoverbergh 1473, R. S. Williams 1952, F. B. 5056 Curran, Ramos & Edaña 65, 69, 102, Merrill 8075.

EXCLUDED OR SYNONYMOUS SPECIES

ROBIQUETIA SPATHULATA Ames Orch. 5 (1915) 237, non J. J. Sm. = *Robiquetia pantherina* (Kränzl.) Ames.

95. SCHOENORCHIS Blume

SCHOENORCHIS MICRANTHA Blume Bijdr. (1825) 362; Lindl. Gen. & Sp. Orch. (1833) 243; J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 100 (reprint 30), Orch. Jav. Fig.-Atlas (1914) fig. 453; Ames Orch. 5 (1915) 239, 6 (1920) 224, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 200.

Saccolabium chionanthum Lindl. in Journ. Linn. Soc. 3 (1859) 35; Miq. Fl. Ind. Bat. 3 (1859) 692; Naves Novis. App. (1882) 241; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 635; Ames in Philip. Journ. Sci. 4 (1909) Bot. 675.

Saccolabium perpusillum Hook. f. Fl. Brit. Ind. 6 (1890) 56, in Hook. Icon. Pl. 22 (1893) t. 2129A; Ridl. in Journ. Linn. Soc. 32 (1896) 358, Mat. Fl. Mal. Penin. 1 (1907) 164.

Gastrochilus chionanthus Ktze. Rev. Gen. Pl. 2 (1891) 661.

Gastrochilus perpusillus Ktze. Rev. Gen. Pl. 2 (1891) 661.

Luzon (Rizal), Leyte, Palawan, Mindanao (Agusan, Davao), *B. S.* 3055, 41637 *Ramos*, *Wenzel* 05, 0122, 0378, *Weber* 319, 603, *Copeland* 685. Epiphyte at sea level to 1,300 m altitude. In Malay Peninsula, Sumatra, Java, and Borneo.

SCHOENORCHIS MINDANAENSIS Ames Orch. 5 (1915) 240.

Luzon (Sorsogon), Mindanao (Agusan), *Elmer* 16669, *Weber* 155. Epiphyte at 15 m altitude in Mindanao.

SCHOENORCHIS PHILIPPINENSIS Ames Orch. 5 (1915) 241.

Luzon (Nueva Vizcaya), Mindanao (Cotabato, Agusan), *B. S.* 20117 *McGregor*, *Weber* 212, *Elmer* 13969. Epiphyte at 550 to 1,000 m altitude.

SCHOENORCHIS VANOVERBERGHII Ames Orch. 5 (1915) 242.

Luzon (Cagayan, Bontoc, Pampanga, Bataan), *F. B.* 16781, 19307 *Curran*, *Vanoverbergh* 1761, 3644, *Whitford* 115 p. p. Epiphyte at 1,700 m altitude in Bontoc.

96. **CAMAROTIS** Lindley

CAMAROTIS PHILIPPINENSIS Lindl. in Journ. Linn. Soc. 3 (1859) 37;

Miq. Fl. Ind. Bat. 3 (1859) 685; *Naves Novis. App.* (1882) 242;

Ames in Philip. Journ. Sci. 8 (1913) Bot. 438, Orch. 5 (1915) 244.

Sarcochilus philippinensis Vidal Phan. Cuming. Philip. (1885) 150,

Rev. Pl. Vasc. Filip. (1886) 271.

Luzon (Laguna, Sorsogon), Leyte, *Cuming s. n.*, *B. S.* 16582 *Ramos*, *Serrato s. n.*, *Wenzel* 04, 0174, *Elmer* 17225. Epiphyte in forest.

CAMAROTIS UTRICULOSA Ames Orch. 5 (1915) 244.

Luzon, Mindanao (Cotabato), *B. S.* 5608 coll.? (flowered in Manila), 11701 *Robinson*.

EXCLUDED OR SYNONYMOUS SPECIES

CAMAROTIS PURPUREA Boxall ex *Naves Novis. App.* (1882) 242, *nomen*.

97. **SARCANTHUS** Lindley

SARCANTHUS BIFIDUS (Lindl.) Ames Orch. 5 (1915) 245.

Saccolabium bifidum Lindl. in Bot. Reg. 24 (1838) Misc. 3; Paxt.

Mag. Bot. 6 (1839) 262; *Reichb. f. in Walp. Ann.* 6 (1864) 886;

Naves Novis. App. (1882) 241.

Gastrochilus bifidus Ktze. *Rev. Gen. Pl.* 2 (1891) 661.

Luzon (Apayao), Mindoro, Leyte, Mindanao (Cotabato, Agusan). A not uncommon species. Epiphyte at 5 to 500 m altitude.

SARCANTHUS DEALBATUS (Lindl.) *Reichb. f. in Walp. Ann.* 6 (1864)

892; *J. J. Sm. in Nat. Tijdschr. Ned. Ind.* 72 (1912) 85 (reprint 15); *Ames Orch.* 5 (1915) 247; *Merr. Sp. Blancoanae* (1918) 115.

Cleisostoma dealbatum Lindl. in Bot. Reg. 29 (1843) Misc. 5.

Cypripedium lineari-subulatum Llanos *Fragm. Pl. Filip.* (1851) 99;

*F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4*¹ (1880) 76; *Naves Novis. App.* (1882) 244.

Luzon (Bontoc, Bulacan, Rizal, Camarines, Sorsogon), Leyte, Palawan, Bancalan, Mindanao (Lanao). A fairly common and widely distributed species. Epiphyte at sea level to 900 m altitude.

SARCANTHUS MALLEIFER J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 90 (reprint 19).

Cleisostoma sagittata Blume Bijdr. (1825) 363, fig. 27; Lindl. Gen. & Sp. Orch. (1833) 226; Miq. Fl. Ind. Bat. 3 (1859) 682; Naves Novis. App. (1882) 238.

Sarcanthus sagittatus J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 607, Fig.-Atlas (1914) fig. 451; Ames Orch. 5 (1915) 251, non King & Pantl.

Palawan, Weber 601. At Alfonso XIII, epiphyte at sea level.

SARCANTHUS MICRANTHUS Ames Orch. 5 (1915) 248.

Luzon (Bataan, Laguna, Tayabas), Leyte, Mindanao (Lanao, Davao), Basilan. A not uncommon species of rather wide distribution. Epiphyte at sea level to 815 m altitude.

SARCANTHUS PACHYPHYLLUS Ames Orch. 5 (1915) 250.

Luzon (Albay, Rizal), *Schultze s. n.*, Lyon 65.

SARCANTHUS RINGENS (Reichb. f.) J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 93 (reprint 22); Ames Orch. 5 (1915) 251.

Cleisostoma ringens Reichb. f. in Gard. Chron. III 4 (1888) 724; Edit. in Kew Bull. (1889) 83.

Reichenbach states: "Ex insulis Philippinis." No specimens of this species have been seen.

SARCANTHUS STRIOLATUS Reichb. f. in Gard. Chron. II 18 (1882) 168; Ames in Philip. Journ. Sci. 2 (1907) Bot. 336 (excl. spec.), Orch. 5 (1915) 252.

Reichenbach states that this species was imported from the Philippines by Low. No specimens certainly referable to it have been seen.

SARCANTHUS WEBERI Ames Orch. 5 (1915) 252.

Luzon (Bataan), Palawan, Mindanao (Cotabato, Davao), *B. S. 5607 Foxworthy*, Weber 252, 604, *Schultze s. n.*, *F. B. 11825 Whitford*. Epiphyte at sea level, from data at hand.

EXCLUDED OR SYNONYMOUS SPECIES

SARCANTHUS CUMINGII (Reichb. f.) J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 85 (reprint 15).

Cleisostoma cumingii Reichb. f. in Allg. Berl. Gartenz. 24 (1856) 218.

The Cuming material on which this species was based may have originated in the Philippines, but no material referable to the description has been seen in the Philippine collections of twenty years.

SARCANTHUS PUGIONIFORMIS Reichb. f. in Allg. Berl. Gartenz. 24 (1856) 219.

According to Dr. Rudolf Schlechter, in a letter, this species was in cultivation in Erlangen from material sent from the Philippines by Loher. No other record of its occurrence in the Islands is known.

SARCANTHUS SAGITTATUS J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 607, Fig.-Atlas (1914) fig. 451; Ames Orch. 5 (1915) 251, non King & Pantl. = *Sarcanthus malleifer* J. J. Sm.

SARCANTHUS SECUNDUS Griff. apud Usteri Beitr. Ken. Philip. Veg. (1905) 128.

Usteri refers a Guimaras specimen here on the basis of an identification made by Schlechter. There is no other evidence to show that this species occurs in the Philippines.

98. ANOTA Schlechter

ANOTA VIOLACEA (Lindl.) Schltr. Orchideen (1914) 588.

Vanda violacea Lindl. in Bot. Reg. 27 (1841) Misc. 12, 33 (1847) t. 30; Paxt. Mag. Bot. 9 (1842) 119; Van Houtte in Fl. des Serres. 17 (1868-69) 92.

Rhynchostylis violacea Reichb. f. in Bonpl. 2 (1854) 93, in Walp. Ann. 6 (1864) 888; Van Houtte in Fl. des Serres 17 (1868-69) 91; Stein Orchideenb. (1892) 537; Ames Orch. 5 (1915) 219.

Saccolabium violaceum Reichb. f. in Bonpl. 2 (1854) 93, in syn.; Warner Sel. Orch. I (1862-65) t. 14; Puydt Orch. (1880) 317, t. 39; Naves Novis. App. (1882) 241; Veitch Man. Orch. Pl. pt. 7 (1891) 117; Ceron Cat. Pl. Herb. Manila (1892) 168.

Luzon (Nueva Ecija, Rizal), *Lyon 50*, *Native collector s. n.*, *B. S. 3090 Curran*, *21334 Ramos*.

99. POMATOCALPA Breda

POMATOCALPA BICOLOR (Lindl.) J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 103 (reprint 33); Ames Orch. 5 (1915) 254.

Epidendrum lineare Blanco Fl. Filip. (1837) 644, ed. 2 (1845) 449, ed. 3, 3 (1879) 44, non Linn.

Cleisostoma bicolor Lindl. in Paxt. Flow. Gard. 2 (1851-52) 99, fig. 185; Reichb. f. in Walp. Ann. 6 (1864) 889; Naves Novis. App. (1882) 239; Ames Orch. 2 (1908) 236; Merr. Fl. Manila (1912) 167, Sp. Blancoanae (1918) 115.

Luzon (Nueva Vizcaya, Bataan, Rizal, Laguna), Romblon, Negros (Occidental Negros), Bohol, Leyte, Palawan, Busuanga, Mindanao (Agusan, Butuan). A common and widely distributed species. Epiphyte or on rocks at sea level to 500 m altitude.

POMATOCALPA LATIFOLIUM (Lindl.) J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 105 (reprint 35), Orch. Jav. Fig.-Atlas (1914) fig. 458; Ames Orch. 5 (1915) 254, in Merr. in Journ. Roy. Asiat. Soc. Straits Branch, Special No. (1921) 202.

Cleisostoma latifolium Lindl. in Bot. Reg. 26 (1840) Misc. 60; Reichb. f. in Walp. Ann. 6 (1864) 889; Hook. f. Fl. Brit. Ind. 6 (1890) 71; Ridl. in Trans. Linn. Soc. II 3 (1893) 370, in Journ. Linn. Soc. 32 (1896) 365; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 612; Ridl. Mat. Fl. Mal. Penin. 1 (1907) 165.

Luzon (Rizal), Palawan, Bancalan, Mindanao (Cotabato, Butuan), *Lyon 23*, *151*, *Reillo s. n.*, *Weber 08*, *600*, *B. S. 11702 Robinson*, *12129 Celestino*. Epiphyte apparently at or near sea level. In Malay Peninsula, Sumatra, Java, and Borneo.

POMATOCALPA MERRILLII Schltr. in Fedde Repert. Beihefte 1 (Orch. Deutsch-Neu-Guin.) (1913) 988; Ames Orch. 6 (1920) 309.

Cleisostoma kunstleri Ames in Philip. Journ. Sci. 6 (1911) Bot. 54, non Hook. f.

Pomatocalpa kunstleri Ames Orch. 5 (1915) 254, non J. J. Sm.

Polillo, Negros, B. S. 10444 McGregor, F. B. 17409 Curran. Epiphyte at 460 m altitude in Negros.

POMATOCALPA VITELLINUM (Reichb. f.) Ames comb. nov.

Cleisostoma vitellinum Reichb. f. in Linnaea 41 (1876) 76; J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 112 (reprint 42); Ames Orch. 5 (1915) 253, in textu.

Cleisostoma expansum Reichb. f. Otia Bot. Hamb. 1 (1878) 52, Xen. Orch. 3 (1881) 29; Naves Novis. App. (1882) 239; Ames Orch. 2 (1908) 237.

Cleisostoma densum Schltr. in Perk. Frag. Fl. Philip. (1904) 43, non Ridl.

Pomatocalpa expansum J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 103 (reprint 33); Ames Orch. 5 (1915) 254.

Luzon (Bataan, Bulacan, Rizal), Leyte, Busuanga, Palawan, Mindanao (Agusan, Davao). A common and rather widely distributed species. Epiphyte or in limestone at sea level to 1,000 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

POMATOCALPA EXPANSUM J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 103 (reprint 33); Ames Orch. 5 (1915) 254 = *Pomatocalpa vitellinum* (Reichb. f.) Ames.

POMATOCALPA KUNSTLERI Ames Orch. 5 (1915) 254, non J. J. Sm. = *Pomatocalpa merrillii* Schltr.

POMATOCALPA ? ROSEUM J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 106 (reprint 36) = *Trichoglottis rosea* (Lindl.) Ames.

100. **TRICHOGLOTTIS** Reichenbach f.

TRICHOGLOTTIS ATROPURPUREA Reichb. f. in Linnaea 41 (1876) 30; Ames Orch. 5 (1915) 254.

Reichenbach (l. c.) states: "Philippinens. insulae." No specimens have been seen.

TRICHOGLOTTIS BATAANENSIS Ames Orch. 1 (1905) 105, *text cut*, 5 (1915) 255.

Cleisostoma subviolaceum Reichb. f. in Bonpl. 10 (1862) 335; Ames Orch. 5 (1915) 253, in nota, non *Trichoglottis subviolacea* (Llanos) Merr.

Trichoglottis rigida Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 271, nec Blume nec Naves.

Trichoglottis subviolacea Merr. Sp. Blancoanae (1918) 116, quoad planta (excl. syn.).

Luzon (Nueva Vizcaya, Pangasinan, Nueva Ecija, Zambales, Bataan, Rizal, Tayabas), Palawan. A common species, especially in Luzon. Epiphyte at about 200 m altitude.

TRICHOGLOTTIS BRACHIATA Ames Orch. 7 (1922) 136.

Biliran, Mindanao (Agusan, Davao), *B. S. 18931 McGregor, Weber 134, Elmer 12021.*

TRICHOGLOTTIS LATISEPALA Ames in Philip. Journ. Sci. 4 (1909) Bot. 675, Orch. 5 (1915) 255.

Luzon (Nueva Vizcaya, Rizal, Tayabas), Palawan, Bancalan, Lum-bucan, Mindanao (Zamboanga, Bukidnon, Agusan, Davao). A fairly common species of rather wide distribution. Epiphyte at sea level to 76 m altitude.

TRICHOGLOTTIS MINDANAENSIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 439, Orch. 5 (1915) 257.

Luzon (Camarines, Sorsogon), Panay (Capiz), Samar, Mindanao (Zamboanga, Lanao, Agusan, Surigao, Davao). A common species, epiphyte at 60 to 150 m altitude.

TRICHOGLOTTIS PHILIPPINENSIS Lindl. in Ann. & Mag. Nat. Hist. 15 (1845) 386; Naves Novis. App. (1882) 243; Vidal Phan. Cuming. Philip. (1885) 150; Usteri Beitr. Ken. Philip. Veg. (1905) 128; Ames Orch. 7 (1922) 137.

Stauroopsis philippinensis Reichb. f. in Hamb. Gartenz. 16 (1860) 117, Xen. Orch. 2 (1862) 8, in Walp. Ann. 6 (1864) 882; Vidal Rev. Pl. Vasc. Filip. (1886) 270; Ames Orch. 2 (1908) 223, 5 (1915) 224.

Luzon (Bulacan, Rizal, Cavite, Camarines), Negros, Mindanao (Misamis, Surigao, Davao). A not uncommon species of fairly wide distribution, at sea level to 300 m altitude.

TRICHOGLOTTIS ROSEA (Lindl.) Ames comb. nov.

Cleisostoma rosea Lindl. in Bot. Reg. 24 (1838) Misc. 80; Paxt. Mag. Bot. 6 (1839) 189, 12 (1845) 165; Reichb. f. in Walp. Ann. 6 (1864) 888; Naves Novis. App. (1882) 239; Ames Orch. 5 (1915) 253, in nota.

Trichoglottis flexuosa Rolfe ex Ames Orch. 1 (1905) 107, 2 (1908) 257, 5 (1915) 255.

Pomatocalpa ? *roseum* J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 106 (reprint 36).

Luzon (Camarines), Panay (Capiz), Bohol, Leyte, Mindanao (Bukidnon, Agusan, Surigao), *Cuming s. n., Loher 6012, 6029, Fortune s. n., Copeland 199, B. S. 38958 Ramos & Edaño, 42085 Edaño, Weber 185, Bolster 230.* Epiphyte at sea level to 1,300 m altitude.

TRICHOGLOTTIS SOLEREDERI Kränzlin in Fedde Repert. 8 (1910) 98; Ames Orch. 5 (1915) 257.

Philippines, probably Luzon, *Loher s. n.*

No specimens of this species have been seen. Kränzlin (l. c.) mentions its relationship to *Trichoglottis bataanensis* Ames.

TRICHOGLOTTIS WENZELII Ames in Philip. Journ. Sci. 8 (1913) Bot. 440, Orch. 5 (1915) 257.

Trichoglottis retusa Ames in Philip. Journ. Sci. 4 (1909) Bot. 676, non Blume.

Leyte, Mindanao (Zamboanga, Agusan, Surigao, Davao), Basilan, *Lyon* 127, *Wenzel* 015, 0176, *Merrill* 8206, *Weber* 189, *Bolster* 225, *Black* s. n., *F. B.* 6103 *Hutchinson*. Epiphyte at sea level to 60 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

TRICHOGLOTTIS BICRURIS Kränzl. in *Ann. k. k. Nat. Hofmus. Wien* 30 (1916) 64.

Based on a Loher plant which Kränzlin questions as being of Philippine origin. No material seen.

TRICHOGLOTTIS BIMAE Reichb. f. apud Naves Novis. *App.* (1882) 243.

TRICHOGLOTTIS FLEXUOSA Rolfe ex Ames *Orch.* 1 (1905) 107, 2 (1908) 257, 5 (1915) 255 = *Trichoglottis rosea* (Lindl.) Ames.

TRICHOGLOTTIS IONOSMA J. J. Sm. in *Nat. Tijdschr. Ned. Ind.* 72 (1912) 108 (reprint 38) (excl. syn. *Epidendrum lineare* Blanco); Ames *Orch.* 5 (1915) 255 = *Staurochilus ionosma* (Lindl.) Schltr.

TRICHOGLOTTIS LANCEOLARIA Blume apud Naves Novis. *App.* (1882) 243.

TRICHOGLOTTIS LUZONENSIS Ames *Orch.* 5 (1915) 255 = *Staurochilus luzonensis* Ames.

TRICHOGLOTTIS PALLENS Lindl. in *Journ. Hort. Soc.* 5 (1850) 34, in *Pact. Flow. Gard.* 1 (1850) 15 = *Phalaenopsis pallens* (Lindl.) Reichb. f., spec. excl.

TRICHOGLOTTIS PUSILLA Reichb. f. apud Naves Novis. *App.* (1882) 243.

TRICHOGLOTTIS RETUSA Ames in *Philip. Journ. Sci.* 4 (1909) Bot. 676, non Blume = *Trichoglottis wenzelii* Ames.

TRICHOGLOTTIS RETUSA Blume apud Naves Novis. *App.* (1882) 243.

TRICHOGLOTTIS RIGIDA Blume apud Naves in Blanco *Fl. Filip.* ed. 3, *Laminas* (1877-83) t. 348; Naves Novis. *App.* (1882) 243.

TRICHOGLOTTIS RIGIDA Vidal Phan. Cuming. *Philip.* (1885) 150, *Rev. Pl. Vasc. Filip.* (1886) 271, nec Blume nec Naves = *Trichoglottis bataanensis* Ames.

TRICHOGLOTTIS SUBVIOLACEA (Llanos) Merr. *Sp. Blancoanae* (1918) 116, quoad syn. (excl. planta).

TRICHOGLOTTIS SUBVIOLACEA Merr. *Sp. Blancoanae* (1918) 116, quoad planta (excl. syn.) = *Trichoglottis bataanensis* Ames.

101. STAUROCHILUS Ridley

STAUROCHILUS IONOSMA (Lindl.) Schltr. in *Fedde Repert. Beihefte* 4 (*Orch. Sino-Jap. Prodr.*) (1919) 287 (excl. syn. *Epidendrum lineare* Blanco).

Cleisostoma ionosmum Lindl. in *Bot. Reg.* 33 (1847) t. 41; *Pact. Mag. Bot.* 14 (1847) 164; G. K. in *Bot. Zeit.* 5 (1847) 633; Reichb. f. in *Walp. Ann.* 1 (1849) 791; *Miq. Fl. Ind. Bat.* 3 (1859) 682; Naves Novis. *App.* (1882) 238; Ames *Orch.* 1 (1905) 103, 2 (1908) 238 (excl. syn. *Epidendrum lineare* Blanco); Merr. in

Philip. Journ. Sci. 1 Suppl. (1906) 39; Hayata Icon. Pl. Formosa 2 (1912) 134, 4 (1914) 96.

Trichoglottis ionosma J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 108 (reprint 38) (excl. syn. *Epidendrum lineare* Blanco); Ames Orch. 5 (1915) 255.

Luzon (Bataan, Rizal, Tayabas, Camarines), Mindanao (Agusan), *F. B. 737 Borden, Lyon 60, B. S. 20852 Escritor, 33626 Ramos & Edaña, Elmer 13977*. Epiphyte at 330 to 1,300 m altitude. Possibly in Formosa; cf. Schltr. l. c.

STAUROCHILUS LUZONENSIS Ames comb. nov.

Trichoglottis luzonensis Ames Orch. 5 (1915) 255.

Vandopsis kupperiana Kränzl. in Fedde Repert. 17 (1921) 390.

Luzon (Tayabas), Mindanao (Agusan), *Loher s. n., B. S. 21199 Escritor, Weber 103*. Epiphyte up to about 400 m altitude in Mindanao.

102. MICROSACCUS Blume

MICROSACCUS BREVIFOLIUS J. J. Sm. in Icon. Bogor. 3 (1906) 63, t. 225, in Bull. Dept. Agr. Ind. Néerl. 13 (1907) 77, in Bull. Jard. Bot. Buitenz. II 26 (1918) 111; Ames Orch. 6 (1920) 309.

Mindanao (Davao), *Wenzel 010003*. Madaum, epiphyte in forest at sea level. In Java.

MICROSACCUS WENZELII Ames Orch. 5 (1915) 257.

Leyte, *Wenzel 029, 0159*. Dagami, epiphyte in forest at 60 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

MICROSACCUS JAVENSIS Blume apud Naves Novis. App. (1882) 244.

103. TAENIOPHYLLUM Blume

TAENIOPHYLLUM COMPACTUM Ames Orch. 2 (1908) 247, *text cut*, 5 (1915) 258.

Microtatorchis compacta Schltr. in Fedde Repert. 10 (1911) 209, in textu.

Luzon (Bontoc, Benguet), Negros, *Vanoverbergh 2821, R. S. Williams 1937, Merrill 7010*. Epiphyte up to 1,700 m altitude in Bontoc.

TAENIOPHYLLUM COPELANDII Ames in Philip. Journ. Sci. 8 (1913) Bot. 436, Orch. 5 (1915) 258.

Mindanao (Davao), *Copeland 339*.

TAENIOPHYLLUM ELMERI Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1588, Orch. 5 (1915) 258.

Luzon (Sorsogon), Negros (Oriental Negros), *Elmer 10336, 16346*. Epiphyte, apparently in mountain regions.

TAENIOPHYLLUM LEYTENSE Ames Orch. 5 (1920) 310.

Leyte, *Wenzel 0998*. Epiphyte at sea level.

TAENIOPHYLLUM MERRILLII Ames in Philip. Journ. Sci. 6 (1911) Bot. 56, Orch. 5 (1915) 258.

Luzon (Bataan), *Merrill 3878, Whitford 117*. Epiphyte at 800 m altitude.

TAENIOPHYLLUM PHILIPPINENSIS Reichb. f. in *Otia Bot. Hamb.* 1 (1878) 53, *Xen. Orch.* 3 (1881) 29; Naves Novis. App. (1882) 237; Ames *Orch.* 2 (1908) 248, 5 (1915) 258.

Luzon (Apayao, Cagayan, Laguna), Leyte, Palawan, Mindanao (Zamboanga). A reasonably common species of wide distribution. Epiphyte at sea level to 130 m altitude.

EXCLUDED OR SYNONYMOUS SPECIES

TAENIOPHYLLUM ALWISII Lindl. apud Naves Novis. App. (1882) 237.

TAENIOPHYLLUM HASSELTII Blume apud Naves Novis. App. (1882) 237.

TAENIOPHYLLUM OBTUSUM Blume apud Naves Novis. App. (1882) 237.

EXCLUDED OR SYNONYMOUS GENERA AND SPECIES

ADENONCOS VIRENS Blume apud Naves Novis. App. (1882) 244.

ALISMORCHIS Thouars = *Calanthe* R. Brown.

ALISMORCHIS FURCATA Ktze. *Rev. Gen. Pl.* 2 (1891) 650 = *Calanthe furcata* Batem.

ALISMORCHIS LYROGLOSSA Ktze. *Rev. Gen. Pl.* 2 (1891) 650 = *Calanthe lyroglossa* Reichb. f.

ARETHUSA GLUTINOSA Blanco *Fl. Filip.* (1837) 641, ed. 2 (1845) 446, ed. 3, 3 (1879) 38, t. 429bis = *Geodorum nutans* (Presl) Ames.

ARUNDINA PENTANDRA Reichb. f. apud Naves Novis. App. (1882) 246.

ARUNDINA SPECIOSA Blume apud Rolfe in *Journ. Bot.* 23 (1885) 215; Vidal Phan. Cuming. Philip. (1885) 150, *Rev. Pl. Vasc. Filip.* (1886) 269.

The Philippine record of this species was based on *Cuming 2058*, but it is almost certain that Cuming's specimen was from the Malay Peninsula, where the species is common and where Cuming also collected. No representative of the genus has been found in the Philippines.

ARUNDINA WALLICHII Reichb. f. apud Naves Novis. App. (1882) 246.

BARTHOLINA PECTINATA R. Br. apud Llanos ex F.-Vill. & Naves in Blanco *Fl. Filip.* ed. 3, 4¹ (1880) 100.

BROMHEADIA PALUSTRIS Lindl. apud Vidal Phan. Cuming. Philip. (1885) 150, *Rev. Pl. Vasc. Filip.* (1886) 270.

The Philippine record was based on *Cuming 2054*, so labeled; the specimen was almost certainly from Singapore or the Malay Peninsula.

CALLISTA Loureiro = *Dendrobium* Swartz.

CALLISTA AMETHYSTOGLOSSA Ktze. *Rev. Gen. Pl.* 2 (1891) 654 = *Dendrobium amethystoglossum* Reichb. f.

CALLISTA ANOSMA Ktze. *Rev. Gen. Pl.* 2 (1891) 653 = *Dendrobium anosmum* Lindl.

CALLISTA BURSIGERA Ktze. *Rev. Gen. Pl.* 2 (1891) 654 = *Dendrobium secundum* Wall.

- CALLISTA TAURINA Ktze. Rev. Gen. Pl. 2 (1891) 655 = *Dendrobium taurinum* Lindl.
- CEPHALANTHERA ENSIFOLIA Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100, non Spreng.
- CERATOCHILUS BIGLANDULOSUS Blume apud Naves Novis. App. (1882) 244.
- CERATOCHILUS MICRANTHUS Lindl. apud Naves Novis. App. (1882) 244.
- CESTICHIS Pfitzer = *Liparis* L. C. Richard.
- CESTICHIS BENGUETENSIS Ames Orch. 1 (1905) 9, 75, t. 3, fig. I, 1-6, 2 (1908) 132 = *Liparis benguetensis* (Ames) Schltr.
- CESTICHIS CAESPITOSA Ames Orch. 2 (1908) 132, t. = *Liparis caespitosa* (Thou.) Lindl.
- CESTICHIS CLEMENSIAE Ames in Philip. Journ. Sci. 4 (1909) Bot. 666 = *Liparis clemensiae* Ames.
- CESTICHIS COMPRESSA Ames Orch. 1 (1905) 76, text cut, 2 (1908) 134 = *Liparis compressa* (Blume) Lindl.
- CESTICHIS CUMINGII Ames Orch. 1 (1905) 13, in Philip. Journ. Sci. 7 (1912) Bot. 7 = *Liparis cumingii* Ridl.
- CESTICHIS DISTICHA Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1888) 131, quoad Philip.; Ames in Philip. Journ. Sci. 2 (1907) Bot. 320 = *Liparis propinqua* Ames.
- CESTICHIS ELMERI Ames Orch. 1 (1905) 10, 75, t. 3, fig. II, 7-12, 2 (1908) 135 = *Liparis elmeri* (Ames) Schltr.
- CESTICHIS FORBESII Ames Orch. 2 (1908) 135 = *Liparis forbesii* Ridl.
- CESTICHIS FRAGILIS Ames in Philip. Journ. Sci. 8 (1913) Bot. 412 = *Liparis fragilis* Ames.
- CESTICHIS GRACILIS Ames Orch. 2 (1908) 136, text cut, non *Liparis gracilis* Hook. f. = *Liparis amesiana* Schltr.
- CESTICHIS HALCONENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 321, Orch. 3 (1908) 63, t. 47 = *Liparis halconensis* Ames.
- CESTICHIS LINEARIFOLIA Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1563 = *Liparis linearifolia* Ames.
- CESTICHIS LONGIPES Ames Orch. 1 (1905) 75 (excl. Luzon specimen), 2 (1908) 137 = *Liparis longipes* Lindl.
- CESTICHIS LYONII Ames in Philip. Journ. Sci. 6 (1911) Bot. 47 = *Liparis lyonii* Ames.
- CESTICHIS MERRILLII Ames Orch. 1 (1905) 11, 75, t. 3, fig. III, 13-18, 2 (1908) 138, in Philip. Journ. Sci. 2 (1907) Bot. 321 = *Liparis merrillii* (Ames) Schltr.
- CESTICHIS NUTANS Ames in Philip. Journ. Sci. 4 (1909) Bot. 597 = *Liparis nutans* Ames.

- CESTICHIS PHILIPPINENSIS** Ames Orch. 1 (1905) 7, 75, *t.* 2, 2 (1908) 138, in Philip. Journ. Sci. 2 (1907) Bot. 321 = *Liparis philippinensis* (Ames) Schltr.
- CESTICHIS TRICHOGLOTTIS** Ames Orch. 2 (1908) 139, *text cut* = *Liparis trichoglottis* (Ames) Schltr.
- CESTICHIS VESTITA** Ames Orch. 2 (1908) 139, p. p., non *Liparis vestita* Reichb. f. = *Liparis confusa* J. J. Sm.
- CESTICHIS VESTITA** Ames Orch. 2 (1908) 139, p. p., non *Liparis vestita* Reichb. f. = *Liparis parviflora* (Blume) Lindl.
- CHILOPOGON MERRILLII** Ames Orch. 5 (1915) 91 = *Appendicula merrillii* Ames.
- CHLOIDIA CONFUSA** Ames Orch. 2 (1908) 67, *text cut* = *Corymborchis confusa* Ames.
- CIRRHOPETALUM** Lindley = *Bulbophyllum* Thouars.
- CIRRHOPETALUM ANTENNIFERUM** Lindl. in Bot. Reg. 29 (1843) sub *t.* 49 = *Bulbophyllum antenniferum* (Lindl.) Reichb. f.
- CIRRHOPETALUM AURATUM** Lindl. in Bot. Reg. 26 (1840) Misc. 50, 29 (1843) *t.* 61; G. K. in Bot. Zeit. 2 (1844) 103; Moore Ill. Orch. Pl. (1857) Cirrhonet. 5, *t.* 4; = *Bulbophyllum auratum* (Lindl.) Reichb. f.
- CIRRHOPETALUM CHRYSSEUM** Kränzl. in Fedde Repert. 8 (1910) 97 = *Bulbophyllum chryseum* (Kränzl.) Ames.
- CIRRHOPETALUM CUMINGII** Lindl. in Paxt. Mag. Bot. 8 (1841) 165, in Bot. Reg. 29 (1843) sub *t.* 49; G. K. in Bot. Zeit. 1 (1843) 791; Moore Ill. Orch. Pl. (1857) Cirrhonet. 5; Hook. in Bot. Mag. 83 (1857) *t.* 4996; Miq. Fl. Ind. Bat. 3 (1859) 653; Veitch Man. Orch. Pl. pt. 3 (1888) 100; Rodigas in Ill. Hort. 39 (1892) 61, in textu = *Bulbophyllum cumingii* (Lindl.) Reichb. f.
- CIRRHOPETALUM LOHERIANUM** Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 61 = *Bulbophyllum loherianum* (Kränzl.) Ames.
- CIRRHOPETALUM MAXILLARE** Lindl. in Bot. Reg. 29 (1843) sub *t.* 49 = *Bulbophyllum blumei* (Lindl.) J. J. Sm.
- CIRRHOPETALUM NUTANS** Lindl. in Bot. Reg. 25 (1839) Misc. 71, 29 (1843) sub *t.* 49; Hook. in Bot. Mag. 75 (1849) *t.* 4418; Miq. Fl. Ind. Bat. 3 (1859) 653 = *Bulbophyllum othonis* (Ktze.) J. J. Sm.
- CIRRHOPETALUM THOUARSII** Lindl. in Bot. Reg. 24 (1838) *t.* 11, non *C. thouarsii* Lindl. Gen. & Sp. Orch. (1830) 58; Hook. in Bot. Mag. 72 (1846) *t.* 4237 = *Bulbophyllum weberi* Ames.
- CLEISOSTOMA AMABILE** Teijsm. & Binn. apud Naves Novis. App. (1882) 238.
- CLEISOSTOMA BICOLOR** Lindl. in Paxt. Flow. Gard. 2 (1851-52) 99, *fig.* 185; Reichb. f. in Walp. Ann. 6 (1864) 889; Naves Novis. App. (1882) 239; Ames Orch. 2 (1908) 236; Merr. Fl. Manila (1912) 167,

Sp. Blancoanae (1918) 115 = *Pomatocalpa bicolor* (Lindl. J. J. Sm.

CLEISOSTOMA BRACHYSTACHYUM Kränzl. in Ann. k. k. Nat. Hofmus. Wien 30 (1916) 62.

Based on a *Loher* plant said to have come from the Philippines. Kränzl. states that it is allied to *Cleisostoma rosea* Lindl., which is *Trichoglottis rosea* (Lindl.) Ames. No material of this species has been seen.

CLEISOSTOMA CHRYSOCHILUM Kränzl. in Fedde Repert. 8 (1910) 545; Ames Orch. 5 (1915) 253, in nota.

Probably a species of *Pomatocalpa*. Kränzl. (l. c.) refers the species to the alliance of *Cleisostoma bicolor* Lindl., which is *Pomatocalpa bicolor* (Lindl.) J. J. Sm. No specimens have been seen.

CLEISOSTOMA CUMINGII Reichb. f. in Allg. Berl. Gartenz. 24 (1856) 218 = *Sarcanthus cumingii* (Reichb. f.) J. J. Sm., spec. dub.

CLEISOSTOMA DEALBATUM Lindl. in Bot. Reg. 29 (1843) Misc. 5 = *Sarcanthus dealbatus* (Lindl.) Reichb. f.

CLEISOSTOMA DENSUM Schltr. in Perk. Frag. Fl. Philip. (1904) 43, non Ridl. = *Pomatocalpa vitellinum* (Reichb. f.) Ames.

CLEISOSTOMA EXPANSUM Reichb. f. Otia Bot. Hamb. 1 (1878) 52, Xen. Orch. 3 (1881) 29; Naves Novis. App. (1882) 239; Ames Orch. 2 (1908) 237 = *Pomatocalpa vitellinum* (Reichb. f.) Ames.

CLEISOSTOMA GUIBERTII Lindl. apud Naves Novis. App. (1882) 238.

CLEISOSTOMA IONOSMUM Lindl. in Bot. Reg. 33 (1847) t. 41; Paxt. Mag. Bot. 14 (1841) 164; G. K. in Bot. Zeit. 5 (1847) 633; Reichb. f. in Walp. Ann. 1 (1849) 791; Miq. Fl. Ind. Bat. 3 (1859) 682; Naves Novis. App. (1882) 238; Ames Orch. 1 (1905) 103, 2 (1908) 238 (excl. syn. *Epidendrum lineare* Blanco) = *Staurochilus ionosma* (Lindl.) Schltr.

CLEISOSTOMA KUNSTLERI Ames in Philip. Journ. Sci. 6 (1911) Bot. 54, non Hook. f. = *Pomatocalpa merrillii* Schltr.

CLEISOSTOMA LONGIFOLIUM Teijsm. & Binn. apud Naves Novis. App. (1882) 238.

CLEISOSTOMA RINGENS Reichb. f. in Gard. Chron. III 4 (1888) 724; Edit. in Kew Bull. (1889) 83 = *Sarcanthus ringens* (Reichb. f.) J. J. Sm.

CLEISOSTOMA ROSEA Lindl. in Bot. Reg. 24 (1838) Misc. 80; Paxt. Mag. Bot. 6 (1839) 189, 12 (1845) 165; Reichb. f. in Walp. Ann. 6 (1864) 888; Naves Novis. App. (1882) 239; Ames Orch. 5 (1915) 253, in nota = *Trichoglottis rosea* (Lindl.) Ames.

CLEISOSTOMA SAGITTATA Blume Bijdr. (1825) 363, fig. 27; Naves Novis. App. (1882) 238 = *Sarcanthus malleifer* J. J. Sm.

CLEISOSTOMA SPATULATUM Blume apud Naves Novis. App. (1882) 238 = *Robiquetia pantherina* (Kränzl.) Ames.

CLEISOSTOMA SUAVEOLENS Blume apud Naves Novis. App. (1882) 238.

CLEISOSTOMA SUBVIOLACEUM Reichb. f. in Bonpl. 10 (1862) 335; Ames Orch. 5 (1915) 253, in nota, non *Trichoglottis subviolacea* (Llanos) Merr. = *Trichoglottis bataanensis* Ames.

CLEISOSTOMA TRIDENTATUM Lindl. apud Naves Novis. App. (1882) 238.

CLEISOSTOMA VITELLINUM Reichb. f. in Linnaea 41 (1876) 76; J. J. Sm. in Nat. Tijdschr. Ned. Ind. 72 (1912) 112 (reprint 42); Ames Orch. 5 (1915) 253, in textu = *Pomatocalpa vitellinum* (Reichb. f.) Ames.

CORYMBIS Thouars = *Corymborchis* Thouars.

CORYMBIS CONFUSA Schltr. in Engl. Bot. Jahrb. 45 (1911) 395 = *Corymborchis confusa* Ames.

CORYMBIS DISTICHA Lindl. Fol. Orch. Corymbis (1854) 1, quoad Philip.; Thou. apud Miq. Fl. Ind. Bat. 3 (1859) 736, quoad Philip. = *Corymborchis confusa* Ames.

CORYMBIS GRACILIS Vidal Rev. Pl. Vasc. Filip. (1886) 272; Ceron Cat. Pl. Herb. Manila (1892) 168, non Blume = *Ceratostylis subulata* Blume.

CORYMBIS VERATRIFOLIA Reichb. f. apud Naves Novis. App. (1882) 248; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 168.

Probably referable to *Corymborchis confusa* Ames.

CORYSANTHES FORNICATA Lindl. apud Naves Novis. App. (1882) 249.

CORYSANTHES MERRILLII Ames Orch. 1 (1905) 65, text cut = *Corybas merrillii* Ames.

CORYSANTHES PICTA Lindl. apud Naves Novis. App. (1882) 249.

CRINONIA ELMERI Ames Orch. 5 (1915) 67, nomen = *Pholidota carnea* (Blume) Lindl.

CYMBIDIUM FLAVESCENS Llanos Frag. Pl. Filip. (1851) 96; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 74; Merr. Sp. Blancoanae (1918) 113 = ? *Oberonia iridifolia* (Roxb.) Lindl.

CYPRIPEDIUM as applicable to the Philippine flora = *Cordula* Raf.

CYPRIPEDIUM ARGUS Reichb. f. in Gard. Chron. (1873) 608, (1874) 710, III 7 (1890) 259; Hook. f. in Bot. Mag. 101 (1875) t. 6175; Naves Novis. App. (1882) 251; Veitch Man. Orch. Pl. pt. 4 (1889) 11; Sander Reichenbachia I 2 (1890) 77, t. 83; Ceron Cat. Pl. Herb. Manila (1892) 169; Kränzl. Orch. Gen. et Sp. 1 (1897) 56 = *Cordula argus* (Reichb. f.) Rolfe.

CYPRIPEDIUM BARBATUM Lindl. in Bot. Reg. 27 (1841) Misc. 53; Vidal Phan. Cuming. Philip. (1885) 151, Rev. Pl. Vasc. Filip. (1886) 273 = *Cordula barbata* (Lindl.) Rolfe.

CYPRIPEDIUM BULBOSUM Blanco Fl. Filip. (1837) 637, ed. 2 (1845) 444, ed. 3, 3 (1897) 33; Merr. Sp. Blancoanae (1918) 117, non Linn.

This is indeterminable from the wholly inadequate description. Blanco's specimens may have been a *Liparis* or a *Malaxis*.

- CYPRIPEDIUM CANNARTIANUM Hort. ex L. Linden in *Lindenia* 3 (1887) 93, *t.* 141 = *Cordula philippinensis* (Reichb. f.) Rolfe.
- CYPRIPEDIUM CILIOLARE Reichb. f. in *Gard. Chron.* II 18 (1882) 488, II 20 (1883) 46; Rodigas in *Ill. Hort.* 31 (1884) 127, *t.* 530; Veitch *Man. Orch. Pl. pt. 4* (1889) 16; *Kränzl. Orch. Gen. et Sp.* 1 (1897) 64 = *Cordula ciliolaris* (Reichb. f.) Rolfe.
- CYPRIPEDIUM ELLIOTTIANUM O'Brien in *Gard. Chron.* III 4 (1888) 501. Cf. *Paphiopedilum rothschildianum* (Reichb. f.) Stein, spec. excl.
- CYPRIPEDIUM FAIRIEANUM Lindl. apud Naves Novis. App. (1882) 251.
- CYPRIPEDIUM HAYNALDIANUM Reichb. f. *Xen. Orch.* 2 (1874) 222, 3 (1881) 33, *t.* 212, in *Gard. Chron.* II 7 (1877) 272, 752; Hook. f. in *Bot. Mag.* 103 (1877) *t.* 6296; Naves Novis. App. (1882) 251; Veitch *Man. Orch. Pl. pt. 4* (1889) 28; *Kränzl. Orch. Gen. et Sp.* 1 (1897) 71 = *Cordula haynaldiana* (Reichb. f.) Rolfe.
- CYPRIPEDIUM HOOKERAE Reichb. f. apud Naves Novis. App. (1882) 251.
- CYPRIPEDIUM JAVANICUM Reinw. apud Naves Novis. App. (1882) 251.
- CYPRIPEDIUM LAEVIGATUM Batem. in *Bot. Mag.* 91 (1865) *t.* 5508; Dombrain in *Floral Mag.* 5 (1866) *t.* 298; Morr. in *Belg. Hort.* 17 (1867) 102, *t.* 6; Van Houtte in *Fl. des Serres* 17 (1869) 83, *t.* 1760-61; Batem. *Second Cent. Orch. Pl.* (1867) *t.* 101; Naves Novis. App. (1882) 251; Ceron *Cat. Pl. Herb. Manila* (1892) 169 = *Cordula philippinensis* (Reichb. f.) Rolfe.
- CYPRIPEDIUM LINEARI-SUBULATUM Llanos *Frag. Pl. Filip.* (1851) 99; F.-Vill. & Naves in *Blanco Fl. Filip. ed. 3, 4*¹ (1880) 76; Naves Novis. App. (1882) 244 = *Sarcanthus dealbatus* (Lindl.) Reichb. f.
- CYPRIPEDIUM LOWII Lindl. apud Naves Novis. App. (1882) 251.
Paphiopedilum lowii Stein apud Ames *Orch.* 2 (1908) 21, 5 (1915) 11, in nota.
- There is no satisfactory evidence that this is Philippine; it was described from Borneo.
- CYPRIPEDIUM MOENSIANUM Hort. ex L. Linden in *Lindenia* 3 (1887) 69, *t.* 129 = *Cordula argus* (Reichb. f.) Rolfe.
- CYPRIPEDIUM PARDINUM Reichb. f. apud Boxall ex Naves Novis. App. (1882) 251.
- CYPRIPEDIUM PHILIPPINENSE Reichb. f. in *Bonpl.* 10 (1862) 335; Veitch *Man. Orch. Pl. pt. 4* (1889) 42, *text cut*; *Kränzl. Orch. Gen. et Sp.* 1 (1897) 46; Rolfe in *Orch. Rev.* 22 (1914) 253, *fig. 29* = *Cordula philippinensis* (Reichb. f.) Rolfe.
- CYPRIPEDIUM PITCHERIANUM Manda in *Am. Flor.* 3 (1887) 178 = *Cordula argus* (Reichb. f.) Rolfe.
- CYPRIPEDIUM PURPURATUM Lindl. apud Naves Novis. App. (1882) 251.
- CYPRIPEDIUM ROEBBELENII Reichb. f. in *Gard. Chron.* II 20 (1883) 684; *Kränzl. Xen. Orch.* 3 (1893) 113, *t.* 265, *Orch. Gen. et Sp.* 1 (1897) 47 = *Cordula philippinensis* (Reichb. f.) Rolfe.

- CYPRIPEDIUM SUPERBIENS Reichb. f. apud Naves Novis. App. (1882) 251.
- CYRTOPERA Lindl. = *Eulophia* R. Brown.
- CYRTOPERA AUREA Boxall ex Naves Novis. App. (1882) 242, *nomen*.
- CYRTOPERA ENSIFORMIS Lindl. in Journ. Linn. Soc. 3 (1859) 31; Miq. Fl. Ind. Bat. 3 (1859) 686; Naves Novis. App. (1882) 242 = *Eulophia stricta* (Presl) Ames.
- CYRTOPERA MACULATA Boxall ex Naves Novis. App. (1882) 242, *nomen*.
- CYRTOPERA SQUALIDA Reichb. f. in Bonpl. 5 (1857) 38; Lindl. in Journ. Linn. Soc. 3 (1859) 31; Miq. Fl. Ind. Bat. 3 (1859) 686; Reichb. f. in Walp. Ann. 6 (1863) 668; Naves Novis. App. (1882) 242 = *Eulophia squalida* Lindl.
- CYRTOPODIUM ENSIFORME Vidal Phan. Cuming. Philip. (1885) 80, 150, Rev. Pl. Vasc. Filip. (1886) 270; Schltr. in Perk. Frag. Fl. Philip. (1904) 43 = *Eulophia stricta* (Presl) Ames.
- CYRTOPODIUM SQUALIDUM Vidal Phan. Cuming. Philip. (1885) 150, Rev. Pl. Vasc. Filip. (1886) 270 = *Eulophia squalida* Lindl.
- CYRTOSIA JAVANICA Blume apud Naves Novis. App. (1882) 247.
- DESMOTRICHUM Blume = *Dendrobium* Swartz.
- DESMOTRICHUM FIMBRIATUM Blume apud Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 354, quoad Philip. = *Dendrobium plicatile* Lindl.
- DESMOTRICHUM SCOPA Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 349 = *Dendrobium scopa* Lindl.
- DIENIA Lindl. = *Malaxis* Swartz.
- DIENIA CARINATA Reichb. f. in Bonpl. 3 (1855) 223 = *Malaxis acuminata* D. Don var. *biloba* Ames.
- DIPLOCONCHIUM INOCEPHALUM Schauer in Nov. Act. Nat. Cur. 19 Suppl. 1 (1843) 428, t. 12, fig. A = *Agrostophyllum inocephalum* (Schauer) Ames.
- DISSORHYNCHIUM MURICATUM Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 434, t. 1, fig. A = *Habenaria hystrix* Ames.
- DORITIS as applicable to the Philippine flora = *Kingiella* Rolfe. Cf. Ames Orch. 6 (1920) 307.
- DORITIS PHILIPPINENSIS Ames Orch. 2 (1908) 235, text cut, 5 (1915) 215 = *Kingiella philippinensis* (Ames) Rolfe.
- DOSSINIA MARMORATA Morr. apud Naves Novis. App. (1882) 250.
- ECHIOGLOSSUM JAVANICUM Blume apud Naves Novis. App. (1882) 244.
- EPIDENDRUM AMETHYSTINUM Reichb. f. apud Naves Novis. App. (1882) 237.
- EPIDENDRUM CARINATUM Linn. Sp. Pl. (1753) 953, ed. 2 (1763) 1350 = *Dendrobium carinatum* (Linn.) Willd.

- EPIDENDRUM EQUITANS Blanco Fl. Filip. (1837) 645, ed. 2 (1845) 449, ed. 3, 3 (1879) 44, non *Dendrobium equitans* Kränzl. = *Dendrobium aporoides* (Lindl.) Merr.
- EPIDENDRUM LINEARE Blanco Fl. Filip. (1837) 644, ed. 2 (1845) 449, ed. 3, 3 (1879) 44, non Linn. = *Pomatocalpa bicolor* (Lindl.) J. J. Sm.
- EPIDENDRUM RHABBARBARUM REDOLENS Blanco Fl. Filip. ed. 3, 3 (1879) 45, t. 389 = *Dendrobium anosmum* Lindl.
- EPIDENDRUM RUCKERAE Reichb. f. apud Naves Novis. App. (1882) 237.
- EPIDENDRUM RUIBARBARUM REDOLENS Blanco Fl. Filip. ed. 2 (1845) 593 = *Dendrobium anosmum* Lindl.
- EPIDENDRUM SOPHRONITIS Lindl. apud Naves Novis. App. (1882) 237.
- EPIDENDRUM ? SPATULATUM Llanos in Mém. Real. Acad. Cienc. Madr. 4 (1857) 499; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100.
- EPIDENDRUM VANILLA Blanco Fl. Filip. (1837) 643, non Linn. = *Vanilla ovalis* Blanco.
- EPHIPPIUM CORNUTUM Blume apud Naves Novis. App. (1882) 244.
- EPHIPPIUM LEPIDIUM Blume apud Naves Novis. App. (1882) 244.
- ESMERALDA SANDERIANA Reichb. f. in Gard. Chron. II 17 (1882) 588, in textu; Kerchove Livre des Orch. (1894) 307, 335, figs. 214, 223; Will. Orch. Grow. Man. ed. 7 (1894) 398, text cut; Ames Orch. 5 (1915) 222 = *Vanda sandariana* Reichb. f.
- EUANTHE Schltr. = *Vanda* Jones.
- EUANTHE SANDERIANA Schltr. Orchideen (1914) 568, fig. 194; Ames Orch. 6 (1920) 308 = *Vanda sandariana* Reichb. f.
- GASTROCHILUS BIFIDUS Ktze. Rev. Gen. Pl. 2 (1891) 661 = *Sarcanthus bifidus* (Lindl.) Ames.
- GASTROCHILUS COMPRESSUS Ktze. Rev. Gen. Pl. 2 (1891) 661 = *Robiquetia compressa* (Lindl.) Schltr.
- GLOSSASPIS TENTACULATA Spreng. apud Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100; Naves Novis. App. (1882) 250.
- GONGORA PHILIPPICA Llanos in Mém. Real. Acad. Cienc. Madr. 4 (1857) 498; F.-Vill. in Blanco Fl. Filip. ed. 3, 4¹ (1880) 100; Merr. Sp. Blancoanae (1918) 117.

The description is very short and imperfect, and no data are given from which it is possible to determine the proper generic position of the species. It certainly is not *Renanthera coccinea* Lour. to which Naves reduced it. Cf. Merrill l. c.

GOODYERA R. Br. = *Epipactis* (Hallier) Boehmer.

GOODYERA DAWSONII Boxall ex Naves Novis. App. (1882) 248, nomen.

- GOODYERA GRANDIS Blume Fl. Jav. Orch. (1858) 36 (Orch. Arch. Ind. 43); Naves Novis. App. (1882) 248 = *Epipactis rubicunda* (Blume) A. A. Eaton.
- GOODYERA LUZONENSIS Ames Orch. 5 (1915) 26 = *Epipactis luzonensis* Ames.
- GOODYERA ORDIANA Boxall ex Naves Novis. App. (1882) 248, *nomen*.
- GOODYERA PARVIFLORA Lindl. apud Naves Novis. App. (1882) 248.
- GOODYERA PHILIPPINENSIS Schltr. in Engl. Bot. Jahrb. 45 (1911) 392; Ames Orch. 5 (1915) 27 = *Epipactis philippinensis* Ames.
- GOODYERA PICTA Boxall ex Naves Novis. App. (1882) 248, *nomen*.
- GOODYERA PROCERA Hook. apud Naves Novis. App. (1882) 248.
- GOODYERA RAMOSII Ames in Philip. Journ. Sci. 8 (1913) Bot. 409, Orch. 5 (1915) 27 = *Epipactis ramosii* Ames.
- GOODYERA REPENS R. Br. apud Naves Novis. App. (1882) 248.
- GOODYERA RUBICUNDA Lindl. in Bot. Reg. 25 (1839) Misc. 61, Gen. & Sp. Orch. (1840) 493, in Journ. Linn. Soc. 1 (1857) 184; Miq. Fl. Ind. Bat. 3 (1859) 728; Naves Novis. App. (1882) 248; J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 121, Fig.-Atlas (1909) fig. 88; Ames Orch. 5 (1915) 27 = *Epipactis rubicunda* (Blume) A. A. Eaton.
- GOODYERA RUBROVENIA Boxall ex Naves Novis. App. (1882) 248, *nomen*.
- GOODYERA VIRIDIFLORA Blume Fl. Jav. Orch. (1858) 34 (Orch. Arch. Ind. 41), t. 9c, fig. 2, 12-16; Ames Orch. 5 (1915) 27 = *Epipactis viridiflora* (Blume) Ames.
- GROSOURDYA APPENDICULATA Reichb. f. Xen. Orch. 2 (1867) 123; Naves Novis. App. (1882) 239 = *Sarcochilus appendiculatus* (Blume) J. J. Sm.
- GROSOURDYA HYSTRIX Reichb. f. apud Naves Novis. App. (1882) 239.
- GROSOURDYA ZOLLINGERI Reichb. f. apud Naves Novis. App. (1882) 239.
- HAEMARIA MERRILLII Ames in Philip. Journ. Sci. 2 (1907) Bot. 315, Orch. 3 (1908) 21, t. 30 = *Kuhlhasseltia whiteheadii* (Rendle) Ames.
- HERPYSMA MERRILLII Ames in Philip. Journ. Sci. 2 (1907) Bot. 313 = *Erythroides merrillii* Ames.
- HEXAMERIA DISTICHA R. Br. in Benn. Pl. Jav. Rar. (1838) 26, t. 7; Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 431 = *Podochilus serphyllifolius* (Blume) Lindl.
- HYLOPHILA RUBRA Ames in Philip. Journ. Sci. 2 (1907) Bot. 315, *nomen*, Orch. 2 (1908) 65, text cut = *Dicerostylis rubra* (Ames) Schltr.
- IRIDORCHIS Thouars = *Oberonia* Lindley.
- IRIDORCHIS ANCEPS Ktze. Rev. Gen. Pl. 2 (1891) 669 = *Oberonia anceps* Lindl.

- IRIDORCHIS APOROPHYLLA Ktze. Rev. Gen. Pl. 2 (1891) 669 = *Oberonia aporophylla* Reichb. f.
- IRIDORCHIS CYLINDRICA Ktze. Rev. Gen. Pl. 2 (1891) 669 = *Oberonia cylindrica* (Lindl.).
- IRIDORCHIS THISBE Ktze. Rev. Gen. Pl. 2 (1891) 669 = *Oberonia thisbe* Reichb. f.
- LEPTORCHIS THOUARS = *Liparis* L. C. Richard.
- LEPTORCHIS FORBESII Ktze. Rev. Gen. Pl. 2 (1891) 671 = *Liparis forbesii* Lindl.
- LEPTORCHIS PECTINATA Ktze. Rev. Gen. Pl. 2 (1891) 671 = *Liparis pectinata* Ridl.
- LIMATODES GRACILIS Lindl. apud Naves Novis. App. (1882) 247.
- LIMATODES GRATA Miq. apud Naves Novis. App. (1882) 247.
- MICROSTYLIS Nutt. = *Malaxis* Swartz.
- MICROSTYLIS ALAGENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 318 = *Malaxis alagensis* Ames.
- MICROSTYLIS BINABAYENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 318 = *Malaxis binabayensis* Ames.
- MICROSTYLIS CARINATA Reichb. f. in Walp. Ann. 6 (1861) 207; Naves Novis. App. (1882) 230; Vidal Phan. Cuming. Philip. (1885) 184, Rev. Pl. Vasc. Filip. (1886) 266; Ridl. in Journ. Linn. Soc. 24 (1888) 338 = *Malaxis acuminata* D. Don var. *biloba* (Lindl.) Ames.
- MICROSTYLIS CONGESTA Reichb. f. in Walp. Ann. 6 (1861) 206; Naves Novis. App. (1882) 230 = *Malaxis latifolia* Sm.
- MICROSTYLIS DENTATA Ames in Philip. Journ. Sci. 2 (July, 1907) Bot. 319 = *Malaxis dentata* Ames.
- MICROSTYLIS HUTCHINSONIANA Ames in Philip. Journ. Sci. 2 (1907) Bot. 319 = *Malaxis hutchinsoniana* Ames.
- MICROSTYLIS MERRILLII Ames in Philip. Journ. Sci. 2 (1907) Bot. 319 = *Malaxis merrillii* Ames.
- MICROSTYLIS MINDORENSIS Rendle in Journ. Bot. 34 (1896) 356 = *Malaxis mindorensis* (Rendle) Ames.
- MICROSTYLIS ODORATA Boxall ex Naves Novis. App. (1882) 230, *nomen*.
- MICROSTYLIS PHILIPPINENSIS Kränzl. in Orchis 2 (Nov. 1907) 16 = *Malaxis dentata* Ames.
- MICROSTYLIS QUADRIDENTATA Ames in Philip. Journ. Sci. 2 (1907) Bot. 320 = *Malaxis quadridentata* Ames.
- MICROSTYLIS RHEEDII Lindl. apud Naves Novis. App. (1882) 230.
- MICROTATORCHIS COMPACTA Schltr. in Fedde Repert. 10 (1911) 209, in textu = *Taeniophyllum compactum* Ames.
- MONOCHILUS CLANDESTINUM Miq. apud Naves Novis. App. (1882) 250.
- MONOCHILUS ZOLLINGERI Reichb. f. apud Naves Novis. App. (1882) 250.

- OPHRYS CERNUA Blanco Fl. Filip. (1837) 639 (as *Ophiris*), ed. 2 (1845) 445, ed. 3, 5 (1879) 35, t. 276, non Linn. = *Grammatophyllum scriptum* (Linn.) Blume.
- ORCHIS LANIGERA Blanco Fl. Filip. (1837) 641, ed. 2 (1845) 446, ed. 3, 3 (1879) 37 = *Rhynchostylis retusa* (Linn.) Blume.
- PANISEA BILAMELLATA Reichb. f. in Walp. Ann. 6 (1861) 240 = *Coelogyne bilamellata* Lindl.
- PAPHIOPEDILUM as applicable to the Philippine flora = *Cordula* Raf.
- PAPHIOPEDILUM ARGUS Stein Orchideenb. (1892) 453; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 40, in Engl. Pflanzenreich IV. 50 (1903) 90; Ames Orch. 1 (1905) 63, 2 (1908) 23 [incl. vars. *moensii* (Hort. ex Veitch) Stein and *nigricans* (Veitch) Hort. ex Stein], 5 (1915) 10 = *Cordula argus* (Reichb. f.) Rolfe.
- PAPHIOPEDILUM BARBATUM Pfitz. in Pringsh. Jahrb. 19 (1888) 159; Ames Orch. 2 (1908) 24, 5 (1915) 10 = *Cordula barbata* (Lindl.) Rolfe.
- PAPHIOPEDILUM CILIOLARE Stein Orchideenb. (1892) 462, fig. 145; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 40, in Engl. Pflanzenreich IV. 50 (1903) 89; Ames Orch. 2 (1908) 23, 5 (1915) 11, in Philip. Journ. Sci. 6 (1911) Bot. 39 = *Cordula ciliolaris* (Reichb. f.) Rolfe.
- PAPHIOPEDILUM ELLIOTTIANUM Stein Orchideenb. (1892) 466. Cf. *Paphiopedilum rothschildianum* (Reichb. f.) Stein, spec. excl.
- PAPHIOPEDILUM HAYNALDIANUM Stein Orchideenb. (1892) 470; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 41, in Engl. Pflanzenreich IV. 50 (1903) 66, fig. 31 A-C; Ames Orch. 2 (1908) 22, 5 (1915) 10, in Philip. Journ. Sci. 6 (1911) Bot. 39; Memml. in Orchis 14 (1920) 18 = *Cordula haynaldiana* (Reichb. f.) Rolfe.
- PAPHIOPEDILUM LAEVIGATUM Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1889) 84 = *Cordula philippinensis* (Reichb. f.) Rolfe.
- PAPHIOPEDILUM LOWII Stein apud Ames Orch. 2 (1908) 21, 5 (1915) 11, in nota. Cf. *Cypripedium lowii* Lindl. apud Naves, spec. excl.
- PAPHIOPEDILUM PHILIPPINENSE Stein Orchideenb. (1892) 480; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 41, in Engl. Pflanzenreich IV. 50 (1903) 61, fig. 28 A-E; Ames Orch. 2 (1908) 19 (incl. vars.), 5 (1915) 10; Memml. in Orchis 14 (1920) 18 = *Cordula philippinensis* (Reichb. f.) Rolfe.
- PAPHIOLEDILUM ROEBELENI Stein Orchideenb. (1892) 480, 482, in syn.; Pfitz. in Engl. Bot. Jahrb. 19 (1894) 41 (as *röbbelenii*), in Engl. Pflanzenreich IV. 50 (1903) 62 = *Cordula philippinensis* (Reichb. f.) Rolfe.
- PAPHIOPEDILUM ROTHSCILDIANUM (Reichb. f.) Stein Orchideenb. (1892) 482; Ames Orch. 2 (1908) 18, 5 (1915) 11, in nota.
Cypripedium elliottianum O'Brien in Gard. Chron. III 4 (1888) 501.
Paphiopedilum elliottianum Stein Orchideenb. (1892) 466.
Paphiopedilum rothschildianum Stein var. *elliottianum* Pfitz. in Engl. Pflanzenreich IV. 50 (1903) 59.

There is no satisfactory evidence that this species is Philippine; the type is from Borneo or from Sumatra. *Cypripedium elliottianum* O'Brien is supposed to have come from the Philippines.

PAXTONIA Lindl. = *Spathoglottis* Blume.

PAXTONIA ROSEA Lindl. in Bot. Reg. 24 (1838) Misc. 61, t. 60; Moore Ill. Orch. Pl. (1857) Paxtonia 4, t. 1; Reichb. f. in Walp. Ann. 6 (1862) 456; Naves Novis. App. (1882) 237 = *Spathoglottis plicata* Blume.

PELEXIA ? FALCATA Llanos Fragm. Pl. Filip. (1851) 95; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 73; Merr. Sp. Blancoanae (1918) 117.

This may be a species of *Oberonia*; the description of Llanos is very imperfect.

PERISTYLUS GOODYEROIDES Lindl. Gen. & Sp. Orch. (1835) 299; Ceron Cat. Pl. Herb. Manila (1892) 169; Kränzl. Orch. Gen. et Sp. 1 (1898) 503 (excl. syn. *P. grandis* Blume); J. J. Sm. Fl. Buitenz. 6 (Orch. Jav.) (1905) 32, Fig.-Atlas (1908) fig. 14 = *Habenaria goodyeroides* D. Don.

PERISTYLUS WARBURGIANUS Kränzl. Orch. Gen. et Sp. 1 (1898) 519.

The type, *Warburg 14987*, is from southern Mindanao. The description suggests close affinity with *Habenaria goodyeroides* D. Don.

PHYLLOECHIS Thouars = *Bulbophyllum* Thouars.

PHYLLOECHIS ANTENNIFERA Ktze. Rev. Gen. Pl. 2 (1891) 677 = *Bulbophyllum antenniferum* (Lindl.) Reichb. f.

PHYLLOECHIS AURATA Ktze. Rev. Gen. Pl. 2 (1891) 677 = *Bulbophyllum auratum* (Lindl.) Reichb. f.

PHYLLOECHIS CHEIRI Ktze. Rev. Gen. Pl. 2 (1891) 676 = *Bulbophyllum blumei* (Lindl.) J. J. Sm.

PHYLLOECHIS CUMINGII Ktze. Rev. Gen. Pl. 2 (1891) 677 = *Bulbophyllum cumingii* (Lindl.) Reichb. f.

PHYLLOECHIS MAXILLARIS Ktze. Rev. Gen. Pl. 2 (1891) 677 = *Bulbophyllum blumei* (Lindl.) J. J. Sm.

PHYLLOECHIS OTHONIS Ktze. Rev. Gen. Pl. 2 (1891) 677 = *Bulbophyllum othonis* (Ktze.) J. J. Sm.

PHYSURUS BLUMEI Lindl. apud Naves Novis. App. (1882) 250.

PINALIA Buch.-Ham. = *Eria* Lindley.

PINALIA APORODES Ktze. Rev. Gen. Pl. 2 (1891) 678 = *Dendrobium aporoides* (Lindl.) Merr.

PINALIA BRACHYSTACHYA Ktze. Rev. Gen. Pl. 2 (1891) 678 = *Eria brachystachya* Reichb. f.

PINALIA POLYURA Ktze. Rev. Gen. Pl. 2 (1891) 679 = *Eria polyura* Lindl.

PINALIA PROREPENS Ktze. Rev. Gen. Pl. 2 (1891) 679 = *Phreatia prorepens* Reichb. f.

PINALIA RETROFLEXA Ktze. Rev. Gen. Pl. 2 (1891) 679 = *Eria retroflexa* Lindl.

PINALA VULPINA Ktze. Rev. Gen. Pl. 2 (1891) 679 = *Eria vulpina* Reichb. f.

PLATANThERA L. C. Richard = *Habenaria* Willdenow.

PLATANThERA AMESIANA Schltr. in Fedde Repert. 9 (1911) 212 = *Habenaria angustata* (Blume) Ktze.

PLATANThERA ANGUSTATA Lindl. Gen. & Sp. Orch. (1835) 290; Naves Novis. App. (1882) 251 = *Habenaria angustata* (Blume) Ktze.

PLATANThERA GOODYEROIDES Vidal Phan. Cuming. Philip. (1885) 151 = *Habenaria goodyeroides* D. Don.

PLATANThERA GRACILIS (Blume) Lindl. apud Naves Novis. App. (1882) 251.

PLATANThERA HALCONENSIS Schltr. in Fedde Repert. 9 (1911) 212 = *Habenaria halconensis* Ames.

PLATANThERA HORSFIELDII (Blume) Naves Novis. App. (1882) 251.

According to J. J. Smith *Peristylus horsfieldii* Blume, upon which Naves based his combination, is a synonym of *Habenaria goodyeroides* D. Don. This conclusion needs further examination before it can be accepted.

PLATANThERA SUSANNAE Lindl. apud Naves Novis. App. (1882) 251.

PLATYCLINIS Benth. = *Acoridium* Nees & Meyen p. p. and *Dendrochilum* Blume p. p.

PLATYCLINIS ARACHNITES Rolfe in Kew Hand-list Orch. (1896) 189, *nomen* = *Dendrochilum arachnites* Reichb. f.

PLATYCLINIS BISTORTA Wendl. & Kränzl. Xen. Orch. 3 (1900) 169, t. 299, fig. I 1-5 = *Dendrochilum bistortum* (Wendl. & Kränzl.) J. J. Sm., spec. excl.

PLATYCLINIS COBBIANA Hemsl. in Gard. Chron. II 16 (1881) 656; Veitch Man. Orch. Pl. pt. 5 (1889) 80 = *Dendrochilum cobbianum* Reichb. f.

PLATYCLINIS CUCUMERINA Rolfe in Kew Hand-list Orch. (1896) 189, *nomen* = *Dendrochilum cucumerinum* Reichb. f. spec. excl.

PLATYCLINIS FILIFORMIS Benth. ex Hemsl. in Gard. Chron. II 16 (1881) 656; Veitch Man. Orch. Pl. pt. 5 (1889) 80; Cogn. in Dict. Icon. Orch. (1902) *Platyclinis* t. 1 = *Dendrochilum filiforme* Lindl.

PLATYCLINIS GLUMACEA Benth. in Journ. Linn. Soc. 18 (1881) 295, by inference only; ex Hemsl. in Gard. Chron. II 16 (1881) 656; Vidal Phan. Cuming. Philip. (1885) 148, Rev. Pl. Vasc. Filip. (1886) 267; Pfitz. in Engl. & Prantl Nat. Pflanzenfam. 2, Abt. 6 (1888) 128, fig. 127 B-F; Veitch Man. Orch. Pl. pt. 5 (1889) 81, t.; Cogn. in Dict. Icon. Orch. (1902) *Platyclinis* t. 2; Ames Orch. 1 (1905) 74 = *Dendrochilum glumaceum* Lindl.


- PLATYCLINIS LATIFOLIA Hemsl. in Gard. Chron. II 16 (1881) 656; Rendle in Journ. Bot. 34 (1896) 355; Ames Orch. 1 (1905) 74 = *Dendrochilum latifolium* Lindl.
- PLATYCLINIS MICROCHILA Schltr. in Bull. Herb. Boiss. II 6 (April, 1906) 302 = *Acoridium microchilum* (Schltr.) Ames.
- PLATYCLINIS UNCATA Rolfe ex Vidal Phan. Cuming. Philip. (1885) 148; Rev. Pl. Vasc. Filip. (1886) 627 = *Dendrochilum uncatum* Reichb. f.
- PLEIONE BILAMELLATA Ktze. Rev. Gen. Pl. 2 (1891) 680 = *Coelogyne bilamellata* Lindl.
- PLEIONE SPARSA Ktze. Rev. Gen. Pl. 2 (1891) 680 = *Coelogyne sparsa* Reichb. f.
- POGONIA A. Jussieu as applicable to the Philippine flora = *Nervilia* Gaudichaud.
- POGONIA CRISPATA Blume Mus. Bot. Lugd.-Bat. 1 (1849) 32; Naves Novis. App. (1882) 248 = *Nervilia discolor* (Blume) Schltr.
- POGONIA DILATATA Blume Fl. Jav. Orch. (1858) 127 (Orch. Arch. Ind. 151) t. 10, fig. 4, 1 & 2; Ames Orch. 1 (1905) 66 = *Nervilia dilatata* (Blume) Schltr.
- POGONIA DISCOLOR Blume Mus. Bot. Lugd.-Bat. 1 (1849) 32; Naves Novis. App. (1882) 248 = *Nervilia discolor* (Blume) Schltr.
- POGONIA FLABELLIFORMIS Lindl. in Wall. Cat. (1832) No. 7400, *nomen*, Gen. & Sp. Orch. (1840) 415; Naves Novis. App. (1882) 248 = *Nervilia aragoana* Gaudich.
- POGONIA JULIANA Wall. apud Naves Novis. App. (1882) 248.
- POGONIA PLICATA Ames Orch. 2 (1908) 43, non Lindl. = *Nervilia discolor* (Blume) Schltr.
- POGONIA PUDICA Ames Orch. 2 (1908) 44, *text cut* = *Nervilia discolor* (Blume) Schltr.
- POGONIA PUNCTATA Blume apud Naves Novis. App. (1882) 248.
- PTYCHOCHILUS SEPTEMNERVIS Schauer in Nov. Act. Nat. Cur. 19, Suppl. 1 (1843) 431, t. 12, fig. B = *Tropidia septemnervis* (Schauer) Reichb. f.
- RHAMPHIDIA GRANDIFLORA Lindl. apud Naves Novis. App. (1882) 248.
- RHAMPHIDIA TENUIS Lindl. in Journ. Linn. Soc. 1 (1857) 182; Miq. Fl. Ind. Bat. 3 (1859) 730; Naves Novis. App. (1882) 248 = *Hetaeria oblongifolia* Blume.
- SALACISTIS NOVEMBRILIS Reichb. f. apud Naves Novis. App. (1882) 249.
- SARCOPODIUM ACUMINATUM Rolfe in Orch. Rev. 18 (August, 1910) 239; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (November, 1910) 329, fig. 32 D, E = *Dendrobium acuminatum* Rolfe.
- SARCOPODIUM CHEIRI Lindl. Fol. Orch. Sarcopod. (1853) 4; Moore Ill. Orch. Pl. (1857) Sarcopod. 4 = *Bulbophyllum blumei* (Lindl.) J. J. Sm.

- SARCOPODIUM CYMBIDIODES** Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (1910) 325, quoad Philip. = *Dendrobium acuminatum* Rolfe.
- SARCOPODIUM LYONII** Rolfe in Orch. Rev. 18 (1910) 240; Ames in Philip. Journ. Sci. 6 (1911) Bot. 53, in Addisonia 8 (1923) 17, t. 265 = *Dendrobium lyonii* Ames.
- SARCOPODIUM STELLA-SILVAE** Loher & Kränzl. in Fedde Repert. 7 (1909) 40; Rolfe in Orch. Rev. 18 (August, 1910) 240; Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21 (November, 1910) 326, fig. 31 A-C; Ames in Philip. Journ. Sci. 8 (1913) Bot. 426 = *Dendrobium stella-silvae* (Loher & Kränzl.) Ames.
- SCHIZMOCERAS** Presl = *Dendrobium* Swartz.
- SCHIZMOCERAS DISTICHA** Presl Rel. Haenk. 1 (1827) 96, t. 13, fig. 2 = *Dendrobium distichum* (Presl) Reichb. f.
- SELENIPEDIUM LAEVIGATUM** May in Rev. Hort. 57 (1885) 301 = *Cordula philippinensis* (Reichb. f.) Rolfe.
- STAUROGLOTTIS EQUESTRIS** Schauer in Nov. Act. Acad. Nat. Cur. 19, Suppl. 1 (1843) 432 = *Phalaenopsis equestris* (Schauer) Reichb. f.
- STAUROPSIS BATEMANII** Nichols. Dict. Gard. 3 (1886) 493 (as *batemanni*) = *Vandopsis lissochiloides* (Gaudich.) Pfitz.
- STAUROPSIS PALLENS** Reichb. f. in Hamb. Gartenz. 16 (1860) 117, Xen. Orch. 2 (1862) 7; Naves Novis. App. (1882) 243 = *Phalaenopsis pallens* (Lindl.) Reichb. f., spec. excl.
- STAUROPSIS PHILIPPINENSIS** Reichb. f. in Hamb. Gartenz. 16 (1860) 117, Xen. Orch. 2 (1862) 8, in Walp. Ann. 6 (1864) 882; Vidal Rev. Pl. Vasc. Filip. (1886) 270; Ames Orch. 2 (1908) 223, 5 (1915) 224 = *Trichoglottis philippinensis* Lindl.
- STAUROPSIS VIOLACEA** Reichb. f. apud Naves Novis. App. (1882) 243.
- STENORHYNCHUS VAGINATUS** Spreng. apud Llanos ex F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 99.
- SYNPTERA SUBVIOLACEA** Llanos Frag. Pl. Filip. (1851) 98; F.-Vill. & Naves in Blanco Fl. Filip. ed. 3, 4¹ (1880) 75.
- TAINIA ELMERI** Ames in Elm. Leaf. Philip. Bot. 5 (1912) 1570 = *Ascotainia elmeri* Ames.
- TAINIA INAMOENA** Kränzl. in Fedde Repert. 17 (1921) 387 = *Ascotainia elmeri* Ames.
- TRICHOTOSIA** Blume = *Eria* Lindley.
- TRICHOTOSIA BINABAYENSIS** Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 141 = *Eria binabayensis* Ames.
- TRICHOTOSIA FASTIGIATIFOLIA** Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 162 = *Eria fastigatifolia* Ames.
- TRICHOTOSIA HALCONENSIS** Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21² (1911) 149 = *Eria halconensis* Ames.

- TRICHOTOSIA MUCRONATA Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21²
(1911) 157 = *Eria mucronata* Lindl. spec. excl.
- TRICHOTOSIA ODORIFERA Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21²
(1911) 153 = *Eria odorifera* Leavitt.
- TRICHOTOSIA RAMOSII Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21²
(1911) 156 = *Eria ramosii* Leavitt.
- TRICHOTOSIA VULPINA Kränzl. in Engl. Pflanzenreich IV. 50. II. B. 21²
(1911) 141 = *Eria vulpina* Reichb. f.
- TYLOSTYLIS sp. Naves Novis. App. (1882) 247.
- VANIFOLIA ALBIDA Blume apud Naves Novis. App. (1882) 248.
- ZEUXINE sp. Vidal Rev. Pl. Vasc. Filip. (1886) 273 = *Adenostylis strateu-*
matica (Linn.) Ames.
- ZEUXINE LUZONENSIS Ames in Philip. Journ. Sci. 2 (1907) Bot. 314, *nomen*
= *Adenostylis luzonensis* Ames.
- ZEUXINE PURPURASCENS Blume apud Naves Novis. App. (1882) 249.
- ZEUXINE WHITEHEADII Rendle in Journ. Bot. 34 (1896) 358 = *Kuhlhassel-*
tia whiteheadii (Rendle) Ames.

ERRATA

- Page 22, line 28, for *Scirpiodendron ghaeri* read *Scirpodendron ghaeri*.
Page 25, line 39, for *Linnophyton obtusifolium* Miq. read *Lymnophyton obtusifolium* Miq.
Page 33, line 23, for Brongh. read Brongn.
Page 41, line 22, for Vizcaya, read Vizcaya),
Page 71, line 44, for Vor. read Var.
Page 75, line 31, for **CHAMAERAPHIS MURICATA** (Linn. f.) comb. nov. read **CHAMAERAPHIS SQUARROSA** (Linn. f.) comb. nov.
Page 84, last line, for Hainan, Indo-China. read Hainan, Indo-China, Celebes, Ternate, Gilolo.
Page 116, after line 37, insert *Fuirena philippinensis* Gandoger in Bull. Soc. Bot. France 66 (1919) 197.
Page 132, line 10, for *Lipironia* read *Lepironia*.
Page 182, after line 6, insert *Schismatoglottis plurivenia* v. A. v. R. in Bull. Jard. Bot. Buitenzorg III 4 (1922) 209.
Page 187, line 27, for **TYPHONIUM DIVARCATUM** (Linn.) Decne. read **TYPHONIUM DIVARICATUM** (Linn.) Decne.
Page 198, line 42, for *Tradescantia critata* Linn. read *Tradescantia cristata* Linn.
Page 201, line 7, for Presl. read Presl
Page 215, line 36, for (Tag.), read (Tag.,
Page 277, line 21, for (Ames) Schltr. & Ames. read (Ames) Ames & Schltr.



INDEX

A

Acampe, 429.
 Acanthophippium, 330.
 Acoridium, 284.
 Acorus, 175.
 Acostainia, 336
 Acriopsis, 398.
 Actinorhynchis, 163.
 Adelmeria, 236.
 Adenostylis, 275.
 Adonidia, 164.
 Aërides, 420.
 Agathis, 5.
 Agave, 212.
 Aglaonema, 182.
 Agrostis, 81.
 Agrostophyllum, 319.
 Aletris, 207.
 Alismaceae, 25.
 Allium, 204.
 Alloteropsis, 70.
 Alocasia, 184.
 Aloe, 204.
 Amaryllidaceae, 210.
 Amomum, 238.
 Amorphophallus, 179.
 Anadendron, 174.
 Ananas, 193.
 Andropogon, 42.
 Aneilema, 196.
 Angraecum, 432.
 Aniselytron, 82.
 Anoectochilus, 274.
 Anota, 438.
 Anthoxanthum, 78.
 Aphyllorchis, 264.
 Apluda, 36.
 Apostasia, 252.
 Apostasiaceae, 252.
 Appendicula, 323.
 Araceae, 172.
 Arachnis, 427.
 Areca, 168.
 Arenga, 159.
 Arisaema, 188.
 Aristida, 78.
 Arthraxon, 41.
 Arundinaria, 94.
 Arundinella, 52.
 Ascocentrum, 434.
 Asparagus, 206.
 Asthenochloa, 49.
 Atamosco, 210.
 Avena, 83.

B

Bambusa, 94.
 Belamcanda, 220.
 Blyxa, 27.
 Boesenbergia, 243.
 Boottia, 27.
 Brachypodium, 93.
 Bromeliaceae, 193.
 Bromus, 92.
 Bulbophyllum, 378.
 Bulbostylis, 127.
 Burmannia, 251.
 Burmanniaceae, 251.

C

Caladium, 187.
 Calamagrostis, 82.
 Calamus, 147.
 Calanthe, 332.
 Camarotis, 436.
 Canna, 247.
 Cannaceae, 247.
 Carex, 136.
 Caryota, 158.
 Cenchrus, 74.
 Centotheca, 91.
 Centrolepidaceae, 191.
 Centrolepis, 191.
 Cephalostachyum, 97.
 Ceratostylis, 317.
 Chamaeanthus, 404.
 Chamaeraphis, 75.
 Cheirostylis, 275.
 Chionachne, 29.
 Chloris, 84.
 Chrysoglossum, 280.
 Cladium, 129.
 Cocos, 170.
 Coelachne, 82.
 Coelococcus, 145.
 Coelogyne, 280.
 Coix, 29.
 Colocasia, 186.
 Commelina, 195.
 Commelinaceae, 194.
 Cordula, 253.
 Cordylina, 206.
 Corybas, 261.
 Corymborchis, 280.
 Corypha, 143.
 Costus, 246.
 Crinum, 210.
 Cryptocoryne, 189.
 Cryptostylis, 261.

Curculigo, 213.
 Curcuma, 243.
 Cyanotis, 198.
 Cycadaceae, 1.
 Cycas, 1.
 Cymbidium, 403.
 Cymodocea, 24.
 Cynodon, 83.
 Cyperaceae, 102.
 Cyperus, 103.
 Cyrtosperma, 179.
 Cystopus, 274.
 Cystorchis, 273.

D

Dacrydium, 4.
 Dactyloctenium, 86.
 Daemonorops, 155.
 Dendrobium, 342.
 Dendrocalamus, 97.
 Dendrochilum, 292.
 Deschampsia, 83.
 Dianella, 202.
 Dicerostylis, 272.
 Didymoplexis, 267.
 Digitaria, 53.
 Diglyphosa, 280.
 Dilochia, 317.
 Dimeria, 30.
 Dinochloa, 100.
 Dioscorea, 215.
 Dioscoreaceae, 215.
 Diplachne, 87.
 Diplacrum, 136.
 Diplanthera, 24.
 Dipodium, 402.
 Disperis, 260.
 Disporum, 206.
 Donax, 248.

E

Eichornia, 200.
 Eleais, 170.
 Eleocharis, 119.
 Eleusine, 85.
 Eleutherine, 220.
 Enhalus, 27.
 Epipactis, 271.
 Epipogon, 265.
 Epipremnopsis, 177.
 Epipremnum, 177.
 Eragrostis, 88.
 Eremochloa, 39.
 Eria, 364.

Eriachne, 82.
 Eriocaulon, 192.
 Eriocaulonaceae, 192.
 Eriochloa, 58.
 Errata, 459.
 Erythroides, 273.
 Eucharis, 211.
 Eulophia, 339.
 Eurycles, 211.

F

Festuca, 92.
 Fimbristylis, 121.
 Flagellaria, 191.
 Flagellariaceae, 190.
 Floscopa, 199.
 Forrestia, 197.
 Freycinetia, 9.
 Fuirena, 116.

G

Gahnia, 131.
 Galeola, 262.
 Garnotia, 81.
 Gastrochilus, 434.
 Gastrodia, 268.
 Geodorum, 341.
 Gigantochloa, 96.
 Globba, 244.
 Glomera, 320.
 Gnetaceae, 7.
 Gnetum, 7.
 Gramineae, 28.
 Grammatophyllum, 399.
 Guadua, 96.

H

Habenaria, 255.
 Halophila, 25.
 Hanguana, 191.
 Hedychium, 241.
 Heliconia, 228.
 Herminium, 255.
 Hetaeria, 278.
 Heterospathe, 162.
 Hippeastrum, 212.
 Hippeophyllum, 310.
 Homalomena, 180.
 Hydrilla, 26.
 Hydrocharitaceae, 25.
 Hymenocallis, 211.
 Hypolytrum, 102.
 Hypoxis, 214.

I

Ichnanthus, 70.
 Imperata, 30.
 Iphigenia, 202.
 Iridaceae, 220.
 Isachne, 58.
 Ischaemum, 37.

J

Joinvillea, 190.
 Juncaceae, 201.
 Juncellus, 114.
 Juncus, 201.

K

Kaempferia, 242.
 Kingiella, 410.
 Kolowratia, 235.
 Korthalsia, 146.
 Kuhlhasseltia, 273.
 Kyllinga, 114.

L

Languas, 230.
 Lecanorchis, 264.
 Leersia, 78.
 Lemna, 190.
 Lemnaceae, 190.
 Lepidogyne, 272.
 Leptaspis, 76.
 Leptochloa, 86.
 Leptosolena, 236.
 Licuala, 143.
 Liliaceae, 202.
 Lilium, 204.
 Liparis, 310.
 Lipocarpa, 102.
 Liriope, 206.
 Livistona, 144.
 Lophatherum, 91.
 Luisia, 426.
 Luzula, 201.

M

Macodes, 275.
 Malaxis, 302.
 Malleola, 434.
 Manisuris, 41.
 Mapania, 131.
 Maranta, 250.
 Marantaceae, 248.
 Mariscus, 112.
 Metroxylon, 145.
 Microchloa, 83.
 Microlaena, 78.
 Microsaccus, 442.
 Microtis, 261.
 Miscanthus, 31.
 Monerma, 93.
 Monochoria, 200.
 Monophrynium, 250.
 Monostachya, 92.
 Muehlenbergia, 80.
 Musa, 221.
 Musaceae, 221.
 Myrmechis, 277.

N

Najadaceae, 24.
 Najas, 24.

Nephelaphyllum, 280.
 Nervilia, 265.
 Neuwiedia, 252.
 Nypa, 170.

O

Oberonia, 307.
 Octarrhena, 392.
 Omoea, 434.
 Oncosperma, 162.
 Ophiopogon, 207.
 Ophiurus, 41.
 Oplismenus, 71.
 Orania, 161.
 Orchidaceae, 253.
 Oryza, 77.
 Ottelia, 27.

P

Pachystoma, 338.
 Palmae, 142.
 Pancreatium, 212.
 Pandanaceae, 9.
 Pandanus, 15.
 Panicum, 61.
 Paspalum, 56.
 Patersonia, 220.
 Pennisetum, 75.
 Perotis, 51.
 Phacelophrynium, 250.
 Phaeomeria, 241.
 Phajus, 331.
 Phalaenopsis, 410.
 Phalaris, 78.
 Philippinaea, 275.
 Phoenix, 142.
 Pholidota, 300.
 Phragmites, 87.
 Phreatia, 395.
 Phrynium, 249.
 Phyllocladus, 4.
 Pinaceae, 5.
 Pinanga, 164.
 Pinus, 6.
 Pistia, 189.
 Plagiostachys, 237.
 Plectocomia, 146.
 Pleomele, 205.
 Plocoglottis, 338.
 Poa, 92.
 Podocarpus, 2.
 Podochilus, 320.
 Pogonatherum, 35.
 Polianthes, 212.
 Pollia, 194.
 Pollinia, 33.
 Polystachya, 329.
 Polytoa, 29.
 Polytrias, 35.
 Pomatocalpa, 438.
 Pontederiaceae, 200.
 Potamogeton, 22.
 Potamogetonaceae, 22.
 Pothodium, 174.

Pothos, 172.
Pseudacoridium, 284.
Ptychoraphis, 163.
Pycereus, 110.

R

Ramosia, 92.
Ravenala, 228.
Remirea, 130.
Renanthera, 428.
Rhaphidophora, 175.
Rhoeo, 199.
Rhynchostylis, 423.
Robiquetia, 434.
Rottboellia, 39.
Ruppia, 24.
Rynchospora, 130.

S

Saccharum, 32.
Saccolabium, 432.
Sagittaria, 25.
Sararanga, 22.
Sarcanthus, 436.
Sarcochilus, 406.
Schismatoglottis, 181.
Schizostachyum, 97.
Schoenorchis, 435.
Schoenus, 128.
Sciaphila, 28.
Scindapsus, 178.
Scirpodendron, 131.
Scirpus, 116.
Scleria, 133.

Setaria, 72.
Smilax, 207.
Spathiphyllum, 178.
Spathoglottis, 336.
Spinifex, 76.
Spiranthes, 268.
Spirodela, 190.
Sporobolus, 80.
Staurochilus, 441.
Stemona, 202.
Stemonaceae, 202.
Stenomeris, 219.
Stereosandra, 267.

T

Tacca, 214.
Taccaceae, 214.
Taeniophyllum, 442.
Taetsia, 205.
Taxaceae, 2.
Taxus, 5.
Thalassia, 27.
Thecostele, 399.
Thelasis, 393.
Thelymitra, 260.
Themeda, 50.
Thoracostachyum, 132.
Thrixspermum, 404.
Thuarea, 75.
Thysanolaena, 53.
Thysanotus, 202.
Torulinium, 114.
Trichoglottis, 439.
Tripogon, 85.
Triticum, 93.

Triuridaceae, 28.
Tropidia, 279.
Typha, 8.
Typhaceae, 8.
Typhonium, 187.

U

Uncinia, 136.

V

Vallisneria, 26.
Vanda, 429.
Vandopsis, 427.
Vanilla, 263.
Vanoverberghia, 238.
Vrydagzynea, 278.

W

Wolffia, 190.

X

Xyridaceae, 192.
Xyris, 192.


Y


Yucca, 204.

Z

Zalacca, 146.
Zea, 28.
Zebrina, 199.
Zingiber, 228.
Zingiberaceae, 228.
Zoisia, 51.









New York Botanical Garden Library

QK3 .M42 v.15

Merrill, Elmer Drew/Botanical publicatio

gen



3 5185 00100 1872

